

VECTORCARDIOGRAPHIC CHANGES DURING EXTENDED SPACE FLIGHT

Raphael F. Smith, M.D.; Kevin Stanton, M.D.†; David Stoop, M.D.†;
Donald Brown, M.D.†; Walter Janusz, M.D.† and Paul King, Ph.D.**

ABSTRACT

To assess the effects of space flight on cardiac electrical properties, vectorcardiograms were taken on the 9 Skylab astronauts during the flights of 28, 59, and 84 days. The Frank lead system was used and observations were made at rest; during 25%, 50%, and 75% of maximum exercise; during a short pulse of exercise (150 watts, 2 minutes); and after exercise. Data from 131 in-flight tests were analyzed by computer and compared to preflight and postflight values. Statistically significant increase in QRS vector magnitude (six of nine crewmen); T vector magnitude (five of nine crewmen); and resting PR interval duration (six of nine crewmen) occurred. During exercise the PR interval did not differ from preflight. Exercise heart rates in-flight were the same as preflight, but increased in the immediate postflight period. No major changes in QRS, T, or ST vector direction occurred. There were sporadic (usually isolated) ectopic ventricular beats in-flight and one astronaut had a brief episode of ventricular tachycardia 21 days after the first mission. Conclusions: with the exception of the arrhythmias, no deleterious vectorcardiographic changes were observed during the Skylab missions. The increase in QRS and T magnitude resembles the electrocardiographic changes associated with athletic conditioning and may be related to increased ventricular volume secondary to centripetal shifts of fluid and/or the in-flight isotonic exercise program. Prolongation of the PR interval at rest with normalization by exercise suggest that there was increased vagal tone in those crewmen exhibiting this response.

INTRODUCTION

The objectives of Skylab Experiment M093 were to measure electrocardiographic signals during space flight, to elucidate the electrophysiological basis for the changes observed, and to assess the effect of the change on the human cardiovascular system. Vectorcardiographic methods were used to quantitate changes, standardize data collection,

*Vanderbilt University, School of Medicine, Nashville, Tennessee 37232.

†Naval Aerospace Medical Institute, Pensacola, Florida 32512.

and to facilitate reduction and statistical analysis of data. Since the Skylab missions provided a unique opportunity to study the effects of prolonged weightlessness on human subjects, an effort was made to construct a data base that contained measurements taken with precision and in adequate number to enable conclusions to be made with a high degree of confidence. Standardized exercise loads were incorporated into the experiment protocol to increase the sensitivity of the electrocardiogram for effects of deconditioning and to detect susceptibility for arrhythmias.

Vectorcardiography provides a comprehensive, three dimensional approach to the analysis of electrocardiographic data which has proven to be useful in both clinical (1, 2) and research applications (3, 4). Vectorcardiographic techniques have been utilized to quantitate electrocardiographic changes during bedrest experiments (4) and Keplerian parabola flights (5). In-flight vectorcardiograms were not obtained during the Mercury, Gemini, or Apollo missions, although preflight and postflight vectorcardiograms were obtained during Apollo 15, 16, and 17.

The purpose of this report is to describe the M093 experiment design, the data transmission system, data reduction methods, and to report the analysis of data from the three Skylab missions.

METHODS

Experiment Design

Vectorcardiograms were taken at rest, during exercise, and after exercise in each crewman during preflight, in-flight, and postflight phases of the Skylab missions. Experiment M093 was designed primarily to obtain electrocardiographic data. In a second Skylab experiment, M171 Metabolic Activity, the Frank lead system (6) was applied for the purpose of obtaining electrocardiographic data during more strenuous exertion. In both experiments vectorcardiograms were obtained from the crewmen at rest for five minutes. In experiment M093 the subject exercised on the bicycle ergometer at a work load of 150 watts for two minutes; during M171 the subject exercised on the ergometer at levels equivalent to 25, 50, and 75% of his maximum aerobic capacity determined prior to the flight. The subject exercised for five minutes at each work load for a total of 15 minutes. After the single exercise load in experiment M093, vectorcardiograms were obtained for 10 minutes. In experiment M171, postexercise vectorcardiograms were obtained for five minutes. The exercise profiles are depicted in figure 1 and the

ergometer is shown in figure 2. Mechanical problems in the orbital workshop during the early portion of Skylab 2 caused scheduling conflicts which resulted in the deletion of the M093 protocol during that flight. However, the M093 protocol was performed throughout Skylab 3 and Skylab 4.




Figure 1. Ergometer exercise profiles for Experiment M093 and Experiment M171. For the M171 protocol, the maximum aerobic capacity of each astronaut was determined prior to the flight and the work levels were 25, 50, and 75 percent of this value.

Instrumentation

Eight electrodes were applied to the crewmen in a modified Frank lead configuration. To lessen muscle noise during exercise, the lead system was modified by transferring the left leg electrode to the left sacral region since the potential difference between the left leg and the left sacrum is negligible. The resistor network proposed by Frank (6) was utilized to correct for the distortion of the cardiac dipole field that




Figure 2. Orbital Workshop bicycle ergometer. The Experiments Support System and Vectorcardiographic subpanel can be seen in the background. The arrows show the positions of two electrodes of the Frank lead system.

results from the shape of the torso and the eccentric location of the heart in the chest. The output of the network is theoretically proportional to the orthogonal components of the cardiac dipole. In the preflight period the electrode sites were marked on each crewman's body by a small tattoo and immediately before the run the electrode site was prepared with benzalkonium chloride. The electrodes were well-type disks and a sponge impregnated with a conductive electrolyte gel served as an interface between the silver-silver chloride electrode and the skin. After attaching the electrodes to the body, the ground reference electrode was tested to determine if there was proper isolation of the subject from the spacecraft ground, then each electrode was tested in sequence to determine the impedance of the skin-electrode interface. The electrode contact was considered to be satisfactory if the impedance was less than 100 000 ohms. To prevent the electrocardiographic signals from exceeding the dynamic range of the recording system, the proper signal conditioner gain for each crewman was determined prior to the flight and the appropriate switch position selected for the individual at the start of the experiment run.

The signal conditioners had differential input with input impedance greater than 40 megohms. The frequency response of the signal conditioners was flat from 0.14 Hz to 90 Hz, at 0.05 Hz and at 100 Hz the frequency response was less than 3 dB down from the flat portion of the frequency response curve. The harmonic distortion of the signal conditioners was less than one percent over the frequency range of the unit. The phase angle difference between vectorcardiographic amplifiers did not exceed one degree over the frequency range of the unit. The three vectorcardiographic channels were simultaneously calibrated by a 10 Hz square wave. The Experiment Support System conditioned and distributed electrical power to experiment equipment, received experiment data in analog and digital form, displayed heart rate data for the crewmen, and routed analog signals to the Airlock Module telemetry system.

The spacecraft recording and telemetry system consisted of two pulse code modulation programmers, pulse code modulation interface box, a data storage and playback system, and remote multiplexers and signal conditioners. This equipment was located primarily in the Airlock Module. The system accepted analog signals from the vectorcardiographic amplifiers and arranged these data into binary coded words at a rate of 320 samples per second. Tape recorders were used to record data for delayed transmission to Lyndon B. Johnson Space Center. The units recorded reduced bit rate segments of the pulse code modulation outputs from the pulse code modulation programmer together with voice data from the crewmen. The recording speed was 1-7/8 inches per second (0.048 meters/second) and the tapes were played back 4 1/4 inches per second (1.05 meters/second) thus allowing four hours of data to be

transmitted in less than 11 minutes. Data were transmitted at "greater-than-real time" rates during passes over receiving stations, a procedure referred to as data "dumps". Due to the volume of data from Skylab experiments it was necessary to compress the vectorcardiographic data and eliminate redundant samples. A zero order predictor algorithm was selected as the data compression technique. In essence, a digital sample of a parameter was tested to determine if it differed from the value of the sample last transmitted. If there was no difference between the current sample and the previous sample, the value was considered to be redundant and not transmitted to L. B. Johnson Space Center.

Computer Analysis Program

The pattern recognition logic of the M093 program is based on a statistical method of identifying components of the vectorcardiogram rather than utilizing empirically derived fiducial values. The program consists of a main program and 10 subprograms that scale and analyze the data. The main program initializes constants, enters identifying information, enters data pertaining to the length of calibration and length of experimental data, generates a digital filter, and serves as a control program for the subroutines. The subroutines compute scale factors from the calibration pulses, apply digital filtering, define the baseline, determine onset and end of waves and segments, and generate a tabular and graphical output of vectorcardiographic items. An optional subroutine derives the 12 conventional electrocardiograms from the three orthogonal vector leads or derives any lead for which spatial coordinates are given. The following vectorcardiographic items are measured or calculated:

- P, QRS, T duration; start and end time
- P, QRS, T maximum voltage X, Y, Z leads; time of occurrence
- P, QRS, T vector loop length
- P, QRS, T maximum vector magnitude, azimuth, elevation; time of occurrence
- P, QRS, T maximum vector velocity; time of occurrence
- P, QRS, T, ST area X, Y, Z leads
- P, QRS, T, ST spatial mean vector magnitude, azimuth, elevation
- Ventricular gradient magnitude, azimuth, elevation

- QRS instantaneous vector magnitude, azimuth, elevation at 10 millisecond intervals
- Angle between spatial mean QRS-T vectors
- Slope and curvature ST segment
- Heart rate

A program for statistical analysis of intra-experiment data has been used in series with the M093 analysis program. The statistical program provides tabular output and graphic displays of both standard statistical parameters and special statistical metrics for directional measurements.

Data Management

For crew safety during the flights, the electrocardiographic signals from each experiment were examined within 24 hours for changes of clinical importance. Occasionally when the orbital workshop was in communications range and an experiment was in progress, electrocardiographic signals were available for immediate analysis. However, due to the gaps in ground station coverage, in most cases the complete data from the experiment were not available until 12 to 24 hours following the run. An analog version of the lead transformation algorithm was available to convert the three orthogonal vectorcardiographic leads to a conventional 12 lead electrocardiogram. Combinations of electrical resistance were chosen to provide the best match between electrocardiographic signals obtained from the standard clinical leads and the derived electrocardiographic signals. Circuit boards were constructed for each astronaut and inserted into the synthesizing unit when an experiment was in progress. Figure 3a is an actual 12-lead electrocardiogram from the Commander of Skylab 2. Figure 3b shows the 12 leads that were derived from the vectorcardiographic signal. Microfilm copies of the computer analysis of experiment data were available within 48 hours after the experiment was performed. These reduced data were reviewed and after inspection of an analog reconstruction, spurious values were deleted from the data base.

RESULTS

Vectorcardiographic parameters from 131 in-flight tests were analyzed by digital computer and compared to pre-flight and postflight values. The vectorcardiographic items examined were heart rate, QRS duration, QRS maximum vector magnitude and direction, T maximum vector magnitude and direction, PR interval, QT interval, area of ST segment X lead, and the spatial angle between QRS and T mean vectors.




Figure 3a. Conventional 12-lead electrocardiogram from Skylab 2 Commander.




Figure 3b. Twelve lead electrocardiogram derived from Frank orthogonal leads.

A statistically significant increase in QRS maximum vector magnitude occurred in six of the nine crewmen. Crew trends plotted as a percentage change from the mean preflight value are shown for the three Skylab missions in figures 4, 5, and 6 respectively. Although crew trends were similar during the three Skylab missions, there were interesting individual differences in the time course of the magnitude changes.

SL 2 CREW TRENDS : QRS VECTOR MAGNITUDE




Figure 4. QRS vector magnitude during Skylab 2 mission. Data plotted as percentage change from the mean preflight value.

SL 3 CREW TRENDS : QRS VECTOR MAGNITUDE




Figure 5. QRS vector magnitude during Skylab 3.




Figure 6. QRS vector magnitude during Skylab 4.

For example, in some astronauts the increase in QRS maximum vector magnitude began in the preflight period as depicted in figures 7 and 8 and in other crewmen the preflight increase was not evident, as shown in figure 9. It should be noted that the data points are spaced equally on the abscissa of figures 7, 8, and 9 although the actual time intervals between the experiments were not equal. The preflight data collection period for Skylab 2 was approximately six months; thus, the rate of QRS maximum vector magnitude increase was considerably greater during the flight than in the preflight period. The magnitude of the spatial T vector increased in five of the nine Skylab crewmen and although the increase was statistically significant, variation in the measurements was large. The angle between the spatial QRS and T mean vectors did not increase significantly in any crewmen during the flights and there were no major changes in QRS, T, or ST vector direction.

The duration of the PR interval measured at rest increased in six of nine crewmen during the three flights and the crew trends for Skylab 3 and Skylab 4 are shown in figures 10 and 11. However, the average




Figure 7. Evolution of QRS vector magnitude change in the Commander of Skylab 2. Experiments are spaced equally on abscissa although the time intervals between the experiments were not equal.




Figure 8. Evolution of QRS vector magnitude change in the Pilot of Skylab 2.




Figure 9. Evolution of QRS vector magnitude changes in the Scientist Pilot of Skylab 3.




Figure 10. PR interval duration during Skylab 3. Data plotted as percentage change from the mean preflight value.




Figure 11. PR interval duration during Skylab 4.

PR interval in-flight did not exceed the clinical standard for the upper limit of normal (0.20 seconds) in any crewmen. During exercise the PR interval did not show a significant difference from the PR interval duration for comparable exercise in the preflight period. A significant decrease in the resting heart rate was observed in the Skylab 2 crew during the flight. However, a significant change in resting heart rate was not a crew trend in the later missions. In general, the average heart rate during the third level of exercise, M171 protocol remained the same as preflight or tended to decrease slightly during the flights. In the immediate postflight period there was a marked increase in the resting heart rate and heart rate response to a given exercise load. As an example, the heart rate responses during Skylab 4 are shown in Figures 12 and 13.

The scalar analog reconstructions of the digital vectorcardiographic signals, the twelve-lead electrocardiograms obtained with the transformation circuitry, and instantaneous vector loop displays were reviewed to check the technical quality of the data and to detect changes of clinical importance. During the three Skylab missions there were no ST segment abnormalities that suggested myocardial ischemia or other changes in the configuration of the electrocardiographic waveforms that were considered to be adverse. During the three flights cardiac arrhythmias were occasionally observed. The Commander

of Skylab 2 had multiple ventricular ectopic beats during the third level of exercise on the initial in-flight M171 test but no arrhythmias were evident in the exercise tests that the Pilot of Skylab 2 performed in the preflight period and during the mission. However, during the third level of exercise (M171 protocol) 21 days after recovery he had salvos of ectopic ventricular beats for approximately 1-1/2 minutes. A representative segment of the arrhythmia is shown in figure 14. The ectopic beats were considered to be ventricular in origin because because the initial beat in the salvo was often a ventricular fusion beat, the isolated ectopic beats did not alter the sinus rhythm, and the degree of QRS aberration was not clearly related to the coupling interval of the premature beat. Furthermore, the ectopic complexes had a monophasic configuration in lead V₁ which suggests a ventricular origin for the arrhythmia. He was monitored for 72 hours, no arrhythmias were detected and he has had no difficulty on subsequent heavy-load exercise tests.

The Scientist Pilot of Skylab 3 had premature ventricular beats sporadically during the second Skylab mission. On mission day 8 during a long extravehicular activity period he was noted to have 80 premature ventricular beats over a 6-1/2 hour period of observation. These ectopic beats were isolated in occurrence and had a configuration suggesting a unifocal origin. This astronaut also had intermittent periods of atrioventricular junctional rhythm at rest throughout the flight. On mission day 21 the Commander of Skylab 3 had a three-beat run of atrioventricular dissociation presumably due to advanced atrioventricular block. The atrial rate was 50 and the junctional escape rate was approximately 39. The episode occurred during the recovery phase of experiment M092 and was not observed in later tests. The crew of Skylab 4 had premature ventricular beats sporadically throughout the mission. On mission day 43 the Commander had two consecutive ectopic ventricular beats during the third level of exercise M171 protocol and on mission day 83 he had three successive ventricular fusion beats during the first exercise level of an M171 test. The Pilot of Skylab 4 had atrioventricular junctional rhythms at rest and after release of lower body negative pressure. There was no impairment of function during the arrhythmia.

DISCUSSION

Elucidation of the mechanisms that underly the cardiac electrical changes is made difficult by the large number of uncontrolled environmental and physiological variables that were operative during the Skylab flights. It is known that electrocardiographic changes occur when there are shifts in the anatomical position of the heart, with hypokalemia, with perturbations of the autonomic nervous system, with changes in the volume of intracavitary blood, and with physical




Figure 14. ECG during arrhythmia experienced 21 days postflight by the Pilot of Skylab 2. The ECG was taken during exercise and six frontal plane leads are shown. The analysis of the arrhythmia is discussed in the text.




Figure 14. Concluded.

conditioning and "deconditioning". These factors have either been shown to vary during space flight or alterations in these factors would intuitively be expected in a weightless environment.

The increase in the magnitude of the QRS maximum vector is an especially interesting change because this has been a crew trend in each Skylab flight. In the majority of the astronauts the QRS maximum vector magnitude has progressively increased during the flight and in several the upward trend began prior to the flight. The T maximum vector magnitude also tended to increase but variation between measurements was greater. The changes observed during the Skylab flights differ from left ventricular hypertrophy encountered clinically in that the angle between the spatial QRS and T vectors was unchanged or decreased in the astronauts and with pathological left ventricular hypertrophy the angle characteristically increases. The QRS and T magnitude increase and the directional relationship between the QRS and T vectors resemble those changes seen in athletes whose electrocardiograms are followed during a physical conditioning program (7). Similarly in dogs given heavy exercise loads over a 12 week period, Wyatt and Mitchell observed a decrease in resting heart rate, a decrease in heart rate response to a standard work load, and an increase in QRS spatial vector magnitude (8).

Increased intracavitary blood due to the centripetal shift of volume during weightlessness may be another mechanism that contributed to the increase in QRS maximum vector magnitude. From a theoretical analysis Brody (9) predicted that an increase in intracavitary blood would augment potentials from radially oriented cardiac dipoles and attenuate those from tangentially oriented dipoles. Since the radially oriented dipoles have the most marked influence on the QRS vector, the net effect of increased diastolic volume would be to increase QRS vector magnitude. Millard, Hodgkin, and Nelson (10) using a series of physiological interventions in experimental animals have confirmed the validity of the Brody effect. Morganroth *et al.* (11) determined left ventricular volumes, wall thickness, and mass by echocardiograph in 26 actively competing college athletes. Athletes competing in events requiring strenuous isotonic exercise had increased left ventricular volume without increased wall thickness and athletes competing in isometric events had increased wall thickness without increased left ventricular volume. Thus, centripetal shift of fluid and isotonic exercise may have had an additive effect in causing the increased QRS vector magnitude that has been observed during the Skylab flights. Measurement of cardiac diastolic dimensions by echocardiography during the preflight and postflight period of Skylab 4 suggested that there was a decrease in the transverse cardiac dimension on the first day after recovery in two of the three crewmen (12). However, after each mission the QRS maximum vector magnitude has remained increased for five to ten days.

An increase in the PR interval duration was a common observation during the three Skylab missions. The PR interval duration is a composite of the conduction time through intra-atrial pathways, the atrioventricular node, and the bundle of His. Since conduction in the atrioventricular node is longer than in the other components, for clinical purposes the PR interval duration serves as an estimate of atrioventricular node conduction. Although drugs such as digitalis, beta-adrenergic blockade, and nodal ischemia can cause prolongation of atrioventricular conduction time, an increase in vagal tone is a more likely explanation of the prolongation of the PR intervals seen in the Skylab astronauts. Further support for this explanation comes from the observation that the PR duration during exercise in-flight was the same as the PR interval duration measured in the preflight period during comparable exercise. Thus the adrenergic influence of exercise tended to overcome the increased vagal influence observed when the men were at rest.

Ventricular ectopy occurred throughout the three Skylab missions. In general this was sporadic, did not alter hemodynamic function in a detectable manner, and electrocardiographic signs of myocardial ischemia were not associated. On three occasions the crewman involved was under extraordinary stress: the first in-flight M171 exercise test of Skylab 2, during a long extravehicular activity in Skylab 3, and during the last in-flight M171 test in Skylab 4. In the case of the more serious ventricular ectopy observed in the Skylab 2 Pilot on the twenty-first postflight day, the relationship of the arrhythmia to the flight is conjectural. He had been in another city on the evening prior to the test and had returned to Houston early on the day of the test. No arrhythmias were observed during the 72 hours following the test and the exercise protocol has been repeated on multiple occasions and the arrhythmia has not recurred during the testing.

With the exception of the arrhythmias, no adverse electrocardiographic changes were observed in the Skylab crews that could be attributed to long exposure to a weightless environment or to the other stresses of extended space flight. Specifically, there was no evidence of myocardial ischemia or changes in the electrocardiogram that would suggest vasoregulatory abnormalities or the emergence of patterns that have been observed in deconditioning experiments (4). The vectorcardiographic techniques utilized in the M093 experiment added both accuracy and precision to the data acquisition and facilitated both scientific investigation and monitoring for crew safety.

REFERENCES

1. Massie, E. and T. J. Walsh. 1969. Clinical Vectorcardiography and Electrocardiography. The Year Book Publishers, Inc., Chicago.

2. Smith, R. F. and R. J. Wherry, Jr. 1966. Quantitative interpretation of the exercise electrocardiogram. Use of computer techniques in the cardiac evaluation of aviation personnel. *Circulation* 34: 1044-1055.
3. Smith, R. F., K. C. Stanton and P. H. King. 1974. Applications of the Frank lead system in clinical and aerospace medical research. H. A. Miller and D. C. Harrison, Eds., Biomedical Electrode Technology, Theory and Practice, pp. 387-404. Academic Press, Inc., New York, N. Y.
4. Saltin, B., G. Blomqvist, J. H. Mitchell, R. L. Johnson, Jr., K. Wildenstahl and C. B. Chapman. 1963. Response to exercise after bedrest and after training. *Circulation* 38, suppl. 7, pp. 1-78.
5. Allebach, N. W. 1966. The Frank lead system as an electrophysiological monitor at 1g, 2g, and 4g. Proceedings of the Second Annual Biomedical Research Conference, NASA Publication (unnumbered) pp. 321-338.
6. Frank, E. 1956. An accurate clinically practical system for spatial vectorcardiography. *Circulation* 13: 737-749.
7. Rautaharju, P. M. and M. J. Karvonen. 1967. Electrophysiological consequences of the adaptive dilatation of the heart. M.J. Karvonen and A. J. Barry. Eds. Physical Activity and the Heart, pp. 159-183. Charles C. Thomas, Springfield, Ill.
8. Wyatt, H. L. and J. H. Mitchell. Influences of physical training on the heart in dogs. *Circulation Research* (in press).
9. Brody, D. A. 1956. A theoretical analysis of intracavitary blood mass influence on the heart lead relationship. *Circulation Research* 4: 731-738.
10. Millard, R. W., B. C. Hodgkin and C. V. Nelson. End-diastolic volume changes reflected in the electrocardiogram. *Circulation* (supplement) (in press).
11. Morganroth, J., B. J. Maron, W. L. Henry and S. E. Epstein. 1974. The athlete's heart: Comparative left ventricular dimensions of collegiate athletes participating in sports requiring isotonic or isometric exertion. *Clinical Research* 22: 291.
12. Henry, W. L., S. E. Epstein, J. M. Griffith, R. E. Goldstein and D. R. Redwood. The effect of prolonged weightlessness on cardiovascular function in Skylab 4 astronauts. Skylab Life Sciences Symposium, Lyndon B. Johnson Space Center, Houston, TX. August 27 to 29, 1974 (in press).