Can Chlorine Anion Catalyze the Reaction of HOCI with HCI?

S. L. Richardson, J. S. Francisco², A. M. Mebel³, and K. Morokuma³

 ¹Department of Electrical Engineering, Howard University, Washington, DC 20059;
 ²Department of Chemistry and Department of Earth and Atmospheric Sciences, Purdue University, West Lafayette, Indiana 47907-1393;
 ³C. L. Emerson Center for Scientific Computation and Department of Chemistry, Emory University, Atlanta, Georgia 30322

Abstract

The reaction of $HOCl + HCl \rightarrow Cl_2 + H_2O$ in the presence of CI has been studied using *ab initio* methods. This reaction has been shown to have a high activation barrier of 46.5 kcal mol⁻¹. The chlorine anion, CI is found to catalyze the reaction, viz. two mechanisms. The first involves CI interacting through the concerted four-center transition state of the neutral reaction. The other mechanism involves the formation of a HCl·HOCl · CI intermediate which dissociates into $Cl_2 + Cl^2 + H_2O$. The steps are found to have no barriers. The overall exothermicity is 15.5 kcal mol⁻¹.

Understanding the underlying chemistry behind stratospheric ozone depletion is central to finding ways of curtailing further loss of ozone in the Antarctic Ozone Hole. One of the major reaction processes implicated in the catalytic removal of ozone is the heterogeneous reaction of HCl with $ClONO_2$ to give HNO_3 and Cl_2 .^{1,2,3} The Cl_2 can subsequently photolyze to produce Cl atoms which can then participate in homogeneous catalytic cycles for ozone destruction through

$$C1 + O_3 \rightarrow ClO + O_2 \tag{1}$$

$$\underline{\text{ClO} + \mathbf{O}} \rightarrow \underline{\text{Cl} + \mathbf{O}_2} \tag{2}$$

net:
$$\mathbf{0} + \mathbf{03} \rightarrow 2O_2$$
 (3)

Recent field measurements of HCl and ClO concentrations in the stratosphere show that the reaction of HCl with ClONO₂ is critical in determining the winter and spring chlorine budget

within the polar vortex. ^{4,5} However, extensive experimental studies suggest that the heterogeneous reaction of HCl with ClONO₂,

may not occur in one step but two viz.^{6,7,8}

$$CIONO_2 + H_2O \rightarrow HOC1 + HNO_3$$
(5)

$$HOCl + HCl + Cl_2 + H_2O$$
(6)

The net reaction of steps 5 and 6 is reaction 3.

A key step is the reaction involving HOCl with HCl (reaction 6). Experimentally,⁹ this reaction is known to be slow in the gas-phase; but little is known about the details of the energetic that could explain the slow rate. Abbatt et al.^{6,7} have asserted that reaction 6 can occur on polar stratospheric cloud (PSC) particles. However, the existence of solvated chlorine in the form of Clon PSC particles has been suggested.^{5,6,7} This has raised the question of whether Cl⁻ can catalyze the reaction of HOCl with HCl (reaction 6). In this paper we use *ab initio* calculations to ascertain if Cl⁻ can lower the activation barrier for the HOCl -t HCl reaction.

The geometries of reactants, products and transition states have been optimized using the *ab initio* MP2 method.¹⁰ Two basis sets have been used for optimization: 6-3 1G(d) and 6-31 1++G(2d,2p) which includes diffuses- and p-functions and an extra set of d-functions on all heavy atoms. The diffuse functions are added because we believe these to be essential for accurate description of geometries and energetic for anions. Single point energies have been calculated using the CCSD(T) method with the extend 6-31 1++G(3df,3pd) basis set.

Geometries of the transition state for the reaction of $HOCl + HCl \rightarrow Cl_2 + H_2O$ is shown in Figure 1a. The number without the asterisk are geometric parameters that have been calculated at the MP2/6-31G(d) level, and the numbers with the asterisk are at the MP2/6-311++G(2d,2p). The transition state is a four-center reaction. The HCl approaches HOCl orienting the hydrogen of the HCl toward the oxygen of HOCl. The HCl bond length increases from the length of the HCl bond in isolated hydrogen chloride. The ClO bond length also increases from that in HOCl. The transition state is a non-planar structure. The four atoms involved in the four-center ring is out-of-plane by 5.9°. Moreover, the hydrogen on HOCl is out of the plane of the ClOH' group by 98°. The transition state is a true first-order saddle point, with one negative frequency.¹¹ The relative energetic are given in Table 1. The experimental heat of reaction¹² at OK for the HOCl + HCl reaction is -18.0 kcal mol⁻¹. At the CCSD(T)/6-311++G(3df,3pd) level we predict a value of -15.4 kcal mol⁻¹, which is in reasonable agreement with the experimental results. The calculated barrier of the same level of theory is 46.5 kcal mol⁻¹. Such a high barrier suggest that the homogeneous reaction will be very slow which is consistent with the experimental observations.

In the presence of Cl-, the anion prefers completing through the hydrogen not involved in the four-center ring. The anion has a dramatic effect on the transition state. The transition state along with structural parameters are given in Figure lb. The transition state is also a first-order saddle point with one imaginary frequency.¹³ Comparing the HCl bond, this bond is longer in the transition state involving the anion than without the anion present. The ClO bond in the transition is significantly elongated as a result of the Cl-. It is also interesting to note that the atoms in the four center ring is more out-of-plane than in the neutral reaction; 24.0° versus 5.9° , respectively. These changes in the transition state for HOCl + HCl involving Cl compared to without the Cl⁻ indicate that the transition state occurs earlier in the entrance channel. The structural changes in the transition state is easier by the presence of the Cl-. The activation barrier for HOCl + HCl + Cl⁻ formation is 8.3 kcal mol⁻¹ at the CCSD(T)/6-311++G(3df,3pd) level. The activation barrier for the neutral reaction (3) is found to be very high and the reaction is slow. However,

involvement of CI anion dramatically changes the potential energy surface. These calculations shows that the involvement of the anion lowers the neutral barrier by $38.2 \text{ kcal mol}^{-1}$. Moreover, these calculations suggest that the anion could play a significant role in reducing the energetic of the HOCl + HCl reaction.

While exploring the potential energy surface of the HOCl + HCl reaction in the presence of Cl-, we found an alternate mechanism for the formation of the $Cl_2 + H_2O$ products. A stepwise mechanism involving the formation of anion complexes such as Cl·HOCl and Cl·H₂O was found. The CI HOCl anionic complex reacts with HCl in a barrier-less process to form a HCl -HOC1. Cl intermediate. This intermediate is shown in Figure 2. At the MP2/6-31G(d) level, the ClO bond in the HCl · HOCl · Cl intermediate is elongated nearly 0.6 Å from the ClO bond in HOCl, and the H'Cl bond is elongated by 0.7 Å from the uncomplexed HCl. We have performed a vibrational frequency analysis to determine if the intermediate is a saddle or minimum energy point on the potential energy surface.¹⁴ The vibrational energy calculations reveal that the structure has all real and positive frequencies. The energy of the intermediate formation from the reaction of HCl+HOCl and Cl is calculated to be -37.3 kcal mol⁻¹ at the MP2/6-31G(d) levels, and reduces to -38.0 kcal mol⁻¹ at the CCSD(T)6-311++G(3df, 3pd) level. The formation of Cl_2 + $H_2O + CI$ requires 22.5 kcal mol⁻¹ at the CCSD(T) level. As a result, one can view an alternate mechanism for HCl + HOCl catalysis by Cl as involving a step-wise mechanism first involving the formation of a Cl. HOCl complex which then attaches to HCl forming an HCl. HOCl. Cl intermediate. The next step involves the detachment of Cl_2 and Cl_2 . H_2O complex. At low temperatures this mechanism involving the HCl · HOCl . Cl intermediate may be preferable. We should point-out that such an attachment/detachment mechanisms has been suggested for the reaction of HCl with ClONO₂ catalyzed by NO₃.

More importantly, the present calculations suggest that anions, such as Cl-, can play a major role in promoting the reaction of HOCl with HCl. The energetic effects on the HOCl + HCl reaction are considerable. Moreover, anions such as nitrate and/or sulfate which are important ions in atmospheric aerosols may play similar roles in the energetic of the HOCl with HCl reaction.

Acknowledgments

One of us (SLR) wishes to thank the Cherry L. Emerson Center for Scientific Computation at Emory University for its support and hospitality as a Emerson Fellow at Emory during the 1995-1996 academic year. One of us (JSF) would like to thank the JPL S upercomputing Project for support of this computing research. The JPL Supercomputing Project is sponsored by JPL and the NASA Office of Space Science and Application. We also wish to acknowledge support from NASA Grant NAGW-2950 for this research.

References

- ¹Solomon, S.; Garcia, R. R.; Rowland, F. S.; Wuebbles, D. J. Nature, 1986,321,755.
- ²Toon, O. B.; Turco, R. P. Sci. Am. 1991,264,68.
- ³Brune, W. H.; Anderson, J. G.; Toohey, D. W. Science, 1991,252, 1260.
- ⁴Webster, C. R.; May, C. D.; Toohey, D. W.; Avallone, L. M.; Anderson, J. G.; Newman, P.; Lait, L.; Schoeberl, M. R.; El King, J. W.; Chan, K. R. Science, 1993,261, 1130.
- ⁵Haas, B. M.; Crellin, K. C.; Kuwata, K. T.; and Okumura, M. J. Phys. Chem. 1994,98,6740.
- ⁶Abbatt, J. P. D.; Molina, M. J. J. Phys. Chem. 1992,96,7074.
- ⁷Abbatt, J. P. D.; Beyer, K. D.; Fucaloro, A. F.; McMahon, J. R.; Wooldridge, P. J.; Zhang, R.; and Molina, M. J. J. Geophys. Res., 1992, 97, 15819.
- ⁸Chu, L. T.; Gen, M. T.; Keyser, L. F. J. Phys. Chem, 1993,97, 12798.
- ⁹DeMore, W. B.; Sander, S. P.; Golden, D. M.; Hampson, R. F.; Kurylo, M. J.; Howard, C. J.; Ravishankara, A. R.; Kolb, C. E.; Molina, M. J. Chemical Kinetics and Photochemical Data for Use in Stratospheric Modeling, Evaluation No. 10, NASA, Jet Propulsion Laboratory, Pasadena, CA.
- ⁶⁰Frisch, M. J.; Trucks, G. W.; Head-Gordon, M.; Gill, P. M. W.; Wong, M. W.; Foresman, J. B.; Johnson, B. G.; Schlegel, H. B.; Robb, M. A.; Replogle, E. S.; Gomperts, R.; Andres, J. L.; Raghavachari, K.; Binkley, J. S.; Gonzalez, C.; Martin, R. L.; Fox, D. J.; DeFrees, D. J.; Baker, J.; Stewart, J. J. P. and Pople, J. A., GAUSSIAN 92/DFT (Gaussian, Inc. Pittsburgh, PA 1993).
- ¹¹ Vibrational frequencies at the MP2/6-31G(d) level of theory for the neutral HCl + HOCl \rightarrow Cl₂ + H₂O reaction are: 3604,2041, 1598,948, 811,365,318,222, 1288 *i* cm⁻¹.
- ¹² Chase, M. W., Jr.; Davies, C. A.; Downey, J. R., Jr.; Frurip, D. J.; McDonald, R. A.; and Syverud, A. N. JANAF Thermochemical Table, 3rd cd., J. Phys. Chem. Ref. Data, 1985, vol. 14.
- ³ Vibrational frequencies at the MP2/6-31G(d) level of theory for the [HCl + HOCl + Cl⁻][‡] transition state are: 3273, 1751, 1544, 1046, 891, 552, 352, 271, 232, 67, 49, 643 *i* cm-l.
- ¹⁴ Vibrational frequencies at the MP2/6-31G(d) level of theory for the HCl. HOCl. Cl⁻ intermediate are: 3793, 2999, 1746, 959, 606, 384, 370, 321, 201, 162, 136, 39 cm-*.
- ¹⁵ Mebel, A. M.; Morokuma, K. J. Phys. Chem. 1956, 100,2985.

				-	
				$CI + HUCI + HCI \rightarrow$	HCI · HUCI · CI →
	HOCI + HCI -	\rightarrow Cl ₂ + H ₂ O	$[HOCI + HCI + CI]^{\ddagger}$	HCI · HOCI · CI	$Cl_2 + H_2O + Cl_2$
Level of Theory	$\Delta H_{r,o}$	H [‡]	H [‡]	$\Delta H_{r,\bullet}^{0}$	$\Delta H^0_{r,o}$
MP2/6-31G(d)	-10.1	52.3	8.1	-37.3	27.2
MP2/6-311++G(2d,2p)	-16.8	49.7	10.3	-41.4	24.6
CCSD(T)/6-311++G(3df,3pd) ^a	-15.5	46.5	8.3	-38.0	22.5
Experiment	0 8 1				

The HOP I HOP Catalward hur OF reative (in boal mal⁻¹) with Dae Tahla 1 Ralativa En

^a Geometries from the MP2/6-311++G(2d,2p) level are used.

10000

Figure Captions

- Figure 1. (a) Transition state for HCl + HOCl → Cl₂ + H₂O for the neutral reaction. Numbers without asterisk are calculated at the MP2/6-31G(d) level, numbers with the asterisk are at the MP2/6-311++G(2d,2p) level of theory.
 (b) Transition state for HCl + HOCl + Cl + Cl₂ + H₂O + Cl-with Cl interacting through the four-center HCl + HOCl transition state. Numbers without asterisk are calculated at the MP2/6-31G(d) level, numbers with the asterisk are calculated at the MP2/6-31G(d) level, numbers with the asterisk are calculated at the MP2/6-31G(d) level, numbers with the asterisk are the MP2/6-31G(d) level, numbers with the asterisk are the MP2/6-31G(d) level, numbers with the asterisk are at the MP2/6-31 1++G(2d,2p) level of theory.
- Figure 2. HCl·HOCl·Cl intermediate. Numbers without asterisk are calculated at the MP2/6-31G(d) level, numbers with the asterisk are at the MP2/6-311++G(2d,2p) level of theory.







-

Í