

the current estimate of the exit pupil wavefront. This estimated PSF is then deconvolved from the image data to provide an estimate of the object. At this point, one has an estimate of the object,

and the estimated wavefront. The process is then repeated with the object included in the model in VSM. The entire process is repeated until a convergence criterion is met.

This work was done by Jeffrey Smith for Goddard Space Flight Center. Further information is contained in a TSP (see page 1). GSC-15963-1

➤ 3D Drop Size Distribution Extrapolation Algorithm Using a Single Disdrometer

Multiple sensors are not required for successful implementation of the 3D interpolation/extrapolation algorithm.

John F. Kennedy Space Center, Florida

Determining the Z-R relationship (where Z is the radar reflectivity factor and R is rainfall rate) from disdrometer data has been and is a common goal of cloud physicists and radar meteorology researchers. The usefulness of this quantity has traditionally been limited since radar represents a volume measurement, while a disdrometer corresponds to a point measurement. To solve that problem, a 3D-DSD (drop-size distribution) method of determining an equivalent 3D Z-R was developed at the University of Central Florida and tested at the Kennedy Space Center, FL. Unfortunately, that method required a minimum of three disdrometers clustered together within a microscale network (≈ 1 -km separation). Since most commercial disdrometers used by the radar meteorology/cloud physics community are high-cost instru-

ments, three disdrometers located within a microscale area is generally not a practical strategy due to the limitations of these kinds of research budgets.

A relatively simple modification to the 3D-DSD algorithm provides an estimate of the 3D-DSD and therefore, a 3D Z-R measurement using a single disdrometer. The basis of the horizontal extrapolation is mass conservation of a drop size increment, employing the mass conservation equation. For vertical extrapolation, convolution of a drop size increment using raindrop terminal velocity is used. Together, these two independent extrapolation techniques provide a complete 3D-DSD estimate in a volume around and above a single disdrometer. The estimation error is lowest along a vertical plane intersecting the disdrometer position in the direction of wind advection.

This work demonstrates that multiple sensors are not required for successful implementation of the 3D interpolation/extrapolation algorithm. This is a great benefit since it is seldom that multiple sensors in the required spatial arrangement are available for this type of analysis.

The original software (developed at the University of Central Florida, 1998–2000) has also been modified to read standardized disdrometer data format (Joss-Waldvogel format). Other modifications to the software involve accounting for vertical ambient wind motion, as well as evaporation of the raindrop during its flight time.

This work was done by John Lane of ASRC Aerospace Corporation for Kennedy Space Center. Further information is contained in a TSP (see page 1). KSC-13302

➤ Social Networking Adapted for Distributed Scientific Collaboration

Sci-Share provides scientists with a set of tools for e-mail, file sharing, and information transfer.

Goddard Space Flight Center, Greenbelt, Maryland

Sci-Share is a social networking site with novel, specially designed feature sets to enable simultaneous remote collaboration and sharing of large data sets among scientists. The site will include not only the standard features found on popular consumer-oriented social networking sites such as Facebook and MySpace, but also a number of powerful tools to extend its functionality to a science collaboration site.

A Virtual Observatory is a promising technology for making data accessible from various missions and instruments through a Web browser. Sci-Share aug-

ments services provided by Virtual Observatories by enabling distributed collaboration and sharing of downloaded and/or processed data among scientists. This will, in turn, increase science returns from NASA missions. Sci-Share also enables better utilization of NASA's high-performance computing resources by providing an easy and central mechanism to access and share large files on users' space or those saved on mass storage.

The most common means of remote scientific collaboration today remains the trio of e-mail for electronic communication, FTP for file sharing, and per-

sonalized Web sites for dissemination of papers and research results. Each of these tools has well-known limitations. Sci-Share transforms the social networking paradigm into a scientific collaboration environment by offering powerful tools for cooperative discourse and digital content sharing. Sci-Share differentiates itself by serving as an online repository for users' digital content with the following unique features:

- Sharing of any file type, any size, from anywhere;
- Creation of projects and groups for controlled sharing;