


# Enhancement of the natural Earth satellite population through meteoroid aerocapture



Althea Moorhead, Bill Cooke  
NASA Meteoroid Environment Office  
April 28, 2014




1972 Great Daylight Fireball


*Image credit: ANSMET, James M. Baker*

# Earth-grazing meteoroids


## Questions:

- How many re-exit?
- How many are aerocaptured?
  - Ratcliff et al. (1993)
  - Hills & Goda (1997)
  - Hunten (1997)
- Can they become natural Earth satellites?
  - Granvik et al. (2012)







# Number and Size



# Velocity




# Directionality




Data from NASA All-Sky Fireball Network  
and Southern Ontario Meteor Network

# Directionality





# The trajectory in the atmosphere




- Compute trajectory under Earth's gravity and atmospheric drag

$$\vec{F}_g = -\frac{GM_{\oplus}m}{r^2}\hat{r}$$


$$\vec{F}_D = -\frac{1}{2}\rho_a v^2 C_d A \hat{v}$$

# Model Earth

Earth as ellipsoid



MSIS-E 90 model



“in-atmosphere” = within 100 km of the surface

# Ablation

- Meteoroid ablation is a function of mass and velocity (Ceplecha, 2000):

$$\dot{m} = -\sigma C_d A \rho_d^{-2/3} \rho m^{2/3} v^3$$



- Meteoroids modeled as spheres ( $C_d = 0.47$ ,  $A = 1.21$ )
- Effective ablation coefficient ( $\sigma$ ) includes some fragmentation

# Catastrophic fragmentation

- Occurs when ram pressure exceeds material strength:

$$S < \rho v^2$$

- Model fragmentation:
  - 20% chance of fragmentation every 0.1 s
  - 0-50% reduction in primary mass



Ceplecha & ReVelle, 2005

# Meteoroid composition

Fireball meteoroid properties by type:


Type	%	$\rho_d$	$\sigma$	$S$
Iron	3	7.8	0.07	200
Stony	29	3.7	0.014	30
Carbonaceous	33	2.0	0.042	10
Cometary	30	0.75	0.1	1

g/cc     $\text{s}^2/\text{km}^2$     MPa

Ceplecha (2001)

Hills & Goda (1993)

# Surviving population



Out of  $10^7$  incident meteoroids:

	<b>Grazers</b>	<b>Aerocaptures</b>
	27,250	10,589
Ablation	18,307	5,040
Frag.	16,754	5,146

Numbers are per year for meteoroids greater than 1 cm in diameter

# Simulations






Image credit: NASA

- Simulations include Sun, Earth, and Moon
- Used *Mercury*, Bulirsch-Stoer method
- Random start between J2K and J2K + 19 years (Metonic cycle; Granvik et al., 2012)

## End states




Leaving Earth's Hill sphere




Atmospheric re-entry



## End states




Re-entry or escape  
after several orbits



Re-entry or escape  
after many orbits


# Steady-state population




Using time spent in near-Earth space and in orbit:

	<b>Number</b>
Outbound	3,000
Orbiting	1,600

# Period distribution



# Size dependence



## Conclusions and future work

- 0.2% of large meteoroids re-exit the atmosphere
- Meteoroid aerocapture can maintain a population of small (cm-sized) NES's
  - $\sim 3,000$  in near-Earth space
  - $\sim 1,500$  orbiting in near-Earth space
- Gravitational capture (Granvik et al., 2012) dominates for meter-sized bodies
- Future work:
  - Convolve results with meteoroid directionality