Division of Engineering
BROWN UNIVERSITY
PROVIDENCE, R. I.

GPO PRICE $ _____

CFSTI PRICE(S) $ _____

Hard copy (HC) 3.00

Microfiche (MF) .65

ff 653 July 65

ON RADIATIVE TRANSFER IN THE LOW REYNOLDS NUMBER BLUNT BODY STAGNATION REGION AT HYPERSONIC SPEEDS PART I. EMISSION DOMINATED CASE

J. T. C. LIU

MAY 1968
RECEIVED
NASA STI FACILITY
INPUT BRANCH

National Aeronautics and Space Administration
Research Grant NASA NGR 40-002-009

Advanced Research Projects Agency
Ballistic Missile Defense Office
and the Fluid Dynamics Branch
Office of Naval Research

Report No.
Nonr 562(35)/21

March 1968

Distribution of This Document is Unlimited

(THRU) (CODE) (CATEGORY)
(ACCESSION NUMBER) (PAGES) (NASA CR OR TMX OR AD NUMBER)
FACILITY FORM 602

On Radiative Transfer in the Low Reynolds Number Blunt Body Stagnation Region at Hypersonic Speeds. Part 1. Emission Dominated Case†

J. T. C. Liu*

Division of Engineering and Center for Fluid Dynamics

Brown University, Providence, Rhode Island

Abstract

One objective of this paper is to contribute to an understanding of radiative transfer effects in the low Reynolds number or merged layer regime of hypersonic flow about axisymmetric blunt bodies. The other objective is to illustrate how the concept of thin shock layer theory can be extended to the radiative case, where the shock structure and shock layer are radiatively coupled, through the introduction of a psuedo-jump condition across the shock structure, accompanied by an iteration technique. The radiative transfer is simplified to the emission dominated case. The gas is taken as calorically and thermally perfect, the viscosity taken as a linear function of the temperature, and the absorption coefficient

†This research was partially supported by the Advanced Research Projects Agency (Ballistic Missile Defense Office) and the Fluid Dynamics Branch of the Office of Naval Research through Contract Nonr 562(35), Task NR 061-132, and by the National Aeronautics and Space Administration through Grant NGR 40-002-009. The assistance of J. Dail and E. Sogame in performing some of the calculation is acknowledged.

*Assistant Professor of Engineering.

of the gas is assumed to be gray. The problem is formulated for the flow about a blunt body but the detailed calculations are carried out for the stagnation region in order to illustrate the techniques used. Radiative cooling decreases the convective heat transfer from the corresponding radiationless case. The additional contributions to the heat transfer due to radiative transfer come from both the shock structure and the shock layer. The relative importance of these contributions are assessed in terms of the rarefaction parameter. Radiative contributions become less important toward the free-molecular range. Radiative cooling decreases the overall shock structure and shock layer thicknesses.

Nomenclature

a = nose radius

a,b = see Eq. (3.3a)

C = see Eq. (3.3a)

C_H = $q_w/\rho_\infty U_\infty(H_\infty - H_w)$, heat transfer coefficient: (C_H^C - convective, C_H^R - radiative, $C_H^{R,SL}$ - radiative, from shock layer, $C_H^{R,STZ}$ - radiative, from shock transition zone)

c_p, c_v = specific heat at constant pressure and at constant volume, respectively

H = $c_p T + \frac{1}{2}(u^2+v^2)$, total enthalpy

ϑ_1, ϑ_2 = see Eq. (3.27)

I = see Eq. (3.28)

K^2 = $\varepsilon\rho_\infty U_\infty a/\mu_o$, rarefaction parameter

k = $(2Pr/3)\ \{[(4/K^2) + 1]^{1/2} - 1\}^{-1}$, also thermal conductivity

Pr = $c_p\mu/k$, Prandtl number

p = pressure

Q = radiative energy loss per unit mass

q_w = rate of energy transferred to body surface (per unit time and per unit area)

q^R = radiative energy flux

R = gas constant

Re_b = $\rho_\infty U_\infty a/\mu_o$, body Reynolds number

T = temperature

u,v = velocity components in the x,y directions, respectively

x,y = orthogonal coordinates along and normal to reference surface (see Fig. 2)

U_∞ = free-stream velocity

Z = distance of reference surface from axis of symmetry

α = absorption coefficient

β = angle between reference surface and free-stream direction

Γ = radiative energy emission to energy convection ratio [see Eq. (3.9)]

γ = c_p/c_v, heat capacity ratio; also incomplete gamma function [see Eq. (3.28)]

ε = $(\gamma-1)/2\gamma$

ζ = $[\psi/(\rho_\infty U_\infty \pi\ Z^2)]^{1/2}$

$\bar{\eta}$ = $(3/4)\ Re_b\ \sin\beta \int_0^y (\mu_o/\mu)\ d(y/a)$

Θ = $(H-H_w)/(H_\infty-H_w)$, dimensionless enthalpy function

θ_o = dimensionless enthalpy function for radiationless case

κ = mass absorption coefficient

κ_c = $d\beta/dx$, logitudinal curvature of reference surface

μ = viscosity coefficient

ξ = x/a

π = 3.1416

ρ = density

σ = Stephen-Boltzmann constant

ψ = stream function

Subscripts

o = stagnation condition in the free stream

1,∞ = conditions in front of the shock and in the free stream, respectively

2,s = conditions behind the shock

w = body surface conditions

1. Introduction

The study of radiative energy transfer in the hypersonic flow over blunt-nosed bodies is relevant to the design of entry vehicles in a flight regime in which the consideration of such an energy transfer mechanism is important to thermal protection systems. Recent fluid mechanical studies of such radiating flows have, in general, fallen into two categories: the inviscid and the fully viscous radiating shock layers.

Recently Cheng and Vincenti[1] presented solutions for the indirect problem of multidimensional, inviscid radiating flow behind a given paraboloidal shock wave. A review of the various earlier works on inviscid radiating shock layers is also given in this paper.

Howe and Viegas[2] considered a completely viscous, radiating shock layer in the stagnation region of a blunt body, using a local similarity approach. In a series of papers, which contain successive improvements on the radiative transfer model, absorption coefficient and gas properties, Hoshizaki and Wilson[3,4] presented solutions for the direct problem of a viscous, radiating shock layer by an integral technique. Burggraf[5] considered the stagnation region of such a shock layer for the emission-dominated case. He obtained asymptotic expansions for large Reynolds numbers, and clarified the radiative interaction of the viscous boundary layer with the inviscid region. In all these viscous radiating shock layer studies, the shock wave is considered to be infinitesimally thin and the usual Rankine-Hugoniot jump conditions are applied, thereby confining the range of validity up to the viscous-layer regime.[6]

In more recent studies of the viscous hypersonic flow over blunt bodies, Bush[7] and Cheng[8] both rule out the conceptual possibility of a flow regime in which transport effects are important throughout the shock layer but not important immediately behind the shock wave. However, Cheng[8] points out the advantage of using the viscous layer model for numerical computations in the moderately high Reynolds number range corresponding to the nonlinear vorticity interaction regime. Use of the viscous layer model would then eliminate the necessity of matching the inner and outer solutions and is thus particularly advantageous when more realistic gas properties are taken into account.

Numerous recent studies, then, have included the high and moderately-high Reynolds number ranges of radiating flow over blunt bodies. However, the low Reynolds number range of radiating hypersonic flows is virtually unexplored. The ability of the viscous hypersonic thin-shock layer theory to describe the essential features of the transition from continuum to free-molecule flow is well known.[8,9,10] It is thus natural to explore the possibility of extending such an approach to radiating, low Reynolds number hypersonic flow.

In the flow regime in which transport effects are important in the shock layer and in modifying the Rankine-Hugoniot jump conditions, the stagnation enthalpy behind the shock structure or at the outer edge of the shock layer decrease from the free stream stagnation value with decreasing body Reynolds number.[9,11] Any preliminary estimate of the importance of energy transfer by radiation relative to energy transfer by convection must take into account this decrease in stagnation enthalpy.

From the standpoint of radiative heat transfer, it would thus be expected that relatively higher flight speeds could be tolerated in the low Reynolds number regime.

This paper presents discussions of the effect of radiative transfer in the low Reynolds number hypersonic flow about a blunt body. The calculations are specialized to the stagnation region to better illustrate the technique used. The radiation model is simplified to that of the emission-dominated case. The radiation parameter for this case is $\Gamma_o \alpha_o \varepsilon a$, where Γ_o is the radiative emission to energy convection ratio, α_o the absorption coefficient and εa gives the order of the shock layer thickness. The subscript o indicates free-stream stagnation conditions. However, as was previously discussed, the free-stream stagnation conditions are not a good indication of the importance of radiative transfer because of the "shock-slip" effects. The appropriate reference condition is one based on the condition at the outer edge of the shock layer, $\Gamma_s \alpha_s \varepsilon a$. A preliminary estimate of this parameter for various values of the rarefaction parameter, K^2, can be made through the use of Cheng's[9] solution $\theta_o(1)$. The parameter $\Gamma_s \alpha_s \varepsilon a$ is shown in Figure 1, where $\Gamma_s \alpha_s \varepsilon a \approx \theta_o^8(1) \Gamma_o \alpha_o \varepsilon a$ is indicated by the dashed lines.* The $K^2 \to \infty$ limit occurs when shock-slip effects are absent and $\Gamma_s \alpha_s = \Gamma_o \alpha_o$ identically. But for $K^2 = O(1)$, for instance, $\Gamma_o \alpha_o \varepsilon a$ can be as high as 10^2 while $\Gamma_s \alpha_s \varepsilon a$ is only $O(1)$. The true value of $\Gamma_s \alpha_s \varepsilon a$, which is not known a priori and is obtained only after the solution has been found, is shown by the solid lines in Figure

*The term $\alpha_s T_s^4$ may be replaced by $\theta_o^8(1) \alpha_o T_o^4$ (see Section 3).

1. Because radiative transfer augments the shock-slip due to transport effects behind the shock wave structure, the dashed lines overestimate this parameter (as is evident from Figure 1).

2. Basic Equations

The basic equations of the two-thin-layer formulation for the description of the (non-radiating) low Reynolds number hypersonic flow over blunt bodies are obtained by Cheng.[8,10] The equivalence of this description to Bush's[7] four-layer formulation is pointed out by Cheng.[8]

In this section Cheng's two-layer formulation will be augmented by the inclusion of the effect of radiative transfer in the energy equations. This presupposes that radiative transfer will not alter the order-of-magnitude analyses in the thin-layer theory. Ultimate justification of this assumption must be sought a posteriori from the actual solutions obtained.*

Throughout this paper, the simplifying assumptions of a perfect gas having constant specific heat and a linear viscosity-temperature law will be retained. It will also be assumed that the gas ahead of the shock structure is cold and neither absorbing nor emitting, that the out going radiation escapes to infinity, that the free stream is uniform, and that the wall is cold. If the smallest local photon mean free path is large compared to the thickness of the shock-transition

*In fact, for the radiative effects in the "order-unity" and the "weak" regimes, the thin-layer approximation is made more valid because of the decrease of shock layer and shock-transition zone thicknesses. The strong radiation case requires separate consideration.

zone and of the shock layer, then the effect of radiation can be represented by the emission-dominated case. The gas is assumed to be gray with a frequency-averaged absorption coefficient which depends on the local pressure and temperature. The contribution of radiation to the pressure and internal energy of the gas are neglected.

2.1 Shock-layer equations

The orthogonal coordinates used are x and y, where x is the distance along and y the distance normal to a reference surface. Following Cheng[8], this reference surface is taken to be the outer edge of the shock layer. (See Figure 2).

The thin-layer approximation then yields the following set of equations for the axially symmetric shock-layer:

$$\frac{\partial}{\partial x}(\rho u Z) + \frac{\partial}{\partial y}(\rho v Z) = 0 \tag{2.1}$$

$$\rho\left(u\frac{\partial u}{\partial x} + v\frac{\partial u}{\partial y}\right) = -\frac{\partial p}{\partial x} + \frac{\partial}{\partial y}\left(\mu\frac{\partial u}{\partial y}\right) \tag{2.2}$$

$$\frac{\partial p}{\partial y} = -\kappa_c \rho u^2 \tag{2.3}$$

$$\rho\left(u\frac{\partial H}{\partial x} + v\frac{\partial H}{\partial y}\right) = \frac{\partial}{\partial y}\left\{\frac{\mu}{Pr}\frac{\partial}{\partial y}\left[H-(1-Pr)\frac{u^2}{2}\right]\right\} - \rho Q \tag{2.4}$$

$$p = \rho R T \tag{2.5}$$

The radiative heat loss term, $-\rho Q$, is the energy lost per unit volume. For the emission-dominated case

$$-\rho Q = -4\rho\kappa\sigma T^4 \tag{2.6}$$

and $-\rho Q$ then has the form of local heat sinks.

2.2 Shock-transition zone equations

Here y is measured from and normal to the outer edge of the shock layer and x is the distance along this layer. The thin-layer approximation then yields the following set of equations for the shock-transition zone:

$$\rho v = \rho_1 v_1 \tag{2.7}$$

$$p + \rho_1 v_1 v - \frac{4}{3}\mu\frac{\partial v}{\partial y} = \rho_1 v_1^{\ 2} \tag{2.8}$$

$$\rho_1 v_1 u - \mu\frac{\partial u}{\partial y} = \rho_1 v_1 u_1 \tag{2.9}$$

$$\rho_1 v_1 \frac{\partial H}{\partial y} = \frac{\partial}{\partial y}\left\{\frac{\mu}{Pr}\frac{\partial}{\partial y}\left[H - (1-Pr)\frac{u^2}{2} - \left(1-\frac{4Pr}{3}\right)\frac{v^2}{2}\right]\right\} - \rho Q \tag{2.10}$$

$$p = \rho RT \tag{2.11}$$

Equations (2.1) through (2.3) are obtained through a simple integration, while (2.4) remains a second-order differential equation. The radiative heat loss, $-\rho Q$, is again given by (2.6). The subscript 1 indicates free stream conditions, where $\rho_1 = \rho_\infty$, $u_1 = U_\infty \cos\beta$, $v_1 = -U_\infty \sin\beta$, and β is the local inclination of the reference surface.

2.3 The unreduced problem

In the nonradiating problem (i.e., for $-\rho Q = 0$) the set of equations (2.7) through (2.10), with Eq. (2.10) integrated once, can be reduced to give a set of shock conservation relations that provide the outer boundary condition for the shock layer.[8,9,10] In the present problem, only Eqs. (2.7) through (2.9) may be reduced; such reduction yields

$$\rho_2 v_2 = \rho_1 v_1 \tag{2.12}$$

$$p_2 = \rho_1 v_1^2 \tag{2.13}$$

$$\rho_1 v_1 (u_2 - u_1) = \left(\mu \frac{\partial u}{\partial y}\right)_2 \tag{2.14}$$

Here the subscript 2 indicates the condition at the outer edge of the shock layer where v and $\partial v/\partial y$ are higher-order quantities. Equations (2.12) through (2.14) are the proper jump conditions, independent of the shock structure problem.

Integration of Eq. (2.10) once across the shock-transition zone yields

$$\rho_1 v_1 (H_2 - H_1) = \left\{\frac{\mu}{Pr} \frac{\partial}{\partial y}\left[H-(1-Pr)\frac{u^2}{2}\right]\right\}_2 - \int_1^2 \rho Q dy \tag{2.15}$$

The last term on the right-hand side of Eq. (2.14) represents the accumulated radiative heat loss through the shock-transition zone. Because of this history-dependent term, a jump condition for the total enthalpy is no longer available. That is, the shock layer can no longer be described independently of the details of the shock-transition zone. The two layers are thus radiatively coupled.

3. Solution of the Two Radiatively-Coupled Layers

Although the radiative coupling of the shock-transition zone to the shock layer is apparent, the method of solution to be described in this section will adhere to the uncoupled, non-radiating two-layer formulation. Thus, an attempt shall be made to derive a "psuedo-jump" condition from the radiating shock-transition zone energy equation. This would provide the boundary condition for the outer edge of the radiating shock layer.

However, such a condition contains the as yet unknown cumulative radiative heat loss through the shock-transition zone. The solution of the shock layer energy equation, using such a psuedo-jump condition, would then provide the boundary condition for the shock-transition zone at the interface. An overall iterative scheme can then be devised.

For a given boundary condition at the shock-interface, the shock-transition zone solution can be calculated and the cumulative heat loss through this layer obtained. For a given cumulative heat loss at the shock interface, the shock layer solution can be calculated and the boundary condition for the shock-transition zone obtained. The final solution is obtained when consistency is achieved in such a "phase-plane".

The solutions for the individual layers used in the overall iteration scheme can be obtained by actually solving the differential equations appropriate to the respective layers. However, the technique of solution to be described below is, again, through iteration.

3.1 Transformation of the shock-transition zone equations

Following Cheng[8.10], we introduce the following dimensionless variables for the shock-transition zone:

$$\bar{u} = u/u_1, \quad \bar{v} = v/v_1, \quad \bar{p} = p/\rho_1 v_1^2, \quad \bar{T} = T/T_o \tag{3.1}$$

$$\bar{\rho} = \rho/\rho_1$$

The transformed normal distance measured from the shock interface, y_2, is defined as

$$\bar{\eta} = \frac{3}{4} \sin\beta \, Re_b \int_{y_2=0}^{y} \left(\frac{\mu_o}{\mu}\right) d\left(\frac{y}{a}\right) \tag{3.2}$$

where $Re_b = \rho_\infty U_\infty a/\mu_o$.

The gray volumetric absorption coefficient, $\rho\kappa$, is assumed to be of the form

$$\alpha = \rho\kappa = Cp^a T^b \tag{3.3a}$$

where C, a and b are constants. For instance, Traugott[12] uses $a = 1$, $b = 4$ and $C = 5.37 \times 10^{-25}$ $sec^2/gm\,^oK$, where p is in units of $dynes/m^2$ and T is in ^oK. The dimensionless form of the absorption coefficient $\bar{\alpha}$ is then written

$$\alpha = \alpha_o(\bar{\alpha}) = \alpha_o(\bar{p}^a \bar{T}^b) \tag{3.3b}$$

where

$$\alpha_o = C(\rho_\infty U_\infty^2)^a T_o^b \tag{3.3c}$$

is based on free stream stagnation conditions.

Substituting Eqs. (3.1) through (3.3) into Eqns. (2.7) through (2.11), one obtains

$$\bar{\rho}\,\bar{v} = 1 \tag{3.4}$$

$$\bar{p} + \bar{v} + \frac{\partial \bar{v}}{\partial \bar{\eta}} = 1 \tag{3.5}$$

$$\bar{u} + \frac{3}{4}\frac{\partial \bar{u}}{\partial \bar{\eta}} = 1 \tag{3.6}$$

$$\frac{\partial \bar{H}}{\partial \bar{\eta}} + \frac{3}{4Pr}\frac{\partial^2}{\partial \bar{\eta}^2}[\bar{H}-(1-Pr)\cos^2\beta\,\bar{u}^2-(1-\frac{4Pr}{3})\sin^2\beta\,\bar{v}^2]$$
$$= (\frac{\Gamma_o \alpha_o a}{Re_b})\,\frac{4}{3}\,\bar{p}^a \bar{T}^{b+5} \tag{3.7}$$

$$\bar{p} = \varepsilon\bar{\rho}\,\bar{T}\csc^2\beta \tag{3.8}$$

The dimensionless total enthalpy is defined as

$$\overline{H} = \overline{T} + \cos^2\beta \ \overline{u}^2 + \sin^2\beta \ \overline{v}^2$$

The linear temperature viscosity law, $\mu = \mu_o T/T_o$, is used to arrive at the right-hand side of Eq. (3.7). The radiative-emisison to convection ratio is defined as

$$\Gamma_o = \frac{4\sigma T_o^4}{\rho_\infty U_\infty c_p T_o} \tag{3.9}$$

where the subscript o indicates its evaluation at free stream stagnation conditions.

Because of the elimination of the coefficient of viscosity through the transformation given by Eq. (3.2), the two momentum equations, (3.5) and (3.6), are explicitly uncoupled from the energy equation. Thus Cheng's[8] forms of the solutions for $\overline{v}$ and $\overline{u}$ are available.

3.2 Transformation of the shock layer equations

Again following Cheng[8,9,10], the shock layer equations are transformed from the (x,y) coordinates to the (x,ψ) Von Mises coordinates. With the introduction of the stream function, ψ, which is related to the velocity components as

$$2\pi Z \ \rho u = \frac{\partial\psi}{\partial y}, \quad 2\pi Z \ \rho v = -\frac{\partial\psi}{\partial x}$$

Eq. (2.1) is then identically satisfied. Equations (2.2) through (2.4), with the use of Eq. (2.6), then become

$$\frac{\partial u}{\partial x} = -\frac{1}{\rho u}\frac{\partial p}{\partial x} + (2\pi Z)^2 \frac{\partial}{\partial\psi}(\mu\rho u \frac{\partial u}{\partial\psi}) \tag{3.10}$$

$$\frac{\partial p}{\partial\psi} + \frac{\kappa_c u}{2\pi Z} = 0 \tag{3.11}$$

$$\frac{\partial H}{\partial x} = (2\pi Z)^2 \frac{\partial}{\partial\psi}\{ \frac{\mu\rho u}{Pr}\frac{\partial}{\partial\psi}[H+(Pr-1)\frac{u^2}{2}]\} - \frac{\alpha 4\sigma T^4}{\rho u} \tag{3.12}$$

The no-slip boundary conditions at the wall, a good approximation in the case of cold walls, are

$$\psi = 0: \quad u = 0, \quad H = H_w$$

The boundary conditions at the outer edge of the shock layer are obtained from Section 2.3:

$$\left.\begin{aligned} \psi = \rho_\infty U_\infty \pi Z^2: \qquad & p = \rho_\infty U_\infty^2 \sin^2\beta \\ & u = U_\infty \cos\beta - \frac{2\pi Z \mu \rho u}{\rho_\infty U_\infty \sin\beta} \frac{\partial u}{\partial \psi} \\ & H = H_\infty - \frac{2\pi Z \mu \rho u}{Pr \rho_\infty U_\infty \sin\beta} \frac{\partial}{\partial \psi}\left[H-(1-Pr)\frac{u^2}{2}\right] + \frac{q_2^R}{\rho_\infty U_\infty \sin\beta} \end{aligned}\right\} \quad (3.13)$$

The outer boundary condition for H is the "psuedo-jump" condition, where

$$q_2^R = \int_\infty^{y_2} \rho \kappa 4 \sigma T^4 \, dy \qquad (3.14)$$

is an integral across the shock-transition zone.

With the introduction of the dimensionless dependent variables

$$\left.\begin{aligned} \bar{p} &= p/(\rho_\infty U_\infty^2 \sin^2\beta) \\ \bar{u} &= u/(U_\infty \cos\beta) \\ \Theta &= (H-H_w)/(H_\infty - H_w) \end{aligned}\right\} \quad (3.15)$$

and the independent variables

$$\xi = x/a, \quad \zeta = [\psi/(\rho_\infty U_\infty \pi Z^2)]^{1/2} \qquad (3.16)$$

Eqs. (3.10) through (3.13) can then be transformed into a form suitable for the calculation of flow over (smooth) blunt bodies.[10] However, the stagnation region, which is the zeroth order term in a series expansion in

ξ , must be considered first, in order to provide initial information for calculations around the body.

In this paper, only the stagnation region will be discussed; in this region the application of the "psuedo-jump" condition and the overall iteration scheme can be illustrated simply.

3.3 The stagnation region

In the stagnation region, $\sin\beta \approx 1$, $\cos\beta \approx \xi$ and $Z/a \approx \xi$. Then, in the leading approximation the following ordinary differential equations are obtained:

$$\bar{u}^2 - \zeta\bar{u}\,\bar{u}' = [\bar{u}/K^2\zeta)]\,[(\bar{u}/\zeta)\bar{u}']' \tag{3.16}$$

$$\bar{u}(0) = 0, \quad \bar{u}(1) = 1 - \bar{u}(1)\bar{u}'(1)/K^2 \tag{3.17}$$

$$-\zeta\bar{u}\,\Theta' = [\bar{u}/(PrK^2\zeta)]\,[(\bar{u}/\zeta)\Theta']' - \Gamma_o\alpha_o\varepsilon a\Theta^9 \tag{3.18}$$

$$\Theta(0) = 0, \quad \Theta(1) = 1-\bar{u}(1)\,\Theta'(1)/(PrK^2)+\bar{q}_2^R \tag{3.19}$$

where $\bar{q}_2^R = q_2^R/(\rho_\infty U_\infty^3/2)$, and primes indicate differentiation with respect to ζ.

The momentum equation (3.16) is uncoupled from the energy equation (3.18) through the use of a linear viscosity-temperature law and the neglect of the tangential pressure gradient term. This term is of order K^2 and is not of importance in the $K^2 = O(1)$ or merged layer regime of flow. (For further discussion of this point, see Cheng.[8]) The solution obtained by Cheng[9] is

$$\bar{u} = PrK^2\zeta/(3k) \tag{3.20}$$

Hence, with the use of (3.20), the final form of the shock layer energy equation and its boundary conditions becomes

$$\left.\begin{aligned} &\Theta'' + 3k\zeta^2\Theta' = (\Gamma_o\alpha_o\varepsilon a)[3k/(PrK^2)]\Theta^9 \\ &\Theta(0) = 0, \quad \Theta(1) = 1 - \Theta'(1)/3k + \bar{q}_2^R \end{aligned}\right\} \quad (3.21)$$

The shock-transition zone energy equation, when specialized to the stagnation region, becomes, with primes indicating differentiation with respect to $\bar{\eta}$

$$\left.\begin{aligned} &\bar{H}'' + \bar{H}' = (4/3)(\Gamma_o\alpha_o\varepsilon a/K^2)\ \varepsilon(\bar{H}-\bar{v}^2)^{10}\ /\bar{v} \\ &\bar{H}(\infty) = 1, \quad \bar{H}(0) = \Theta(1) \end{aligned}\right\} \quad (3.22)$$

where the shock-interface is taken as located at $\bar{\eta} = 0$. Cheng's solution[8,10] for the shock-transition zone momentum equations are

$$\bar{u} = 1 - (1 - \bar{u}_2)e^{-4\bar{\eta}/3} \quad (3.23)$$

and

$$(1-2\bar{v}_*)\bar{\eta} = \ln[(\bar{v}-\bar{v}^*)^{\bar{v}^*}/(1-\bar{v})] + \text{constant} \quad (3.24)^{\dagger}$$

where $\bar{v}^* = \bar{v}_*/(1-\bar{v}_*)$ and $\bar{v}_* = \varepsilon\bar{H}_2$. See Cheng[8] for details of the approximate solution for $\bar{v}$.

In terms of the shock-transition zone variables, $\bar{q}_2^R$ in Eq. (3.21) is

$$\bar{q}_2^R = (4/3)\ (\Gamma_o\alpha_o\varepsilon a/K^2)\int_{\infty}^{0} [\varepsilon(\bar{H} - \bar{v}^2)^{10}/\bar{v}]d\bar{\eta} \quad (3.25)$$

---

†The exact value of the constant of integration is not important. The constant could be fixed by matching $\partial\bar{v}/\partial\bar{\eta}$ with that of the shock layer solution at $\bar{\eta}=0$ or it could be entirely omitted. In both cases, the normal velocity and temperature profiles are expected to be discontinous at the interface to order ε^2. (See Reference 8). In this paper, for simplicity, the constant is omitted.

Since the integrand in Eq. (3.25) is positive and the integration is taken from ∞ to 0, $\bar{q}_2^R$ is a negative number, representing a further mode of heat loss besides conduction at the shock interface.

The overall iteration scheme is as follows: The parameters $\Gamma_o\alpha_o\varepsilon a$, ε , K^2 and Pr are first specified. We decouple Eq. (3.21) from Eq. (3.22) by assuming various values of $\bar{q}_2^R$ to obtain the solution for $\Theta(\zeta)$. From these solutions, $\Theta(1)$ (and hence $\Theta'(1)$) is obtained as a function of the assumed values of $\bar{q}_2^R$.

On the other hand, Eq. (3.22) may be decoupled from Eq. (3.21) by assuming various values of $\Theta(1)$ in the solutions for $\bar{H}$. From these solutions $\bar{q}_2^R$ can be calculated from Eq. (3.25) and obtained as a function of the assumed values of $\Theta(1)$. The intersection of the two "independent" solution curves in the $\bar{q}_2^R - \Theta(1)$ phase plane then gives the final consistent overall solution.

Because of the nonlinearity of Eqs. (3.21) and (3.22), analytical solutions are not expected. The technique of solving the individual decoupled Eqs. (3.21) and (3.22) is a matter of choice. They could be numerically integrated, subject to their respective two-point boundary conditions. However, here their solutions will be obtained by iteration.

To proceed, it is assumed that the right-hand sides of the differential equations in Eqs. (3.21) and (3.22) can be treated as known nonhomogenous terms. Upon applying the respective boundary conditions, we obtain for the shock layer

$$\Theta(z) = \Gamma_o\alpha_o\varepsilon a(3k^{1/3}/PrK^2)\ \{\mathcal{J}_2(z)-\theta_o(z)[\mathcal{J}_2(k^{1/3})+e^{-k}\mathcal{J}_1(k^{1/3})/3k^{2/3}]\}$$

$$+ (1 + \bar{q}_2^R)\theta_o(z) \qquad (3.26)$$

where $z = k^{1/3}\zeta$

$$\left.\begin{aligned} \mathcal{J}_1(z) &= \int_0^z e^{\bar{z}^3}\,\theta^9(\bar{z})d\bar{z} \\ \mathcal{J}_2(z) &= \int_0^z e^{-\bar{z}^3}\,\mathcal{J}_1(\bar{z})d\bar{z} \end{aligned}\right\} \quad (3.27)$$

and

$$\theta_o(z) = I(z)/[I(k^{1/3}) + e^{-k}/(3k^{2/3})]$$

is the solution for the nonradiating stagnation region shock layer.[9] The integral $I(z)$ is defined by

$$I(z) = \int_0^z e^{-z^3}\,dz = \frac{1}{3}\,\gamma(\tfrac{1}{3}, z^3), \quad (3.28)$$

which is related to the incomplete gamma function of order $1/3$ and argument z^3.

For the shock-transition zone, we obtain

$$\bar{H} = 1 - [1-\Theta(1) - \int_0^{\bar{\eta}} e^{\eta'}\bar{q}^R(\eta')\,d\eta']e^{-\bar{\eta}} \quad (3.29)$$

where

$$\bar{q}_2^R(\bar{\eta}) = (\Gamma_o\alpha_o\varepsilon a/K^2)(4/3)\int_\infty^{\bar{\eta}} [\varepsilon(\bar{H}-\bar{v}^2)^{10}/\bar{v}]d\eta \quad (3.30)$$

In Eq. (3.29), $\Theta(1)$ is the value of the dimensionless shock layer enthalpy function at the interface $\zeta = 1$ (or $z = k^{1/3}$), which is assumed known. The shock-transition zone velocity, $\bar{v}$, is in the form given by Eq. (3.24) and is consistent with the assumed $\Theta(1)$.

An alternate form of Eq. (3.29), in terms of $\bar{T}$, can be obtained:

$$\bar{T} = 1 - \bar{v}^2 - [1 - \Theta(1) - \int_0^{\bar{\eta}} e^{\eta'}\bar{q}^R(\eta')\,d\eta']e^{-\bar{\eta}} \quad (3.31)$$

with

$$\bar{q}^R(\bar{\eta}) = (\Gamma_o \alpha_o \varepsilon a/K^2)(4/3) \int_{\infty}^{\bar{\eta}} (\varepsilon \bar{T}^{10}/\bar{v}) d\eta \tag{3.32}$$

4. Results and Discussion

As anticipated in Section 3.3, the intersection of the shock layer and shock-transition zone solution curves in the $\Theta(1) - \bar{q}_2^R$ phase plane would give the final consistent solution. Each point on the respective solution curves is here obtained by iteration, using the energy equations in the form given by Eq. (3.26) and either (3.29) or (3.31). The calculations begin as follows: first, the parameters $\Gamma_o \alpha_o \varepsilon a$, K^2, ε and Pr are fixed. Then the radiationless shock layer solution[9] (i.e. fixing $\Theta(1)$) corresponding to these values of K^2, ε, and Pr, is used to obtain $\bar{v}$ in the shock-transition zone. Then $\bar{T}$ in the shock-transition zone is obtained from Eq. (3.31) as accurately as desired through iteration, the zeroth approximation used being the radiationless solution[8,10] for $\bar{T}$. Thus $\bar{q}_2^R$ can be obtained from Eq. (3.25) for the particular value of $\Theta(1)$; this corresponds to the first circle in Figure 3. The next step is to use this value of $\bar{q}_2^R$ in Eq. (3.16). The iteration begins by using the radiationless solution, θ_o, as the zeroth approximation in the integrals $\mathcal{J}_1$ and $\mathcal{J}_2$. Again, Θ is obtained to the desired accuracy for the particular value of $\bar{q}_2^R$ used, and this yields a new $\Theta(1)$ which corresponds to the first square in Figure 3. In this figure the horizontal paths indicate fixed $\Theta(1)$ and the vertical paths indicate fixed $\bar{q}_2^R$. Such a procedure produces a spiral path in approaching the final consistent solution, as

is dictated by physical reasoning from the implications of such a procedure.

The particular values of parameters for the trajectory shown in the phase plane of Figure 3 are $\Gamma_o \alpha_o \varepsilon a = 100$, $K^2 = 1$, $\varepsilon = 0.1428$ ($\gamma = 1.40$) and $Pr = 0.75$. From Figure 1 we see that this combination of $\Gamma_o \alpha_o \varepsilon a$ and K^2 falls in the regime where radiative emmision effects are of "order one". When calculations are performed in the "weak radiation" regime, only a signle vertical path in the $\Theta(1) - \bar{q}_2^R$ plane is necessary, which corresponds to a small perturbation away from the radiationless case. Since for the "strong radiation" regime, radiation becomes the dominant mode of energy transfer, this regime must be considered separately. The techniques described here then encompass both the order-one and weak radiation cases, but not the strong radiation case.

Of particular interest are the contributions of the various modes of heat transfer to the body surface. In this discussior the local surface heat transfer rates will be made dimensionless with resepct to $\rho_\infty U_\infty (H_\infty - H_w)$ ($\tilde{=} \rho_\infty U_\infty^3/2$ for a highly cooled wall and a hypersonic free stream) to give the coefficients of heat transfer C_H. The contributions from the radiative heat flux to the wall are obtained by the following integrals:

$$q_w^R = q_w^{R,SL} + q_w^{R,STZ} = \frac{1}{2} \int_o^{y_s} \rho \kappa 4 \sigma T^4 dy_{SL} + \frac{1}{2} \int_{y_s}^{\infty} \rho \kappa 4 \sigma T^4 dy_{STZ} \qquad (4.1)$$

The radiative heat flux to the wall is taken as half of the total energy radiated from the shock layer and the shock-transition zone. The thin

radiating slab approximation is implicit in this consideration. This would be consistent with the thin-layer approximations already used in treating the shock layer and shock-transition zone. In terms of the heat transfer coefficient and of the respective variables used for the two layers, then:

$$C_H^{R,SL} = [(\Gamma_o \alpha_o \varepsilon a) 3k^{2/3}/(2PrK^2)] \int_0^{k^{1/3}} \Theta^9(\bar{z}) d\bar{z} \tag{4.2}$$

$$C_H^{R,STZ} = [(\Gamma_o \alpha_o \varepsilon a) 4/(6K^2)] \int_0^{\infty} (\varepsilon \bar{T}^{10}/\bar{v}) d\bar{\eta} = - \bar{q}_2^R/2 \tag{4.3}$$

The form of the absorption coefficient given by Eq. (3.3) shows its dependence on pressure. In the shock layer $\bar{p} = 1$ in the stagnation region. In the shock-transition zone. $\bar{p} = \varepsilon\bar{T}/\bar{v}$. As with the $\bar{p}$ term in the momentum equation (3.5), the contribution of $\varepsilon\bar{T}/\bar{v}$ does not become important until the region near the shock interface is approached, where $\bar{v} = O(\varepsilon)$. In general, because of this pressure dependence, it is expected that the contribution to C_H^R from the shock layer, $C_H^{R,SL}$, is more important than that from the shock transition zone, $C_H^{R,STZ}$. However, for decreasing K^2, the shock layer thickness, $y_s/\varepsilon a$, decreases (for the case of a cold wall) while the shock transition zone stretches out. In this case, $C_H^{R,STZ}$ overtakes $C_H^{R,SL}$. This is shown in Figure 4 for $\Gamma_o \alpha_o \varepsilon a = 10$, $\varepsilon = 0.1428$, $Pr = 0.75$, where C_H^R is broken down into the contributions from the two regions. Since radiative emission decreases the shock layer temperature, and hence its gradient, the convective heat transfer coefficient, C_H^C, becomes lower than in the

radiationless case C_{H_o}. The total heat transfer coefficient, C_H^{R+C}, is asymptotic to C_H^R at the higher end of K^2 and asymptotic to C_H^C at the lower end of K^2. Radiative effects become negligible for small K^2 for a fixed $\Gamma_o\alpha_o\varepsilon a$, since the lower temperature level in this range of K^2 puts the radiative energy transfer in the "weak" range, (see Figure 1). The effect of changes in $\Gamma_o\alpha_o\varepsilon a$ on C_H^{R+C} is shown in Figure 5.

The dimensionless enthalpy function at the shock interface, $\Theta(1)$, is shown as a function of K^2 in Figure 6, and the shock detachment distance, or the thickness of the shock layer, is shown in Figure 7. Because the overall temperature level in the shock layer decreases for increasing $\Gamma_o\alpha_o\varepsilon a$, the shock layer thickness decreases correspondingly.

Some typical stagnation region profiles are shown in Figure 8 for the case of $K^2 = 1$. The discontinuity in the temperature profile at the shock interface is as expected in the thin shock layer treatment [see Footnote to Eq. (3.24)]. The explicit form of the tangential and normal velocity profiles are identical to the radiationless case in their respective transformed plane. In particular, the tangential velocity differs from the corresponding radiationless case in the physical plane because of coordinate stretching only. The normal velocity differs from the corresponding radiationless case because of the modifications upon $\overline{H}_2$ in addition to coordinate stretching. These differences become more pronounced with increasing $\Gamma_o\alpha_o\varepsilon a$. The effects of radiative cooling on the temperature profile is illustrated. The temperature profiles become less steep near the wall and the entire

shock layer and shock-transition zone decrease in thickness with increasing importance of radiation.

Throughout this paper, comparisons of the present solutions have been made only with the corresponding radiationless solutions. Comparisons with other radiative solutions at the higher Reynolds number range have not been made. Because of the various differing assumptions on the gas properties, the equation of state and the radiation model, such comparisons are not found to be meaningful.

Since the gas near the cold wall and in the outer portion of the shock-transition zone is relatively cool, reabsorption effects would become important. An extension to the corresponding low Reynolds number hypersonic flow, including reabsorption effects, is under way.

References

1. Cheng, P. and Vincenti, W. G., "Inviscid Radiating Flow Over A Blunt Body", Journal of Fluid Mechanics, Vol. 27, Part 4, 1967, pp. 625-646.

2. Howe, J. T. and Viegas, J. R., "Solutions of the Ionized Radiating Shock Layer Including Reabsorption and Foreign Species Effects and Stagnation Region Heat Transfer," TR R-159, 1963, NASA.

3. Hoshizaki, H. and Wilson, K. H., "Viscous, Radiating Shock Layer about a Blunt Body," AIAA Journal, Vol. 3, No. 9, Sept. 1965, pp. 1614-1622.

4. Hoshizaki, H. and Wilson, K. H., "Convective and Radiative Heat Transfer During Superorbital Entry," AIAA Journal, Vol. 5, No. 1, Jan. 1967, pp. 25-35.

5. Burggraf, O. R., "Asymptotic Solution for the Viscous Radiating Shock Layer," AIAA Journal, Vol. 4, No. 10, October 1966, pp. 1725-1734.

6. Hayes, W. D. and Probstein, R. F., Hypersonic Flow Theory, 1st ed., Academic Press, New York, 1959, Chap. 10.

7. Bush, W. B., "On the Viscous Hypersonic Blunt-Body Problem," Journal of Fluid Mechanics, Vol. 20, Part 3, 1964, pp. 353-368.

8. Cheng, H. K., "Viscous Hypersonic Blunt-Body Problems and the Newtonian Theory," Proceedings of International Symposium on Fundamental Phenomena in Hypersonic Flow, 1964, Hall, J. G. ed., Cornell University Press, 1966, pp. 90-131.

9. Cheng, H. K., "Hypersonic Shock-Layer Theory of the Stagnation Region at Low Reynolds Number," Proceedings of the 1961 Heat Transfer and Fluid Mechanics Institute, Stanford University Press, 1961, pp. 161-175.

References (cont.)

10. Cheng, H. K., "The Blunt-Body Problem in Hypersonic Flow at Low Reynolds Number," Report No. AF 1285-A-10, June 1963, Cornell Aeronautical Laboratory, Buffalo, New York.

11. Probstein, R. F. and Kemp, N., "Viscous Aerodynamic Characteristics in Hypersonic Rarefied Gas Flow," Journal of Aerospace Science, Vol. 27, No. 3, March 1960, pp. 174-192, 218.

12. Traugott, S. C., "A Differential Approximation for Radiative Transfer with Application to Normal Shock Structure," Proceedings of the 1963 Heat Transfer and Fluid Mechanics Institute, Stanford University Press, 1963, pp. 1-13.

JTCL/cl



Figure 1. The radiation parameter.



Figure 2. Thin shock structure and shock layer notation. Schematic.



Figure 3. The $\theta(1) - \overline{q}_2^R$ phase plane.



Figure 4. Contributions to the heat transfer coefficient: convective heat transfer and radiative heat transfer from shock layer and shock-transition zone as functions of the rarefaction parameter.



Figure 5. Overall heat transfer coefficient as a function of the rarefaction parameter.



Figure 6. Dimensionless enthalpy at the outer-edge of the shock layer as a function of the rarefaction parameter.



Figure 7. Shock layer thickness as a function of the rarefaction parameter.



Figure 8. Dimensionless temperature, tangential velocity and normal velocity profiles for various values of the radiation parameter.

DISTRIBUTION LIST

INDIVIDUAL

DR H N ABRAMSON ABCFGHI
SOUTHWEST RESEARCH INST
8500 CULEBRA ROAD
SAN ANTONIO TEXAS
78228

DR R A ALLEN
AVCO MFG CORPORATION
RESEARCH LABORATORY
2385 REVERE BEACH PARKWAY
EVERETT MASSACHUSETTS
02149

PROFESSOR I AMDUR B
MASSACHUSETTS INSTITUTE
OF TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

PROFESSOR T L BAILEY B
DEPARTMENT OF PHYSICS
UNIVERSITY OF FLORIDA
GAINESVILLE FLORIDA
32603

MR MYLES B BERG AB
AEROSPACE CORPORATION
P O BOX 1308
SAN BERNARDINO CALIFORNIA
92402

PROF D BERSHADER B D
DEPT OF AERONAUTICAL
ENGINEERING
STANFORD UNIVERSITY
STANFORD CALIFORNIA
94305

DR MARTIN H BLOOM ABCDFI
POLYTECH INST OF BROOKLYN
GRADUATE CENTER
DEPT OF AEROSPACE
ENG & APPLIED MECHANICS
FARMINGDALE N Y

PROFESSOR S BODGONOFF B
GAS DYNAMICS LABORATORY
FORRESTAL CAMPUS
PRINCETON UNIVERSITY
PRINCETON NEW JERSEY
08540

DR B N PRIDMORE BROWN
ABDG
NORTHROP CORPORATION
NORAIR DIV
HAWTHORNE CALIFORNIA
90250

PROF ODUS R BURGGRAF
OHIO STATE UNIVERSITY
COLUMBUS OHIO
43210

DR STANLEY BYRON AB
AERONUTRONICS APPLIED
RESEARCH LAB
FORD ROAD
NEWPORT BEACH CALIFORNIA
92660

PROF J E CERMAK ABDFH
PROF IN CHARGE
FLUID MECHS PRO
COLLEGE OF ENGINEERING
COLORADO STATE UNIVERSITY
FT COLLINS COLORADO
80521

PROF R D CESS
STATE UNIV OF NEW YORK
STONY BROOK LONG ISLAND
NEW YORK
11790

PROF ALLEN CHAPMANN
CHAIRMAN BHI
MECHANICAL ENGIN DEPT
WILLIAM M RICE INSTITUTE
BOX 1892
HOUSTON TEXAS
77001

ATT L I CHASEN
GENERAL ELECTRIC COMPANY
MISSILE & SPACE DIVISION
MGR MSD LIB B
P O BOX 8555
PHILADELPHIA PENNSYLVANIA
19104

DR W J CHRISTIAN BF
RESEARCH FOUNDATION
10 WEST 35TH STREET
CHICAGO ILLINOIS
60616

DR PING CHENG
NASA AMES
MOFFETT FIELD CALIFORNIA
94035

PROFESSOR J F CLARKE
COLLEGE OF AERONAUTICS
CRANFIELD BUCKINGHAMSHIRE
ENGLAND

DR C COOK B
STANFORD RESEARCH
INSTITUTE
MENLO PARK CALIFORNIA
94025

MR L G COOPER BG
THE MARTIN COMPANY
DEPARTMENT 2550
MAIL NO J 3072
BALTIMORE MARYLAND
21203

DR L F CRABTREE BG
MINSTRY OF AVIATION
ROYAL AIRCRAFT ESTABLISH
AERPDYNAMICS DEPARTMENT
FARNBOROUGH HANTS ENGLAND

MR R H CRAMER
APPLIED PHYSICS LAB
JOHN HOPKINS UNIV
8621 GEORGIA AVENUE
SILVER SPRING MARYLAND
21218

PROFESSOR G E DUVALL AB
DEPARTMENT OF PHYSICS
WASHINGTON STATE UNIV
PULLMAN WASHINGTON
99164

MR BILL EAST BG
LOCKHEED AIRCRAFT CORP
CALIFORNIA DIVISION
HYDRODYNAMICS RESEARCH
BURBANK CALIFORNIA
91503

PROFESSOR E R G ECKERT B
241 MECHANICAL
ENGINEERING BLDG
UNIVERSITY OF MINNESOTA
MINNEAPOLIS MINNESOTA
55455

DR GEORGE EMANUEL
AEROPHYSICS LABORATORY
AEROSPACE CORPORATION
2400 E EL SEGUNDO BLVD
EL SEGUNDO CALIFORNIA
90246

PROF J A FAY
DEPARTMENT OF
MECHANICAL ENGINEERING
M I T
CAMBRIDGE MASSACHUSETTS
02139

DR SAUL FELDMAN B
HELIODYNE CORPORATION
7810 BURNET AVENUE
VAN NUYS CALIFORNIA
91405

PROFESSOR CARLO FERRARI
INSTITUTO DI
MECCANICA APPLICATA
POLITECNICO DI TORINO
TORINO ITALY

GAIL T FLESHER 44 AB
GM DEFENSE RESEARCH LAB
BOX T
SANTA BARBARA CALIFORNIA
93102

DR N C FREEMAN B
AERONAUTICS DEPARTMENT
IMPERIAL COLLEGE
LONDON S W 7 ENGLAND

PROFESSOR D GILBARG B C
STANFORD UNIVERSITY
STANFORD CALIFORNIA
94305

DR ROBERT GOULARD
SCHOOL OF AERONAUTICS &
ENGINEERING SCIENCE
PURDUE UNIVERSITY
LAFAYETTE INDIANA
47907

PROFESSOR R GREIF
DEPARTMENT OF
MECHANICAL ENGINEERING
UNIVERSITY OF CALIFORNIA
BERKELEY CALIFORNIA
94704

PROFESSOR H R GRIEM
UNIVERSITY OF MARYLAND
COLLEGE PARK MARYLAND
20742

PROF ROBERT A GROSS
DEPARTMENT OF
MECHANICAL ENGINEERING
COLUMBIA UNIVERSITY
NEW YORK NEW YORK
10027

DR J S GRUSXCZYNSKI
SPACE SCIENCES LABORATORY
GENERAL ELECTRIC
MISSILES & SPACE DIVISION
KING OF PRUSSIA PENN
19406

DR R J HAKKINEN
CHIEF SCIENTIST
PHYSICAL SCIENCES DEPT B
RESEARCH AND DEVELOPMENT
DOUGLAS MISS & SPACE DIV
SANTA MONICA CALIFORNIA
90406

DR G HALL BD
CORNELL AERONAUTICAL
LABORATORY INC
P O BOX 235
BUFFALO NEW YORK
14221

DR C F HANSEN
GENERAL MOTORS
DEFENSE LABORATORIES
SANTA BARBARA CALIFORNIA
93101

PROF R P HARRINGTON
HEAD B
DEPT OF AERONAUTICAL
ENGINEERING
UNIVERSITY OF CINCINNATI
CINCINNATI OHIO
45221

DR ANTHONY G HEARNE
UKAEA CULHAM LABORATORY
AILHAM ABINGDON
BERSHIRE ENGLAND

MR A HERTZBERG
UNIVERSITY OF WASHINGTON
SEATTLE WASHINGTON
98105

DR R HILLENDAHL
PHYSICAL SCIENCES LAB
LOCKHEED MISS & SPACE CO
3251 HANOVER AVENUE
PALO ALTO CALIFORNIA
94304

DR J S HOLDHUSEN B
FLUIDYNE ENGINEERING
CORPORATION
5900 OLSON MEMORIAL HGWY
MINNEAPOLIS MINNESOTA
55455

DR ROBERT N HOLLYER JR B
RESEARCH LABORATORIES
GENERAL MOTORS CORP
12 MILE AND MOUND ROAD
WARREN MICHIGAN
48090

PROFESSOR M HOLT ABCH
DIVISION OF AERONAUTICAL
SCIENCES
UNIVERSITY OF CALIFORNIA
BERKELEY CALIFORNIA
94720

DR H HOSHIZAKI
AEROSPACE SCIENCES LAB
LOCKHEED MISS & SPACE CO
RESEARCH LABORATORIES
3251 HANOVER STREET
PALO ALTO CALIFORNIA
94304

PROFESSOR H C HOTTEL AB
DEPT OF CHEMICAL ENGINEER
MASSACHUSETTS INSTITUTE
OF TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

MR C C HUDSON AB
SANDIA CORPORATION
SANDIA BASE
ALBUQUERQUE NEW MEXICO
87115

R G JORDAN BCE
OAK RIDGE GASEOUS
DIFFUSION PL
UNION CARBIDE CORP
NUCLEAR DIV
P O BOX P
OAK RIDGE TENNESSEE

DR ARTHUR KANTROWITZ
DIRECTOR B
AVCO EVERETT RESEARCH
LABORATORY
2385 REVERE BEACH PARKWAY
EVERETT MASSACHUSETTS
02149

PROFESSOR K KARAMCHETI B
DEPT OF AERONAUTICAL
ENGINEERING
STANFORD UNIVERSITY
STANFORD CALIFORNIA
94305

DR J KULANDER
LOCKHEED MISS & SPACE CO
3251 HANOVER AVENUE
PALO ALTO CALIFORNIA
94304

A R KUHLTHAU DIRECTOR B
RESEARCH LABORATORIES FOR
THE ENGINEERING SCIENCES
THORNTON HALL
UNIVERSITY OF VIRGINIA
CHARLOTTESVILLE VIRGINIA
22903

PROF E V LAITONE BFGHI
UNIVERSITY OF CALIFORNIA
BERKELEY CALIFORNIA
94720

PROFESSORS HARVEY LAM
FORRESTAL RESEARCH CENTER
DEPT OF AERO ENGIN &
MECHANICAL SCIENCE
PRINCETON UNIVERSITY
PRINCETON NEW JERSEY
08541

DR FRANK LANE ABF
GENERAL APPLIED
SCIENCE LAB
MERRICK & STEWART AVENUES
WESTBURY LONG ISLAND N Y
11590

PROFESSOR L LEES
GUGGENHEIM AERONAUTCL LAB
CALIFORNIA INST OF TECH
PASADENA CALIFORNIA
91109

PROFESSOR M LESSEN HEAD
DEPARTMENT OF MECHANICAL
ENGINEERING
THE UNIV OF ROCHESTER
RIVER CAMPUS STATION
ROCHESTER NEW YORK
14627

MR PHILIP LEVINE
2 POWELL ROAD
DANVERS MASSACHUSETTS
01923

DR S L LEVY DIRECTOR B
MIDWEST RESCH INSTITUTE
PHYSICS AND MATHEMATICS
DIVISION
425 VOLKER BOULEVARD
KANSAS CITY MISSOURI
64110

DR H G LEW B
MANAGER GAS DYNAMICS
SPACE SCIENCE LABORATORY
SPACE TECHNOLOGY CENTER
KING OF PRUSSIA PENN
19406

PROFESSOR TING YI LI B
DEPT OF AERO &
ASTRONAUTICAL ENGR
OHIO STATE UNIVERSITY
2036 NEIL AVENUE
COLUMBUS OHIO
43210

PROFESSOR WILBERT LICK
CASE WEST UNIV.
CLEVELAND, OHIO

PROF M J LIGHTHILL
UNIVERSITY OF LONDON
LONDON ENGLAND

PROFESSOR C C LIN ABDH
MASSACHUSETTS INSTITUTE
OF TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

PROFESSOR MAEDER B
DIVISION OF ENGINEERING
BROWN UNIVERSITY
PROVIDENCE RHODE ISLAND
02912

PROFESSOR D MINTZER B
DEPARTMENT OF MECHANICAL
ENGINEERING
NORTHWESTERN UNIVERSITY
TECHNOLOGICAL INSTITUTE
EVANSTON ILLINOIS
60201

DR F K MOORE
DIRECTOR
THERMAL ENGINEERING DEPT
208 UPSON HALL
CORNELL UNIVERSITY
ITHACA NEW YORK
14850

DR H T NAGAMATSU AB
GENERAL ELECTRIC COMPANY
RES & DEV CENTER K 1
P O BOX 8
SCHENECTADY NEW YORK
12301

DR YOSHINARI NAKAGAWA
NCAR
UNIVERSITY OF COLORADO
BOULDER COLORADO
80304

PROFESSOR R M NEREM
DEPT AERO & ASTRO ENGIN
OHIO STATE UNIVERSITY
COLUMBUS OHIO
43210

PROF D B OLFE
UNIV OF CALIFORNIA
AT SAN DIEGO
DEPARTMENT OF
AEROSPACE ENGINEERING
LA JOLLA CALIFORNIA
92037

PROF A K OPPENHEIM AB
DIVISION OF MECHANICAL
ENGINEERING
UNIVERSITY OF CALIFORNIA
BERKELEY CALIFORNIA
94720

PROFESSOR PAI
INSTITUTE FOR
FLUID MECHANICS AND
APPLIED MATHEMATICS
UNIVERSITY OF MARYLAND
COLLEGE PARK MARYLAND
20740

DR ADRIAN PALLONE
AVCO MISSILE SYSTEMS DIV
201 LOWELL STREET
WILMINGTON MASSACHUSETTS
01887

PROF G N PATTERSON B
INSTITUTE FOR AEROSPACE
STUDIES
UNIVERSITY OF TORONTO
TORONTO 5 CANADA

PROF S S PENNER
UNIV OF CALIFORNIA
AT SAN DIEGO
DEPARTMENT OF
AEROSPACE ENGINEERING
LA JOLLA CALIFORNIA
92037

DR GLEN PIPPERT B
MASS INST OF TECHNOLOGY
M I T LINCOLN LABORATORY
BOX 73
LEXINGTON MASSACHUSETTS
02173

PROF R F PROBSTEIN
ABCDFHI
DEPT OF MECHANICAL ENGIN
MASSACHUSETTS INST OF
TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

PROF E L RESLER ABCDHI
GRADUATE SCHOOL OF
AERONAUTICAL ENGINEERING
CORNELL UNIVERSITY
ITHACA NEW YORK
14851

DR ALFRED RITTER ABGHI
THERM ADVANCED RESCH INC
100 HUDSON CIRCLE
ITHACA NEW YORK
14850

MISS MARY ROMIG
THE RAND CORP
1700 MAIN STREET
SANTA MONICA CALIFORNIA
90401

DR PETER ROSE
AVCO MFG CORPORATION
RESEARCH LABORATORY
2385 REVERE BEACH PARKWAY
EVERETT MASSACHUSETTS
02149

SINCLAIR M SCALA MGR AB
SPACE SCIENCES LAB
GENERAL ELECTRIC CO
P O BOX 8555
PHILADELPHIA PENNSYLVANIA
19101

PROFESSOR W R SEARS
ABCDFI
4927 PACIFICA DRIVE
SAN DIEGO CALIFORNIA
92109

DR E E SECHLER ABCDEFGHI
EXECUTIVE OFFIC FOR AERO
CALIFORNIA INSTITUTE OF
TECHNOLOGY
PASADENA CALIFORNIA
91109

PROFESSOR A H SHAPIRO BDH
DEPART OF MECHANICAL
ENGINEERING
MASSACHUSETTS INST OF
TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

DR MARTIN P SHERMAN
HIGH ALTITUDE
AERODYNAMICS LABORATORY
GENERAL ELECTRIC
MISSILES & SPACE DIVISION
KING OF PRUSSIA PENN
19406

PROFESSOR P SHERMAN B
AERONAUTICAL ENGIN DEPT
UNIVERSITY OF MICHIGAN
ANN ARBOR MICHIGAN
48108

DR J D SHREVE JR BH
SANDIA CORPORATION
SANDIA BASE
ALBUQUERQUE NEW MEXICO
81115

PROF E M SPARROW
DEPARTMENT OF
MECHANICAL ENGINEERING
UNIV OF MINNESOTA
MINNEAPOLIS MINNESOTA
55455

BARBARA SPENCE
TECHNICAL LIB AB
AVCO EVERETT RESEARCH LAB
2385 REVERE BEACH PARKWAY
EVERETT MASSACHUSETTS
02149

PROF H B STILLWELL CHMN B
DEPT OF AERONAUTICAL AND
ASTRONAUTICAL ENGINEERING
UNIVERSITY OF ILLINOIS
URBANA ILLINOIS
61803

PROFESSOR R G STONER
DEPARTMENT OF PHYSICS
ARIZONA STATE UNIVERSITY
TEMPE ARIZONA
85281

DR L TALBOT ABI
UNIVERSITY OF CALIFORNIA
DEPARTMENT OF ENGINEERING
BERKELEY CALIFORNIA
94720

DR R N THOMAS B
UNIVERSITY OF COLORADO
BOULDER COLORADO
80302

M J THOMPSON ABDH
DEFENSE RESEARCH LAB
UNIVERSITY OF TEXAS
P O BOX 8029
AUSTIN TEXAS
78712

DR S C TRAUGOTT
MAIL H-3033
THE MARTIN COMPANY
BALTIMORE MARYLAND
21203

DR L TRILLING B
DEPT OF AERONAUTICS AND
ASTRONAUTICS
MASSACHUSETTS INST OF
TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

PROF MILTON VAN DYKE B
DEPT OF AERONAUTICAL
ENGINEERING
STANFORD UNIVERSITY
STANFORD CALIFORNIA
94305

PROFESSOR J K VENNARD BCD
DEPT OF CIVIL ENGINEERING
STANDARD UNIVERSITY
STANFORD CALIFORNIA
94305

PROF WALTER G VINCENTI B
DEPT OF AERONAUTICS
AND ASTRONAUTICS
STANFORD UNIVERSITY
STANFORD CALIFORNIA
94305

DR M VINOKUR
AEROSPACE SCIENCES LAB
LOCKHEED MISS & SPACE CO
3251 HANOVER AVENUE
PALO ALTO CALIFORNIA
94304

PROF R VISKANTA
DEPARTMENT OF
MECHANICAL ENGINEERING
PURDUE UNIVERSITY
LAFAYETTE INDIANA
47907

PROFESSOR P P WEGENER B
DEPT OF ENGINEERING AND
APPLIED SCIENCES
YALE UNIVERSITY
NEW HAVEN CONNECTICUT
06520

CHARLES A WHITNEY
SMITHSONIAN ASTROPHYSICAL
OBSERVATORY
HARVARD COLLEGE OBSERVTY
HARVARD UNIVERSITY
CAMBRIDGE MASSACHUSETTS
02138

DR K H WILSON
AEROSPACE SCIENCES LAB
LOCKHEED MISSILES AND
SPACE COMPANY
3251 HANOVER STREET
PALO ALTO CALIFORNIA
94304

DR W H WURSTER
CORNELL AERO LABORATORY
BUFFALO NEW YORK
14221

DR S L ZIEBERG HEAD AB
GAS DYNAMICS SECT
FLUID MECHANICS DEPT
BLDG B 1 ROOM 1320
AEROSPACE CORPORATION
SAN BERNARDINO CALIFORNIA
92402

PROFESSOR F ZWICKY ABCFH
DEPARTMENT OF PHYSICS
CALIFORNIA INSTITUTE OF
TECHNOLOGY
PASADENA CALIFORNIA
91109

AERO DEPARTMENT B
IMPERIAL COLLEGE
LONDON S W 7 ENGLAND

BATTELLE DEFENDER
INFORMATION
ANALYSIS CENTER B
BATTELLE MEMORIAL INST
505 KING AVENUE
COLUMBUS OHIO
43201

CHIEF OF AERODYNAMICS B
AVCO MISSILE SYSTEMS DIV
201 LOWELL STREET
WILMINGTON MASSACHUSETTS
01887

COMMANDING OFFICER ABCDGH
NROTC & NAVAL
ADMINISTRATIVE UNIT
MASS INST OF TECHNOLOGY
CAMBRIDGE MASSACHUSETTS
02139

CONDUCTRON CORPO
3475 PLYMOUTH ROAD
P O BOX 614
ANN ARBOR MICHIGAN
48107

EDITOR ABCDEFGHI
APPLIED MECHANICS REVIEW
SOUTHWEST RESEARCH INST
8500 CULEBRA ROAD
SAN ANTONIO TEXAS
78206

ENGINEERING DIVISION B
CALIFORNIA INSTITUTE OF
TECHNOLOGY
PASADENA CALIFORNIA
91109

ENGINEERING LIBRARY B
UNIVERSITY OF SOUTHERN
CALIFORNIA
BOX 77929
LOS ANGELES CALIFORNIA
90007

LOCKHEED MISSILES & SPACE
COMPANY
TECHNICAL INFORMATION CTR
ABDEFH
3251 HANOVER STREET
PALO ALTO CALIFORNIA
94301

NIELSEN ENGINEERING &
RESEARCH COMPANY BDG
P O BOX 595
LOS ALTOS CALIFORNIA
94022

NORTH AMER AVIATION INC B
SPACE & INFORMATION
SYSTEMS DIV
12214 LAKEWOOD BLVD
DOWNEY CALIFORNIA
90240

SCHOOL OF APPLIED MATH
INDIANA UNIVERSITY
ABCDFGHI
BLOOMINGTON INDIANA
47401

SUPERSONIC GASDYNAMICS
RES LAB B
DEPT OF MECHANICAL
ENGINEERING
MC GILL UNIVERSITY
MONTREAL 12 QUEBEC CANADA

TECHNICAL REPORTS CTR B
BELL TELEPHONE LABS INC
WHIPPANY LABORATORIES
WHIPPANY NEW JERSEY
07981

GOVERNMENT AGENCIES

AFAPL (APRC)
AB
WRIGHT PATTERSON AFB OHIO
45433

AFOSR (SREM) ABCDEFG
1400 WILSON BOULEVARD
ARLINGTON VA
22209

ARL (ARN) BUILDING 450
ABDH
WRIGHT PATTERSON AFB OHIO
45433

ADVANCED RESH PROJ AGENCY
RESEARCH & DEVELOPMENT
FIELD UNIT
ATTN MR TOM BRUNDAGE B
APO 146 BOX 271
SAN FRANCISCO CALIFORNIA
96346

MR H J ALLEN
NASA
AMES RESEARCH CTR NASA
MT VIEW CALIFORNIA
94040

DR B S BALDWIN
NASA
AMES RESEARCH CTR NASA
MT VIEW CALIFORNIA
94040

DR F D BENNETT AB
EXTERIOR BALLISTICS LAB
BALLISTICS RESEARCH LABS
ABERDEEN PROV GROUND MD
21005

CHIEF OF RESEARCH &
DEVELOPMENT
OFFICE OF CHIEF OF STAFF
ABDEFH
DEPARTMENT OF THE ARMY
THE PENTAGON
WASHINGTON D C

CHIEF OF NAVAL RESEARCH
DEPARTMENT OF THE NAVY
ATTN CODE 421 ABI
WASHINGTON D C
20360

CHIEF OF NAVAL RESCH 03
DEPARTMENT OF THE NAVY
ATTN CODE 438 ABCDEFGHI
WASHINGTON D C
20360

CHIEF OF NAVAL RESEARCH
DEPARTMENT OF THE NAVY
ATTN CODE 461 ABDFGHI
WASHINGTON D C
20360

COMMANDER R & T DIVISION
AIR FORCE SYSTEMS COMMAND
ATTN L M MEDGEPETH
(APIP 1) BCE
WRIGHT PATTERSON AFB OHIO
45433

COMMANDER
NAVAL AIR SYSTEMS COMMAND
DEPARTMENT OF THE NAVY
ATTN CODE AIR 3032 B
WASHINGTON D C
20360

COMMANDER
NAVAL ORDNANCE LABORATORY
ATTN AEROPHYSICS DIVISION
ABDG
WHITE OAK
SILVER SPRING MD
20910

COMMANDER
NAVAL ORDNANCE LABORATORY
ATTN LIBRARIAN ABCDEFGHI
WHITE OAK
SILVER SPRING MD
20910

COMMANDER
NAVAL ORDNANCE LABORATORY
ATTN DR A E SEIGEL ABCD
WHITE OAK
SILVER SPRING MD
20910

COMMANDER
NAVAL ORDNANCE SYSTEMS
COMMAND
ATTN ORD 035 ABDGI
WASHINGTON D C
20360

COMMANDER
NAVAL ORDNANCE SYSTEMS
COMMAND
ATTN ORD 913 (LIBRARY)
ABCDEFGHI
WASHINGTON D C
20360

COMMANDER
NAVAL ORDNANCE TEST STA
ATTN CODE 753 ABCDEG
CHINA LAKE CALIFORNIA
93557

COMMANDER
NAVAL ORDNANCE TEST STA
ATTN CODE 5014 AB
CHINA LAKE CALIFORNIA
93557

COMMANDER
NAVAL MISSILE CENTER
ATTN TECHNICAL LIBRARY
ABCDEFGHI
POINT MUGU CALIFORNIA
93041

COMMANDER
NAVAL WEAPONS LABORATORY
ATTN TECHNICAL LIBRARY
ABCDEFGHI
DAHLGREN VA
22418

COMMANDING GENERAL
ARMY ENGIN R & D LABS
TECH DOCUMENTS CENTER
ABCDEFGH
FORT BELVOIR VIRGINIA
22060

COMMANDING OFFICER ABCEGH
ARMY RESEARCH OFFICE
BOX CM DUKE STATION
DURHAM NORTH CAROLINA
27706

COMMANDING OFFICER
ARMY RESEARCH OFFICE
ATTN ESD ARUD ABCDEH
BOX CM DUKE STATION
DURHAM NORTH CAROLINA
27706

COMMANDING OFFICER
AND DIRECTOR
DAVID TAYLOR MODEL BASIN
ATTN CODE 042 ABCDEFGHI
WASHINGTON D C
20007

COMMANDING OFFICER AND
DIRECTOR
DAVID TAYLOR MODEL BASIN
ATTN CODE 046 B
WASHINGTON D C
20007

COMMANDING OFFICER AND
DIRECTOR
DAVID TAYLOR MODEL BASIN
ATTN CODE 108 ABCDHI
WASHINGTON D C
20007

COMMANDING OFFICER AND
DIRECTOR
DAVID TAYLOR MODEL BASIN
ATTN CODE 800 ABCDGH
WASHINGTON D C
20007

COMMANDING OFFICER
ATTN TECH LIB (BLDG) 313
ABCDEFGHI
ABERDEEN PROV GROUND MD
21005

COMMANDING OFFICER &
DIRECTOR BCDHI
NAVAL CIVIL ENGINEERING
LABORATORY
PORT HUENEME CALIFORNIA
93041

COMMANDING OFFICER
OFFICE OF NAVAL RESEARCH
ABCDEFGHI
BRANCH OFFICE (25)
BOX 39
FPO NEW YORK NEW YORK
09510

COMMANDING OFFICER
OFFICE OF NAVAL RESEARCH
ABCDEFGHI
BRANCH OFFICE
1030 E GREEN STREET
PASADENA CALIFORNIA
91101

COMMANDING OFFICER
OFFICE OF NAVAL RESEARCH
ABCDEFGHI
BRANCH OFFICE
219 SOUTH DEARBORN STREET
CHICAGO ILLINOIS
60604

COMMANDING OFFICER
OFFICE OF NAVAL RESEARCH
ABCDEFGHI
BRANCH OFFICE
495 SUMMER STREET
BOSTON MASSACHUSETTS
02210

DEFENSE DOCUMENT CTR 20
ABCDEFGHI
CAMERON STATION
ALEXANDRIA VIRGINIA
22314

DIRECTOR B
WEAPONS SYSTEMS
EVALUATION GR
WASHINGTON D C
20305

DIRECTOR OF
RESEARCH CODE RR BCD
NATIONAL AERONAUTICS &
SPACE ADMINISTRATION
600 INDEPENDENCE AVE S W
WASHINGTON D C
20546

DIRECTOR
NATIONAL BUREAU OF STDS
ATTN LIBRARY AB
WASHINGTON D C
20234

DIRECTOR
NATIONAL BUREAU OF STDS
ATTN DR D H TSAI AB
WASHINGTON D C
20234

DIRECTOR 06
U S NAVAL RESEARCH LAB
ATTN CODE 2027
ABCDEFGHI
WASHINGTON D C
20390

DR A J EGGERS JR
NASA
AMES RESEARCH CTR NASA
MT VIEW CALIFORNIA
94040

DR J H FRAZER CHIEF B
INTERNAL BALLISTICS LAB
BALLISTIC RESEARCH LABS
ABERDEEN PROV GROUND MD
21005

I E GARRICK MS 242 AB
NASA LANGLEY RESEARCH CTR
HAMPTON VIRGINIA
23365

MR W T HAMILTON ABDGI
CHIEF SST TECHNOLOGY
SUPERSONIC TRANSPORT DIV
P O BOX 733
RENTON WASHINGTON
98055

DR M HEASLET
NASA
AMES RESEARCH CTR NASA
MT VIEW CALIFORNIA
94040

DR JOHN T HOWE
NASA AMES
MOFFETT FIELD CALIFORNIA
94035

ELMER G JOHNSON DIRECTOR
(ARF) BDG
FLUID DYNAMICS FACIL LAB
AEROSPACE RESEARCH LABS
WRIGHT PATTERSON AFB OHIO
45433

MR LUKASIEWICZ ABDEG
CHIEF
GAS DYNAMICS FACILITY
ARNOLD AIR FORCE STATION
TENNESSEE
37389

DR J MENKES ABCDEFGHI
INST FOR DEFENSE ANALYSES
400 ARMY NAVY DRIVE
ARLINGTON VIRGINIA
22204

NASA
DR D R CHAPMAN
AMES RESEARCH CTR NASA
MT VIEW CALIFORNIA
94040

NASA LANGLEY RESEARCH CTR
LANGLEY STATION
ATTN LIBRARY MS185
ABCDEFGHI
HAMPTON VIRGINIA
23365

NASA LEWIS RESEARCH CTR
ATTN LIBRARY MS 60 3
ABCDEFGI
21000 BROOKPARK ROAD
CLEVELAND OHIO
44135

NASA SCIENTIFIC & TECH
INFO FAC
ABCDEFGHI
ATTN ACQUISITIONS BR
(S AK/DL)
P O BOX 33
COLLEGE PARK MARYLAND

NATIONAL SCIENCE FOUND
ENGINEERING DIVISION
ABCDEI
1800 G STREET N W
WASHINGTON D C
20550

OFFICE OF NAVAL RESEARCH
NEW YORK AREA OFFICE
ABCDEFGHI
207 W 24TH STREET
NEW YORK NEW YORK
10011

OFFICE OF NAVAL RESEARCH
SAN FRANCISCO AREA
OFFICE ABCDEFGHI
1076 MISSION STREET
SAN FRANCISCO CALIFORNIA
94103

DR W A PAGE
NASA AMES
MOFFETT FIELD CALIFORNIA
94035

J L POTTER BDG
MANAGER, VKF AP
ARNOLD AIR FORCE STATION
TENNESSEE
37389

REDSTONE SCIENTIFIC
INFORMATION CENTER
ATTN CHIEF DOCUMENT SECT
ABCDEFGHI
ARMY MISSILE COMMAND
REDSTONE ARSENAL ALABAMA
35809

SCIENCE & TECHNOLOGY DIV
ABCDEFGHI
LIBRARY OF CONGRESS
WASHINGTON D C
20540

SHIPYARD TECHNIC LIBRARY
CODE 130L7 BLDG 746
ABCDEFGHI
SAN FRANCISCO BAY NAVAL
SHIPYARD
VALLEJO CALIFORNIA
94592

SUPERINTENDENT
NAVAL ACADEMY
ATTN LIBRARY ABCDEFGHI
ANNAPOLIS MD
21402

UNITED STATES ATOMIC
ENERGY COMMISSION
DIV OF TECHNICAL INFO
EXTENSION
P O BOX 62 BCE
OAK RIDGE TENNESSEE
37830

DR R E WILSON ABCDFGH
ASSOC TECH DIR
(AEROBALLISTICS)
NAVAL ORDNANCE LABORATORY
WHITE OAK
SILVER SPRING MD
20910

LIBRARIES

AMER INST OF AERONAUTICS
AND ASTRONAUTICS
1290 SIXTH AVENUE
NEW YORK NEW YORK
ATTN INTL AEROSPACE
ABSTRACTS LIBRARY
10019

AVCO CORPORATION B
RESEARCH LIBRARY
201 LOWELL STREET
WILMINGTON MASSACHUSETTS
01887

CONVAIR DIVISION OF
GENERAL DYNAMICS
P O BOX 12009
ATTN LIBRARY (128 00)
ABCDFGHI
SAN DIEGO CALIFORNIA
92112

DEPARTMENT LIBRARIAN B
DEPT OF AERONAUTICAL
ENGINEERING
UNIVERSITY OF WASHINGTON
SEATTLE WASHINGTON
98105

ENGINEERING LIBRARY
ABCDEFGHI
DEPT 218 BLDG 101
MC DONNEL AIRCRAFT CORP
P O BOX 516
ST LOUIS MISSOURI
63166

ENGINEERING SOCIETIES
LIBRARY ABCDEFGHI
345 EAST 47TH STREET
NEW YORK NEW YORK
10017

LIBRARY ABCDEFGHI
AEROJET GENERAL CORP
6352 N IRWINDALE AVENUE
AZUSA CALIFORNIA
91702

LIBRARY ABDFH
THE RAND CORPORATION
1700 MAIN STREET
SANTA MONICA CALIFORNIA
90401

LIBRARY ABDH
MITHRAS INC
701 CONCORD AVENUE
CAMBRIDGE MASSACHUSETTS
02138

LIBRARY B
JET PROPULSION LAB
4800 OAK GROVE DRIVE
PASADENA CALIFORNIA
91103

LIBRARY BF
MIDWEST RESEARCH
INSTITUTE
425 VOLKER BOULEVARD
KANSAS CITY MISSOURI
64110

LIBRARY &
INFORMATION SERVICES BE
GENERAL DYNAMICS CONVAIR
P O BOX 1128
SAN DIEGO CALIFORNIA
92112

LTV AEROSPACE CORPORATION
TECHNICAL LIBRARY
2 51131 B
P O BOX 5907
DALLAS TEXAS
75222

NATIONAL RESEARCH COUNCIL
ABCDEFGHI
AERONAUTICAL LIBRARY
ATTN MISS O M LEACH LIBR
MONTREAL ROAD
OTTAWA 7 CANADA

RADIO CORPORATION OF
AMERICA
ENGINEERING LIBRARY
127 223 B
ATTN MRS E K DALY LIBRAR
MOORESTOWN NEW JERSEY
08057

SUPERINTENDENT
NAVAL POSTGRADUATE SCHOOL
ATTN LIBRARY ABCDEFGHI
MONTEREY CALIFORNIA
93940

SUPERVISOR
TECH LIB SEC BDE
THIOKOL CHEMICAL CORP
WASATCH DIVISION
BRIGHAM CITY UTAH
84302

THE LIBRARY B
INSTITUTE FOR AEROSPACE
STUDIES
UNIVERSITY OF TORONTO
TORONTO 5 CANADA

UNITED AIRCRAFT CORP
RESEARCH LABORATORIES
ATTN LIBRARY B
SILVER LANE
EAST HARTFORD CONN
06103

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY *(Corporate author)* Brown University, Providence, R. I. 02912	2a. REPORT SECURITY CLASSIFICATION Unclassified 2b. GROUP

3. REPORT TITLE

ON RADIATIVE TRANSFER IN THE LOW REYNOLDS NUMBER BLUNT BODY STAGNATION REGION AT HYPERSONIC SPEEDS PART 1. EMISSION DOMINATED CASE

4. DESCRIPTIVE NOTES *(Type of report and inclusive dates)*

5. AUTHOR(S) *(First name, middle initial, last name)*

Liu, Joseph T. C.

6. REPORT DATE March 1968	7a. TOTAL NO. OF PAGES 35	7b. NO. OF REFS 12
8a. CONTRACT OR GRANT NO. Nonr 562(35) b. PROJECT NO. Task NR 061-132 c. d. ARPA Project Code Number 2740	9a. ORIGINATOR'S REPORT NUMBER(S) Nonr 562(35)/21 9b. OTHER REPORT NO(S) *(Any other numbers that may be assigned this report)*	

10. DISTRIBUTION STATEMENT

11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY Advanced Research Projects Agency and Office of Naval Research

13. ABSTRACT

One objective of this paper is to contribute to an understanding of radiative transfer effects in the low Reynolds number or merged layer regime of hypersonic flow about axisymmetric blunt bodies. The other objective is to illustrate how the concept of thin shock layer theory can be extended to the radiative case, where the shock structure and shock layer are radiatively coupled, through the introduction of a psuedo-jump condition across the shock structure, accompanied by an iteration technique. The radiative transfer is simplified to the emission dominated case. The gas is taken as calorically and thermally perfect, the viscosity taken as a linear function of the temperature, and the absorption coefficient of the gas is assumed to be gray. The problem is formulated for the flow about a blunt body but the detailed calculations are carried out for the stagnation region in order to illustrate the techniques used. Radiative cooling decreases the convective heat transfer from the corresponding radiationless case. The additional contributions to the heat transfer due to radiative transfer come from both the shock structure and the shock layer. The relative importance of these contributions are assessed in terms of the rarefaction parameter. Radiative contributions become less important toward the free-molecular range. Radiative cooling decreases the overall shock structure and shock layer thicknesses.

DD FORM 1 NOV 65 1473

Unclassified
Security Classification

Security Classification

14. KEY WORDS	LINK A ROLE	LINK A WT	LINK B ROLE	LINK B WT	LINK C ROLE	LINK C WT
Radiative Transfer						
Low Reynolds Number Hypersonic Flow						
Blunt-Body Flow						

Unclassified
Security Classification