


# PLASMA PHYSICS GROUP



DEPARTMENT OF PHYSICS  
UNIVERSITY OF CALIFORNIA  
LOS ANGELES 90024

FACILITY FORM 502

N66-15440	
(ACCESSION NUMBER)	(THRU)
7	1
(PAGES)	(CODE)
CL 69387	25
(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

GPO PRICE \$ \_\_\_\_\_

CFSTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) 1.00

Microfiche (MF) .50

ff 653 July 65

Propagation of Ion Acoustic Waves  
Along Cylindrical Plasma Columns

A. Y. WONG

July 1965

R - 1

Plasma Physics Group  
Department of Physics  
University of California  
Los Angeles, California

This research was partially supported by the  
National Aeronautics and Space Administration  
under grant NGR-05-007-066.

# Propagation of Ion Acoustic Waves Along Cylindrical Plasma Columns

A. Y. Wong

Department of Physics, University of California, Los Angeles

It has been found by Wong, Motley and D'Angelo<sup>1</sup> that the propagation of ion acoustic waves along an external axial magnetic field between 4kg and 14 kg in highly ionized plasmas is not affected by the size and shape of the cylindrical plasma column. The experimental results indeed agree with a plane wave analysis even though the plasma radius  $R$  was of the order of the wavelength  $\lambda$ . Neither changing the plasma diameter by a half nor altering the density profile through varying the axial magnetic field (4-14kg) seemed to have any effect on the phase velocity and damping. On the other hand, Little<sup>2</sup> has observed the effects of boundaries on ion acoustic waves in a weakly ionized plasma in the presence of a weak magnetic field (less than 50 gauss) with  $\lambda \approx R$ . Malmberg and Wharton<sup>3</sup> also found that in the propagation of electrostatic electron oscillations the size of the plasma column must be taken into account when  $kR \approx 1$ . These experimental results led us to re-examine the theory of ion acoustic waves to include transverse boundary effects.

Making the electrostatic approximation for ion acoustic waves  $\underline{E} = -\nabla\phi$  and  $\nabla \cdot \underline{D} = 0$ , we can derive a wave equation for the wave potential  $\phi$ :

$$\epsilon_{\perp} \nabla_{\perp}^2 \phi - \epsilon_{zz} k^2 \phi = 0 \quad (1)$$

where  $\epsilon_{\perp}$  and  $\epsilon_{zz}$  are elements of the dielectric tensor  $\underline{\epsilon}$

$$\underline{\epsilon} = \begin{vmatrix} \epsilon_{\perp} & \epsilon_2 & 0 \\ -\epsilon_2 & \epsilon_{\perp} & 0 \\ 0 & 0 & \epsilon_{zz} \end{vmatrix}$$

and  $k$  = the wave number in the direction of propagation.

From a theory taking the ions as collisionless and the electrons as a fluid (as appropriate to WMD's experiment<sup>1</sup>) we obtained

$$\epsilon_{zz} = 1 - \frac{k_D^2}{k^2} Z' \left( \frac{\omega}{k v_i} \right) + 2 \frac{T_i}{T_e} \frac{k_D^2}{k^2}$$

$$\epsilon_{\perp} \cong 1 + k_D^2 \rho_i^2 \quad \text{For } \omega < \omega_{ci} \ll \omega_{ce} \quad \text{and} \quad \frac{B^2}{8\pi K T} \gg 1$$

$$\text{where } k_D^2 = \frac{4\pi n e^2}{K T_i} \quad \rho_i = \text{ion cyclotron radius}$$

and  $\omega_{ci}$   $\omega_{ce}$  the ion and electron cyclotron frequency respectively.

Neglecting the electron inertia in the ion wave, it can be shown that

$$\frac{e\phi}{K T} = \frac{n}{n_0}$$

where  $n_0$  = the unperturbed plasma density

$n$  = the perturbed plasma density

Equation (1) can thus be written as

$$\nabla_{\perp}^2 n - k^2 \frac{\epsilon_{zz}}{\epsilon_{\perp}} n = 0 \quad (2)$$

We shall make use of the following experimental observation in our selection of the appropriate solution for equation (2):

$$1) \quad n_o(r) \approx J_o \left( \frac{\chi_1 r}{R} \right), \text{ where } \chi_1 = \text{1st root of the zeroth order Bessel function}$$

$R = \text{plasma radius.}$

$$2) \quad \frac{n}{n_o} \approx \text{constant along the radial direction.}$$

3) azimuthal symmetry.

The solution of equation (2) will then be  $n(r) \propto J_o \left( \sqrt{\frac{-k^2 \epsilon_{zz}}{\epsilon_{\perp}}} r \right)$

with the condition that  $k^2 \epsilon_{zz} = - \frac{\epsilon_{\perp} \chi_1^2}{R^2}$

This condition can be rewritten as

$$1 - \frac{k_D^2}{k^2} Z' \left( \frac{\omega}{k v_i} \right) + 2 \frac{T_i}{T_e} \frac{k_D^2}{k^2} + \frac{\epsilon_{\perp} \chi_1^2}{k^2 R^2} = 0$$

Under the experimental conditions of WMD

$$k \ll k_D \quad \text{and} \quad T_i = T_e$$

$$\text{we have } Z' \left( \frac{\omega}{k v_i} \right) \approx 2 + \frac{\epsilon_{\perp} \chi_1^2}{k_D^2 R^2}$$

The effects of transverse boundary conditions are important only if

$$\left| \frac{\epsilon_{\perp} \chi_1^2}{k_D^2 R^2} \right| \approx 2 \quad (3)$$

A similar condition is also arrived at when both ions and electrons are treated as fluids.

In WMD's experiment

$$T_e = T_i = 2500^\circ\text{K}$$

$$n \approx 10^{11} \text{ cm}^3/\text{sec}$$

$$R = 1.5 \text{ cm},$$

$$k_D \approx 6 \times 10^2 \text{ cm}^{-1}$$

$$4 \times 10^{-2} \text{ cm} < \rho_i < 2 \times 10^{-1} \text{ cm} \quad \text{for } 14 \text{ kg} > B > 4 \text{ kg}$$

we have  $\epsilon_{\perp} \approx k_D^2 \rho_i^2$  since  $k_D \rho_i \gg 1$

Condition (3) reduces to  $\frac{\rho_i^2}{R^2} = \frac{2}{X_1^2}$  (4)

Thus we come to the conclusion that the transverse boundary conditions are important if the ion cyclotron radius (rather than the wavelength of the ion wave) is comparable to the plasma radius. This agrees with the experimental finding of WMD: the independence of propagation characteristics on the size of the plasma column in the presence of a magnetic field such that  $\rho_i \ll R$ . Examination of condition (3) shows that the transverse boundary effects on ion acoustic waves can be vitiated by the application of a sufficiently high magnetic field which reduces the boundary contribution with respect to the electron contribution in ion acoustic waves. In contrast, the longitudinal electron plasma oscillations are sensitively dependent upon the boundary conditions since the corresponding ion contribution to the electron oscillations is zero. The importance of such dependence has been shown experimentally<sup>3</sup> and theoretically.<sup>4,5</sup> In summary, the independence of ion acoustic waves on

transverse boundary conditions is attainable under typical laboratory conditions and further simplifies its laboratory investigation and increases its role in plasma diagnosis.<sup>6</sup>

#### Acknowledgements

It is a pleasure to acknowledge useful discussions with Dr. D. Sachs, Dr. A. Kaufman and Dr. B. Fried. This work was supported by the National Aeronautics and Space Administration under grant NGR 05-007-066.

References

1. A. Y. Wong, R. W. Motley and N. D'Angelo, Phys. Rev. 133, A436 (1964).
2. P. F. Little, Nature 194, 1137 (1962).
3. J. H. Malmberg and C. B. Wharton, Phys. Rev. Letters 13, 184.
4. R. W. Gould, "Propagation of Electrostatic Waves Along a Plasma Column,"  
Culham Laboratory Report (1964).
5. Knehl, Stewart and Yeh, Phys. of Fluids (to be published).
6. I. Alexeff and R. Neidigh, Phys. Rev. 129, 516 (1963).  
A. Y. Wong, Appl. Phys. Letters 6, 147, 1965.