

JAN 14 1966
UCSD-SP-65-5
December 21, 1965

FACILITY FORM 802

N66 24683
(ACCESSION NUMBER)
12
(PAGES)
CR-74808
(NASA CR OR TMX OR AD NUMBER)

(THRU)
29
(CODE)
(CATEGORY)

The Spectrum of Crab Nebula X-Rays to 120 Kev

By

Laurence E. Peterson, Allan S. Jacobson, and R. M. Pelling

Physics Department

University of California at San Diego

La Jolla, California

Hard copy (HC) 1.00

Microfiche (MF) 1.50

653 July 65

In this letter, we wish to report a balloon determination of the Crab Nebula X-ray spectrum over the range 16 to 120 kev. X-rays from the Crab were first discovered from a rocket⁽¹⁾, and were observed from a balloon to 60 kev by Clark⁽²⁾. In this work the spectrum has been extended and determined to a much higher precision.

The detector⁽³⁾ consists of a NaI crystal 5 mm thick by 24 mm diameter. Collimation is provided by a cylindrical CsI scintillation cup 9.3 cm O.D. by 22.9 cm long with a 2 cm wall which surrounds the central detector and its phototube. This collimator, which is in anti-coincidence with the central detector also rejects background events produced by cosmic-rays. The detector area is 9.4 cm^2 , its geometry factor is 1.5 cm^2 -steradian, and its resolution, FWHM, 54%, at 30 kev. Post flight and pre-flight energy calibrations were identical. Inflight calibrations during a previous flight with the same apparatus showed no tendency for gain changes. Data was telemetered event by event in a digital format from a 128 channel pulse-height analyzer. The detector was mounted at a fixed

GPO PRICE \$

CFSTI PRICE(S) \$

elevation of 80° , and servo-controlled in azimuth to point true south. The reference was obtained from a magnetometer set to compensate for the average magnetic declination over Texas. The azimuth was independently verified to within about a degree by two additional crossed magnetometers.

The balloon was launched from Palestine, Texas at 2234 CST, September 23, 1965, reached ceiling at 0050 CST, and floated level at 3.3 gm/cm^2 until 0610 CST, when the instrument was inadvertently released by radio command. The Crab transit on this date, corrected for the drift in longitudes of the balloon, was 0545 CST.

Figure 1 shows the counting rates as a function of time for various energy ranges, averaged over 12 minute intervals. The peak on ascent is due to the cosmic ray maximum. The rate increase at lower energies as the balloon approaches ceiling is apparently due to a diffuse component of cosmic photons. This flux, when corrected to zero depth seems in general agreement with the results reported previously.^(4,5) This component contributes about half of the measured background in the 20 - 50 kev range. About one-half is due to cosmic ray production in the overlying atmosphere and only about 5 or 10% is true detector background.

The increase near the end of the flight is attributed to the meridian passage of an X-ray source. The dashed line in Figure 1(a) shows the expected time dependent response for a point object of the observed peak intensity at the position of the Crab. The observed source must be within about $\pm 2^{\circ}$ in R.A. and $\pm 4^{\circ}$ in declination of the Crab Nebula.

Figure 2 shows the pulse height distribution of the background, and of the Crab during its transit. The incident flux is obtained by

subtracting these spectra and correcting for atmospheric absorption, efficiency, deadtime, channel width, and geometry factor. The spectrum obtained by carrying out this procedure is shown in Figure 3, averaged over suitable channel groupings. No statistically significant fluxes above background were found above 120 kev. Atmospheric absorption removes X-rays below 16 kev. A least square fit to the spectrum gives a power law exponent of $1.91 \pm .10$. Therefore α , the index of the differential energy flux is $0.91 \pm .10$ over this range.

Clark⁽²⁾ quotes a spectral index $\alpha = 2$ with an upper limit at 60 kev, as shown in Figure 3. Our flux seems significantly above his upper limit. His observations were made through a considerably higher background and with less energy resolution. Inspection of his results reveals that an α of one is also consistent with his data, if the upper limit at 60 kev is excluded. There may possibly be secular variations in the Crab Nebula. Also shown in Figure 3 are the rocket observations reported by the NRL group⁽⁶⁾ and the upper limits obtained during a search for solar γ -rays.⁽⁷⁾

Various mechanisms for the production of X-rays in astronomical sources have been described by Gould and Burbidge⁽⁸⁾. Thermal Bremsstrahlung from a hot gas produces an exponential spectrum. A model temperature and density distribution to reproduce the power law observed here requires a temperature on the order of 10^9 °K. Bremsstrahlung from non-thermal electrons gives a $1/E$ number spectrum. A distribution of slow electrons which will give the observed photon spectrum seems somewhat artificial, and would be quickly cooled by collision loss. The synchrotron

mechanism, with an appropriate time dependent source of 10^4 Bev electrons in a 10^{-4} gauss field, can produce the observed photon spectrum. (9)

Clayton and Craddock⁽¹⁰⁾ have estimated γ -ray line emission from the Crab Nebula on the basis of heavy element production by the r-process in supernova explosions. They predict a line at 60 kev due to Am^{241} decays with an intensity about 5.7×10^{-5} photons cm^2 -sec. Our upper limit on this line is about 1.1×10^{-2} photons cm^2 -sec.

Acknowledgements:


The authors are indebted to R. J. Gould, W. H. Tucker and G. R. Burbidge for helpful discussion. The apparatus was constructed by Rod Jerde, who also assisted with the flight details. Balloon flight services were provided by the National Center for Atmospheric Research. This research was supported under Grant NsG 318 from the National Aeronautics and Space Administration.

References:

1. R. Giacconi, H. Gursky, F. R. Paolini, B. R. Rossi, Phys. Rev. Letters 9, 439 (1962).
2. G. W. Clark, Phys. Rev. Letters 14, 91 (1965)
3. D. B. Hicks, L. Ried, Jr., L. E. Peterson, Transactions 11th Nuclear Science Symposium, IEEE (1965)
4. R. Rothenflug, R. Rocchia, L. Koch, presented at 9th Int. Conf. on Cosmic Rays (1965)
5. A. E. Metzger, E. C. Anderson, M. A. Van Dilla, J. R. Arnold, Nature 204, 766 (1964)
6. S. Bowyer, E. T. Byram, T. A. Chub, H. Friedmann, Nature 201, 1307 (1964)
7. K. J. Frost, E. D. Rothe, L. E. Peterson, J. Geophys. Res. (to be published)
8. R. J. Gould, G. R. Burbidge, Annales d'Astrophysique 28 171 (1965); Handbuch der Physik 46,2 (to be published)
9. W. H. Tucker (Private Communication)
10. D. D. Clayton, W. L. Craddock, Ap. J. 142, 189 (1965)


Figure Captions:

1. The counting rates of various energy ranges as a function of time.
Background data was obtained at ceiling before the meridian transit of the X-ray source. The dashed line shows the expected transit profile for a point source at the position of the Crab Nebula.
2. The pulse height distributions obtained during the background interval, and during the transit of the Crab. No significant difference is apparent above 120 kev.
3. Observations of the spectrum of the Crab Nebula in the X-ray range.
The previous measurements of Clark were made with a detector of higher background and may not be inconsistent with this determination.



COUNTING RATE SPECTRA CRAB NEBULA OBSERVATION

23 SEPT. 1965



17000 2-10-62

