

SPACE RESEARCH COORDINATION CENTER



EXTENSIONS OF THE MOMENTUM  
TRANSFER THEOREM

BY

E. GERJUOY

DEPARTMENT OF PHYSICS

SRCC REPORT NO. 26

UNIVERSITY OF PITTSBURGH  
PITTSBURGH, PENNSYLVANIA

28 JANUARY 1966


GPO PRICE \$ \_\_\_\_\_

CFSTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) 1.00

Microfiche (MF) .50

# 653 July 65



33469

FACILITY FORM 602	N66 32254	
	(ACCESSION NUMBER)	(THRU)
	15	(CODE)
	CR-76651	(CATEGORY)
	(NASA CR OR TMX OR AD NUMBER)	

## Extensions of the Momentum Transfer Theorem\*

E. Gerjuoy

University of Pittsburgh

Recently Lippmann,<sup>1</sup> in this Journal, has discussed the extension of the momentum transfer theorem<sup>2,3</sup> to systems more complicated than the (elastic or inelastic) collisions of electrons with atomic hydrogen. Lippmann<sup>1</sup> also has discussed extensions of the theorem to other observables, so as to derive, e.g., an energy transfer theorem. In his discussion, Lippmann took exception to some remarks concerning the validity of the symbolic methods customarily employed in scattering theory. These remarks, from a preprint version of the paper which proved the momentum transfer theorem for e-H collisions, were accurately quoted by Lippmann, but do not appear in the actually published paper,<sup>3</sup> because I already had decided the remarks were not wholly defensible.<sup>4</sup> Nevertheless there remain some differences between Lippmann's and my views of the status of the momentum transfer theorem and its extensions. Making these differences explicit is the primary objective of this Letter.

\*Supported by the Office of Naval Research

In Lippmann's derivation of the momentum transfer theorem, the starting point is

$$(\Psi^{(+)}, [p_{1z} H - H p_{1z}] \Psi^{(+)}) \quad (1a)$$

the "expectation value" of the commutator between the Hamiltonian  $H$  and  $p_{1z}$ , the momentum operator (along its incident direction) of the incident particle. Lippmann relates (1a) to the momentum transfer cross section via symbolic methods. My starting point has been much the same as Lippmann's, namely the identity

$$(\Psi^{(+)}, p_{1z} H \Psi^{(+)}) - (H \Psi^{(+)}, p_{1z} \Psi^{(+)}) = 0 \quad (1b)$$

On the other hand, I have chosen to evaluate the integrals on the left side of (1b) in a particular representation, the coordinate representation. In this representation, the terms in (1b) independent of the potential  $V$  are related, via Green's Theorem, to a surface integral at infinity, which then can be evaluated from the known asymptotic behavior of  $\Psi^{(+)}$  at large interparticle distances.

For the case of potential scattering, the aforementioned surface integral reduces immediately<sup>2</sup> to the physical momentum transfer cross section, thus yielding the momentum transfer cross section theorem. The situation is less simple in e-H collisions, however, wherein  $\Psi^{(+)}$  must be symmetric (singlet scattering) or antisymmetric (triplet scattering) under interchange of  $r_1$  and  $r_2$ , the coordinates respectively of the indistinguishable "initially incident" and "initially bound" electrons. In this event the surface integral (now over the five-dimensional boundary of the sphere at infinity in the six-dimensional space of  $(r_1, r_2)$ ) reduces to the momentum transfer cross section plus terms proportional to

$\int d\mathbf{r} \phi_j^*(\mathbf{r}_1) p_{1z} \phi_j(\mathbf{r}_1)$  integrated over all  $\mathbf{r}_1$ , where  $\phi_j$  is the eigenfunction of atomic hydrogen in its  $j$ th bound state. Such terms, discussed in connection with Eqs. (G-42) - (G-44b), apparently are absent from<sup>1</sup> Eq. (L-6). But, because  $p_z$  has odd parity, these terms proportional to  $(\phi_j, p_z \phi_j)$  vanish. At least superficially, therefore, Lippmann's version of the momentum transfer theorem for e-H collisions agrees with mine.

Next, let  $p_1^2$  replace  $p_{1z}$  in (1a) and (1b). Then proceeding from (1b) just as in Eqs. (G-36) - (G-49), one finds for singlet or triplet e-H collisions that the energy transfer cross section  $\sigma_E$  is given by

$$\begin{aligned} \sigma_E = & \frac{1}{2ik_0} \frac{2m}{\hbar^2} \int d\mathbf{r} \psi^{(+)*} [\psi^{(+)} \nabla_1^2 V + 2\nabla_1 V \cdot \nabla_1 \psi^{(+)}] \\ & - \frac{\sigma}{\hbar^2 k_0^2} \int d\mathbf{r}_1 \phi_0^*(\mathbf{r}_1) p_1^2 \phi_0(\mathbf{r}_1) \\ & + \sum_j \frac{\sigma_j}{\hbar^2 k_0^2} \int d\mathbf{r}_1 \phi_j^*(\mathbf{r}_1) p_1^2 \phi_j(\mathbf{r}_1), \end{aligned} \quad (2)$$

where the definition of  $\sigma_E$  is

$$\sigma_E = \frac{1}{k_0^2} \left[ \sum_j (k_0^2 - k_j^2) \sigma_j + (k_0^2 - k_1'^2 - k_2'^2) \sigma_{ion} \right] \quad (3)$$

In Eqs. (2) - (3),  $k_0$  is the wave number of the incident electron;  $k_j$  is the wave number of the outgoing electron after a collision leaving the atom in its  $j$ th bound state;  $k_1'$ ,  $k_2'$  are the wave vectors of the outgoing electrons when ionization occurs;  $\sigma$  is the total cross section, including ionization;  $\sigma_j$  is the cross section (including direct and exchange processes) for collisions producing outgoing electrons with wave number  $k_j$ ;  $\sigma_{ion}$  is the cross section for ionization, integrated over all allowed values of  $k_1'$ ,  $k_2'$ ; the sums over  $j$  include elastic scattering,  $j = 0$ . Of course,

$$k_0^2 = k_j^2 = -\frac{2m}{\hbar^2}(\epsilon_0 - \epsilon_j) \quad (4)$$

$$k_0^2 - k_1'^2 - k_2'^2 = -\frac{2m}{\hbar^2} \epsilon_0$$

where  $\epsilon_j$  is the (negative) energy of the  $j$ th bound state.

Eq. (L-7) apparently lacks the last two terms in Eq. (2) above. These terms, which correspond to the terms proportional to  $(\phi_j, p_z \phi_j)$  in the momentum transfer theorem, now do not vanish because  $p_1^2$  has even parity. Thus, in the case of e-H collisions, Lippmann's result for the energy transfer theorem disagrees (superficially, at least) with the result of a detailed calculation in the coordinate representation.

This apparent disagreement between Lippmann's and my version of the energy transfer theorem persists even when the particles 1 and 2 are considered distinguishable, i.e., when  $\psi^{(+)}$  is not symmetrized. To be specific, in this situation

$$\begin{aligned} \sigma_E = & \frac{1}{2ik_0^3} \frac{2m}{\hbar^2} \int d\mathbf{r} \psi^{(+)*} [\psi^{(+)} \nabla_1^2 V + 2\nabla_1 V \cdot \nabla_1 \psi^{(+)}] \\ & - \frac{1}{\hbar^2 k_0^2} \sum_j \sigma_j^{\text{exch}} [\hbar^2 k_0^2 - \int d\mathbf{r}_1 \phi_j^*(\mathbf{r}_1) p_1^2 \phi_j(\mathbf{r}_1)] \end{aligned} \quad (5)$$

where  $\sigma_j^{\text{exch}}$  is the exchange cross section for production of free particles 2, leaving the initially free incident particle 1 in the  $j$ th bound state. In Eq. (5), because the particles now are distinguishable,  $\sigma_E$  is defined not by Eq. (3) but rather by

$$\sigma_E = \frac{1}{k_0^2} \sum_j [(k_0^2 - k_j^2)(\sigma_j - \sigma_j^{\text{exch}}) + \int d\mathbf{k}_1' d\mathbf{k}_2' (k_0^2 - k_1'^2) \sigma_{\text{ion}}(\mathbf{k}_1', \mathbf{k}_2')] \quad (6)$$

where the total ionization cross section  $\sigma_{\text{ion}}$  satisfies

$$\sigma_{ion} = \int dk_1^i dk_2^i \sigma_{ion}(k_1^i, k_2^i)$$

Evidently Eq. (6) supposes the kinetic energies of outgoing free particles 1 only--not of particles 2-- will be measured and compared with the initially incident kinetic energy. Eq. (6) is not the only physically sensible possible definition of  $\sigma_E$  in (5), but no definition of  $\sigma_E$  will eliminate the expectation values  $(\phi_j, p_j^2 \phi_j)$  in the energy transfer theorem unless such expectation values explicitly are incorporated into the definition of  $\sigma_E$ . For actual e-H collisions, involving indistinguishable particles, Eq. (3) provides the only physically sensible definition of  $\sigma_E$ .

The presence of the expectation values  $(\phi_j, p_j^2 \phi_j)$  in Eq. (5) is understandable. Whether or not the particles are indistinguishable, i.e., whether or not  $\Psi^{(+)}$  is symmetrized, the surface integral arising from Eq. (1b) (with  $p_1^2$  replacing  $p_{1z}$ ) represents the net flux of probability current--weighted by  $p_1^2$ --across the sphere at infinity in  $r_1, r_2$  space; the presence of forces, contained in the first term on the right side of (5) or (2), causes this net weighted flux to differ from zero. All collision processes, including those which convert 1 from a free to a bound particle, are included in the net probability current flux; but any physically sensible definition of  $\sigma_E$ , e.g., Eqs. (6) or (3), corresponding to actually feasible measurements, should involve the kinetic energy fluxes of free (unbound) particles only. Consequently, only in the circumstances that particle 1 is always free, or that  $p_1^2$  is expected to vanish whenever particle 1 is not free, does one expect  $\sigma_E$  of Eq. (6) to equal exactly the force terms involving  $V$  on the right side of (5). In fact, the extra terms in (5), proportional to  $\sigma_j^{exch}$ , have precisely the form one expects (in terms of the cross sections) for the rate at which the forces are causing a flow of  $p_1^2$  from unbound to bound states of 1.

For actual e-H collisions, where the particles are indistinguishable, the precise form of the extra terms involving  $(\phi_j, p^2 \phi_j)$  is less readily interpreted physically, but it is clear that the genesis of these extra terms in (2) is essentially the same as in (5). The preceding paragraph also clarifies the fact that the vanishing expectation values  $(\phi_j, p_z \phi_j)$  appear in the derivation of the momentum transfer theorem, and suggests that extra terms involving the expectation values  $(\phi_j, A \phi_j)$  will have to be included in the transfer theorem for any even parity operator A, e.g., the angular momentum transfer theorem, Eq. (L-8). However, I have not examined the angular momentum transfer theorem, or the transfer theorem for any other even parity operator A, in the detail that I have examined the energy transfer theorem.

It is to be noted that the presence of extra terms involving  $(\phi_j, A \phi_j)$  implies the transfer theorem for A--unlike the momentum transfer theorem--has little chance of being generally useful. For instance, granting exact knowledge of V, prediction of  $\sigma_E$  from (2) or (5) requires accurate knowledge of  $\sigma_j$  and the associated expectation values  $(\phi_j, p^2 \phi_j)$ . Hence use of (2) or (5) to estimate  $\sigma_E$  generally will be no easier or more accurate than direct employment of the corresponding defining equations (3) or (6). For this reason the energy transfer theorem and similar obvious extensions of the momentum transfer theorem were not included in my e-H collisions paper.<sup>3</sup>

On the other hand, it is possible to eliminate the extra terms involving  $(\phi_j, A \phi_j)$  in special cases. One important such case is the energy transfer theorem for Coulomb interactions, i.e. just the case for which (2) was derived. In this case we know from the virial theorem that

$$\frac{1}{2m} \int d\mathbf{r} \phi_j^*(\mathbf{r}) p_1^2 \phi_j(\mathbf{r}) = -\epsilon_j \quad (7)$$

Recalling Eqs. (3) - (4), using (7) converts Eq. (2) to

$$2\sigma_E = \frac{1}{2ik_0^3} \frac{2m}{\hbar^2} \int d\mathbf{r} \psi^{(+)*} [\psi^{(+)} \nabla_1^2 V + 2\nabla_1 V \cdot \nabla_1 \psi^{(+)}] \quad (8)$$

Thus in the special case of e-H collisions there is a useful energy transfer theorem, but (superficially at least) it differs by exactly a factor 2 from Lippmann's version. The result (8) suggests the energy transfer theorem remains useful--though differing by a numerical factor from Lippmann's version--in the collisions of many-particle systems interacting via Coulomb forces, e.g., to atom-atom collisions. The same comment should hold for any collisions wherein the virial theorem is applicable, e.g., to the collisions of many-particle systems interacting via homogeneous potentials of any degree  $n$  (if any case other than the Coulomb  $n = -1$  actually exists).

Admittedly the coordinate representation proofs--of the momentum transfer theorem published previously,<sup>3</sup> and of the energy transfer theorem outlined here--become awkward and inelegant when extended to collisions more complicated than e-H. By finding the route to short elegant proofs for arbitrarily complicated colliding systems, Lippmann has made an important contribution therefore. This Letter has indicated, however, that the symbolic methods he employs must be made more precise before the extensions of the momentum transfer theorem to arbitrarily complicated colliding systems, and to other observables, can be regarded as more than "plausible". In particular (concentrating now on the energy transfer theorem), for many-particle systems including both distinguishable and indistinguishable particles, it is at least necessary to establish: (a) the connection



between the right side of Eq. (L-7) and the physically sensible  $\sigma_E$ ;

(b) the presence of the extra terms involving  $(\phi_j, p^2 \phi_j)$ , which are not obviously explicitly manifested (though very likely contained) in Eqs. (L-7) and (L-9).

References

1. B. A. Lippmann, Phys. Rev. Letters 15, 11 (1965). Equation numbers referring to this paper are preceded by L.
2. E. Gerjuoy, J. Math. Phys. 6, 993 (1965).
3. E. Gerjuoy, J. Math. Phys. 6, 1396 (1965). Equation numbers referring to this paper are preceded by G.
4. Regrettably, Lippmann did not check with me before publishing his Letter.