

270 X-641-67-303 END

291A NASA-TM-X-55818

# 3 ION ACOUSTIC WAVES IN A COLLISIONAL PLASMA 6

BY

6 B. BUTI 9

N67-29994

(ACCESSION NUMBER)

1029K52-111

(PAGES)

TMX-55818

(NASA CR OR TMX OR AD NUMBER)

(THRU)

1

(CODE)

25

(CATEGORY)

GPO PRICE \$ \_\_\_\_\_

CFSTI PRICE(S) \$ \_\_\_\_\_

Hard copy (HC) 3.00

Microfiche (MF) .65

9 JUNE 1967 10

ff 653 July 65



1. NASA

GODDARD SPACE FLIGHT CENTER

GREENBELT, MARYLAND 3

ION ACOUSTIC WAVES IN A COLLISIONAL PLASMA

by

B. Buti\*

National Aeronautics and Space Administration

Laboratory for Theoretical Studies

June 1967

GODDARD SPACE FLIGHT CENTER

Greenbelt, Maryland

---

\*National Academy of Sciences - National Research Council Resident Research Associate

# ION ACOUSTIC WAVES IN A COLLISIONAL PLASMA

by

B. Buti\*

National Aeronautics and Space Administration,

Laboratory for Theoretical Studies,

Goddard Space Flight Center,

Greenbelt, Maryland

---

\*National Academy of Sciences - National Research Council Resident Research Associate

## ABSTRACT

The damping of the ion-acoustic waves by Coulomb collisions is studied by using the Fokker-Planck equation of Rosenbluth et al. for both the species constituting the plasma, namely the electrons and the ions. In a plasma with weak collisions and in the absence of any external field, one finds that irrespective of the ratio of the ion temperature to the electron temperature,  $T$ , the characteristic frequency gets affected only by the electron-ion collisions. However, as far as the collisional damping is concerned, both the ion-ion and the electron-ion collisions play a somewhat equally important role; the electron-electron and the ion-electron contributions are negligible compared to the other two. The damping increases with increase in the collision frequency but decreases with increase in  $T$ .

## I. INTRODUCTION

Ion acoustic waves, which were first predicted by Tonks and Langmuir<sup>1</sup> using the fluid analysis, have been studied by a number of authors<sup>2-5</sup> on the basis of collisionless Boltzmann equation. The collisionless theory shows that when the electrons and the ions have the same temperatures, these waves are very heavily Landau-damped. Experimentally, the ion waves were first observed by Revans<sup>6</sup> and their Landau-damping was measured by Wong et al.<sup>7</sup> Wong et al. studied the space damping rather than the time damping in Cesium and Potassium and observed that the damping constant depends on the magnitude of the ion drift.

The collisional damping of the ion waves was studied by Bhadra and Varma;<sup>8</sup> they used Krook's model<sup>9</sup> to describe the ion-ion collisions. Kulsrud and Shen<sup>10</sup> used a slightly more realistic model for the collisional process; in their model, the ions were described by the Fokker-Planck equation of Rosenbluth et al.,<sup>11</sup> however, the electrons were treated by a fluid equation with the further assumption that the electrons were isothermal. This assumption of electrons being isothermal is equivalent to considering the Vlasov equation for the electrons, thereby neglecting the electron-electron and the electron-ion collisions. We remove this restriction on the electrons and treat both the

electrons and the ions by the corresponding Fokker-Planck equations. We indeed find that the electron-ion contribution to the collision term can not be neglected as compared to the ion-ion contribution.

Similar calculations for high-frequency plasma waves have been done by Comisar<sup>12</sup> and Buti,<sup>13-15</sup> The effect of strong collisions on the low-frequency electrostatic plasma oscillations with and without an external magnetic field has been studied by Kuckes<sup>16</sup> by using fluid equations. Following Comisar and Buti, we solve the Fokker-Planck equation on the assumption that the Coulomb collisions are weak i.e.,  $\omega\tau \gg 1$ , where  $\omega$  is the characteristic frequency of the wave and  $\tau$  is the mean collision time for the collision process under consideration. We find that in the absence of any external field, irrespective of the ratio of the ion temperature to the electron temperature, the characteristic frequency gets affected only by the electron-ion collisions; however, as far as the collisional damping is concerned, both the electron-ion and the ion-ion collisions play a role: the electron-electron and the ion-electron contributions are negligible compared to the other two.

## II. GENERAL THEORY

Consider an unbounded fully ionized hot plasma consisting only of electron and ions without any external field. In equilibrium both the electrons and the

ions obey Maxwellian distribution of velocities i.e.,

$$f_{oa}(v) = (2\pi v_a^2)^{-3/2} e^{-v^2/(2v_a^2)}, \quad (1)$$

where the subscript  $a$  stands either for the electron or for the ion and  $V_a^2 = KT_a/m_a$  with  $m_e = m$  and  $m_i = M$ .  $f_{oa}$  is the equilibrium distribution function for species  $a$ . For small perturbations, the linearized Fokker-Planck equation for longitudinal oscillations is given by

$$\frac{\partial f_a}{\partial t} + \mathbf{v} \cdot \frac{\partial f}{\partial \mathbf{x}} + \frac{Ne_a}{m_a} \mathbf{E} \cdot \frac{\partial f_{oa}}{\partial \mathbf{v}} = \left( \frac{\partial f_a}{\partial t} \right)_c, \quad (2)$$

where  $f_a(\mathbf{x}, \mathbf{v}, t)$  is the perturbed distribution function and  $(\partial f_a / \partial t)_c$  takes into account the collisions between like and unlike particles and is represented by

$$\begin{aligned} \left( \frac{\partial f_a}{\partial t} \right)_c = & - \frac{\partial}{\partial \mathbf{v}} \cdot \left[ f_{oa} \langle \Delta \rangle_a + f_a \langle \Delta \rangle_{oa} \right] \\ & + \frac{1}{2} \frac{\partial^2}{\partial \mathbf{v} \partial \mathbf{v}} \cdot \left[ f_{oa} \langle \Delta \Delta \rangle_a + f_a \langle \Delta \Delta \rangle_{oa} \right] \end{aligned} \quad (3)$$

with

$$\langle \Delta \rangle_a = N\Gamma_a \sum_{J=e,i} \left( 1 + \frac{m_a}{m_J} \right) \frac{\partial}{\partial \mathbf{v}} \int d\mathbf{v}' \frac{f_J(\mathbf{v}')}{|\mathbf{v} - \mathbf{v}'|}, \quad (4)$$

$$\langle \Delta \rangle_{oa} = N\Gamma_a \sum_{J=e,i} \left( 1 + \frac{m_a}{m_J} \right) \frac{\partial}{\partial \mathbf{v}} \int d\mathbf{v}' \frac{f_{oJ}(\mathbf{v}')}{|\mathbf{v} - \mathbf{v}'|}, \quad (5)$$

$$\langle \Delta \Delta \rangle_a = N \Gamma_a \sum_{J=e,i} \frac{\partial^2}{\partial \mathbf{v} \partial \mathbf{v}'} \int d\mathbf{v}' f_J(\mathbf{v}') |\mathbf{v} - \mathbf{v}'| \quad (6)$$

and

$$\langle \Delta \Delta \rangle_{oa} = N \Gamma_a \sum_{J=e,i} \frac{\partial^2}{\partial \mathbf{v} \partial \mathbf{v}'} \int d\mathbf{v}' f_{oJ}(\mathbf{v}') |\mathbf{v} - \mathbf{v}'| \quad (7)$$

In these equations  $\Gamma_a = (4\pi e_a^2 / m_a^2) \ln(4\pi N \lambda_D^3)$  with  $\lambda_D^2 = KT_{av} / (4\pi Ne^2)$ ;  $T_{av}$  is the average of the electron and the ion temperatures. The perturbed electric field in Eq. (2) is given by the Poisson's equation

$$\text{div } \mathbf{E} = 4\pi \sum_J e_J \int d\mathbf{v} f_J \quad (8)$$

To solve the pair of coupled equations (2) and (8), we take the Fourier transforms in space and Laplace transforms in time of all the perturbed quantities; Eq. (2) then becomes

$$(s + i \mathbf{k} \cdot \mathbf{v}) f'_a(\mathbf{k}, \mathbf{v}, s) - g_a(\mathbf{k}, \mathbf{v}) + \frac{Ne_a}{m_a} \mathbf{E}'_k \cdot \frac{\partial f_{oa}}{\partial \mathbf{v}} = \left( \frac{\partial f'_a}{\partial t} \right)_c \quad (9)$$

where  $f'_a$  and  $\mathbf{E}'_k$  are the Fourier-Laplace transforms of  $f_a$  and  $\mathbf{E}$  and  $g_a(\mathbf{k}, \mathbf{v})$  is the Fourier transform of the initial perturbation in the distribution function. To ensure the convergence of the integrals, we can have  $s$  in Eq. (9) such that  $\text{Re } s > 0$ . On further taking the Fourier transforms in velocity space, thus

defining

$$F_a(\mathbf{k}, \sigma, s) = \int d\mathbf{v} e^{-i\sigma \cdot \mathbf{v}} f_a'(\mathbf{k}, \mathbf{v}, s) , \quad (10)$$

Eq. (9) takes the form

$$\left( \frac{s}{k} - \frac{\partial}{\partial \sigma_z} \right) F_a - \frac{1}{k} G_a(\mathbf{k}, \sigma) + \frac{i e_a N \sigma_z E_z'}{m_a k} e^{-\sigma^2 v_a^2 / 2} = \frac{1}{k} \left( \frac{\partial F_a}{\partial t} \right)_c , \quad (11)$$

where

$$E_z' = - \frac{4i\pi}{k} \sum_J e_J F_J(\mathbf{k}, \mathbf{o}, s) \quad (12)$$

In writing Eqs. (11) and (12), we have assumed that  $\mathbf{k}$  is along the  $z$ -axis. Following the procedure outlined in references 12-14, we can easily evaluate the collision term and on integrating Eq. (11), we get

$$\begin{aligned} F_a(\mathbf{k}, \sigma, s) = & e^{s\sigma_z/k} \left[ Q_a(\sigma) + \frac{i e_a N E_z'}{m_a k} P_a(\sigma) \right] \\ & + \frac{v_a^4}{k L_a} e^{s\sigma_z/k} \int_{\sigma_z}^{\infty} d\sigma_z' e^{-s\sigma_z'/k} \int d\eta \left[ F_a(\eta) (K_1^b + K_2^a + K_3^a) \right. \\ & \left. + F_b(\eta) K_4^a \right] ; \quad a \neq b , \quad (13) \end{aligned}$$

where  $\sigma' = (\sigma_x, \sigma_y, \sigma_z')$ ,

$$Q_a(\sigma) = \frac{1}{k} \int_{\sigma_z}^{\infty} d\sigma_z' G_a(\mathbf{k}, \sigma') e^{-s\sigma_z'/k}, \quad (14)$$

$$P_a(\sigma) = - \int_{\sigma_z}^{\infty} d\sigma_z' \sigma_z' \exp \left[ - \frac{s\sigma_z'}{k} - \frac{v_a^2 \sigma'^2}{2} \right], \quad (15)$$

$$K_1^b(\sigma, \eta) = \frac{1}{2\pi^2} \left[ \left( 1 + \frac{m_a}{m_b} \right) \frac{\sigma \cdot (\sigma - \eta)}{(\sigma - \eta)^2} - \frac{\{\sigma \cdot (\sigma - \eta)\}^2}{(\sigma - \eta)^4} \right] \exp \left[ - \frac{v_b^2}{2} (\sigma - \eta)^2 \right], \quad (16)$$

$$K_2^a(\sigma, \eta) = \frac{1}{\pi^2} \left[ \frac{\sigma \cdot (\sigma - \eta)}{(\sigma - \eta)^2} - \frac{\{\sigma \cdot (\sigma - \eta)\}^2}{2(\sigma - \eta)^4} \right] \exp \left[ - \frac{v_a^2}{2} (\sigma - \eta)^2 \right], \quad (17)$$

$$K_3^a(\sigma, \eta) = \frac{1}{\pi^2} \left[ \frac{(\sigma \cdot \eta)}{\eta^2} - \frac{(\sigma \cdot \eta)^2}{2\eta^4} \right] \exp \left[ - \frac{v_a^2}{2} (\sigma - \eta)^2 \right] \quad (18)$$

and

$$K_4^a(\sigma, \eta) = \frac{1}{2\pi^2} \left[ \left( 1 + \frac{m_a}{m_b} \right) \frac{(\sigma \cdot \eta)}{\eta^2} - \frac{(\sigma \cdot \eta)^2}{\eta^4} \right] \exp \left[ - \frac{v_a^2}{2} (\sigma - \eta)^2 \right]. \quad (19)$$

In Eq. (13), we have introduced the Coulomb mean free path  $L_a = v_a^4 / (N\Gamma_a)$ .

If we assume that the collisions are infrequent i.e.,  $kL_a \gg 1$ , then to first

order in  $(kL_a)^{-1}$ , Eq. (13) gives

$$\begin{aligned}
F_a(\mathbf{k}, \boldsymbol{\sigma}, s) = & e^{s\sigma_z/k} \left[ Q_a(\boldsymbol{\sigma}) + \frac{ie_a N E_z'}{m_a k} P_a(\boldsymbol{\sigma}) \right] \\
& + \frac{v_a^4}{kL_a} e^{s\sigma_z/k} \int_{\sigma_z}^{\infty} d\sigma_z' e^{-s\sigma_z'/k} \int d\eta e^{s\eta_z/k} \left[ (K_1^b + K_2^a + K_3^a) \right. \\
& \left. \left\{ Q_a(\eta) + \frac{ie_a E_z' N}{m_a k} P_a(\eta) \right\} + K_4^a \left\{ Q_b(\eta) + \frac{ie_b E_z' N}{m_b k} P_b(\eta) \right\} \right] \quad (20)
\end{aligned}$$

On substituting Eq. (20) in Eq. (12), we obtain

$$E_z' = \frac{\Phi(k, s)}{\Psi(k, s)}, \quad (21)$$

where

$$\begin{aligned}
\Phi(k, s) = & + \frac{4i\pi}{k} \left[ \sum_{J=e, i} e_J Q_J(0) + \frac{v_e^4}{kL_e} \int_0^{\infty} d\sigma_z' e^{-s\sigma_z'/k} \int d\eta e^{s\eta_z/k} \right. \\
& \left\{ Q_e(\eta) (K_1^i + K_2^e + K_3^e)_{\boldsymbol{\sigma}=0} - \frac{m^2}{M^2} Q_i(\eta) (K_1^e + K_2^i + K_3^i)_{\boldsymbol{\sigma}=0} \right. \\
& \left. \left. + Q_i(\eta) K_4^e(\boldsymbol{\sigma}=0) - \frac{m^2}{M^2} Q_e(\eta) K_4^i(\boldsymbol{\sigma}=0) \right\} \right] \quad (22)
\end{aligned}$$

and

$$\begin{aligned} \Psi(k, s) = 1 - \sum_{J=e,i} \frac{\omega_{pJ}^2}{k^2} P_J(\sigma = 0) - \frac{v_e^4 \omega_{pe}^2}{k^3 L_e} \int_0^\infty d\sigma_z' e^{-s\sigma_z'/k} \\ \int d\eta e^{s\eta_z/k} \left[ P_e(\eta) (K_1^i + K_2^e + K_3^e)_{\sigma=0} - \frac{m}{M} P_i(\eta) K_4^e(\sigma = 0) \right. \\ \left. - \frac{m^2}{M^2} P_e(\eta) K_4^i(\sigma = 0) + \frac{m^3}{M^3} P_i(\eta) (K_1^e + K_2^i + K_3^i)_{\sigma=0} \right]. \quad (2) \end{aligned}$$

In Eq. (23),  $\omega_{pJ}^2 = 4\pi N e^2/m_J$  is the plasma frequency for the species J. We may note here that  $\Phi(k, s)$  depends only on the initial perturbation. If we consider only those perturbations for which  $\Phi(k, s)$  is analytic in the complex s-plane, then for the Laplace inversion of Eq. (21), we have to consider only the zeros  $\Psi(k, s)$  which are given by

$$1 = \sum_{J=e,i} \frac{\omega_{pJ}^2}{k^2} P_J(\sigma = 0) + \frac{v_e^3 \nu_c \omega_{pe}^2}{k^3} (\alpha_{ei} + \alpha_{ii} + \alpha_{ee} + \alpha_{ie}), \quad (3)$$

where  $\nu_c = v_e/L_e$  is the effective electron collision frequency and

$$\begin{aligned} \alpha_{ei} = \int_0^\infty d\sigma_z' e^{-s\sigma_z'/k} \int d\eta e^{s\eta_z/k} \left[ P_e(\eta) K_1^i(\sigma = 0) \right. \\ \left. - \frac{m}{M} P_i(\eta) K_4^e(\sigma = 0) \right], \quad (4) \end{aligned}$$

$$\alpha_{ii} = \frac{m^3}{M^3} \int_0^\infty d\sigma_z' e^{-s\sigma_z'/k} \int d\eta e^{s\eta_z/k} \left[ (K_2^i + K_3^i)_{\sigma=0} \right] P_i(\eta) , \quad (26)$$

$$\alpha_{ee} = \int_0^\infty d\sigma_z' e^{-s\sigma_z'/k} \int d\eta e^{s\eta_z/k} P_e(\eta) (K_2^e + K_3^e)_{\sigma=0} \quad (27)$$

and

$$\alpha_{ie} = \frac{m^2}{M^2} \int_0^\infty d\sigma_z' e^{-s\sigma_z'/k} \int d\eta e^{s\eta_z/k} \left[ \frac{m}{M} P_i(\eta) K_1^e(\sigma=0) - P_e(\eta) K_4^i(\sigma=0) \right] . \quad (28)$$

Eq. (24) represents the required dispersion relation which gives the oscillatory behavior of the plasma under consideration.

### III. ANALYSIS OF THE DISPERSION RELATION

From Eq. (15), we have

$$P_J(\sigma=0) = -\frac{1}{v_J^2} \left[ 1 - \sqrt{2} a_J e^{a_J^2/2} \text{Erf}\left(a_J/\sqrt{2}\right) \right] , \quad (29)$$

where  $a_J = s/(kv_J)$  and

$$\text{Erf}(x) = \int_x^\infty dy e^{-y^2} . \quad (30)$$

While evaluating these integrals ( $P_j$  etc.), strictly speaking we should take into account the pole contribution which gives Landau damping but since this has been discussed earlier, we shall not repeat it here. We shall further assume that  $a_i \gg 1$  but  $a_e \ll 1$  which actually is a consequence of the fact that  $v_e^2 \gg v_i^2$ ; thus on using the proper asymptotic and series expansions for the error function,<sup>17</sup> we immediately get

$$P_i (\sigma = 0) = -\frac{1}{v_i^2 a_i^2} \left( 1 - \frac{3}{a_i^2} \right) + O\left(\frac{1}{a_i^6}\right) \quad (31)$$

and

$$P_e (\sigma = 0) = -\frac{1}{v_e^2} \left( 1 - \frac{\pi^{1/2}}{\sqrt{2}} a_e \right) + O(a_e^2) \quad (32)$$

Like in the collisionless theory, we shall neglect terms of the order of  $a_e^2$ . Moreover, in evaluating the collision integrals, we shall retain terms only upto  $(1/a_i^2)$ .

Let us first consider the electron-ion contribution which is given by  $\alpha_{ei}$  in Eq. (25). On substituting Eqs. (16) and (19) in Eq. (25), we obtain

$$\alpha_{ei} = I_1^i + I_4^e, \quad (33)$$

where

$$I_1^i = \frac{1}{2\pi^2} \int_0^\infty d\sigma e^{-s\sigma/k} \int d\eta P_e(\eta) \exp \left[ \frac{s\eta_z}{k} - \frac{v_i^2}{2} (\sigma \hat{e}_z - \eta)^2 \right] \left[ \left( 1 + \frac{m}{M} \right) \frac{\sigma(\sigma - \eta_z)}{(\sigma \hat{e}_z - \eta)^2} - \frac{\sigma^2(\sigma - \eta_z)^2}{(\sigma \hat{e}_z - \eta)^4} \right] \quad (34)$$

and

$$I_4^e = -\frac{m}{2\pi^2 M} \int_0^\infty d\sigma e^{-s\sigma/k} \int d\eta P_i(\eta) \exp \left[ \frac{s\eta_z}{k} - \frac{v_e^2}{2} (\sigma \hat{e}_z - \eta)^2 \right] \left( \frac{\sigma\eta_z}{\eta^2} - \frac{\sigma^2 \eta_z^2}{\eta^4} \right) \cdot \quad (35)$$

On rewriting Eq. (15) as

$$F_a(\eta) = -e^{v_a^2 \eta_z^2 / 2} \int_0^\infty dy (y + \eta_z) \exp \left[ -\frac{s}{k} (y + \eta_z) - \frac{v_a^2}{2} (y + \eta_z)^2 \right] \quad , \quad (36)$$

and on using Eq. (36) for  $P_i$ , Eq. (35) on performing  $\eta$ -integration reduces to

$$I_4^e = \frac{m}{4\pi^{1/2} M} \int_0^\infty \frac{d\nu}{(\nu + b^2)^{5/2}} \int_0^\infty d\sigma \sigma \exp \left( -\frac{s\sigma}{k} - \frac{v_e^2 \sigma^2}{2} \right) \int_0^\infty dy \exp \left[ \frac{\beta^2}{4(\nu + b^2)} - \frac{sy}{k} - \frac{v_i^2 y^2}{2} \right] \left[ 1 - \nu\sigma y + \beta y + \frac{\beta}{2(\nu + b^2)} \left\{ \beta(1 - \nu\sigma y) - 3\nu\sigma \right\} - \frac{\nu\sigma\beta^3}{4(\nu + b^2)^2} \right], \quad (37)$$

where

$$\beta = (v_e^2 \sigma - v_i^2 y) \quad \text{and} \quad b^2 = \frac{1}{2} (v_e^2 + v_i^2). \quad (38)$$

Now to carry out the  $y$ -integration, we shall neglect terms of the order of

$\delta^2 = v_i^2/v_e^2$  as well as terms of the order of  $(1/a_i^4)$ . The resulting expression

can then be put into the dimensionless form by using the variables  $(v_e \sigma) = x$

and  $1/2 v_e^2 (\nu + b^2)^{-1} = z^2$ ; Eq. (37) thus reads

$$I_4^e = \frac{m}{\pi^{1/2} M v_e^6} \int_0^{(1-\delta^2/2)} dz \int_0^\infty dx \frac{x}{\xi} \exp \left[ -a_e x - \frac{x^2}{2} (1 - z^2) \right] \left[ b_0 + \frac{b_1}{\xi} + \frac{1}{\xi^2} \left( b_2 - \frac{\delta^2}{2} b_0 \right) + \frac{1}{\xi^3} \left( b_3 - \frac{\delta^2}{2} b_1 \right) - \frac{\delta^2 b_2}{2\xi^4} - \frac{\delta^2 b_3}{2\xi^5} \right], \quad (39)$$

where

$$\begin{aligned}
\xi &= (a_e + \delta^2 z^2 x) , \\
b_0 &= z^2 \left[ 1 - \frac{x^2}{2} (3 - 5z^2) - \frac{z^2 x^4}{2} (1 - z^2) \right] , \\
b_1 &= -\frac{x}{2} \left[ 1 - 3z^2 + z^2 x^2 (1 - z^2) \right] , \\
b_2 &= -\delta^2 z^2 (1 - x^2 + x^2 z^2)
\end{aligned}$$

and

$$b_3 = -\frac{\delta^4 z^2 x}{2} (1 - z^2) . \quad (40)$$

$I_1^i$  can be evaluated by proceeding on the lines similar to the ones for  $I_4^e$  and we get

$$\begin{aligned}
I_1^i &= \frac{2^{1/2}}{\pi^{1/2} v_e^6} \int_0^{(1-\delta^2/2)} dy \int_0^\infty dz z \int_z^\infty dx \exp \left[ -a_e x - \frac{x^2}{2} (1 - y^2) \right] \\
&\quad \left[ y^2 + (1 - y^2) \left\{ -y^2 x^2 + \frac{xz}{2} (1 + x^2 y^2 - 3y^2 - x^2 y^4) \right\} \right] . \quad (41)
\end{aligned}$$

The ion-ion contribution can be similarly obtained by simplifying Eq. (26); this gives

$$a_{ii} = -8 / (15\pi^{1/2} v_i^6 a_i^5) . \quad (42)$$

This, with  $i$  replaced by  $e$ , is exactly the same as the electron-electron contribution obtained by Comisar.<sup>12</sup> Eq. (28) however, on integration gives

$$\alpha_{ie} = \frac{\sqrt{2} m^2}{\pi^{1/2} v_e^3 v_i^3 a_i^3 M^2} \left[ 1 - \frac{10}{a_i^2} + \frac{8m}{3a_e^2 M} - \frac{2\delta^2 M}{3m} \left( 1 - \frac{6}{a_i^2} \right) - \frac{\sqrt{2} a_e}{\pi^{1/2}} S \right], \quad (43)$$

where

$$S = \int_1^\infty \frac{d\nu}{\nu^{5/2}} \int_0^\infty dy e^{-y^2} \int_{-\infty}^\infty dz \exp \left[ -z^2 - \frac{\sqrt{2} v_e y}{\nu^{1/2} b} z \right] \left[ \left( 1 - \frac{6}{a_i^2} \right) \left( 1 - \frac{M v_i^2}{m b^2} z^2 \right) + \frac{4M v_i^4}{\nu m b^4 a_i^2} z^4 \right]. \quad (44)$$

Now it is easy to see that  $S_{\max}$  is  $1/\delta$ ; so we immediately find that  $\alpha_{ie} \sim (a_e^3 M/m) \alpha_{ii}$  and can thus be neglected. On using the similar argument, we can show that  $\alpha_{ee} \sim (a_e^3 \delta M/m) \alpha_{ii}$ . Hence to analyze the dispersion relation, we have to consider only the ion-ion and the electron-ion collisions.

Let us now introduce the following dimensionless variables:

$$\frac{\omega}{\omega_{pi}} = \omega^*, \quad \frac{\nu_c}{\omega_{pi}} = B, \quad \frac{T_i}{T_e} = T \quad \text{and} \quad a = k\lambda_e,$$

where  $\lambda_e^{-2} = KT_e / (4\pi Ne^2)$ , is the Debye length for the electrons. Eq. (24), with the help of Eqs. (31), (32), (39), (41) and (42) can be written as

$$1 = \frac{1}{\omega^{*2}} \left( 1 + \frac{3a^2 T}{\omega^{*2}} \right) - \frac{1}{a^2} \left( 1 - \frac{\pi^{1/2}}{2^{1/2}} a_e \right) + C \quad (45)$$

where

$$C = (C_{ii} + C_{ei}) \frac{B\epsilon^{1/2}}{a^3} \quad (46)$$

with  $\epsilon = m/M$ ,  $C_{ii} = v_e^6 \alpha_{ii}$  and  $C_{ei} = v_e^6 \alpha_{ei}$ . We shall solve Eq. (45) by the method of successive approximations. To lowest order, on neglecting  $C$  and  $a_e$ , we obtain

$$\omega^{*2} = \frac{1}{2\chi} \left[ 1 + (1 + 12\chi a^2 T)^{1/2} \right] \equiv \omega_0^2 \text{ say} \quad (47)$$

Here  $\chi$  stands for  $(1 + 1/a^2)$ . For  $T \ll 1$ , Eq. (47) further reduces to

$$\omega_0^2 = \frac{1}{\chi} + 3k^2 \lambda_i^2 \quad (48)$$

the result obtained by Bernstein and Trehan.<sup>18</sup>  $\lambda_i$  in Eq. (48) is the ion-Debye length. To next order Eq. (45) takes the form

$$\chi = \frac{1}{\omega^{*2}} \left( 1 + \frac{3a^2 T}{\omega^{*2}} \right) - i \left( \frac{\epsilon\pi}{2} \right)^{1/2} \frac{\omega_0}{a^3} + C (\omega^* = \omega_0) \quad (49)$$

which we shall solve numerically. It is worth pointing out that

$$C_{ii}(\omega^* = \omega_0) = - \frac{8i a^5}{15\pi^{1/2} T^{1/2} \omega_0^5}, \quad (50)$$

being pure imaginary, does not affect the real part of  $\omega^*$ ; however,  $C_{ei}(\omega^* = \omega_0)$  is a complex quantity and thus changes both the real and the imaginary parts of  $\omega^*$ .  $C_{ii}$  and  $C_{ei}$  for various values of  $a$  and  $T$  are given in Table 1.

According to Eq. (49), although  $\text{Re } \omega^*(\omega_r^*)$  increases with an increase in  $a$ ,  $T$  as well as  $B$  but this increase, as shown in Tables (2a) and (2b), is negligible. As illustrated in Figures 1 and 2,  $\text{Im } \omega^*(\omega_i^*)$  increases with increase in  $a$  and  $B$  but decreases with increase in  $T$ . So two-body Coulomb collisions have a tendency to stabilize the ion acoustic waves. As  $T$  increases, Landau damping takes over the collisional damping. For the sake of completeness, in Figure 3, we have shown the thermal effects on the ion waves.

In a system where the collisions are frequent, this model will break down; in this case one should use the kinetic equation which takes into account the correlations between the charged particles.

#### IV. CONCLUSIONS

Independent of the ratio of the ion temperature to the electron temperature  $T$ , the characteristic frequency of the ion acoustic waves in a plasma with weak Coulomb collisions gets affected only by the electron-ion collisions.

Just as in a collisionless plasma, in this case also  $\omega_r$  increases with increase in  $T$ ; moreover it increases with increase in the collision frequency  $\nu_c$  as well as with  $(k\lambda_e)$ . The electron-electron and the ion-electron collisions are negligible compared to the ion-ion and the electron-ion collisions. The latter two play a somewhat equally important role in damping these waves. The collisional damping increases with increase in  $\nu_c$  and  $(k\lambda_e)$  but decreases with increase in  $T$ .

## V. ACKNOWLEDGMENTS

This work was carried out while the author held the Post-doctoral Resident Research Associateship of the National Academy of Sciences, National Research Council. I am thankful to Mrs. E. Glover for assistance with the numerical computations.

## REFERENCES

1. L. Tonks and I. Langmuir, Phys. Rev. 33, 195 (1929).
2. J. D. Jackson, J. Nucl. Energy: Pt. C1, 171 (1960).
3. E. A. Jackson, Phys. Fluids, 3, 786 (1960).
4. I. B. Bernstein, E. A. Frieman, R. M. Kulsrud and M. N. Rosenbluth, Phys. Fluids 3, 136 (1960); I. B. Bernstein and R. M. Kulsrud, Phys. Fluids 3, 937 (1960).
5. B. D. Fried and R. W. Gould, Phys. Fluids 4, 139 (1961).
6. R. W. Revans, Phys. Rev. 44, 798 (1933).
7. A. Y. Wong, R. W. Motley and N. D'Angelo, Phys. Rev. 133, A436 (1964).
8. D. Bhadra and R. K. Varma, Phys. Fluids 7, 1091 (1964).
9. P. L. Bhatnagar, E. P. Gross and M. Krook, Phys. Rev. 94, 511 (1954).
10. R. M. Kulsrud and C. S. Shen, Phys. Fluids 9, 177 (1966).
11. M. N. Rosenbluth, W. M. MacDonald and D. L. Judd, Phys. Rev. 107, 1 (1957).
12. G. G. Comisar, Phys. Fluids 6, 76 (1963) and 6,1660 (1963).
13. B. Buti and R. K. Jain, Phys. Fluids 8, 2080 (1965).
14. B. Buti and S. K. Trehan, Ann. Phys. 40, 296 (1966).
15. B. Buti, Phys. Rev. (to be published).
16. A. F. Kuckes, Phys. Fluids 7, 511 (1964).
17. E. Jahnke and F. Emde, Tables of Higher Functions (McGraw-Hill Book Company, Inc., New York, 1960), 6th ed.
18. I. B. Bernstein and S. K. Trehan, Nuc. Fusion 1, 3 (1960).

Table 1

 $C_{ii}$  and  $C_{ei}$  for Various Values of  $a = k\lambda_e$  and  $T = T_i/T_e$ 

a	T	Im $C_{ii}$	Re $C_{ei}$	Im $C_{ei}$
1.0	0.10	- 2.3183	3.3309	-0.0228
	0.05	- 6.9959	3.6253	0.0217
	0.01	-10.7640	4.0307	0.0686
0.1	0.10	- 0.5656	2.1755	0.0465
	0.05	- 1.0082	2.2879	0.0648
	0.01	- 2.8690	2.4265	0.0825
0.05	0.10	- 0.5568	2.1659	0.0471
	0.05	- 0.9915	2.2770	0.0652
	0.01	- 2.8175	2.4039	0.0826
0.01	0.10	- 0.5540	2.1629	0.0473
	0.05	- 0.9862	2.2736	0.0653
	0.01	- 2.8011	2.3999	0.0827
0.001	0.10	- 0.5539	2.1627	0.0473
	0.05	- 0.9860	2.2734	0.0653
	0.01	- 2.8004	2.3998	0.0827

Table 2a

$\text{Re } \omega^*$  for  $a = 0.1$  and  $T = 0.1, 0.05$  and  $0.01$  for Various Values of  $B$

B	T = 0.1	T = 0.05	T = 0.01
0.0	0.110936	0.105914	0.100927
0.0001	0.110938	0.105916	0.100929
0.001	0.110959	0.105939	0.100953
0.01	0.111167	0.106160	0.101187
0.05	0.112108	0.107161	0.102213
0.10	0.113323	0.108447	0.103456

Table 2b

$\text{Re } \omega^*$  for  $a = 1.0$  and  $T = 0.1, 0.05$  and  $0.01$  for Various Values of  $B$

B	T = 0.1	T = 0.05	T = 0.01
0.0	0.883308	0.858313	0.838316
0.0001	0.883309	0.858313	0.838317
0.001	0.883315	0.858318	0.838322
0.01	0.883376	0.858370	0.838368
0.05	0.883647	0.858599	0.838566
0.10	0.883987	0.858881	0.838807

## ILLUSTRATIONS

Figure 1. Variation of  $\text{Im } \omega^*$  with  $B$  for  $T = 0.1$  is illustrated for  $a = 0.01, 0.1$  and  $1.0$

Figure 2. Variation of  $\text{Im } \omega^*$  with  $B$  for  $a = 0.1$  is illustrated for  $T = 0.01, 0.05$  and  $0.1$

Figure 3. Variation of  $\text{Re } \omega^*$  with  $a$  for  $B = 0$  is illustrated for  $T = 0.01$  and  $0.1$





