

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

CRITICAL CASES FOR NEUTRAL FUNCTIONAL DIFFERENTIAL EQUATIONS*

by

Jack K. Hale

Center for Dynamical Systems
Division of Applied Mathematics
Brown University
Providence, Rhode Island

* This research was supported in part by the National Aeronautics and Space Administration under Grant No. NGL 40-002-015, in part by the Air Force Office of Scientific Research under Grant No. AF-AFOSR-67-0693A and in part by the United States Army - Durham under Contract No. DA-31-124-ARO-D-270.

FACILITY FORM 808

N70-26311

(ACCESSION NUMBER)

42

(PARTS)

CR-86367

(NASA CR OR TMX OR AD NUMBER)

(THRU)

1

(CODE)

19

(CATEGORY)

CRITICAL CASES FOR NEUTRAL FUNCTIONAL DIFFERENTIAL EQUATIONS

Jack K. Hale

1. Introduction. A neutral functional differential equation as defined below includes the scalar differential-difference equation

$$(1.1) \quad \frac{d}{dt} [x(t) + ax(t-1) + G(x(t-1))] = bx(t) + cx(t-1) + F(x(t), x(t-1)),$$

where a, b, c are constants and $G(x), F(y, x)$ are continuous functions of x, y . For any continuous function φ defined on $[-1, 0]$, a solution of (1.1) through φ is a continuous function x defined on some interval $[-1, \alpha)$, $\alpha > 0$, which coincides with φ on $[-1, 0]$ and is such that the expression

$$x(t) + ax(t-1) + G(x(t-1))$$

[not $x(t)$] is continuously differentiable on $(0, \alpha)$ and satisfies (1.1) on $(0, \alpha)$.

It has been shown in [1] that the solution $x = 0$ of (1.1) is asymptotically stable provided the solution $x = 0$ of the linear equation

$$(1.2) \quad \frac{d}{dt} [x(t) + ax(t-1)] = bx(t) + cx(t-1)$$

is asymptotically stable and the functions $G(x), F(y, x)$ as well as their first derivatives vanish at $x = y = 0$. Furthermore, the solution $x = 0$ of (1.2) is asymptotically stable if all roots of the characteristic equation

$$(1.3) \quad \lambda(1+ae^{-\lambda}) = b + ce^{-\lambda}$$

have real parts $\leq -\delta < 0$.

The purpose of this paper is to obtain sufficient conditions for the zero solution of (1.1) to be asymptotically stable even when some roots of (1.3) have zero real parts. Of course, the discussion involves much more general equations, but it is easier to describe the essential ideas for (1.1). A basic hypothesis for (1.1) is that $|a| < 1$ and no roots of (1.3) have positive real parts. This hypothesis eliminates the possibility of a sequence of distinct roots λ_j of (1.3) having $\text{Re } \lambda_j \rightarrow 0$ as $j \rightarrow \infty$. Suppose P is the finite dimensional linear subspace of the space C of continuous functions on $[-1, 0]$ which corresponds to all initial values of solutions of (1.2) of the form $p(t)e^{\lambda t}$ where p is a polynomial and λ is a root of (1.3) with $\text{Re } \lambda = 0$. If P has dimension d , it is shown that there exists a d -dimensional manifold P^* in C with zero in P^* , and an ordinary differential equation on P^* such that the stability properties of the zero solution of this equation on P^* determine the stability properties of the zero solution of (1.1). Also,

constructive methods are given for obtaining this information about P^* . The case of zero roots is discussed in detail and generalizes the paper of Lefschetz [2] for ordinary differential equations. The results about P^* seem to be new even for ordinary differential equations although a partial result of this type appears in Pliss [3]. For retarded equations (that is, $a = 0$, $G = 0$), Shimanov [4,5] has given some sufficient conditions for the stability of (1.1) in special cases. For neutral equations, the presence of the term G introduces many new difficulties in the discussion and, in fact, seems to prevent the use of the converse theorems of Liapunov, a tool systematically employed by Shimanov. The approach used here follows more closely the method of integral manifolds in the spirit of Krylov, Bogoluibov and Mitropolski (see [6]).

2. Notation and background. Let E^n be a real or complex n -dimensional linear vector space with norm $|\cdot|$, $r \geq 0$ a given real number and C be the space of continuous functions mapping $[-r, 0]$ into E^n with $|\varphi| = \sup_{-r \leq \theta \leq 0} |\varphi(\theta)|$ for $\varphi \in C$. If x is a continuous function taking $[\sigma-r, \sigma+A]$, $A \geq 0$, into E^n , then, for each $t \in [\sigma, \sigma+A]$, we let $x_t \in C$ be defined by $x_t(\theta) = x(t+\theta)$, $-r \leq \theta \leq 0$. Suppose $\mu(\theta), \eta(\theta)$, are $n \times n$ matrix functions of bounded variation in θ , $-r \leq \theta \leq 0$, $\varphi \in C$ and define

$$\begin{aligned}
 & \text{a) } L(\varphi) = \int_{-r}^0 [d\eta(\theta)]\varphi(\theta) \\
 (2.1) \quad & \text{b) } g(\varphi) = \int_{-r}^0 [d\mu(\theta)]\varphi(\theta) \\
 & \text{c) } D(\varphi) = \varphi(0) - g(\varphi)
 \end{aligned}$$

for all φ in C . The functions L and D are continuous linear operators. Also, suppose $G: C \rightarrow E^n$, $F: C \rightarrow E^n$, G has a continuous first derivative $G'(\varphi)$, $G(\varphi)$ depends only upon values of $\varphi(\theta)$ for $\theta < 0$ and $G(\varphi), G'(\varphi)$ are uniformly continuous on closed bounded sets in C . Also, suppose there exist continuous scalar functions $r(s), q(s), s \geq 0, r(0) = q(0) = 0$, such that

$$\begin{aligned}
 & \text{a) } \left| \int_{-s}^0 [d_{\theta} \mu(\theta)]\varphi(\theta) \right| \leq r(s)|\varphi| \\
 & \text{b) } F(0) = G(0) = 0 \\
 (2.2) \quad & \text{c) } |F(\varphi) - F(\psi)| \leq q(\sigma)|\varphi - \psi| \\
 & \text{d) } |G(\varphi) - G(\psi)| \leq q(\sigma)|\varphi - \psi|
 \end{aligned}$$

for $s \geq 0, \sigma \geq 0$ and all φ, ψ in C and, furthermore, $|\varphi|, |\psi| \leq \sigma$ in (2.3c), (2.3d).

Our main concern throughout this paper is with the autonomous neutral functional differential equation

$$(2.3) \quad \frac{d}{dt} [D(x_t) - G(x_t)] = L(x_t) + F(x_t).$$

A solution $x = x(\varphi)$ of (2.3) through a point φ in C is a continuous function taking $[-r, A)$, $A > 0$, into E^n such that $x_0 = \varphi$ and $D(x_t)$ is continuously differentiable and satisfies (2.3) for t in $(0, A)$. It is proved in [7, 8] that there is a unique solution $x(\varphi)$ through φ and $x(\varphi)(t)$ is continuous in (t, φ) .

Along with (2.3), we consider the linear system

$$(2.4) \quad \frac{d}{dt} D(y_t) = L(y_t).$$

If the transformation $T(t): C \rightarrow C$ is defined by

$$(2.5) \quad y_t(\varphi) = T(t)\varphi,$$

then it is shown in [9] that $\{T(t), t \geq 0\}$ is a strongly continuous semigroup of linear operators with infinitesimal generator $A: \mathcal{D}(A) \rightarrow C$, $A\varphi(\theta) = \dot{\varphi}(\theta)$, $-r \leq \theta \leq 0$,

$$(2.6) \quad \mathcal{D}(A) = \{\varphi \in C: \dot{\varphi} \in C, \dot{\varphi}(0) = g(\dot{\varphi}) + L(\varphi)\}$$

and the spectrum $\sigma(A)$ of A consists of those λ which satisfy the characteristic equation

$$(2.7) \quad \det \Delta(\lambda) = 0, \quad \Delta(\lambda) = \lambda \left[I - \int_{-r}^0 e^{\lambda \theta} d\mu(\theta) \right] - \int_{-r}^0 e^{\lambda \theta} d\eta(\theta).$$

Let $\{T_D(t), t \geq 0\}$ be the strongly continuous semigroup of linear transformations associated with the solution of the equation

$$(2.8) \quad \frac{d}{dt} D(x_t) = 0.$$

Definition. The order a_D of D is defined by

$$(2.9) \quad a_D = \inf \{ \text{real } a: \text{ there is a constant } K(a) \text{ with} \\ |T_D(t)\varphi| \leq K(a)e^{at}|\varphi|, t \geq 0, \varphi \in C, D(\varphi) = 0 \}.$$

If

$$(2.10) \quad D(\varphi) = \varphi(0) - \sum_{k=1}^N A_k \varphi(-\tau_k)$$

the A_k are $n \times n$ constant matrices, each $\tau_j > 0$ and τ_j/τ_k is rational for $N > 1$, it is shown in [10] that

$$(2.11) \quad a_D = \sup \{ \text{Re } \lambda: \det(I - \sum A_k e^{-\lambda \tau_k}) = 0 \}.$$

Suppose $a_D < 0$ and all roots of (2.7) have nonpositive real parts. If $\Lambda = \{\lambda: \det \Delta(\lambda) = 0, \text{Re } \lambda = 0\}$, then Λ is a

finite set and it follows from [9] that the space C can be de-
composed by Λ as $C = P \oplus Q$ where P, Q are subspaces of C
invariant under $T(t)$ and A , the space P is finite dimensional
and corresponds to the initial values of all those solutions of
(2.4) which are of the form $p(t)e^{\lambda t}$ where $p(t)$ is a polynomial
in t and $\lambda \in \Lambda$.

Let $X(t)$ be the $n \times n$ matrix function defined for all
 $t \in [0, \infty)$, of bounded variation in t and continuous in t from
the right such that

$$(2.12) \quad D(X_t) = \int_0^t L(X_s) ds + I, \quad t \geq 0,$$

$$X_0(\theta) = \begin{cases} 0 & -r \leq \theta < 0 \\ I & \theta = 0 \end{cases} .$$

Since $X(t)$ is a solution (2.4), it is reasonable to let

$$(2.13) \quad X_t = T(t)X_0.$$

Using the same arguments as in [1,9], it is easily shown that the
solution of (2.3) with initial value φ satisfies the equation

$$(2.14) \quad x_t - X_0 G(x_t) = T(t)[\varphi - X_0 G(\varphi)]$$

$$+ \int_0^t \{d_s[-T(t-s)X_0]G(x_s) + T(t-s)X_0 F(x_s)\} ds, \quad t \geq 0.$$

Conversely, any solution of (2.14) satisfies (2.3). The integrals in (2.14) are evaluated at each θ in $[-r, 0]$ as ordinary integrals in E^n . Also, if C is decomposed by Λ as $C = P \oplus Q$, then equation (2.14) is equivalent to

$$\begin{aligned}
 \text{a) } x_t^P - X_0^P G(x_t) &= T(t) [\varphi^P - X_0^P G(\varphi)] \\
 &+ \int_0^t (d_s [-T(t-s) X_0^P] G(x_s) + T(t-s) X_0^P F(x_s)) ds \\
 \text{b) } x_t^Q - X_0^Q G(x_t) &= T(t) [\varphi^Q - X_0^Q G(\varphi)] \\
 &+ \int_0^t (d_s [-T(t-s) X_0^Q] G(x_s) + T(t-s) X_0^Q F(x_s)) ds
 \end{aligned}
 \tag{2.15}$$

where the superscripts P and Q designate the projections of the corresponding functions onto the subspaces P and Q , respectively. Everything is clear in (2.15) except for the meaning of the projections X_0^P, X_0^Q since X_0 is not continuous. These terms will be defined after we have given an explicit way for determining the projections of C onto P and Q .

Projection operators taking C onto P and Q are easily determined by means of the adjoint differential equation

$$(2.16) \quad \frac{d}{d\tau} [z(\theta) - \int_{-r}^0 z(\tau-\theta) d\mu(\theta)] = - \int_{-r}^0 z(\tau-\theta) d\eta(\theta)$$

and the bilinear form

$$(2.17) \quad (\alpha, \varphi) = \alpha(0)D(\varphi) + \int_{-r}^0 \int_0^\theta \alpha(\xi - \theta) [r\mu(\theta)] \varphi(\xi) d\xi \\ - \int_{-r}^0 \int_0^\theta \alpha(\xi - \theta) [d\eta(\theta)] \varphi(\xi) d\xi$$

defined for all $\alpha \in C^* = C([0, r], E^n)$, $\alpha \in C^*$, $\varphi \in C$.

If $\Phi = (\varphi_1, \dots, \varphi_\nu)$ is a basis for the initial values of those solutions of (2.4) of the form $p(t)e^{\lambda t}$ where p is a polynomial and $\lambda \in \Lambda$ and $\Psi = \text{col}(\psi_1, \dots, \psi_\nu)$ is a basis for the initial values of those solutions of (2.16) of the form $p(t)e^{-\lambda t}$, p a polynomial, $\lambda \in \Lambda$, then it is shown in [9] that the $\nu \times \nu$ matrix $(\Psi, \Phi) = ((\psi_i, \varphi_j))$, $i, j = 1, 2, \dots, \nu$ is nonsingular and, therefore, can be assumed to be the identity. If Φ, Ψ are defined in this way, then, for any $\varphi \in C$, we define φ^P, φ^Q by

$$(2.18) \quad \varphi = \varphi^P + \varphi^Q \\ \varphi^P = \Phi(\Psi, \varphi).$$

One can now show that (Ψ, X_0) is well defined and $(\Psi, X_0) = \Psi(0)$.

Therefore, if we put

$$(2.19) \quad X_0^P = \Phi\Psi(0), \quad X_0^Q = X_0 - X_0^P$$

the quantities in (2.15) are well defined.

It is shown in [10] that the hypothesis $a_{D_0} < 0$ implies

there are $K \geq 1$, $\alpha > 0$, such that

$$(2.20) \quad \begin{aligned} \text{a) } & |T(t)\varphi| \leq Ke^{-\alpha t} |\varphi|, \quad t \geq 0, \varphi \in Q, \\ \text{b) } & |T(t)X_0^Q| + \int_0^1 |d_s T(t-s)X_0^Q| \leq Ke^{-\alpha t}, \quad t \geq 0. \end{aligned}$$

3. Integral manifolds in critical cases. Throughout the remainder of this paper, we assume D is given in (2.9), $a_D < 0$ and the space C is decomposed by $\Lambda = \{\lambda: \det \Delta(\lambda) = 0, \operatorname{Re} \lambda = 0\}$ as $C = P \oplus Q$ where P, Q are defined as in the previous section. Our objective in this section is to prove there is a ν -dimensional manifold in C which is an integral manifold of system (2.4) in a neighborhood of zero and the stability properties of the solution $x = 0$ relative to this manifold determine the stability properties of the solution $x = 0$ of (2.4). We remark that if there are some roots of (2.8) with positive real parts, then one could obtain the existence of the manifold corresponding to the roots with zero real parts by using essentially the same procedure as below. It only complicates the notation by forcing one to consider some integrals from 0 to $+\infty$ to take care of the roots with positive real parts.

Let $PC = PC([-r, 0], E^n)$ be the space of functions mapping $[-r, 0]$ into E^n which are right continuous and bounded on $[-r, 0]$ and, for any φ in PC , define $|\varphi| = \sup_{\theta \in [-r, 0]} |\varphi(\theta)|$.

It is not difficult to see that the operator $T(t)$ associated with the linear system (2.5) can be defined on PC. Also, for any $\varphi \in PC$ with $\dot{\varphi} \in PC$, for $\theta < 0$,

$$\lim_{t \rightarrow 0^+} \frac{1}{t} [T(t)\varphi(\theta) - \varphi(\theta)] = \lim_{t \rightarrow 0^+} \frac{1}{t} [\varphi(t+\theta) - \varphi(\theta)] = \dot{\varphi}(\theta)$$

and, for $\theta = 0$,

$$\begin{aligned} \lim_{t \rightarrow 0^+} \frac{1}{t} [T(t)\varphi(0) - \varphi(0)] &= \lim_{t \rightarrow 0^+} \frac{1}{t} [g(T(t)\varphi) - g(\varphi) - \int_0^t L(T(s)\varphi) ds] \\ &= g(\dot{\varphi}) + L(\varphi) \end{aligned}$$

since g satisfies (2.1b). Therefore, if we let $A: \mathcal{D}(A) \rightarrow PC$,

$\mathcal{D}(A) = \{\varphi \in PC: \dot{\varphi} \in PC\}$, be defined by

$$(3.1) \quad A\varphi(\theta) = \begin{cases} \dot{\varphi}(\theta), & -r \leq \theta < 0, \\ g(\dot{\varphi}) + L(\varphi), & \theta = 0, \end{cases}$$

then

$$(3.2) \quad \frac{d}{dt} T(t)\varphi = T(t)A\varphi$$

for all φ in $\mathcal{D}(A)$.

Note that if φ is continuously differentiable on $[-r, 0]$ and $A\varphi$ is continuous, then $A\varphi$ coincides with the action on φ of the infinitesimal generator of the semigroup on C generated by

(2.4). Therefore, we may consider the operator A in (3.2) as an extension of the infinitesimal generator of (2.4) to PC .

The variation of constants formula (2.14) for the solution of (2.3) suggests the change of variables $x_t - X_0 G(x_t) = z_t$ to obtain a new equation for z_t in PC . This transformation is a well defined transformation from PC to PC and preserves stability properties since $G(x_t)$ depends only upon the values of $x_t(\theta)$ for $-r \leq \theta < 0$ and, therefore, $G(x_t) = G(z_t)$. Thus, if, in (2.14),

$$(3.3) \quad \begin{aligned} z_t &= x_t - X_0 G(x_t), \\ x_t &= z_t + X_0 G(z_t) \stackrel{\text{def}}{=} H(z_t), \end{aligned}$$

then (2.14) becomes

$$(3.4) \quad z_t = T(t)z_0 + \int_0^t \{[-d_s T(t-s)X_0]G(z_s) + T(t-s)X_0 F(H(z_s))\} ds.$$

Let Φ, Ψ be the matrices defined in Section 2 for the decomposition $C = P \oplus Q$, $(\Psi, \Phi) = I$, and let E be the $q \times q$ matrix such that $T(t)\Phi = \Phi \exp Et$, $t \in (-\infty, \infty)$. The spectrum of E is Λ . For any $\varphi \in PC$, one can define (Ψ, φ) and, therefore, it is meaningful to put

$$(3.5) \quad \varphi^P = \Phi(\Psi, \varphi), \quad \varphi^Q = \varphi - \varphi^P, \quad \varphi \in PC.$$

Also, the equation (3.4) can be split as (2.15) with appropriate

substitutions from (3.3). Furthermore, if $z_t^P = \Phi u(t)$, then it follows from (2.15a) and the transformation (3.3) that

$$\begin{aligned} \Phi u(t) = & \Phi e^{Et} u(0) + \Phi \int_0^t \{ [-d_s e^{E(t-s)} \Psi(0)] G(z_s) \\ & + e^{E(t-s)} \Psi(C) F(H(z_s)) \} ds. \end{aligned}$$

Therefore, we see that equation (3.4) is equivalent to

$$\begin{aligned} (3.6) \quad z_t &= \Phi u(t) + w_t, \quad w_t \in Q \\ \frac{du(t)}{dt} &= Eu(t) + F_1(u(t), w_t) \\ w_t &= T(t)w_0 + \int_0^t \{ -[d_s T(t-s) X_0^Q] g_0(u(s), w_s) + T(t-s) X_0^Q F_0(u(s), w_s) \} ds, \end{aligned}$$

where

$$\begin{aligned} (3.7) \quad F_1(u, \varphi) &= \Psi(0) F(H(\Phi u + \varphi)) + E \Psi(0) G(\Phi u + \varphi) \\ G_0(u, \varphi) &= G(\Phi u + \varphi) \\ F_0(u, \varphi) &= F(H(\Phi u + \varphi)), \quad u \in E^V, \quad \varphi \in Q. \end{aligned}$$

For any $\rho > 0$, let $\dot{\Omega}_\rho = \{(u, \varphi) \in E^V \times Q: 0 \leq |u| < \infty, 0 \leq |\varphi| \leq \rho\}$ and let F_1^e, F_0^e, G_0^e be functions defined on $\dot{\Omega}_\rho$, which coincide with F_1, F_0, G_0 respectively on $\{(u, \varphi) \in E^V \times Q: 0 \leq |u|, |\varphi| \leq \rho\}$ and

$$\begin{aligned} (3.8) \quad F_j^e(u, \varphi) &= F_j\left(\frac{u\varphi}{|u|}, \varphi\right), \quad j = 0, 1 \\ G_0^e(u, \varphi) &= G_0\left(\frac{u\varphi}{|u|}, \varphi\right), \quad \rho < |u| < \infty. \end{aligned}$$

From the properties of F_1 , F_0 and G_0 , there is a nondecreasing continuous function $v(\rho)$, $\rho \geq 0$, $v(0) = 0$, such that

$$\begin{aligned}
 & F_j^e(0,0) = 0, \quad |F_j^e(u,\varphi)| \leq v(\rho)\rho \\
 & |F_j^e(u,\varphi) - F_j^e(v,\psi)| \leq v(\rho)(|u-v| + |\varphi-\psi|), \quad j = 0,1. \\
 (3.9) \quad & G_0^e(0,0) = 0, \quad |G_0^e(u,\varphi)| \leq v(\rho)\rho \\
 & |G_0^e(u,\varphi) - G_0^e(v,\psi)| \leq v(\rho)(|u-v| + |\varphi-\psi|)
 \end{aligned}$$

for $(u,\varphi), (v,\psi) \in \Omega_\rho$ (see, for example, Chafee [11]).

In order to discuss the local properties of (3.6) near $u = 0$, $w_t = 0$, it is convenient to first discuss the global properties of the system

$$\begin{aligned}
 (3.10) \quad & \frac{du(t)}{dt} = Eu(t) + F_1^e(u(t), w_t) \\
 & w_t = T(t)w_0 - \int_0^t [d_s T(t-s)X_0^Q] G_0^e(u(s), w_s) \\
 & \quad + \int_0^t T(t-s)X_0^Q F_0^e(u(s), w_s) ds, \quad t \geq 0.
 \end{aligned}$$

Theorem 3.1. There is a $\rho_0 > 0$ and a lipschitz continuous function $h: E^V \rightarrow Q$ such that for any $0 < \rho \leq \rho_0$, $(u, h(u)) \in \Omega_\rho$, $0 \leq |u| < \infty$ and the set $M_\rho = \{(u,\varphi) \in \Omega_\rho : \varphi = h(u), 0 \leq |u| < \infty\}$ is an integral manifold of (3.10). Furthermore, any solution of (3.10) with initial value in M_ρ is defined for all $t \in (-\infty, \infty)$.

Proof. For simplicity in the estimates we assume $|\exp Et| = 1$. Only slight changes in the proof are needed since for any $\epsilon > 0$ there is in (2.20) and $v(\rho)$ as in (3.9), for any ρ with $v(\rho) = \alpha/4$, there is a constant $K_1 = K_1(\alpha)$ such that

$$\int_{-\infty}^0 |d_s T(-s) X_0^Q| e^{-2v(\rho)s} \leq K_1.$$

In fact, if we write this integral as an infinite sum of integrals of length one and use (2.20), then

$$\begin{aligned} \int_{-\infty}^0 |d_s T(-s) X_0^Q| e^{-2v(\rho)s} &= \sum_{j=0}^{\infty} \int_0^1 |d_s T(j+1-s) X_0^Q| e^{2v(\rho)(j+1-s)} \\ &\leq \sum_{j=0}^{\infty} e^{2v(\rho)(j+1)} \int_0^1 |d_s T(j+1-s) X_0^Q| \\ &\leq \sum_{j=0}^{\infty} e^{-(\alpha-2v(\rho))(j+1)} \\ &\leq \sum_{j=0}^{\infty} e^{-\alpha(j+1)/2} \stackrel{\text{def}}{=} K_1(\alpha). \end{aligned}$$

Let $K_2 = \max(K_1, K)$, choose ρ_0 so that

$$(3.11) \quad 8K_2(1+\alpha^{-1})v(\rho_0) < 1,$$

and for any $0 < \rho \leq \rho_0$, let

$$(3.12) \quad S = \{h: E^V \rightarrow Q, \text{ continuous, } (u, h(u)) \in \Omega_\rho, u \in E^V, h(0) = 0,$$

$$|h(u) - h(v)| \leq |u - v|, u, v \in E^V\}$$

For $h \in S$, let $|h| = \sup\{|h(u)|, u \in E^V\}$.

For any $h \in S$, let $u(t) = u(t, u_0, h)$, $u(0, u_0, h) = u_0$,
be the solution of the system

$$(3.13) \quad \dot{u}(t) = Eu(t) + F_1^e(u(t), h(u(t))), \quad t \geq 0,$$

and define the mapping $\mathcal{P}: S \rightarrow Q$ by the relation

$$(3.14) \quad (\mathcal{P}h)(u_0) = -\int_{-\infty}^0 [d_s T(-s) X_0^Q] G_0^e(u(s), h(u(s))) \\ + \int_{-\infty}^0 T(-s) X_0^Q F_0^e(u(s), h(u(s))) ds,$$

for $u_0 \in E^V$. Our objective is to show that \mathcal{P} has a fixed point in S and then to show that this fixed point defines an integral manifold M_p .

For any h in S , it follows from (3.9), (3.11) and our estimates on $T(t)$ in (2.20) that

$$|(\mathcal{P}h)(u_0)| \leq K\nu(\rho)\rho(1+\alpha^{-1}) \leq \rho.$$

To estimate the dependence of $(\mathcal{P}h)(u_0)$ upon h and u_0 we need the dependence of $u(t, u_0, h)$ upon the same parameters. From (3.13), one easily obtains from the variation of constants formula and simple estimates that

$$|u(t, u_0, h) - u(t, \bar{u}_0, \bar{h})| \leq e^{-2\nu(\rho)t} (|u_0 - \bar{u}_0| + \frac{1}{2}|h - \bar{h}|), \quad t \leq 0.$$

Since

$$\begin{aligned} & |G_0^e(u(t, u_0, h), h(u(t, u_0, h))) - G_0^e(u(t, \bar{u}_0, \bar{h}), \bar{h}(u(t, \bar{u}_0, \bar{h})))| \\ & \leq 2\nu(\rho)|u(t, u_0, h) - u(t, \bar{u}_0, \bar{h})| + \nu(\rho)|h - \bar{h}| \end{aligned}$$

it follows that

$$\begin{aligned} & \left| \int_{-\infty}^0 [d_s T(-s) X_0^Q] [G_0^e(u(s, u_0, h), h(u(s, u_0, h))) - G_0^e(u(s, \bar{u}_0, \bar{h}), \bar{h}(u(s, \bar{u}_0, \bar{h})))] \right| \\ & \leq \int_{-\infty}^0 |d_s T(-s) X_0^Q| [2\nu(\rho)e^{-2\nu(\rho)s} (|u_0 - \bar{u}_0| + \frac{1}{2}|h - \bar{h}|) + \nu(\rho)|h - \bar{h}|] \\ & \leq K\nu(\rho)|h - \bar{h}| + K_1 2\nu(\rho) (|u_0 - \bar{u}_0| + \frac{1}{2}|h - \bar{h}|) \\ & \leq 2\nu(\rho)K_2 (|u_0 - \bar{u}_0| + |h - \bar{h}|). \end{aligned}$$

In a similar manner, one shows that

$$\begin{aligned} & \left| \int_{-\infty}^0 T(-s) [F_0^e(u(s, u_0, h), h(u(s, u_0, h))) - F_0^e(u(s, \bar{u}_0, \bar{h}), \bar{h}(u(s, \bar{u}_0, \bar{h})))] ds \right| \\ & \leq \frac{4K\nu(\rho)}{\alpha} (|u_0 - \bar{u}_0| + \frac{1}{2}|h - \bar{h}|) + \frac{K\nu(\rho)}{\alpha} |h - \bar{h}| \\ & \leq \frac{4K_2\nu(\rho)}{\alpha} (|u_0 - \bar{u}_0| + |h - \bar{h}|). \end{aligned}$$

Combining these estimates and using (3.11), we obtain

$$|(\mathcal{P}h)(u_0) - (\mathcal{P}\bar{h})(u_0)| \leq |\bar{u}_0 - u_0| + \frac{1}{2}|h - \bar{h}|.$$

Therefore, $\mathcal{P}: S \rightarrow S$ and is a contraction. The unique fixed point h of \mathcal{P} in S is easily shown to define an integral manifold satisfying the properties of the theorem. This completes the proof.

Our next objective is to determine the stability properties of the manifold M_ρ given in Theorem 3.1. To do this, we need

Lemma 3.1. There are a $\rho_1 > 0$, $K_3 > 0$, $\alpha_1 > 0$, and a continuous function $p: \mathbb{R}^+ \times \Omega_{\rho_1} \rightarrow \mathbb{Q}$ such that if $(u(t), w_t)$ is a solution of (3.10) with initial value $(u_0, \varphi) \in \mathbb{R}^+ \times \Omega_{\rho_1}$, then $(u(t), w_t)$ exists for $t \geq 0$ and

$$(3.15) \quad w_t = p(t, u(t), \varphi), \quad t \geq 0.$$

Moreover, p satisfies the inequalities

$$(3.16) \quad \begin{aligned} \text{a) } & |p(t, u, \varphi)| \leq \rho_1 \\ \text{b) } & |p(t, u, \varphi) - p(t, \bar{u}, \bar{\varphi})| \leq |u - \bar{u}| + K_3 e^{-\alpha_1 t} |\varphi - \bar{\varphi}| \end{aligned}$$

for $(t, u, \varphi), (t, \bar{u}, \bar{\varphi}) \in \mathbb{R}^+ \times \Omega_{\rho_1}$.

Proof: Consider the equation

$$\frac{du(\tau)}{d\tau} = Eu(\tau) + F_1^e(u(\tau), p(\tau, u(\tau), \varphi)), \quad t \geq 0,$$

$$u(0) = u_0,$$

$$p(t, u_0, \varphi) = T(t)\varphi - \int_0^t [d_s T(t-s)X_0^Q] G_0^e(u(s), p(s, u(s), \varphi)) \\ + \int_0^t T(t-s)X_0^Q F_0^e(u(s), p(s, u(s), \varphi)) ds,$$

for $(t, u_0, \varphi) \in R^+ \times \Omega_{\rho_0}$. By using arguments very similar to the proof of Theorem 3.1, one proves the existence of a $p(t, u, \varphi)$ satisfying (3.16a) which is lipschitzian in u, φ . To show the precise estimate (3.17a) is more difficult and one can use an argument similar to the one used in the basic stability theorem in [1] to obtain this estimate. It is then easy to verify (3.15). The details are omitted.

Theorem 3.2. With h as in Theorem 3.1 and ρ_1 as in Lemma 3.1, if $(u(t), w_t)$ is a solution of (3.10) with initial value $(u_0, \varphi) \in \Omega_{\rho_1}$, then $(u(t), w_t)$ is defined for all $t \geq 0$ and

$$|w_t - h(u(t))| \leq K_1 e^{-\alpha_1 t} |\varphi - h(u_0)|.$$

Proof: If M_{ρ_1} is the integral manifold in Theorem 3.1, then any solution lying in M_{ρ_1} must satisfy (3.15). If $(u(t), w_t)$ is any solution of (3.10) with initial value in Ω_{ρ_1} , then Lemma 3.1 implies

this solution is defined for all $t \geq 0$. For an arbitrary $\tau \geq 0$, the solution of (3.10) through $(u(\tau), h(u(\tau)))$ is defined for all $t \in (-\infty, \infty)$ and lies on M_{ρ_1} from Theorem 3.1. This solution can therefore be considered as the value of a solution of (3.10) at time τ starting from some point (u^*, φ^*) at $t = 0$. Lemma 3.1, therefore, implies

$$\begin{aligned} |w_\tau - h(u(\tau))| &= |p(\tau, u(\tau), \varphi) - p(\tau, u(\tau), \varphi^*)| \\ &\leq K_3 e^{-\alpha_1 t} |\varphi - \varphi^*| \end{aligned}$$

and this proves the theorem.

The following corollary is immediate from Theorem 3.2.

Corollary 3.1. With h as in Theorem 3.1 and ρ_1 as in Lemma 3.1, any solution $(u(t), w_t) \in \Omega_{\rho_1}$ for $t \leq 0$ must lie on M_{ρ_1} .

Let h be as in Theorem 3.1 and ρ_1 as in Lemma 3.1 and consider the ordinary differential equation

$$(3.17) \quad \frac{du(t)}{dt} = Eu(t) + F_1^e(u(t), h(u(t)))$$

which describes the behavior of the solutions $(u(t), w_t) = (u(t), h(u(t)))$ of (3.10) on the integral manifold M_{ρ_1} .

Theorem 3.3. If the solution $u = 0$ of (3.17) is uniformly asymptotically stable (unstable), then the solution $u = 0, w_t = 0$ of (3.10)

[and therefore the solution $x = 0$ of (2.3)] is uniformly asymptotically stable (unstable).

Proof: For any $\delta > 0$, let $B_\delta = \{(u, \varphi) \in E^V \times Q: |u| + |\varphi| < \delta\}$.

Let $M_{\rho_1} \cap B_{b_0}$ be contained in the region of attraction of the solution $u = 0$ of (3.17). We first prove stability. From the hypothesis on (3.17), for any $\epsilon > 0$, $2\epsilon < \rho_1$, there is a $\delta > 0$, $\delta < b_0$, such that any solution $u(t)$ of (3.17) satisfies $|u(t)| < \epsilon$ for $t \geq 0$ provided that $|u(0)| < \delta$. Also, there is a $t_0 = t_0(\delta)$ such that $|u(t)| < \delta/2$ for $t \geq t_0$ since the zero solution of (3.17) is assumed to be uniformly asymptotically stable. With K_1, α_1 as in Theorem 3.2, further restrict δ so that $K_1 \exp(-\alpha_1 t_0(\delta)) < 1/2$. Such a choice of δ is possible since $t_0(\delta) \rightarrow \infty$ as $\delta \rightarrow 0$.

Since the solutions of (3.10) depend continuously upon the initial data and the set $M_{\rho_1} \cap B_\delta$ is precompact, there is a δ_1 -neighborhood V_{δ_1} of $M_{\rho_1} \cap B_\delta$ such that $(u(0), w_0) \in V_{\delta_1}$ implies the solution $(u(t), w_t)$ of (3.10) belongs to $B_{2\epsilon}$ for $0 \leq t \leq t_0$ and $(u(t_0), w_{t_0})$ is in a $\delta/2$ neighborhood $W_{\delta/2}$ of $M_\rho \cap B_{\delta/2}$. Since $K_1 \exp(-\alpha_1 t_0) < 1/2$, it follows that $|w_{t_0}| < \delta_1/2$ and, thus, $(u(t_0), w_{t_0}) \in V_{\delta_1}$. Therefore, $(u(t), w_t)$ must belong to $B_{2\epsilon}$ for all $t \geq 0$. This proves stability of the zero solution of (3.10).

Suppose V_{δ_1} is chosen as above and $(u(0), w_0) \in V_{\delta_1}$. The solution $(u(t), w_t)$ of (3.10) through $(u(0), w_0)$ is in $B_{2\epsilon}$. Since $2\epsilon < \rho_1$, this defines a solution x of (2.3) which is bounded. Since $a_D < 0$, it follows from [12] that the orbit of this solution

has a nonempty ω -limit set. Theorem 3.2 implies this limit set must be in $M_{\rho_1} \cap B_\delta$. Since the ω -limit set is invariant and the only invariant set in $M_{\rho_1} \cap B_\delta$ is zero [by the hypothesis on system (3.17)] it follows that every solution of (3.10) with initial value in B_{δ_1} approaches zero as $t \rightarrow \infty$. This completes the proof of the asymptotic stability.

If the solution $u = 0$ is unstable, then it is obvious that the zero solution of (3.10) is unstable. This completes the proof of the theorem.

4. Stability in critical cases-zero roots. In the previous section, we proved a result (Theorem 3.3) which stated that the asymptotic stability (or instability) of the zero solution of (2.3) are determined by the asymptotic stability (or instability) of the zero solution of an ordinary differential equation (3.17). It, therefore, remains to analyze the behavior of the solutions of (3.17) near $u = 0$. Of course, this is an extremely difficult task and no general procedure is available to treat all possible situations. Therefore, one is forced to consider particular cases, one of which will be discussed in this section.

Suppose D, L, G, F as before and $a_D < 0$. Also, suppose $x = 0$ is an isolated equilibrium point of (2.3) and if $\varphi(y)$ is an analytic function of a ν -vector y in a neighborhood of $y = 0$, then $G(\varphi(y)), F(\varphi(y))$ are analytic functions of y in a neighbor-

hood of $y = 0$. Finally, $\lambda = 0$ is a root of the characteristic equation (2.7) of multiplicity ν , the dimension of the null space of $\Delta(0)$ is ν and all other roots of (2.7) have negative real parts; that is, Λ of Section 2 consists only of the element 0, and if C is decomposed by Λ as $C = P \oplus Q$, then a basis ϕ for P can be chosen to consist of constant functions. Since $T(t)\phi = \phi$, the matrix E in (3.6) is zero. For notational convenience throughout this section, if ϕ is a constant function in C , then ϕ_0 will denote the value of this function in E^n .

If Ψ is a basis for the constant solutions of the adjoint equation (2.16) with $(\Psi, \phi) = I$, then a direct computation shows that

$$(4.1) \quad I = (\Psi, \phi) = \Psi_0 \dot{\Delta}(0) \phi_0,$$

$$\dot{\Delta}(0) = I - \int_{-r}^0 d\mu(\theta) - \int_{-r}^0 \theta d\eta(\theta).$$

Suppose PC is the space of piecewise continuous functions defined in Section 3 and A is defined by (3.1). Let $\Delta^\#(0)$ be that $n \times n$ matrix which takes the range of $\Delta(0)$ onto the range of the transpose $\Delta'(0)$ of $\Delta(0)$ in a one-to-one manner and $\Delta(0)\Delta^\#(0) = I$.

Lemma 4.1. If $Q^* = \{\varphi \in PC: (\Psi, \varphi) = 0\}$ and A is defined by (3.1), then there is a bounded right inverse A^{-1} of A taking PC into $Q^* \cap \mathcal{D}(A)$; that is, if $\varphi \in PC$, $\psi = A^{-1}\varphi$, then $\psi \in Q^* \cap \mathcal{D}(A)$ and $A\psi = AA^{-1}\varphi = \varphi$. Also, if $\varphi \in PC$, then $A^{-1}\varphi$ is defined by

$$(4.2) \quad \begin{aligned} (A^{-1}\varphi)(\theta) &= \varphi_1(\theta) + \phi_0 a \\ \varphi_1(\theta) &= \int_0^\theta \varphi(s) ds + c_1(\varphi), \quad -r \leq \theta \leq 0, \\ c_1(\varphi) &= \Delta^\#(0)(I, \varphi) \\ a &= -(\Psi, \varphi_1) \end{aligned}$$

where (I, φ) is the bilinear form defined in (2.17) evaluated at the identity I .

Proof: With A^{-1} defined in (4.2), it is clear that A^{-1} is a bounded linear operator taking PC in Q^* . Also $dA^{-1}\varphi/d\theta$ is in PC and, for $-r \leq \theta < 0$,

$$(AA^{-1}\varphi)(\theta) = \frac{d}{d\theta} (A^{-1}\varphi)(\theta) = \varphi(\theta).$$

Since $g(\varphi)$ satisfies (2.2a), $(AA^{-1}\varphi)(\theta) = d(A^{-1}\varphi)(\theta)/d\theta$, $-r \leq \theta < 0$, $f(\phi) = 0$ and $\Delta(0)\Delta^\#(0) = I$, we have

$$\begin{aligned}
g\left(\frac{d}{d\theta} A^{-1}\varphi\right) + f(A^{-1}\varphi) &= g(\varphi) + f(\varphi_1 + \phi a) \\
&= g(\varphi) + f(\varphi_1) \\
&= g(\varphi) + f\left(\int_0^{\cdot} \varphi(s) ds\right) + f(\Delta^{\#}(0)(I, \varphi)) \\
&= g(\varphi) + \int_{-r}^0 [d\eta(\theta)] \int_0^{\theta} \varphi(s) ds + \Delta(0)\Delta^{\#}(0)(I, \varphi) \\
&= \varphi(0).
\end{aligned}$$

This completes the proof of the lemma.

Let X_0 be the $n \times n$ matrix function on $[-r, 0]$ defined by $X_0(\theta) = 0$, $-r \leq \theta < 0$, $X_0(0) = I$, the identity.

Lemma 4.2. Suppose \tilde{G}, \tilde{F} are arbitrary functions satisfying (2.2b) - (2.2d), $\tilde{G}(\varphi)$ depends only on the values of $\varphi(\theta)$ for $\theta < 0$ and $\tilde{G}(\varphi(y)), \tilde{F}(\varphi(y))$ are analytic functions of the v -vector y in a neighborhood of $y = 0$ if $\varphi(y)$ is analytic in y in a neighborhood of $y = 0$. With A^{-1} as in Lemma 4.1, the equation

$$(4.3) \quad \varphi + X_0^{\mathcal{Q}} \tilde{G}(\phi y + \varphi) = -A^{-1} X_0^{\mathcal{Q}} \tilde{F}(\phi y + \varphi)$$

has a unique solution $\varphi^{\mathcal{Q}} = \alpha(y)$ in a neighborhood of $y = 0$, $\varphi^{\mathcal{Q}} = 0$, and the solution is analytic in y in a neighborhood of $y = 0$. Furthermore, if the power series expansion of $\tilde{G}(\phi y + \varphi^{\mathcal{Q}}), \tilde{F}(\phi y + \varphi^{\mathcal{Q}})$

begin with terms of degree k in y , then the power series expansion of $\alpha(y)$ begin with terms of at least degree k in y .

The proof is not given since it is a standard application of the method of successive approximations and the properties of A^{-1} .

We now apply these lemmas to the study of the stability of the zero solution of (2.3). Suppose system (2.3) has been transformed to the system (3.6) through the transformation (3.3). We now make an additional transformation on (3.6) to a more convenient form.

Let $\alpha(u)$ be the solution assured by Lemma 4.2 of the equation

$$(4.4) \quad \phi + X_0^0 G(\phi u + \phi) = -A^{-1} X_0^0 F(H(\phi u + \phi))$$

and consider the transformation of variables

$$(4.5) \quad w_t = v_t + \alpha(u(t))$$

in (3.6). If $\beta(u) = \phi u + \alpha(u)$, $F \circ H = \tilde{F}$, the new equation for v_t is given by

$$\begin{aligned}
(4.6) \quad v_t &= T(t)v_0 + T(t)\alpha(u(0)) - \alpha(u(t)) \\
&+ \int_0^t \{d_s[-T(t-s)X_0^Q]G(\beta(u)) + T(t-s)X_0^Q \tilde{F}(\beta(u))\} ds \\
&+ \int_0^t d_s[-T(t-s)X_0^Q]\{G(\beta(u) + v_s) - G(\beta(u))\} \\
&+ \int_0^t T(t-s)X_0^Q\{\tilde{F}(\beta(u) + v_s) - \tilde{F}(\beta(u))\} ds
\end{aligned}$$

where the function u under the integrals is always evaluated at s . Since $u(t)$ has a continuous first derivative, the function $G(\Phi u(t) + \alpha(u(t)))$ has a continuous first derivative. Therefore, the first term in the first integral in (4.6) can be integrated by parts to obtain

$$(4.7) \quad \int_0^t T(t-s)X_0^Q d_s G(\beta(u(s))) + T(t)X_0^Q G(\beta(u(0))) - X_0^Q G(\beta(u(t))).$$

If $r(u(t)) = \alpha(u(t)) + X_0^Q G(\beta(u(t)))$, then $r(u(t))$ is in $\mathcal{D}(A)$ for each t and $r(u(t))$ is also continuously differentiable in t . Therefore, from (3.2),

$$\frac{d}{dt} T(t-s)r(u(s)) = -T(t-s)Ar(u(s)) + T(t-s) \frac{\partial r(u(s))}{\partial s}$$

and,

$$(4.8) \quad T(t)r(u(0)) - T(t)r(u(t)) = -\int_0^t T(t-s)Ar(u(s))ds + \int_0^t T(t-s) \frac{\partial r(u(s))}{\partial s} ds.$$

If we use these relations (4.7), (4.8) in (4.6) together with the fact that $\alpha(u)$ satisfies (4.4), we obtain

$$(4.9) \quad v_t = T(t)v_0 - \int_0^t [d_s T(t-s)X_0^Q] \{G(\beta(u(s))) + w_s - G(\beta(u(s)))\} \\ + \int_0^t T(t-s)X_0^Q \{\tilde{F}(\beta(u(s)) + v_s) - \tilde{F}(\beta(u(s)))\} ds \\ + \int_0^t T(t-s) \frac{\partial \alpha(u(s))}{\partial u} \dot{u}(s) ds$$

$$\beta(u) = \Phi u + \alpha(u), \quad \tilde{F} = F \circ H,$$

where α satisfies (4.4) and H is defined in (3.3).

We summarize these results in the following

Theorem 4.1. Suppose D, L, G, F satisfy the conditions of section 2, zero is a characteristic value of (2.4) whose geometric and algebraic multiplicities are ν and $G(\varphi(y)), F(\varphi(y))$ are analytic functions of the ν -vector y in a neighborhood of $y = 0$. Suppose PC is the space of functions mapping $[-r, 0]$ into E^n which are right continuous and bounded and $A: \mathcal{D}(A) \rightarrow PC, \mathcal{D}(A) \subset PC$ is defined by (3.1). Let C be decomposed by $\{0\}$ as $C = P \oplus Q$. With H defined as in (3.3), Let $\alpha(u)$ be the solution of (4.4). If

$$(4.10) \quad \begin{array}{ll} \text{a)} & x_t = H(z_t), \beta(u) = \Phi u + \alpha(u) \\ \text{b)} & z_t = \beta(u(t)) + v_t, v_t \in Q \\ \text{c)} & \hat{F}(u, \varphi) = F(H(\beta(u) + \varphi)) \\ \text{d)} & \hat{G}(u, \varphi) = G(\beta(u) + \varphi), \end{array}$$

then the initial problem for (2.3) in a neighborhood of $\varphi = 0$ is equivalent to the equations

$$(4.11) \quad a) \quad \frac{du(t)}{dt} = \Psi(0)\hat{F}(u(t), v_t)$$

$$b) \quad v_t = T(t)v_0 - \int_0^t [d_s T(t-s)X_0^Q][\hat{G}(u(s), v_s) - \hat{G}(u(s), 0)] \\ + \int_0^t T(t-s)X_0^Q[\hat{F}(u(s), v_s) - \hat{F}(u(s), 0)]ds \\ + \int_0^t T(t-s) \frac{\partial \chi(u(s))}{\partial u} \Psi(0)\hat{F}(u(s), v_s)ds.$$

There is a degenerate case of (4.11) which needs to be discussed separately; namely, the case in which F in (2.3) satisfies $F(\varphi) = 0$ for all φ in a neighborhood of $\varphi = 0$. Equations (4.11) for this case are

$$\frac{du(t)}{dt} = 0$$

$$v_t = T(t)v_0 - \int_0^t [d_s T(t-s)X_0^Q][\hat{G}(u(s), v_s) - \hat{G}(u(s), 0)].$$

Using the same type of argument as in the proof of the stability theorem in [1], one can show that the solution $(u, v_t) = (0, 0)$ is uniformly stable and, thus, the solution $x = 0$ of (2.3) is uniformly stable. That is, a perturbation in (2.3) which occurs only in the term which is being differentiated does not affect the zero roots of (2.7) provided that the algebraic and geometric multiplicity of this root are the same.

We now discuss the case when $F(\varphi) \neq 0$. System (4.11) is of the same form as (3.6). The results in section 3 did not depend upon the form of F_1, F_0, G_0 but only upon the estimates (3.9). Therefore, the conclusion of Theorem 3.3 is valid for our situation and it remains only to analyze the behavior of the solutions of (3.17) under our present hypotheses. Now the form of the terms in (4.11) are important since they permit us to determine the qualitative behavior of the integral manifold given in Theorem 3.1 near $u = 0$.

To be more specific, let us define

$$\begin{aligned}
 (4.12) \quad F_1(u, \varphi) &= \Psi_0 \hat{F}(u, \varphi) \\
 G_0(u, \varphi) &= \hat{G}(u, \varphi) - \hat{G}(u, 0) \\
 F_0(u, \varphi) &= X_0^0 [\hat{F}(u, \varphi) - \hat{F}(u, 0)] + \frac{\partial X(u)}{\partial s} \Psi_0 \hat{F}(u, \varphi)
 \end{aligned}$$

and write (4.11) as

$$\begin{aligned}
 (4.13) \quad \frac{du(t)}{dt} &= F_1(u, v_t) \\
 v_t &= T(t)v_0 - \int_0^t [d_s T(t-s)] G_0(u(s), v_s) \\
 &\quad + \int_0^t T(t-s) F_0(u(s), v_s) ds.
 \end{aligned}$$

If the functions F_1, G_0, F_0 in (4.12) are extended as F_1^e, G_0^e, F_0^e in (3.8), then Theorem 3.1 guarantees the existence of

an integral manifold M of the extended system with $M = \{(u, h_0(u)), 0 \leq |u| < \infty\}$ where $h_0: E^V \rightarrow Q$ is lipschitz continuous. Furthermore, the proof of that theorem gave h_0 as the limit of the sequence of successive approximations

$$(4.14) \quad h^0 = 0,$$

$$h^{k+1}(u_0) = -\int_{-\infty}^0 [d_s T(-s) X_0^Q] G_0^e(u^k(s), h^k(u(s))) \\ + \int_{-\infty}^0 T(-s) F_0^e(u^k(s), h^k(u(s))) ds,$$

$$\dot{u}^k(t) = F_1(u^k(t), h^k(u^k(t))), \quad u^k(0) = u_0.$$

Furthermore, for every k ,

$$(4.15) \quad |u^k(t, u_0, h^k)| \leq e^{-2\nu(\rho)t} |u_0|, \quad t \geq 0.$$

Suppose the power series expansion of $F(u, 0)$ begins with terms of degree $m \geq 2$ and ρ_0 in (3.11) is further restricted so that $4(m+1)\nu(\rho_0) < \alpha$. From (4.14)

$$h^1(u_0) = \int_{-\infty}^0 T(-s) F_0^e(u^1(s), 0) ds.$$

From the definition of F_0^e in (3.8) and (4.12) and Lemma 4.2, it

follows that there is a constant k such that

$$|F_0^e(u, 0)| \leq k|u|^{m+1}.$$

Therefore,

$$\begin{aligned} |h^1(u_0)| &\leq \int_{-\infty}^0 K e^{\alpha s} k e^{-2(m+1)v(\rho)s} |u_0|^{m+1} ds \\ &\leq 2kK\alpha^{-1} |u_0|^{m+1}. \end{aligned}$$

A simple induction argument on the sequence h^k allows one to conclude that $h_0(u) = \mathcal{O}(|u|^{m+1})$ as $|u| \rightarrow 0$. This is summarized in

Lemma 4.3. Suppose the hypotheses of Theorem 4.1 are satisfied and $h_0: E^v \rightarrow Q$ is the Lipschitz continuous function assured by Theorem 3.1. If the power series expansion of $\hat{F}(u, 0)$ begins with terms of degree m , then

$$h_0(u) = \mathcal{O}(|u|^{m+1}) \text{ as } |u| \rightarrow 0.$$

With h_0 as in Lemma 4.3, the analogue of system (3.17)

is

$$(4.16) \quad \dot{u} = \Psi_0 \hat{F}(u, h_0(u)).$$

Theorem 4.2. Under the assumptions and notations of Theorem 4.1, let $Q(u)$ designate the homogeneous polynomial of the lowest degree terms in the power series expansion of $\Psi(0)\hat{F}(u,0)$. If the zero solution of the ordinary differential equation

$$(4.17) \quad \dot{u} = Q(u)$$

is asymptotically stable, then the zero solution of (2.3) is asymptotically stable (and therefore, the degree of $Q(y)$ is odd). If there is a homogeneous polynomial $A(u)$ which is positive on some set and

$$B(u) = -[\partial A(u)/\partial u]Q(u)$$

is negative definite, then the zero solution of (2.3) is unstable.

Proof: Suppose the degree of $Q(y)$ is m . If the zero solution of (4.17) is asymptotically stable, it is known from ordinary differential equations (see Zubov [13]) that there are two positive definite quadratic forms $A(y)$, $B(y)$ homogeneous of degree $m+1$, $2m$, respectively such that

$$\dot{A}_{(4.17)}(u) = -B(u),$$

where $\dot{A}_{(4.17)}(u)$ represents the derivative of $A(u)$ along the

solutions of (4.17). There is a $\rho_2 > 0$ such that, for $|u| \leq \rho_2$, $F_1^e(u, h_0(u)) = F_1(u, h_0(u)) = \Psi(0)\tilde{F}(u, h_0(u))$. Therefore, for $|u| \leq \rho_2$,

$$\dot{A}_{(4.16)}(u) = -B(u) + \frac{\partial A(u)}{\partial t} \Psi_0[\hat{F}(u, h_0(u)) - \hat{F}(u, 0)].$$

Lemma 3.3 implies the second term in this expression is at least order $2n + 1$ and therefore $\dot{A}_{(4.16)}(u)$ is negative definite in a neighborhood of $u = 0$. The classical Liapunov theorem implies the zero solution of (4.16) is uniformly asymptotically stable. Theorem 3.3 implies the zero solution of (2.3) is uniformly asymptotically stable.

If there is a homogeneous polynomial $A(u)$ of degree ≥ 1 which is not of constant sign and $B(u) = -[\partial A(u)/\partial t]Q(u)$ is negative definite, then

$$\dot{A}_{(4.16)}(u) = -B(u) + \frac{\partial A(u)}{\partial t} \Psi_0[\hat{F}(u, h_0(u)) - \hat{F}(u, 0)].$$

In a sufficiently small neighborhood of $u = 0$, the right hand side of this expression is a positive definite function. The classical Cetaev theorem implies the solution $u = 0$ of (4.16) is unstable. Theorem 3.3 implies the solution $x = 0$ of (2.3) is unstable and the proof is complete.

Corollary 4.1. Under the assumptions and notations of Theorem 4.1, let $R(u)$ designate the homogeneous polynomial of the lowest

degree terms in the power series expansion of $\Psi(0)F(H(\Phi u))$. If the zero solution of the ordinary differential equation

$$(4.18) \quad \dot{u} = R(u)$$

is asymptotically stable, then the zero solution of (2.3) is asymptotically stable. If there is a homogeneous polynomial which is positive on some set and

$$S(u) = -[\partial A(u)/\partial u]R(u)$$

is negative definite, then the zero solution of (2.3) is unstable.

Proof: Let the degree of $R(u)$ be m . From (4.10), $\Psi(0)\hat{F}(u,0) = F(H(\Phi u + \alpha(u)))$. Furthermore, from Lemma 4.2, the power series expansions of $\alpha(u)$ begins with terms of at least degree 2. This implies that

$$F(H(\Phi u + \alpha(u))) = R(u) + T(u)$$

where the power series expansion of $T(u)$ begins with terms of at least degree $m + 1$. Theorem 4.2 now gives the result.

Corollary 4.1 includes Theorem 3.1 in [5] for retarded functional differential equations.

Corollary 4.2. Suppose the hypotheses of Theorem 4.1 are satisfied

and zero is a simple root of (2.7) and equation (4.17) is

$$(4.19) \quad \dot{u} = au^m, \quad a \neq 0.$$

If $a < 0$ and m is odd, the solution $x = 0$ of (2.3) is asymptotically stable. Otherwise, the solution $x = 0$ of (2.3) is unstable.

Proof: If $a < 0$ and m is odd, $A(u) = u^2/2$, then

$$\dot{A}_{(4.19)}(u) = au^{m+1}$$

is a negative definite function and the solution $u = 0$ of (4.19) is asymptotically stable. Theorem 4.2 implies the solution $x = 0$ of (2.3) is asymptotically stable.

If $a > 0$, m is odd and $A(u) = u^2/2$, then $\dot{A}_{(4.19)} = -B(u)$ is positive definite. Theorem 4.2 implies the solution $x = 0$ of (2.3) is unstable. If m is even and $A(u) = (\text{sgn } a)u$, then $\dot{A}_{(4.19)}(u) = -B(u) = |a|u^m$ is positive definite. Theorem 4.2 implies the solution $x = 0$ of (2.3) is unstable. The proof is complete.

As an example, let us consider the two dimensional system $[x = \text{col } (x_1, x_2)]$

$$(4.20) \quad \begin{aligned} \frac{d}{dt} x_1(t) &= x_2(t) \\ \frac{d}{dt} [x_2(t) - g(x(t-r))] &= \alpha x_2(t-r) + f(x(t), x(t-r)) \end{aligned}$$

where $r > 0$, f, g are analytic in their arguments in a neighborhood of zero and the power series expansions begin with terms of degree ≥ 2 . The associated linear equation is the RFDE

$$(4.21) \quad \begin{aligned} \frac{d}{dt} x_1(t) &= x_2(t) \\ \frac{d}{dt} x_2(t) &= \alpha x_2(t-r) \end{aligned}$$

which has $a_D = -\infty$ and a characteristic equation given by

$$(4.22) \quad \lambda(\lambda - \alpha e^{-\lambda r}) = 0.$$

For $-\pi/2r < \alpha < 0$, equation (4.22) has a simple zero root and all other roots have negative real parts. The bases for the constant solutions of (4.21) and its adjoint may be taken as Φ, Ψ , respectively, with

$$\Phi(0) = \text{col}(\alpha^{-1}, 0), \quad \Psi(0) = (\alpha, -1).$$

Suppose C is decomposed by $\{0\}$ as $P \oplus Q$ where P is the one dimensional subspace spanned by Φ . The function H in Theorem 4.1

is given by

$$(4.23) \quad \begin{aligned} H(\psi)(\theta) &= \psi(\theta), \quad -r \leq \theta < 0 \\ &\psi(0) + G(\psi(-r)), \quad \theta = 0, \end{aligned}$$

where $G = \text{col}(0, g)$. If $x_t = H(z_t)$, $z_t = \phi u(t) + w_t$, then the equations for $u(t)$, w_t are

$$(4.24) \quad \begin{aligned} \text{a) } \dot{u}(t) &= -f(z(t) + G(z(t-r)), z(t-r)) \\ \text{b) } w_t &= T(t)w_0 + \int_0^t T(t-s)X_0^Q f(z(s) + G(z(s-r)), z(s-r)) ds. \end{aligned}$$

For $w_t = 0$, the right hand side of (4.24a) becomes

$$(4.25) \quad \begin{aligned} -f(\phi u + G(\phi u), \phi u) &= -f(\alpha^{-1}u, g(\alpha^{-1}u, 0), \alpha^{-1}u, 0) \\ &\stackrel{\text{def}}{=} au^m + bu^{m+1} + \dots \end{aligned}$$

If $\alpha(u)$ is the solution of (4.4) for this particular case, then the fact that $\alpha(u)$ begins with second order terms in u implies that

$$-f(\phi u + \alpha(u) + G(\phi u + \alpha(u)), \phi u + \alpha(u)) = au^m + cu^{m+1} + \dots ;$$

that is, the lowest order terms in the expansion of $-f$ are not affected by $\alpha(u)$.

Therefore, an application of Corollary 4.1 implies the following result. If $a < 0$ and m is odd, the solution $x = 0$ of (4.20) is uniformly asymptotically stable. Otherwise, for any $a \neq 0$, the solution $x = 0$ of (4.21) is unstable.

As a particular illustration, if

$$f(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = b_1 \alpha_1^3 + b_2 \alpha_1 \alpha_2 + b_3 \alpha_3^3$$

$$g(\beta_1, \beta_2) = c_1 \beta_1^2$$

then m, a in (4.25) are given by

$$m = 3, a = (-\alpha)^3 (b_1 + b_2 c_1 + b_3).$$

Since $\alpha < 0$, it follows that the solution of the equation is asymptotically stable if $b_1 + b_2 c_1 + b_3 < 0$ and unstable if $b_1 + b_2 c_1 + b_3 > 0$.

References

1. Hale, J.K. and M.A. Cruz, Asymptotic behavior of neutral functional differential equations. Arch. Rat. Mech. Ana. 34(1969), 331-353.
2. Lefschetz, S., The critical case in differential equations. Bol. Soc. Mat. Mex., 1961, 5-18.
3. Pliss, V.A., On conditional stability in critical cases. Dokl. Akad. Nauk SSSR 147(1962), 1292-1293.
4. Shimanov, S.N., On stability in the critical case of a zero root with time lag. Prikl. Mat. Meh. 24(1960), 447-457.
5. Prokopenv, V.P. and S.N. Shimanov, On stability in the critical case of two zero roots for systems with lags. Differentialniye Uraveniya 2(1966), 453-462.
6. Hale J.K., Ordinary Differential Equations, Wiley Interscience, 1969.
7. Hale, J.K. and M.A. Cruz, Existence, uniqueness and continuous dependence for hereditary systems. Annali di Mat. Pura Appl. 1970.
8. Hale, J.K., Forward and backward continuation for neutral functional differential equations. J. Differential Eqns. 8(1971).
9. Hale, J.K. and K.R. Meyer, A class of functional equations of neutral type. Mem. Amer. Math. Soc. 76(1967).

10. Cruz, M.A. and J.K. Hale, Exponential estimates and the saddle point property for neutral functional differential equations. J. Math. Ana. Appl., 1970.
11. Chafee, N., The bifurcation of one or more closed orbits from an equilibrium point of an autonomous differential system. J. Differential Eqns. 4(1968), 661-679.
12. Cruz, M.A. and J.K. Hale, Stability of functional differential equations of neutral type. J. Differential Eqns. 7(1970).
13. Zubov, V.I., Methods of A.M. Lyapunov and Their Application, Noordhoff, Groningen, 1964.