

CASE FILE
COPY

THE HYDROLYSIS OF PROTEINS¹

Don Roach² and Charles W. Gehrke³

*University of Missouri
Columbia, Missouri 65201

INTRODUCTION

The particular method used for the hydrolysis of proteins prior to an amino acid analysis is of considerable importance since some amino acids are preferentially destroyed and the hydrolysis of others is incomplete. In view of the high precision attained in the gas-liquid chromatographic analysis of amino acid mixtures, the nature of the hydrolytic conditions plays an increasingly important role and can be easily evaluated. The speed, precision, and accuracy of the gas-liquid chromatographic methods developed by Gehrke et al. (1, 2, 3, 4, 5, 6, 7) make possible a thorough investigation of the various parameters involved in the quantitative hydrolysis of different proteins and their compositional characterization.

¹Contribution from the Missouri Agricultural Experiment Station, Journal Series No. 5993. Approved by the Director. Supported in part by grants from the National Aeronautics and Space Administration (NGR 26-004-011), the National Science Foundation (GB 7182), and the United States Department of Agriculture (12-14-100-9468 (34)).

²Experimental data taken in part from doctoral research, University of Missouri, William Henry Hatch predoctoral research fellow. Now Associate Professor at Miami Dade Jr. College, Miami, Florida.

³Professor. Experiment Station Chemical Laboratories.

A hydrolysis reagent of broad specificity is required to break all of the possible peptide bonds which are found in natural products of varying complexity. The particular reagent used must be capable of cleaving all peptide bonds in a protein. Further, the peptide bonds must be accessible to the hydrolytic agent, however, two features of protein structure present difficulties in this respect. First, there is steric hindrance due to the bulky side chains of the aliphatic amino acids, and second, the macromolecular structure, i.e. that due to secondary and tertiary bonding of the protein, prevent complete hydrolysis. The degree to which a protein molecule can unfold is limited by its secondary and tertiary structure, therefore, the hydrolysis reagent may react rapidly on one part of a protein molecule and slowly on another. This is evidenced by the number of different hydrolysis methods that are reported.

Differences in the stability of the various functional groups of amino acids necessitate a compromise among several experimental conditions in order to achieve the optimum hydrolysis of the protein. Moore and Stein (8) reported that the best "all around" hydrolysis can be achieved by reaction for 24 hours with 6 N HCl at 110⁰C under conditions rigorously excluding oxygen, non protein substances, and metals. Oxygen can be excluded by using a sealed tube hydrolysis technique. Generally, acid hydrolysis will yield over 95% recovery for aspartic and glutamic acids, proline, glycine, alanine, methionine,

leucine, tyrosine, phenylalanine, lysine, histidine, and arginine. However, tryptophan is completely destroyed, whereas 5 to 15% of threonine and serine are destroyed. Extrapolation to "zero-time" of hydrolysis can be done, but requires several different times of hydrolysis for each sample. GLC makes studies of "zero-time" hydrolysis practical. The peptide bonds of valine, isoleucine, and leucine are quite stable and thus a longer hydrolysis time is required to obtain maximum yield for these amino acids. Whitfield (9) has studied this problem and explained it in terms of steric factors. Extending the hydrolysis time to 70 hours gives maximum yields (8) for these three amino acids. This, of course, results in lower yields for the other amino acids as compared to a 24-hour hydrolysis time. As yet, no satisfactory method has been found for tryptophan, except alkaline, or enzymatic hydrolysis (8).

The purpose of this research was to study the rates and yields of protein hydrolysis and to determine the optimum reaction conditions which would give maximum yields of all 20 of the protein amino acids in the shortest possible time, using ribonuclease as a representative protein.

LITERATURE REVIEW

Braconnot, in 1820, (10) first used sulfuric acid for the hydrolysis of a protein. The use of hydrochloric acid as a hydrolytic agent was introduced by Bopp (11) in 1849. The hydrolytic agent commonly used today is hydrochloric acid since the rate of peptide bond cleavage is increased in hydrochloric acid over what it would be in sulfuric acid

of equal concentration. An added advantage of HCl is that it can easily be removed from an amino acid mixture by evaporation. Protein samples are usually hydrolyzed with 2.5 to 5000 times their weight of 6 N HCl under reflux for 18 to 24 hours.

The method of Macpherson (12) is generally recommended for large protein samples (ca. 0.2 g or larger). A protein sample which has been equilibrated under atmospheric conditions is weighed into a suitable round-bottomed flask which is fitted with a condenser. Concentrated Analytical Reagent HCl (36 w/w%) is added (ca. 20 ml/g protein), the protein is dissolved on a water bath at 35 to 40°C, then sufficient hot double distilled water is added to bring the concentration of HCl to 20 w/w%. The solution is boiled gently under reflux for 24 hours. The excess of HCl is removed under a partial vacuum and the sample is diluted to a suitable volume with 0.1 N HCl. An aliquot of this solution is then removed for classical amino acid analysis or GLC amino acid analysis.

The method of Moore and Stein is in common use. A sample of air-dried or lyophilized protein is placed in a 16 x 135 mm heavy-walled Pyrex tube (Corning No. 9860). The protein is suspended in 1 ml of 6 N HCl (a 1:1 v/v dilution of concentrated reagent HCl with double distilled H₂O). The sample is frozen by placing in a bath of ethanol and solid carbon dioxide. After freezing, the sample container is evacuated to below 50 microns then sealed under vacuum. The hydrolysis is conducted at 110° ± 1°C for 20 hours or 70 hours. Excess HCl is removed under vacuum at 40 to 45°C, the sample is diluted to a known volume, and aliquots are

removed for analysis. This technique or some modification of it is presently the preferred method for the hydrolysis of protein samples.

A serious problem associated with the acidic hydrolysis of proteins is the partial decomposition of some of the amino acids. The destruction of tryptophan is almost complete and a considerable loss of cysteine may occur. The breakdown of the other amino acids generally occurs to a lesser degree.

Rees (13) reported in 1946 that hydrolysis with 6 N HCl for 24 hours leads to a recovery of only 89.5% of serine and 94.7% for threonine. Corfield and Robson (14) reported a 14% loss of serine in the hydrolysis of salmine. Rees (13) and Hirs et al. (15) found the rate of decomposition of serine and threonine to vary with the purity of the HCl used in the acidic hydrolysis. However, an accurate determination of the threonine and serine content can be made by extrapolation to "zero-time" of hydrolysis if data are available for several different hydrolysis times. Examples of this technique were included in publications by Harfenist (16) in 1953, by Smith and Stockell (17) in 1954, by Hirs et al. (15) in 1954, and by Noltmann et al. (18) in 1962.

There is a possibility that proline is degraded during acid hydrolysis. Elliot et al. (19) and Zuber and Jagues (20) both suggested an empirical formula of $\text{Arg}_2\text{Phe}_2\text{Pro}_2\text{Gly.Ser}$ for the peptide, bradykinin, from results based on amino acid analyses after acidic hydrolysis. Boissonmas et al. (21), however synthesized bradykinin and found that the actual

structure corresponded to the formula $\text{Arg}_2\text{Phe}_2\text{Pro}_3\text{Gly.Ser}$. The variance between the formula determined from amino acid analysis and the actual formula may be due to the decomposition of proline during the acid hydrolysis prior to analysis.

Lugg (22) observed that pure tyrosine was not affected by heating it in acid at 100°C for 20 to 30 hours. Light and Smith (23), however, reported that tyrosine was completely destroyed during the acid hydrolysis of the peptide, Ala.Val.Gly.Tyr . Shepherd et al. (24) also obtained low recoveries of tyrosine from several peptides. This destruction was reduced but not eliminated when the samples were hydrolyzed under a N_2 atmosphere. The decomposition of tyrosine may involve aspartic acid since tyrosine was quantitatively recovered from the peptide Val.Tyr.Pro but not from Val.Tyr.Pro.Asp . Munier (25) reported that tyrosine may be converted to 3-chlorotyrosine during hydrolysis by reacting with traces of chlorine in the HCl . This reaction could not, however, account entirely for the losses observed by Hirs et al. (15).

A large concentration of carbohydrates in the hydrolysis medium may seriously reduce certain amino acids. Tristran (26) noted that arginine was extensively destroyed during acidic hydrolysis in the presence of carbohydrates with the amount of destruction being proportional to the concentration of carbohydrates, and Bailey (27) reported losses of methionine as high as 20% in samples which were high in carbohydrate content. Osono et al. (28) found that refluxing methionine with 10% HCl

resulted in the production of some homocystine, homocysteine, and glycine. Lugg (29) observed only a slight loss of cystine during acidic hydrolysis in the absence of carbohydrate, however, losses of 6 to 7% were noted in the presence of carbohydrates.

Lysine is considered to be the most stable of the diamino acids, but Ishii (30) reported a loss of 3% when lysine was heated at reflux with 20% HCl. The reported degradation products were aspartic acid, glycine, glutamic acid, and α -amino adipic acid.

Steric hindrance by bulky side chain residues results in the slow release of some amino acids, particularly valine and isoleucine. Kinetic studies, by Synge (31) in 1945, and by Harris et al. (32) in 1956, clearly indicated hindrance by valine, leucine, alanine, and isoleucine and the yields for these amino acids which have been hydrolyzed for varying lengths of time were found to be a function of time. An accurate value for each of these amino acids can be determined by plotting yield as a function of hydrolysis time and extrapolating to infinite time, or by drawing tangents to the maximums in the curve, then extrapolating to "zero-time".

The rates of decomposition of the amino acids during acidic hydrolysis are dependent on several factors including: the concentration of the hydrolyzing acid, the purity of the acid used, the time and temperature of hydrolysis, the presence of carbohydrates, aldehydes or metal impurities. Current methods represent a compromise among the several considerations mentioned above. The most common methods

for the hydrolysis of proteins are outlined in two excellent review articles by Light and Smith (33) and by Moore and Stein (8).

Because of the considerable time required for the hydrolysis of proteins by the reported methods and the losses involved, it was considered highly important to investigate other procedures. From studies directed toward this goal, this research reports on a rapid hydrolysis method which can be completed in 4 hours.

EXPERIMENTAL

Apparatus

A Varian Aerograph Model 2100 gas chromatograph with a four-column oven bath, four flame ionization detectors, two dual differential electrometers, and equipped with a Varian Model 20 recorder, and a Packard Instruments Co. Model 7300 dual column gas chromatograph with hydrogen flame detectors and equipped with a Honeywell Electronik 16 strip chart recorder were used. Peak areas were determined with a digital readout integrator, Infotronics Model CRS 104.

Solvents were removed from the samples with a CaLab rotary evaporator "cold finger" condenser, and a Welch Duo-Seal vacuum pump.

Pyrex 2.5 x 20 cm screw top culture tubes with teflon lined caps (Corning No. 9826) were used as the reaction vessels for the hydrolysis.

Reagents

All reagents used were those specified by Gehrke et al. (2).

The 6 N HCl solution in water was prepared by distilling a 6 N solution in an all-glass system and then adjusting the concentration of the constant boiling HCl to 6 N by the addition of double-distilled water.

The 6 N HCl solution in n-butanol was prepared by bubbling anhydrous HCl into anhydrous n-butanol.

Ribonuclease A from bovine pancreas, crystallized 5X, Type I-A, protease free, essentially salt free, activity of 70 Kunitz units per mg, was obtained from Sigma Chemical Co., St. Louis, Missouri.

Chromatography

The columns and chromatographic conditions were those as specified by Gehrke et al. (2).

HYDROLYSIS METHODS

A. Hydrolysis of samples in 6 N HCl in H₂O for GLC analysis.

1. Accurately weigh 10.0 mg of dry protein (ribonuclease) into the pyrex screw top with teflon lined cap culture tube.
2. Flush tube with filtered N₂ gas.
3. Add 10.0 ml of 6 N HCl in H₂O to each tube.
4. Flush each tube again with N₂ gas.

5. Place teflon lined cap on each tube and heat at $110^{\circ} \pm 1^{\circ}\text{C}$, or $145^{\circ} \pm 2^{\circ}\text{C}$ in an oil bath for the specified time.
6. Dry the samples at 60°C under a partial vacuum with a rotary evaporator.
7. Add an appropriate aliquot (1.0 ml of I.S., 0.50 mg butyl stearate/1.0 ml).
8. Add 1.5 ml of n-butanol.3 N in HCl per 1.0 mg of total amino acids, ultrasonic mix for 15 seconds, esterify at 100°C for 35 minutes, then dry at 60°C under partial vacuum, and acylate as described by Gehrke et al. (2).

B. Hydrolysis of Samples in 6 N HCl in H₂O for Analysis by Both GLC and Classical Ion-Exchange.

1. Accurately weigh 25.0 mg of dry protein (ribonuclease) into a large culture tube.
2. Flush tube with filtered N₂ gas.
3. Add 25.0 ml of 6 N HCl in H₂O to each tube.
4. Flush each tube again with N₂ gas.
5. Place teflon lined cap on each tube and heat at $110^{\circ} \pm 1^{\circ}\text{C}$, or $145^{\circ} \pm 2^{\circ}\text{C}$ in an oil bath for the specified time.
6. Dry the samples at 60°C under a partial vacuum with a rotary evaporator.
7. Accurately pipet 20.0 ml of 0.1 N HCl into each of the samples to dissolve the amino acid residue.
Mix each sample thoroughly.

8. Draw a 5.0 ml aliquot of each sample and place in a 125 ml flat-bottom boiling flask for GLC analysis, or analyze by classical ion-exchange.
9. Dry the samples at 60°C under a partial vacuum with a rotary evaporator.
10. Add an appropriate aliquot (1.0 ml of I.S., 0.50 mg of butyl stearate/1.0 ml).
11. Add 1.5 ml BuOH·3 N in HCl per 1.0 mg of total amino acids, ultrasonic mix for 15 seconds, esterify at 100°C for 35 minutes, then dry at 60°C under a partial vacuum, and acylate as described by Gehrke et al. (2).

C. Hydrolysis of Samples by 6 N HCl in n-Butanol.

1. Accurately weigh 10.0 mg of dry protein (ribonuclease) into a large culture tube.
2. Flush tube with filtered N₂ gas.
3. Add 15.0 ml of n-butanol·6 N HCl (1.5 ml of BuOH·HCl per 1.0 mg of protein).
4. Flush reaction vessel again with filtered N₂ gas.
5. Place teflon lined cap on each tube and heat at 110° ± 1°C, or 145° ± 2°C in an oil bath for the specified time.
6. Add an appropriate quantity of I.S., 0.50 mg butyl stearate, dissolved in BuOH·HCl.
7. Dry the samples at 60°C under a partial vacuum with a rotary evaporator.
8. Acylate as described by Gehrke et al. (2).

RESULTS AND DISCUSSION

According to the direct esterification procedure reported by Roach and Gehrke (6) the amino acids are esterified in n-butanolic HCl prior to their acylation with trifluoroacetic anhydride. The present procedure for the GLC analysis of a protein sample requires that the protein sample be hydrolyzed in 6 N HCl in H₂O, dried, and then esterified with n-butanol·3 N HCl. If complete hydrolysis of the protein were achieved in n-butanolic HCl, one of the steps in the GLC analysis of a protein hydrolysate could be eliminated since the n-butyl esters of the amino acids would be formed during the hydrolysis of the protein. Thus, studies were made to investigate the yields of hydrolysis of a model protein, ribonuclease, in 6 N HCl in n-butanol.

Samples of ribonuclease were hydrolyzed at 110⁰ ± 1⁰C in n-butanol 6 N in HCl for varying lengths of time. The experimental results from the GLC analysis of this hydrolysate are presented in Table I. A sample of ribonuclease was also hydrolyzed at 110⁰ ± 1⁰C in H₂O 6 N in HCl for varying lengths of time. The GLC data are presented in Table II. Much higher yields were obtained for the samples hydrolyzed in aqueous 6 N HCl. Also, decomposition of all of the amino acids became a serious problem at hydrolysis times longer than 44 hours with n-butanol 6 N in HCl.

Since n-butanol 6 N in HCl was found to be unsuitable for the hydrolysis of proteins, it was decided to investigate the

effect of temperature on the hydrolysis reaction in aqueous 6 N HCl. Samples were hydrolyzed at $110^{\circ} \pm 1^{\circ}\text{C}$, $145^{\circ} \pm 2^{\circ}\text{C}$, and $175^{\circ} \pm 5^{\circ}\text{C}$ for 1 to 10 hours. The optimum hydrolysis temperature in terms of rate of hydrolysis and a minimum of decomposition was found to be at $145^{\circ} \pm 2^{\circ}\text{C}$ for 4 hours. The GLC data in Table III were obtained from the analysis of samples of ribonuclease which had been hydrolyzed at $145^{\circ} \pm 2^{\circ}\text{C}$ for periods of time from 2 to 9 hours. These data indicate that approximately equivalent hydrolyses of a protein can be achieved in 4 hours at $145^{\circ} \pm 2^{\circ}\text{C}$ and in 24 hours at 110°C .

Moore and Stein (8) reported that the best "all around" hydrolysis can be achieved with 6 N HCl for 24 hours at 110°C under conditions rigorously excluding oxygen, non protein substances, and metals. These experimental conditions were selected to give maximum total recovery of the amino acids. Certain of the amino acids, however, undergo serious decomposition and the hydrolysis of others is incomplete. Maximum values for isoleucine, leucine, valine, and phenylalanine can be obtained by drawing a tangent to the maximum of a plot of yield versus hydrolysis time to obtain the maximum values. Also, an extrapolation to "zero-time" can be made to determine more accurate values for those amino acids which undergo serious decomposition, i.e. threonine, serine, and methionine.

Plots of yield versus hydrolysis time at $110^{\circ} \pm 1^{\circ}\text{C}$ for valine, isoleucine, phenylalanine, threonine, serine, and

methionine from ribonuclease are shown in Figures 1 and 2. Similar plots for hydrolysis of ribonuclease at $145^{\circ} \pm 2^{\circ}\text{C}$ are included in Figures 3 and 4. Maximum values for threonine, serine, and methionine were obtained by extrapolating to "zero-time" as shown; whereas, the maximum values for valine, isoleucine, and phenylalanine were obtained by drawing tangents to the maximum in the curves. Maximum values for the protein amino acids from ribonuclease obtained in this way are included in Table IV. The agreement of the maximum values from the 110°C curves with the corresponding values at $145^{\circ} \pm 2^{\circ}\text{C}$ indicates that hydrolysis under both sets of conditions gave essentially the same results. Also, both sets of data are in excellent agreement with the literature values.

After developing the set of "hydrolysis conditions" ($145^{\circ} \pm 2^{\circ}\text{C}$) for 4 hours using ribonuclease as a model protein, several other proteins were then hydrolyzed under the same conditions and analyzed. The data for one of the proteins, bovine serum albumin, analyzed by classical ion-exchange are given in Table V. The good agreement of the data for hydrolysis for 4 hours at $145^{\circ} \pm 2^{\circ}\text{C}$ with 24 hours at 110°C shows that the hydrolysis of these proteins can be conducted equally well under these two sets of conditions.

Recovery data for GLC analysis of mixtures of the protein amino acids which had been heated for varying times at $145^{\circ} \pm 2^{\circ}\text{C}$ are included in Table VI. The recovery for all of the amino acids except arginine was excellent with a hydrolysis time of 2 hours. On hydrolysis for four hours, the losses in some cases were significant. Low recoveries were obtained for proline, threonine, serine, methionine, hydroxyproline, and

TABLE I

AMINO ACID ANALYSIS OF RIBONUCLEASE
AS A FUNCTION OF HYDROLYSIS TIME

Amino Acid	w/w% ^b				
	4 hr.	10 hr.	24 hr.	44 hr.	70 hr.
Alanine	2.26	3.61	6.54	9.14	3.31
Valine	0.81	1.72	3.69	5.27	3.48
Glycine	0.73	0.84	1.59	2.06	1.01
Isoleucine	0.27	0.47	0.73	1.23	1.50
Leucine	0.28	0.58	1.95	2.22	1.47
Proline	1.45	1.87	3.62	3.90	2.75
Threonine	1.34	1.59	2.59	2.20	1.24
Serine	3.53	3.48	3.98	3.29	1.81
Methionine ^c	---	---	---	---	---
Phenylalanine	0.61	0.94	2.52	2.90	2.36
Aspartic Acid	2.93	4.33	8.52	7.00	3.80
Glutamic Acid	3.06	4.31	8.15	7.10	4.56
Tyrosine ^c	---	---	---	---	---
Lysine	1.90	3.06	6.94	5.76	2.17

^aHydrolyzed in a closed tube with *n*-butanol 6 N in HCl for the specified time at 110° + 1°C. EGA column only. *n*-butyl stearate as internal standard. Ribonuclease, Type I-A, Sigma Chemical Co. GLC Analyses.

^bEach value represents an average of two determinations at the macro level (10 mg total protein + 15 ml of reagent).

^cThe peaks obtained for methionine and tyrosine were too small to allow an accurate determination.

TABLE II

AMINO ACID ANALYSIS OF RIBONUCLEASE AS A FUNCTION OF HYDROLYSIS TIME

Amino Acid	HYDROLYSIS TIME (HOURS ^a) , AND W/W%					
	9 1/2	26	35	48	81	116
Alanine	7.06	7.12	7.35	7.46	7.57	7.51
Valine	4.33	6.43	6.90	7.28	7.56	7.52
Glycine	1.60	1.65	1.66	1.66	1.69	1.68
Isoleucine	0.81	1.50	1.69	1.99	2.39	2.59
Leucine	1.83	1.91	1.96	1.99	2.02	2.01
Proline	2.78	3.09	3.12	3.14	3.19	3.17
Threonine	6.91	7.35	7.71	7.72	7.58	7.27
Serine	9.91	10.04	9.96	9.80	9.20	8.32
Methionine	3.47	3.39	3.36	3.44	3.17	2.93
Phenylalanine	2.59	3.05	3.17	3.22	3.25	3.29
Aspartic Acid	13.23	13.39	13.74	13.97	13.96	13.99
Glutamic Acid	11.13	12.08	12.43	12.04	12.49	12.33
Tyrosine	5.74	6.56	6.63	6.59	6.26	6.06
Lysine	8.88	10.15	10.44	10.59	10.61	10.75

^aHydrolyzed in a closed tube with 6 N HCl for the specified time at $110^{\circ} \pm 1^{\circ} \text{C}$. EGA column only. *n*-Butyl stearate as internal standard. Ribonuclease, Type I-A, Sigma Chemical Co. GLC Analyses.

^bEach value represents an average of two determinations at the macro level (10 mg total protein + 15 ml 6 N HCl).


TABLE III

AMINO ACID ANALYSIS OF RIBONUCLEASE AS A FUNCTION OF HYDROLYSIS TIME


Amino Acid	HYDROLYSIS TIME (HOURS ^a), AND W/W% ^b						
	2	4	5	6	7	8 1/2	9
Alanine	7.30	7.28	7.24	7.37	7.40	7.54	7.45
Valine	6.25	7.15	7.16	7.31	7.41	7.54	7.55
Glycine	1.69	1.71	1.66	1.72	1.67	1.71	1.71
Isoleucine	1.40	1.94	2.07	2.23	2.26	2.53	2.55
Leucine	1.99	1.98	1.98	2.04	2.00	2.02	2.07
Proline	3.18	3.19	3.15	3.16	3.14	3.15	3.16
Threonine	7.73	7.62	7.52	7.36	7.11	7.18	7.17
Serine	10.25	9.66	9.40	8.82	8.52	8.15	8.01
Methionine	3.54	3.36	3.29	3.11	3.06	2.77	2.24
Phenylalanine	3.16	3.26	3.22	3.29	3.22	3.42	3.38
Aspartic Acid	13.83	13.60	13.66	13.76	13.60	13.81	13.97
Glutamic Acid	11.79	11.88	11.97	12.00	11.75	12.05	12.29
Tyrosine	6.57	6.80	7.00	7.26	7.29	6.17	6.47
Lysine	10.29	10.37	10.45	10.42	10.10	10.27	10.65

^aHydrolyzed in a closed tube with 6 N HCl for the specified time at $145^{\circ} \pm 2^{\circ}\text{C}$. EGA column only. *n*-Butyl stearate as internal standard. Ribonuclease, Type I-A, Sigma Chemical Co. GLC Analyses.


^bEach value represents an average of two determinations at the macro level (10 mg total protein + 15 ml 6 N HCl).



**YIELDS OF AMINO ACIDS FROM RIBONUCLEASE
AS A FUNCTION OF HYDROLYSIS TIME**




**YIELDS OF AMINO ACIDS FROM RIBONUCLEASE
AS A FUNCTION OF HYDROLYSIS TIME**



**YIELDS OF AMINO ACIDS FROM RIBONUCLEASE
AS A FUNCTION OF HYDROLYSIS TIME**

145° ± 2°C
6N HCl



**YIELDS OF AMINO ACIDS FROM RIBONUCLEASE
AS A FUNCTION OF HYDROLYSIS TIME**

TABLE IV
SELECTED AMINO ACIDS FROM THE
AMINO ACID ANALYSIS OF RIBONUCLEASE

Amino Acid	W/W%, AND HYDROLYSIS TEMPERATURE		
	110 ⁰ C-24 hrs.	145 ⁰ C-4 hrs.	Literature Value ^a
Valine	7.54 ^b	7.55 ^b	7.49
Isoleucine	2.54 ^b	2.59 ^b	2.67
Threonine	8.00 ^c	8.00 ^c	8.90
Serine	10.70 ^c	11.05 ^c	11.40
Methionine	3.54 ^c	3.83 ^c	4.00
Phenylalanine	3.30 ^b	3.40 ^b	3.51

^aC. H. W. Hirs, W. H. Stein, and S. Moore. J. Biol. Chem., 211, 941, (1954).

^bValues obtained by drawing a tangent to the maximum in a plot of yield of amino acid versus time of hydrolysis to obtain the maximum values.

^cValues obtained by extrapolation to zero time a plot of yield of amino acid versus time of hydrolysis.

TABLE V

AMINO ACID ANALYSIS OF BOVINE SERUM ALBUMIN AS A FUNCTION OF HYDROLYSIS TIME

Amino Acid	HYDROLYSIS TIME (HOURS ^a) AND W/W%					
	2	4	6	8	9	24 ^b
Alanine	4.95	5.00	5.14	4.83	4.91	4.76
Valine	4.15	4.46	5.10	4.79	4.89	4.56
Glycine	2.05	2.25	2.11	2.02	1.97	1.97
Isoleucine	1.81	2.12	2.27	2.16	2.22	2.04
Leucine	11.45	11.91	12.30	11.91	11.70	11.42
Proline	4.72	4.23	4.34	4.73	3.84	3.76
Threonine	4.97	5.07	5.05	4.91	4.93	4.97
Serine	3.64	3.73	3.47	3.08	3.40	3.55
Methionine	0.74	0.81	0.68	0.72	0.72	0.70
Phenylalanine	6.50	6.88	6.83	6.53	6.45	6.59
Aspartic Acid	9.43	9.49	9.62	9.42	9.18	9.02
Glutamic Acid	15.82	15.79	16.13	15.91	15.56	15.49
Tyrosine	5.57	5.92	5.34	5.38	5.15	5.53
Lysine	11.51	11.88	12.01	11.73	11.76	11.56
Histidine	4.00	4.43	4.28	4.16	4.06	4.09
Arginine	5.84	6.10	6.00	5.73	5.87	5.83
Tryptophan	---	---	---	---	---	---
Half-Cystine	5.93	6.08	5.84	5.96	5.51	5.86

^aHydrolyzed with 6 N HCl for the specified time at $145^{\circ} \pm 2^{\circ}$ C in a closed tube, with norleucine as internal standard. Analyzed by classical ion-exchange.

^bHydrolyzed with 6 N HCl for 24 hours at $110^{\circ} \pm 1^{\circ}$ C in a closed tube, norleucine as internal standard. Analyzed by classical ion-exchange.

TABLE VI
RECOVERY OF AMINO ACIDS FROM A STANDARD MIXTURE^a

Amino Acid	HYDROLYSIS TIME (HOURS) , AND RECOVERY, %					
	2	4	6	7	8	9
Alanine	100.6	101.8	96.9	95.6	95.5	93.1
Valine	101.2	100.9	94.1	92.9	91.8	89.4
Glycine	100.6	99.7	95.5	92.9	90.3	87.7
Isoleucine	101.8	102.9	95.8	92.3	89.9	87.6
Leucine	101.2	98.8	94.9	92.8	91.5	89.2
Proline	99.4	94.6	89.9	88.7	89.9	88.9
Threonine	98.6	93.2	90.4	86.3	84.9	84.6
Serine	99.3	90.5	87.6	84.7	81.8	78.8
Methionine	98.1	90.6	79.2	73.6	71.7	75.5
Hydroxyproline	97.4	94.7	89.5	85.5	84.2	83.8
Phenylalanine	102.3	97.7	95.5	94.1	93.2	90.9
Aspartic Acid	101.9	98.9	97.6	96.3	95.5	95.4
Glutamic Acid	101.6	100.7	100.6	99.7	99.4	97.2
Tyrosine	102.6	100.0	98.7	94.2	93.5	90.9
Lysine	100.0	97.4	94.8	93.4	89.5	92.1
Arginine	95.2	88.1	85.7	76.2	77.1	65.2
Histidine	101.1	97.5	97.4	97.3	97.3	88.4
Cystine	102.8	97.1	100.0	85.7	77.1	74.3

^a0.4 mg of each amino acid + 10 ml 6N HCl heated for the specified time at 145⁰ + 2⁰C. Each value an average of two independent analyses. Analyzed by gas-liquid chromatography.

arginine. Plots of yield versus hydrolysis time are required to obtain accurate results for these amino acids.

SUMMARY AND CONCLUSIONS

The use of n-butanol 6 N in HCl as a protein hydrolysis reagent would obviate one of the steps in the reported (2) GLC analysis of proteins since the n-butyl esters of the amino acids would be formed during the hydrolysis. Thus, these studies were initiated to investigate the yields on hydrolysis of a model protein, ribonuclease, in 6 N HCl in n-butanol. However, this reagent was found to be unsatisfactory since the rate of hydrolysis was much slower in this medium than it was in 6 N HCl in water, and the rates of decomposition of the amino acids were considerably faster.

Since n-butanol 6 N in HCl was found to be unsuitable for the hydrolysis of proteins, experiments were made to investigate the effect of temperature on the hydrolysis reaction in aqueous 6 N HCl with a view to developing a rapid hydrolysis procedure. The maximum yield for all of the protein amino acids was obtained at $145^{\circ} \pm 2^{\circ}\text{C}$ for the minimal time of 4 hours. Essentially equivalent hydrolysis of ribonuclease was achieved at the two different hydrolysis conditions, i.e. $110^{\circ} \pm 1^{\circ}\text{C}$ for 26 hours, or $145^{\circ} \pm 2^{\circ}\text{C}$ for 4 hours. The yields obtained were in good agreement.

Plots of yield versus hydrolysis time must be made for each amino acid to obtain the best possible values for the amino acid composition of a protein. These plots are then

extrapolated to "zero-time" for those amino acids which undergo decomposition; and tangents are drawn to the maximum in the curves to obtain the maximum values for those amino acids which are difficult to hydrolyze. Several protein samples must be hydrolyzed at different times to obtain all the data necessary to construct these plots. Therefore, a rapid hydrolysis method is needed which gives maximum values for the amino acids with a minimum of decomposition. Recovery studies as a function of hydrolysis time at 145°C were made using standard amino acid mixtures; essentially complete recovery was obtained on hydrolyzing for 2 to 4 hours. Pro., Thr., Ser., Met., and Arg. were the most sensitive to heat. This reported procedure allows one to rapidly hydrolyze several samples of a protein at $145^{\circ} \pm 2^{\circ}\text{C}$ and then to conveniently obtain the data by GLC for plotting these curves. The agreement among the data for the two different hydrolysis temperatures and times conclusively demonstrates that such plots can be made. Of special interest is a comparison of the data obtained for ribonuclease and bovine serum albumin for the 4 hour hydrolysis at $145^{\circ} \pm 2^{\circ}\text{C}$ with the 24 hour hydrolysis at $110^{\circ} \pm 1^{\circ}\text{C}$. In almost every case a higher recovery was obtained for the $145^{\circ} \pm 2^{\circ}\text{C}$ hydrolysis.

With this method a protein can be essentially completely hydrolyzed in 4 hours with a minimum of decomposition of the amino acids. Rapid hydrolysis of proteins coupled with quantitative gas-liquid chromatographic analysis of amino

acids provides a powerful tool in protein research, biochemical, and nutritional investigations.

REFERENCES

1. Gehrke, C. W., Lamkin, W. M., Stalling, D. L., and Shahrokhi, F., Biochem. Biophys. Res. Commun., 13, 328, (1965).
2. Gehrke, C. W., Roach, D., Zumwalt, R. W., Stalling, D. L., and Wall, L. L., Quantitative Gas-Liquid Chromatography of Amino Acids in Proteins and Biological Substances, Analytical Biochemistry Laboratories, Inc., P.O. Box 1097, Columbia, Mo., 65201 (1968).
3. Gehrke, C. W., and Shahrokhi, F., Anal. Biochem., 15 97 (1966).
4. Gehrke, C. W., and Stalling, D. L., Sep. Sci., 2, 101 (1967).
5. Gehrke, C. W., Zumwalt, R. W., and Wall, L. L., J. of Chromatog., 37, 398 (1968).
6. Roach, D., and Gehrke, C. W., J. Chromatog., 43, 303, (1969).
7. Roach, D., Gehrke, C. W., and Zumwalt, R. W., J. Chromatog., 43, 311, (1969).
8. Moore, S., and Stein, W. H., Methods in Enzymology, S. P. Colwick and N. O. Kaplan, eds., Academic Press, New York (1963) Vol. VI, p. 819.
9. Whitfield, A., Science, 142, 577, (1963).
10. Braconnot, H., Ann. Chim. Phys., 13, 113, (1820).
11. Bopp, F., Ann., 69, 16, (1849).
12. Macpherson, H. T., Biochem. J., 40, 470 (1946).
13. Rees, M. W., Biochem. J. (London), 40, 632, (1946).
14. Corfield, M. C., and Robson, A., Biochem. J. (London), 55, 517, (1953).
15. Hirs, C. H. W., Stein, W. H., and Moore, S., J. Biol. Chem., 211, 941 (1954).
16. Harfenist, E. L., J. Am. Chem. Soc., 75, 5528 (1953).
17. Smith, E. L., and Stockell, A., J. Biol. Chem., 207, 501 (1954).

18. Noltmann, E. A., Mahowald, T. A., and Kuby, S. A., J. Biol. Chem., 237, 1146 (1962).
19. Elliot, D. F., Lewis, G. P., and Horton, E. W., Biochem. J., 74, 15P (1960).
20. Zuber, H., and Jaques, R., Helv. Chim. Acta, 43, 1128 (1960).
21. Boissonnas, R. A., Guttman, St., and Jaquenoud, P. A., Helv. Chim. Acta, 43, 1349 (1960).
22. Lugg, J. W. H., Biochem. J. (London), 32, 775 (1938).
23. Light, A., and Smith, E. L., J. Biol. Chem., 237, 2537 (1962).
24. Shepherd, R. G., Willson, S. D., Howard, K. S., Davies, D. S., Davies, S. B., Eigner, E. A., and Shakespeare, N. E., J. Am. Chem. Soc., 78, 5067 (1958).
25. Munier, R. J., J. Chromatog., 1, 524 (1958).
26. Tristran, G. R., Biochem. J. (London), 33, 1271 (1939).
27. Bailey, K., Biochem. J. (London), 31, 1396 (1937).
28. Osono, K., Mukai, I., and Tominaga, F., Met. Nagasaki Igakkai Zasshi., 30, 156 (1955).
29. Lugg, J. W. H., Biochem. J. (London), 27, 1022 (1933).
30. Ishii, K., Nagasaki Igakkai Zasshi., 29, 218 (1954) from the Chem. Abs., 48, 8837 (1954).
31. Synge, R. L. M., Biochem. J., 39, 351 (1945).
32. Harris, J. I., Cole, R. D., and Pon, N. G., Biochem. J., 62, 154 (1956).
33. Light, A., and Smith, E. L., The Proteins, 2nd ed., H. Neurath, ed., Academic Press, New York (1963) Vol. I, p. 32.