


## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

29

CR0-3662-3



# A Facsimile Report

Reproduced by  
**UNITED STATES**  
**ATOMIC ENERGY COMMISSION**  
Division of Technical Information  
P.O. Box 62 Oak Ridge, Tennessee 37830

FACILITY FORM 602

|                               |            |
|-------------------------------|------------|
| N71-13500                     |            |
| (ACCESSION NUMBER)            |            |
| 55                            | (THRU)     |
| (PAGES)                       | G3         |
| CR-115810                     | (CODE)     |
| (NASA CR OR TMX OR AD NUMBER) | 08         |
|                               | (CATEGORY) |

505-63935R



Technical Report 69-91  
AT-(40-1)-3662

May, 1969  
ORO-3662-3

Noise cleaning in binary-valued  
digital pictures  
using "propagation" processes

Aziel Rosenfeld\*, Chan M. Park\*,  
and James P. Strong\*\*

Abstract

Methods of cleaning noise in a binary-valued digital picture by using "propagation" processes are described, with emphasis on the possibility of preserving elongated parts of objects while deleting small non-elongated parts.

The support of the U. S. Atomic Energy Commission, under Contract AT-(40-1)-3662 with the Computer Science Center of the University of Maryland, is gratefully acknowledged.

- \* Computer Science Center, University of Maryland
- \*\* Goddard Space Flight Center, National Aeronautics and Space Administration

Table of Contents

|                                                |    |
|------------------------------------------------|----|
| 1. Introduction                                | 1  |
| 2. Neighbor-counting methods and modifications | 2  |
| 3. Elongation analysis                         | 7  |
| 4. Isolatedness                                | 13 |



Technical Report 69-91  
AT-(40-1)-3662

May, 1969  
ORO-3662-3

Noise cleaning in binary-valued  
digital pictures  
using "propagation" processes

Azriel Rosenfeld\*, Chan M. Park\*,  
and James P. Strong\*\*

#### Abstract

Methods of cleaning noise in a binary-valued digital picture by using "propagation" processes are described, with emphasis on the possibility of preserving elongated parts of objects while deleting small non-elongated parts.

The support of the U. S. Atomic Energy Commission, under Contract AT-(40-1)-3662 with the Computer Science Center of the University of Maryland, is gratefully acknowledged.

- \* Computer Science Center, University of Maryland
- \*\* Goddard Space Flight Center, National Aeronautics and Space Administration

#### Table of Contents

|                                                |    |
|------------------------------------------------|----|
| 1. Introduction                                | 1  |
| 2. Neighbor-counting methods and modifications | 2  |
| 3. Elongation analysis                         | 7  |
| 4. Isolatedness                                | 13 |

# LIST OF FIGURES

| <u>Fig. No.</u> | <u>Caption</u>                                                                                   | <u>Follows page</u> |
|-----------------|--------------------------------------------------------------------------------------------------|---------------------|
| 1               | Test picture                                                                                     | 2                   |
| 2-8             | Results of applying $F_{r,\lambda}$ to Figure 1 for various values of $r$ and $\lambda$          | 3                   |
| 9-15            | Results of applying $G_{r,\lambda}$ to Figure 1 for the same values of $r$ and $\lambda$         | 4                   |
| 16              | Results of applying $H_1$ to Figure 1                                                            | 6                   |
| 17              | Results of one-step contraction of the 1's in Figure 1                                           | 11                  |
|                 | a) Connected components which disappeared                                                        |                     |
|                 | b) Table of their areas                                                                          |                     |
|                 | c) Result of deleting "noise" components                                                         |                     |
| 18              | Results of one-step contraction of the 0's in Figure 1 [(a-c) as in Figure 17]                   | 11                  |
| 19              | Results of two-step contraction and reexpansion of the 1's in Figure 18c                         | 11                  |
|                 | a) Connected components which failed to reappear                                                 |                     |
|                 | b) Table of their areas (divided by 4)                                                           |                     |
|                 | c) Result of deleting "noise" components                                                         |                     |
| 20              | Results of two-step contraction and reexpansion of the 0's in Figure 17c [(a-c) as in Figure 19] | 11                  |
| 21              | Fifteen-step elongatedness analysis of a cloud cover picture                                     | 12                  |
| 22              | Isolatedness analysis of Figure 1                                                                | 13                  |
|                 | a) Components after one-step expansion                                                           |                     |
|                 | b) Components after two-step expansion                                                           |                     |
|                 | c) Table of areas for (a), divided by 9                                                          |                     |
|                 | d) Table of areas for (b), divided by 25                                                         |                     |

## 1. Introduction

One of the earliest papers on picture processing by computer [1] introduced a simple method of "cleaning up" noise in a binary-valued digital picture : 1's are changed to 0's if they have more than a certain number of 0's as neighbors, and vice versa. Since that time, many other methods of noise cleaning have been developed, including refinements of the neighbor-counting method [e.g., 2-3] and methods based on selective spatial frequency filtering [e.g., 4-6]. In addition, many investigators have developed noise cleaning techniques appropriate to particular classes of pictures, and have used them as part of the "preprocessing" for specific pictorial pattern recognition tasks.

The purpose of the present report is to describe an approach to noise cleaning based on applying "propagation" (or "distance"-computing) operations to the given picture. In particular, it is shown how this approach can be used to clean noise while preserving thin, elongated "signals".

# LIST OF FIGURES

| <u>Fig. No.</u> | <u>Caption</u>                                                                                   | <u>Follows page</u> |
|-----------------|--------------------------------------------------------------------------------------------------|---------------------|
| 1               | Test picture                                                                                     | 2                   |
| 2-8             | Results of applying $F_{r,\lambda}$ to Figure 1 for various values of $r$ and $\lambda$          | 3                   |
| 9-15            | Results of applying $G_{r,\lambda}$ to Figure 1 for the same values of $r$ and $\lambda$         | 4                   |
| 16              | Results of applying $H_1$ to Figure 1                                                            | 6                   |
| 17              | Results of one-step contraction of the 1's in Figure 1                                           | 11                  |
|                 | a) Connected components which disappeared                                                        |                     |
|                 | b) Table of their areas                                                                          |                     |
|                 | c) Result of deleting "noise" components                                                         |                     |
| 18              | Results of one-step contraction of the 0's in Figure 1 [(a-c) as in Figure 17]                   | 11                  |
| 19              | Results of two-step contraction and reexpansion of the 1's in Figure 18c                         | 11                  |
|                 | a) Connected components which failed to reappear                                                 |                     |
|                 | b) Table of their areas (divided by 4)                                                           |                     |
|                 | c) Result of deleting "noise" components                                                         |                     |
| 20              | Results of two-step contraction and reexpansion of the 0's in Figure 17c [(a-c) as in Figure 19] | 11                  |
| 21              | Fifteen-step elongatedness analysis of a cloud cover picture                                     | 12                  |
| 22              | Isolatedness analysis of Figure 1                                                                | 13                  |
|                 | a) Components after one-step expansion                                                           |                     |
|                 | b) Components after two-step expansion                                                           |                     |
|                 | c) Table of areas for (a), divided by 9                                                          |                     |
|                 | d) Table of areas for (b), divided by 25                                                         |                     |

## 1. Introduction

One of the earliest papers on picture processing by computer [1] introduced a simple method of "cleaning up" noise in a binary-valued digital picture: 1's are changed to 0's if they have more than a certain number of 0's as neighbors, and vice versa. Since that time, many other methods of noise cleaning have been developed, including refinements of the neighbor-counting method [e.g., 2-3] and methods based on selective spatial frequency filtering [e.g., 4-6]. In addition, many investigators have developed noise cleaning techniques appropriate to particular classes of pictures, and have used them as part of the "preprocessing" for specific pictorial pattern recognition tasks.

The purpose of the present report is to describe an approach to noise cleaning based on applying "propagation" (or "distance"-computing) operations to the given picture. In particular, it is shown how this approach can be used to clean noise while preserving thin, elongated "signals".

## 2. Neighbor-counting methods and modifications


To illustrate the limitations of the classical approach to noise cleaning described in [1], consider the binary-valued digital picture shown in Figure 1. Here it might be reasonable to regard the circles (which look like ellipses because the vertical and horizontal spacing on the printer are unequal) as "signal", and the dots as "noise". the streaks might be regarded as signal for some purposes, as noise for others. In Figure 1a, the probability that an element is a noise dot, or the center of a streak, is 0.025; in Figure 1b, 0.05; in Figure 1c, 0.10; and in Figure 1d, 0.20.

Let  $D_{ij}^{(r)}$  be the set of elements at distance  $< r + \frac{1}{2}$  from  $(i, j)$ ; let  $n_r$  be the number of elements in this set. Let  $F_{r,\lambda}$  (where  $0 \leq \lambda \leq 1$ ) be the operation which takes the binary-valued digital picture  $(a_{ij})$  into the binary-valued digital picture  $(b_{ij})$  defined as follows:

$$b_{ij} = \begin{cases} 0 & \text{if the number of 1's in } D_{ij}^{(r)} \text{ is } \leq \lambda n_r \\ 1 & \text{if this number is } > (1-\lambda)n_r \\ a_{ij} & \text{otherwise} \end{cases}$$

In other words,  $F_{r,\lambda}$  leaves  $(a_{ij})$  unchanged except (possibly) for elements having very many 0's or very many 1's as neighbors; such elements themselves become 0's or 1's, respectively.

Figures 2-8 show the results of applying  $F_{r,\lambda}$  to Figure 1 for the following values of  $r$  and  $\lambda$ :






| Figure | $r$ | $\lambda$ |
|--------|-----|-----------|
| 2      | 1   | 0.2       |
| 3      | 1   | 0.3       |
| 4      | 1   | 0.4       |
| 5      | 2   | 0.1       |
| 6      | 2   | 0.2       |
| 7      | 2   | 0.3       |
| 8      | 2   | 0.4       |

Note that for  $r = 1$ , the set  $D_{ij}^{(r)}$  is a 3-by-3 "square" consisting of  $(i, j)$  and its eight neighbors; while for  $r = 2$ ,  $D_{ij}^{(r)}$  is an "octagon" having 21 elements.

It is seen from these results that except for the lowest values of  $\lambda$ , these operations treat the streaks as noise; however, only for high  $\lambda$  are they eliminated entirely, while for lower  $\lambda$ 's they shrink but do not completely disappear. (Note that for  $r = 2$ , diagonal streaks are treated more harshly than horizontal and vertical streaks, since the set  $D_{ij}^{(r)}$  is five elements high and wide, but only three elements across in the diagonal directions.) Evidently, neighbor-counting methods cannot be effective for high levels of "dot" noise without also treating streaks as noise.

A further difficulty with these operations, as seen in Figures 2-8, is that noise which is close to the edge of the "signal" is usually not eliminated, since when  $(i, j)$  is such a noise element,  $D_{ij}^{(r)}$  contains many "signal" elements. (This problem becomes increasingly troublesome as  $r$  increases.) To eliminate this objection, one can modify the neighbor-counting method along the following lines (compare [7]):








Figure 5




Figure 6




Figure 7




Figure 8

Let  $c_{ij} = 0$  if  $(i, j)$  is in some  $D_{hk}^{(r)}$  in which the number of 1's is  $\leq \lambda n_r$ ;  $c_{ij} = 1$  otherwise

$d_{ij} = 1$  if  $(i, j)$  is in some  $D_{hk}^{(r)}$  in which the number of 1's is  $\leq (1-\lambda)n_r$ ;  $d_{ij} = 0$  otherwise

and let  $G_{r,\lambda}$  be the operation which takes  $(a_{ij})$  into  $(e_{ij})$  defined by

$$\begin{aligned} e_{ij} &= 0 \text{ if } c_{ij} = d_{ij} = 0 \\ e_{ij} &= 1 \text{ if } c_{ij} = d_{ij} = 1 \\ e_{ij} &= a_{ij} \text{ otherwise.} \end{aligned}$$

In other words,  $e_{ij} = 0$  if  $(i, j)$  is in some neighborhood consisting mostly of 0's, and in no neighborhood consisting mostly of 1's;  $e_{ij} = 1$  if the opposite is true; and  $e_{ij} = a_{ij}$  if  $(i, j)$  is in neither type or in both types of neighborhood. Evidently, this operation can, in principle, eliminate noise even if it is close to the edge of the signal. For example, it can turn a "1" into a "0" if some neighborhood containing it consists mostly of 0's, even though the neighborhood centered at it does not.

Figures 9-15 show the results of applying  $G_{r,\lambda}$  to Figure 1, for the same values of  $r$  and  $\lambda$  as in Figures 2-8. It will be noted that the noise cleaning power of  $G_{r,\lambda}$  is generally higher than that of the corresponding  $F_{r,\lambda}$ . (Even the weakest  $G$ 's treat streaks as noise; note, however, that for  $r = 1$ , diagonal streaks are treated more harshly, since a 3-by-3 square can intersect such a streak in only one element, while it must intersect a horizontal






Figure 9






Figure 12




Figure 13




Figure 14




Figure 15

or vertical streak in at least three elements.) Note also that for  $r = 2$  and  $\lambda = 0.4$ ,  $G$  is actually poorer than  $F$ , since for large  $r$  and  $\lambda$  it becomes increasingly easy to have  $c_{ij} = 0$  and  $d_{ij} = 1$  at the same point, so that  $e_{ij} = a_{ij}$  and noise is not cleaned.


We can also consider another approach to noise cleaning in which "propagation" processes are used. Specifically, let  $H_r$  be the operation which changes 0's to 1's if any of their eight neighbors are 1's, and let  $\bar{H}$  similarly change 1's to 0's if any of their eight neighbors are 0's. Let  $(f_{ij}) = H(\bar{H}(r+1)(H(r)(a_{ij})))$ ,  $(g_{ij}) = \bar{H}(H(r+1)(\bar{H}(r)(a_{ij})))$ , and let  $H_r$  be the operation which takes  $a_{ij}$  into  $(h_{ij})$  defined by

$$\begin{aligned} h_{ij} &= 0 \text{ if } f_{ij} = g_{ij} = 0 \\ h_{ij} &= 1 \text{ if } f_{ij} = g_{ij} = 1 \\ h_{ij} &= a_{ij} \text{ otherwise.} \end{aligned}$$

In other words, we obtain  $(f_{ij})$  from  $(a_{ij})$  by "expanding" (or "propagating") the 1's  $r$  times, then "recontracting" them (or equivalently, expanding the 0's)  $r + 1$  times, and finally re-expanding the 1's once. Evidently, an isolated "1" first expands under this treatment into a square of radius  $r$ , but then contracts and vanishes completely. However, if the "1" belongs to a cluster of 1's which "fuses" (see [8]) under the expansion, it will not vanish under re-contraction. Similarly, we obtain  $(g_{ij})$  from  $(a_{ij})$  by first expanding the 0's (or equivalently, contracting the 1's)  $r$  times, then contracting them  $r + 1$  times, and finally re-expanding them once. Clearly an isolated "1" will vanish

immediately under this treatment, so that for such a "1" we have  $h_{ij} = f_{ij} = g_{ij} = 0$ , and analogously for isolated 0's; but in a region where 1's and 0's are mixed together,  $f_{ij}$  and  $g_{ij}$  will usually differ, so that we will have  $h_{ij} = a_{ij}$ . Note that in general, if there are many 1's in a neighborhood of radius  $2r$  of a given "1", expansion of the 1's by  $r$  will cause "fusion" to take place, so that we will have  $f_{ij} = 1$  and the given "1" will not be changed to a "0" by  $H_r$ . Figure 16 shows the result of applying  $H_1$  to Figure 1.

Figure 16



### 3. Elongation analysis

The "propagation" scheme just described, like the neighbor counting schemes, treats both dots and streaks as noise. In this section, a more complex scheme is described, based on propagation, which treats the dots (and eventually, even the large circles) as noise, and the streaks as signal.

To see the basic idea underlying this scheme [8], consider the effect of contracting an object (e.g., the set of 1's in a digital picture) and then re-expanding it by the same amount. It is easily seen that the re-expanded object cannot contain any elements which were not in the original object. On the other hand, it need not contain every element of the original; for example, if an object is contracted by an amount greater than its radius, it disappears completely, so that there is nothing left to re-expand.

Suppose that the object  $S$  is contracted by  $k$  steps (i.e., the operation which turns 1's into 0's if they have 0's as neighbors is applied  $k$  times to the picture) and then re-expanded by  $k$  steps (0's having 1's as neighbors are turned into 1's,  $k$  times). Let  $S_k$  be the result of this contraction and re-expansion (so that  $S_k$  is a subset of  $S$ ), and let  $\bar{S}_k = S - S_k$  be the set of elements of  $S$  which are not in  $S_k$ . Let  $\bar{A}_k$  be a connected component of  $\bar{S}_k$ . Since every element of the subset  $\bar{A}_k$  of  $S$  disappeared under contraction and re-expansion by  $k$  steps,  $\bar{A}_k$  cannot be more than  $2k$  "wide". Suppose that the area of  $\bar{A}_k$  is (say)  $10k^2$  or greater. Since its "width"

is at most  $2k$ , its "length" must thus be at least  $5k$ , i.e.,  $2\frac{1}{2}$  times its "width", making it elongated. Here "length" and "width" can be taken literally if  $\tilde{A}_x$  is a rectangle; but in general too, it is reasonable to call a connected set elongated if its area is more than  $2\frac{1}{2}$  times the square of its greatest "width" (where the "width" of a set at a point might be defined as twice the distance from the point to the set's border).

The criterion just described requires refinement before it can be used as reasonable definition of "elongated". Note, in fact, that the object

1111

disappears under contraction by one step, but since its area is only 4, it would not be called elongated. This is because the criterion was designed on the basis of objects which are elongated in spite of being as wide as possible. Indeed, the object

11  
11

also disappears under contraction by one step, and has area 4, but is certainly not elongated. If an object disappears under one step of contraction, the criterion calls it elongated only if its area is at least 10, since it may be 2 wide, but this criterion is too severe for objects which are only 1 wide. In general, the criterion must be too severe for objects of width  $2k-1$ , if it is not to be too lenient for objects of width  $2k$ ; but as  $k$  increases, this difficulty diminishes.

The objection just raised can be overcome if the definition of the contraction process can be modified so that objects of width  $2k-1$  and  $2k$  require different numbers of contraction steps before they disappear.

In particular, suppose that the process is redefined as follows: on odd-numbered steps, change 1's to 0's if they have 6 or more 0's as neighbors; on even-numbered steps, change 1's to 0's if they have any 0's as neighbors. It is easily seen that a rectangle  $n$  elements wide takes  $n$  steps to disappear under this modified process (rather than  $\frac{n}{2}$  or  $\frac{n+1}{2}$ , depending on whether  $n$  is even or odd, as under the process described originally). If an object is contracted by  $k$  steps using the new procedure, and then re-expanded (by changing 0's to 1's if they have any 1's as neighbors,  $\frac{k}{2}$  times, where  $\frac{k}{2}$  is the largest integer not greater than  $\frac{k}{2}$ ), one can again consider connected components of what fails to reappear, and define a component to be elongated if its area is (say)  $2\frac{1}{2}k^2$  or greater.

A further objection to this class of elongatedness criteria is their sensitivity to noise. Suppose, for example, that a  $(2k+1)$ -by- $(2k+1)$  square of 1's contains a single "0" at its center; then it disappears under contraction by only  $k+1$  steps (rather than  $2k+1$  steps), so that (for  $k \geq 2$ ) it would be called elongated by the criterion just described, which (at least for large  $k$ ) seems unreasonable. On the other hand, if the "hole" in the center of the square is large with respect to  $k$ , it is not unreasonable to call the "square" elongated, since it is really an annulus.

This last objection can be overcome by carrying out a more elaborate propagation procedure, in which elongatedness criteria are applied, alternatingly, to both the 1's and the 0's in the given picture  $P_0$ , and connected components which are not found to be elongated are treated as noise. Specifically, the 1's in  $P_0$  are contracted by one step as described earlier\*, and the small (=non-elongated) connected components which disappear (=fail to reappear) remain as 0's, while the elongated components are changed back to 1's. Conversely, the 0's in  $P_0$  are contracted by one step, and their non-elongated components which disappear become 1's, while their elongated components are changed back to 0's. Let the resulting pictures be  $P_1$  and  $P_1'$ , respectively. At the second stage, the 1's in  $P_1$  are contracted by two steps and re-expanded, and their non-elongated components which fail to reappear remain as 0's, while the elongated components are changed back to 1's; and analogously for the 0's in  $P_1$ . Let the results of this stage be  $P_2$  and  $P_2'$ , respectively, and repeat the process, contracting the 1's in  $P_2$  (the 0's in  $P_2'$ ) by three steps and re-expanding; and so on. In the example of the square with a hole at its center considered in the preceding paragraph, it is easily seen that if the width of the hole is less than  $1/3$  that of the square, the hole will disappear under shrinking and re-expansion of 0's before the square (annulus) disappears under shrinking and re-expansion of 1's; so that

\* Note that for  $k = 1$  we have  $\left\lfloor \frac{k}{2} \right\rfloor = 0$ , so that at this step there is actually no re-expansion.

if the hole itself is not elongated, the square will not be called elongated. More generally, the presence of holes will not make an object elongated unless they are at least as wide as the pieces of object which separate them.

In practice, it turns out that the threshold used to decide whether a component is elongated should not be the same as that used to decide whether a component is "noise". In the examples which follow, a component is allowed to disappear only if its area is  $\leq k^2$  (rather than  $2k \cdot k^2$  as would be required for an elongatedness decision).

Figures 17-18 show the results of one-step contraction and re-expansion of both the 1's and the 0's in Figure 1: (a) Connected components of what failed to reappear, successively labelled 1, 2, ..., 9, A, B, ..., V, 1, 2, ..., etc. (b) Table of the areas of the components (c) Result of deleting "noise" components which failed to reappear. Figures 19-20 show the next stage (two step contraction and re-expansion). If the process were repeated further, little would change until the 9th stage, when the circles themselves would disappear. The results of applying the process to a cloud cover picture, for 15 stages, are shown in Figure 21: (a) Original picture (cloud elements denoted by blanks, noncloud by 1's); (b) Noncloud elements labelled 1, 2, ... according to the stage at which they fail to reappear and are deleted ("x" denotes elements which have not been deleted even at the 15th stage); (c) Noncloud




Figure 17a

| PARAMETER | SYMBOL | ARFA | RATIO |
|-----------|--------|------|-------|
| 2         | 2      | 7    | 7.000 |
| 3         | 3      | 3    | 3.000 |
| 4         | 4      | 3    | 2.000 |
| 5         | 5      | 4    | 4.000 |
| 6         | 6      | 4    | 4.000 |
| 7         | 7      | 1    | 1.000 |
| 8         | 8      | 1    | 1.000 |
| 9         | 9      | 1    | 1.000 |
| 10        | A      | 3    | 3.000 |
| 11        | B      | 3    | 3.000 |
| 12        | C      | 1    | 1.000 |
| 13        | D      | 1    | 1.000 |
| 14        | E      | 1    | 1.000 |
| 15        | F      | 1    | 1.000 |
| 16        | G      | 1    | 1.000 |
| 17        | H      | 1    | 1.000 |
| 18        | I      | 1    | 1.000 |
| 19        | J      | 1    | 1.000 |
| 20        | K      | 5    | 5.000 |
| 21        | L      | 5    | 5.000 |
| 22        | M      | 1    | 1.000 |
| 23        | N      | 1    | 1.000 |
| 24        | O      | 1    | 1.000 |
| 25        | P      | 1    | 1.000 |
| 26        | Q      | 1    | 1.000 |
| 27        | R      | 1    | 1.000 |
| 28        | S      | 1    | 1.000 |
| 29        | T      | 1    | 1.000 |
| 30        | U      | 1    | 1.000 |
| 31        | V      | 1    | 1.000 |
| 32        | W      | 1    | 1.000 |
| 33        | X      | 1    | 1.000 |
| 34        | Y      | 1    | 1.000 |
| 35        | Z      | 1    | 1.000 |
| 36        | 0      | 1    | 1.000 |
| 37        | 1      | 1    | 1.000 |
| 38        | 2      | 1    | 1.000 |
| 39        | 3      | 1    | 1.000 |
| 40        | 4      | 1    | 1.000 |
| 41        | 5      | 1    | 1.000 |
| 42        | 6      | 1    | 1.000 |
| 43        | 7      | 1    | 1.000 |
| 44        | 8      | 1    | 1.000 |
| 45        | 9      | 1    | 1.000 |
| 46        | A      | 1    | 1.000 |
| 47        | B      | 1    | 1.000 |
| 48        | C      | 1    | 1.000 |
| 49        | D      | 1    | 1.000 |
| 50        | E      | 1    | 1.000 |
| 51        | F      | 1    | 1.000 |
| 52        | G      | 1    | 1.000 |
| 53        | H      | 1    | 1.000 |
| 54        | I      | 1    | 1.000 |
| 55        | J      | 1    | 1.000 |
| 56        | K      | 1    | 1.000 |
| 57        | L      | 1    | 1.000 |
| 58        | M      | 1    | 1.000 |
| 59        | N      | 1    | 1.000 |
| 60        | O      | 1    | 1.000 |
| 61        | P      | 1    | 1.000 |
| 62        | Q      | 1    | 1.000 |
| 63        | R      | 1    | 1.000 |
| 64        | S      | 1    | 1.000 |
| 65        | T      | 1    | 1.000 |
| 66        | U      | 1    | 1.000 |
| 67        | V      | 1    | 1.000 |
| 68        | W      | 1    | 1.000 |
| 69        | X      | 1    | 1.000 |
| 70        | Y      | 1    | 1.000 |
| 71        | Z      | 1    | 1.000 |
| 72        | 0      | 1    | 1.000 |
| 73        | 1      | 1    | 1.000 |
| 74        | 2      | 1    | 1.000 |
| 75        | 3      | 1    | 1.000 |
| 76        | 4      | 1    | 1.000 |
| 77        | 5      | 1    | 1.000 |
| 78        | 6      | 1    | 1.000 |
| 79        | 7      | 1    | 1.000 |
| 80        | 8      | 1    | 1.000 |
| 81        | 9      | 1    | 1.000 |
| 82        | A      | 1    | 1.000 |
| 83        | B      | 1    | 1.000 |
| 84        | C      | 1    | 1.000 |
| 85        | D      | 1    | 1.000 |
| 86        | E      | 1    | 1.000 |
| 87        | F      | 1    | 1.000 |
| 88        | G      | 1    | 1.000 |
| 89        | H      | 1    | 1.000 |
| 90        | I      | 1    | 1.000 |
| 91        | J      | 1    | 1.000 |
| 92        | K      | 1    | 1.000 |
| 93        | L      | 1    | 1.000 |
| 94        | M      | 1    | 1.000 |
| 95        | N      | 1    | 1.000 |
| 96        | O      | 1    | 1.000 |
| 97        | P      | 1    | 1.000 |
| 98        | Q      | 1    | 1.000 |
| 99        | R      | 1    | 1.000 |
| 100       | S      | 1    | 1.000 |

Figure 17b

|     |   |   |       |
|-----|---|---|-------|
| 109 | S | 1 | 1.000 |
| 108 | T | 1 | 1.000 |
| 107 | U | 2 | 2.000 |
| 106 | V | 2 | 2.000 |
| 105 | W | 1 | 1.000 |
| 104 | X | 1 | 1.000 |
| 103 | Y | 4 | 5.000 |
| 102 | Z | 1 | 4.000 |
| 101 | 1 | 1 | 1.000 |
| 100 | 2 | 1 | 7.000 |
| 99  | 3 | 4 | 4.000 |
| 98  | 4 | 1 | 1.000 |
| 97  | 5 | 1 | 1.000 |
| 96  | 6 | 1 | 1.000 |
| 95  | 7 | 1 | 1.000 |
| 94  | 8 | 1 | 1.000 |
| 93  | 9 | 1 | 1.000 |
| 92  | 0 | 1 | 1.000 |
| 91  | A | 1 | 1.000 |
| 90  | B | 1 | 1.000 |
| 89  | C | 1 | 1.000 |
| 88  | D | 1 | 1.000 |
| 87  | E | 1 | 1.000 |
| 86  | F | 1 | 1.000 |
| 85  | G | 1 | 1.000 |
| 84  | H | 1 | 1.000 |
| 83  | I | 1 | 1.000 |
| 82  | J | 1 | 1.000 |
| 81  | K | 1 | 1.000 |
| 80  | L | 1 | 1.000 |
| 79  | M | 1 | 1.000 |
| 78  | N | 1 | 1.000 |
| 77  | O | 1 | 1.000 |
| 76  | P | 1 | 1.000 |
| 75  | Q | 1 | 1.000 |
| 74  | R | 1 | 1.000 |
| 73  | S | 1 | 1.000 |
| 72  | T | 1 | 1.000 |
| 71  | U | 1 | 1.000 |
| 70  | V | 1 | 1.000 |
| 69  | W | 1 | 1.000 |
| 68  | X | 1 | 1.000 |
| 67  | Y | 1 | 1.000 |
| 66  | Z | 1 | 1.000 |
| 65  | 1 | 1 | 1.000 |
| 64  | 2 | 1 | 1.000 |
| 63  | 3 | 1 | 1.000 |
| 62  | 4 | 1 | 1.000 |
| 61  | 5 | 1 | 1.000 |
| 60  | 6 | 1 | 1.000 |
| 59  | 7 | 1 | 1.000 |
| 58  | 8 | 1 | 1.000 |
| 57  | 9 | 1 | 1.000 |
| 56  | A | 1 | 1.000 |
| 55  | B | 1 | 1.000 |
| 54  | C | 1 | 1.000 |
| 53  | D | 1 | 1.000 |
| 52  | E | 1 | 1.000 |
| 51  | F | 1 | 1.000 |
| 50  | G | 1 | 1.000 |
| 49  | H | 1 | 1.000 |
| 48  | I | 1 | 1.000 |
| 47  | J | 1 | 1.000 |
| 46  | K | 1 | 1.000 |
| 45  | L | 1 | 1.000 |
| 44  | M | 1 | 1.000 |
| 43  | N | 1 | 1.000 |
| 42  | O | 1 | 1.000 |
| 41  | P | 1 | 1.000 |
| 40  | Q | 1 | 1.000 |
| 39  | R | 1 | 1.000 |
| 38  | S | 1 | 1.000 |
| 37  | T | 1 | 1.000 |
| 36  | U | 1 | 1.000 |
| 35  | V | 1 | 1.000 |
| 34  | W | 1 | 1.000 |
| 33  | X | 1 | 1.000 |
| 32  | Y | 1 | 1.000 |
| 31  | Z | 1 | 1.000 |
| 30  | 1 | 1 | 1.000 |
| 29  | 2 | 1 | 1.000 |
| 28  | 3 | 1 | 1.000 |
| 27  | 4 | 1 | 1.000 |
| 26  | 5 | 1 | 1.000 |
| 25  | 6 | 1 | 1.000 |
| 24  | 7 | 1 | 1.000 |
| 23  | 8 | 1 | 1.000 |
| 22  | 9 | 1 | 1.000 |
| 21  | A | 1 | 1.000 |
| 20  | B | 1 | 1.000 |
| 19  | C | 1 | 1.000 |
| 18  | D | 1 | 1.000 |
| 17  | E | 1 | 1.000 |
| 16  | F | 1 | 1.000 |
| 15  | G | 1 | 1.000 |
| 14  | H | 1 | 1.000 |
| 13  | I | 1 | 1.000 |
| 12  | J | 1 | 1.000 |
| 11  | K | 1 | 1.000 |
| 10  | L | 1 | 1.000 |
| 9   | M | 1 | 1.000 |
| 8   | N | 1 | 1.000 |
| 7   | O | 1 | 1.000 |
| 6   | P | 1 | 1.000 |
| 5   | Q | 1 | 1.000 |
| 4   | R | 1 | 1.000 |
| 3   | S | 1 | 1.000 |
| 2   | T | 1 | 1.000 |
| 1   | U | 1 | 1.000 |

Figure 17b (continued)




Figure 17c

Figure 18a








Figure 18b

| NUMBER | SYMBOL | APFA | RATIO |
|--------|--------|------|-------|
| 1      | 1      | 1    | 1.000 |
| 2      | 2      | 1    | 1.000 |
| 3      | 3      | 3    | 3.000 |
| 4      | 4      | 3    | 3.000 |
| 5      | 5      | 1    | 1.000 |
| 6      | 6      | 1    | 1.000 |
| 7      | 7      | 1    | 1.000 |
| 8      | 8      | 4    | 4.000 |
| 9      | 9      | 1    | 1.000 |
| 10     | 10     | 4    | 4.000 |
| 11     | 11     | 2    | 2.000 |
| 12     | 12     | 1    | 1.000 |
| 13     | 13     | 1    | 1.000 |
| 14     | 14     | 1    | 1.000 |
| 15     | 15     | 1    | 1.000 |
| 16     | 16     | 1    | 1.000 |
| 17     | 17     | 1    | 1.000 |
| 18     | 18     | 1    | 1.000 |
| 19     | 19     | 1    | 1.000 |
| 20     | 20     | 1    | 1.000 |
| 21     | 21     | 1    | 1.000 |
| 22     | 22     | 1    | 1.000 |
| 23     | 23     | 2    | 2.000 |
| 24     | 24     | 1    | 1.000 |
| 25     | 25     | 1    | 1.000 |
| 26     | 26     | 1    | 1.000 |
| 27     | 27     | 4    | 4.000 |
| 28     | 28     | 1    | 1.000 |
| 29     | 29     | 1    | 1.000 |
| 30     | 30     | 1    | 1.000 |
| 31     | 31     | 1    | 1.000 |



| NUMBER | SYMBOL | AGE | HAIR  |
|--------|--------|-----|-------|
| 2      | 2      | 7   | 1.750 |
| 3      | 3      | 13  | 3.250 |
| 4      | 4      | 2   | .500  |
| 5      | 5      | 5   | 1.250 |
| 6      | 6      | 2   | .500  |
| 7      | 7      | 2   | .500  |
| 8      | 8      | 1   | .250  |
| 9      | 9      | 7   | 1.750 |
| 10     | 10     | 2   | .500  |
| 11     | 11     | 2   | .500  |
| 12     | 12     | 2   | .250  |
| 13     | 13     | 13  | 3.250 |
| 14     | 14     | 4   | 1.000 |
| 15     | 15     | 1   | .250  |
| 16     | 16     | 5   | 1.250 |
| 17     | 17     | 2   | .500  |
| 18     | 18     | 2   | .500  |
| 19     | 19     | 6   | 1.500 |
| 20     | 20     | 1   | .250  |
| 21     | 21     | 5   | 1.250 |
| 22     | 22     | 5   | 1.250 |
| 23     | 23     | 1   | .250  |
| 24     | 24     | 1   | .250  |
| 25     | 25     | 1   | .250  |
| 26     | 26     | 1   | .250  |
| 27     | 27     | 5   | 1.250 |
| 28     | 28     | 39  | 9.500 |
| 29     | 29     | 1   | .250  |
| 30     | 30     | 1   | .250  |
| 31     | 31     | 10  | 2.400 |
| 32     | 32     | 1   | .250  |
| 33     | 33     | 11  | 2.750 |
| 34     | 34     | 5   | 1.250 |
| 35     | 35     | 1   | .250  |
| 36     | 36     | 1   | .250  |
| 37     | 37     | 1   | .250  |
| 38     | 38     | 2   | .500  |
| 39     | 39     | 4   | 2.000 |
| 40     | 40     | 1   | .250  |
| 41     | 41     | 1   | .250  |
| 42     | 42     | 1   | .250  |
| 43     | 43     | 17  | 4.250 |
| 44     | 44     | 6   | 1.500 |
| 45     | 45     | 7   | 1.750 |
| 46     | 46     | 3   | .750  |
| 47     | 47     | 1   | .250  |
| 48     | 48     | 1   | .250  |
| 49     | 49     | 1   | .250  |
| 50     | 50     | 2   | .500  |
| 51     | 51     | 9   | 2.250 |
| 52     | 52     | 1   | .250  |
| 53     | 53     | 1   | .250  |
| 54     | 54     | 7   | 1.750 |
| 55     | 55     | 11  | 4.500 |
| 56     | 56     | 1   | .250  |
| 57     | 57     | 1   | .250  |
| 58     | 58     | 1   | .250  |

Figure 19b

|    |   |    |        |
|----|---|----|--------|
| 59 | S | 1  | .250   |
| 60 | T | 9  | 2.250  |
| 61 | U | 4  | 2.250  |
| 62 | V | 5  | 1.250  |
| 63 | 1 | 3  | .750   |
| 64 | 2 | 2  | .500   |
| 65 | 3 | 20 | 7.250  |
| 66 | 4 | 1  | .250   |
| 67 | 5 | 1  | .250   |
| 68 | 6 | 1  | .250   |
| 69 | 7 | 53 | 13.250 |
| 70 | 8 | 2  | .500   |
| 71 | 9 | 1  | .250   |
| 72 | A | 11 | 2.750  |
| 73 | H | 1  | .250   |
| 74 | C | 1  | .250   |
| 75 | D | 1  | .250   |
| 76 | E | 7  | 1.750  |
| 77 | F | 14 | 4.500  |
| 78 | G | 1  | .250   |
| 79 | H | 1  | .250   |
| 80 | I | 2  | .500   |
| 81 | J | 1  | .250   |
| 82 | K | 1  | .250   |

Figure 19b (continued)



Figure 19c




Figure 20a

| NUMBER | SYMBOL | AREA | DATA  |
|--------|--------|------|-------|
| 1      | 1      | 1    | .750  |
| 2      | 2      | 1    | .250  |
| 3      | 3      | 4    | 1.000 |
| 4      | 4      | 1    | .250  |
| 5      | 5      | 1    | .250  |
| 6      | 6      | 4    | 1.000 |
| 7      | 7      | 4    | 1.000 |
| 8      | 8      | 3    | .750  |
| 9      | 9      | 1    | .250  |
| 10     | 10     | 1    | .250  |
| 11     | 11     | 1    | .250  |
| 12     | 12     | 2    | .500  |
| 13     | 13     | 2    | .500  |
| 14     | 14     | 16   | 4.000 |
| 15     | 15     | 3    | .750  |
| 16     | 16     | 3    | .750  |
| 17     | 17     | 3    | 2.250 |
| 18     | 18     | 1    | .250  |
| 19     | 19     | 2    | .500  |
| 20     | 20     | 6    | 1.500 |
| 21     | 21     | 1    | .250  |
| 22     | 22     | 5    | 1.250 |
| 23     | 23     | 2    | .500  |
| 24     | 24     | 2    | .500  |
| 25     | 25     | 11   | 2.750 |
| 26     | 26     | 4    | 1.000 |
| 27     | 27     | 4    | 1.000 |
| 28     | 28     | 4    | 1.000 |
| 29     | 29     | 1    | .250  |
| 30     | 30     | 4    | 1.000 |
| 31     | 31     | 3    | .750  |
| 32     | 32     | 4    | 1.000 |
| 33     | 33     | 11   | 2.750 |
| 34     | 34     | 2    | .500  |
| 35     | 35     | 1    | .250  |
| 36     | 36     | 2    | .500  |
| 37     | 37     | 3    | .750  |
| 38     | 38     | 1    | .250  |
| 39     | 39     | 1    | .250  |
| 40     | 40     | 1    | .250  |
| 41     | 41     | 2    | .500  |
| 42     | 42     | 1    | .250  |
| 43     | 43     | 1    | .250  |
| 44     | 44     | 1    | .750  |
| 45     | 45     | 4    | 1.000 |
| 46     | 46     | 2    | .500  |
| 47     | 47     | 4    | 1.250 |
| 48     | 48     | 1    | .250  |
| 49     | 49     | 1    | .500  |
| 50     | 50     | 2    | .500  |
| 51     | 51     | 1    | .250  |
| 52     | 52     | 1    | .250  |

Figure 20b




Figure 20c

elements labelled 1, 2,... according to the stage at which they fail to reappear and are found to belong to elongated components; (d-e) analogously for cloud elements. [Note that this noise cleaning method may occasionally make a "cut" across a region at a point of constriction; this is probably aggravated by the fact that the expansion and contraction operations being used are "non-Euclidean" (a "circle" of radius  $k$  is actually a square), so that the constrictions look narrower to the operations than they actually are.]






Figure 21b





Figure 21c



#### 4. Isolatedness

A propagation scheme can also be used to define isolated elements in a binary-valued digital picture [8]. Suppose, in fact, that the 1's in such a picture are "expanded" (in the simple sense of Section 2) by  $k$  steps, and let  $R_k$  be a connected component of 1's in the resulting picture. If  $R_k$  arises from a single "1" in the original picture which was at distance greater than  $2k$  from the nearest other "1", it will be a square of side  $2k + 1$  (assuming that in the expansion, an element is regarded as having eight neighbors), and so will have area  $(2k+1)^2$ . Conversely, if a component has area close to  $(2k+1)^2$ , it must have arisen from a small, closely clustered set of 1's in the original picture which had no other 1's within distance  $2k$  of them -- in short, such a component must have arisen from an isolated set of 1's. Under some circumstances, one might want to preserve small objects in a picture when they are isolated, but not otherwise; the procedure just outlined can be used for this purpose. Figure 22 shows, and tabulates the areas of, the connected components resulting when the 1's in Figure 1 are expanded by 1 and by 2 steps. Note that an elongated object will never be called elongated by this scheme (e.g., components 2, 7, 8, A in Figure 22a, and 4 in Figure 22b).





| NUMBER | SYMBOL | AREA | RATIO   |
|--------|--------|------|---------|
| 1      | 1      | 9    | 1.000   |
| 2      | 2      | 27   | 3.000   |
| 3      | 3      | 45   | 4.500   |
| 4      | 4      | 145  | 16.111  |
| 5      | 5      | 9    | 1.000   |
| 6      | 6      | 139  | 15.444  |
| 7      | 7      | 29   | 3.222   |
| 8      | 8      | 29   | 3.222   |
| 9      | 9      | 1434 | 159.669 |
| 10     | A      | 24   | 2.667   |
| 11     | H      | 9    | 1.000   |

Figure 22c



| NUMBER | SYMBOL | AREA | RATIO  |
|--------|--------|------|--------|
| 1      | 1      | 630  | 25.560 |
| 2      | 2      | 521  | 20.920 |
| 3      | 3      | 25   | 1.000  |
| 4      | 4      | 61   | 2.440  |
| 5      | 5      | 2014 | 80.560 |

Figure 22d

References

- [1] G. P. Dinneen, "Programming pattern recognition", Proc. Western Joint Computer Conf., March 1955, 94-100.
- [2] R. E. Graham, "Snow removal - a noise-stripping process for picture signals", I.R.E. Trans. IT-8, February 1962, 129-144.
- [3] S. M. Pizer and H. G. Vetter, "Perception and processing of medical radioisotope scans", Pictorial Pattern Recognition, Thompson 1968, 147-156.
- [4] H. Tairy, "Some qualitative and quantitative results on spatial filtering of granularity", Appl. Opt. 3, January 1964, 39-43.
- [5] P. G. Roetling, "Effects of signal-dependent granularity", J. Opt. Soc. Amer. 55, January 1965, 67-71.
- [6] R. Nathan, "Picture enhancement for the moon, Mars, and man", Pictorial Pattern Recognition, Thompson 1968, 239-266.
- [7] A. Rosenfeld, C. Fried and J. N. Orton, "Automatic cloud interpretation", Photogram. Eng. 31, November 1965, 991-1002.
- [8] A. Rosenfeld and J. L. Pfaltz, "Distance functions on digital pictures", Pat. Recog. 1, July 1968, 33-61.