

N71-21470
NASA CR-114931

MULTIPLE ZEROS OF POLYNOMIALS

CASE FILE
COPY

by

CRAIG A. WOOD

Department of Mathematics
Oklahoma State University
Stillwater, Oklahoma

A National Aeronautics and Space Administration

Research Grant

Grant Number NASA NGR 37-002-084

MULTIPLE ZEROS OF POLYNOMIALS

by

CRAIG A. WOOD

Department of Mathematics
Oklahoma State University
Stillwater, Oklahoma

A National Aeronautics and Space Administration

Research Grant

Grant Number NASA NGR 37-002-084

Abstract: MULTIPLE ZEROS OF POLYNOMIALS

Grant: The National Aeronautics and Space Administration

Grant Number: NASA NGR 37-002-084

Grantee: Professor Craig A. Wood
Department of Mathematics
Oklahoma State University
Stillwater, Oklahoma

Various classical methods exist for extracting the zeros of a polynomial

$$P(X) = a_1 X^N + a_2 X^{N-1} + \dots + a_{N+1}$$

where $a_1 \neq 0$ and $a_1, a_2, \dots, a_{N+1}$ are complex numbers, when $N=1,2,3,4$. For polynomials of higher degree, iterative numerical methods must be used. In this material four iterative methods are presented for approximating the zeros of a polynomial using a digital computer. Newton's method and Muller's method are two well known iterative methods which are presented. They extract the zeros of a polynomial by generating a sequence of approximations converging to each zero. However, both of these methods are very unstable when used on a polynomial which has multiple zeros. That is, either they fail to converge to some or all of the zeros, or they converge to very bad approximations of the polynomial's zeros.

This material introduces two new methods, the greatest common divisor (G.C.D.) method and the repeated greatest common divisor (repeated G.C.D.) method, which are superior methods for numerically approximating the zeros of a polynomial having multiple zeros.

The above methods were all programmed in FORTRAN IV and comparisons in time and accuracy are given. These programs were executed on the

IBM 360/50 computer as well as the UNIVAC 1108 and the CDC 6600 computer.

This material also contains complete documentations for six FORTRAN IV programs. Flow charts, program listings, definition of variables used in the program, and instructions for use of each program are included.

PREFACE

Four iterative methods for approximating the zeros of a polynomial using a digital computer are presented in this material. Chapter I is an introduction. Chapters II and III contain Newton's and Muller's methods, respectively. Chapters IV and V present two new methods which depend upon finding the greatest common divisor of two polynomials. Chapter VI contains a comparison of the four methods. Flow charts, FORTRAN IV programs, and complete program documentations for these four methods are presented in appendices A through H.

I would like to express my appreciation to the National Aeronautics and Space Administration, specifically the Manned Spacecraft Center in Houston, Texas, for their financial support in making this work possible under grant number NASA NGR 37-002-084. I would also like to thank Randy Snider, a graduate assistant supported by this grant, for the great deal of work he put in on the FORTRAN programs. In particular, the material on Newton's and Muller's Methods included in this paper is part of his masters thesis at Oklahoma State University.

TABLE OF CONTENTS

Chapter	Page
I. INTRODUCTION.	1
II. NEWTON'S METHOD	4
Derivation of the Algorithm.	4
Convergence of Newton's Method	8
Procedure for Newton's Method.	9
Geometrical Interpretation of Newton's Method. . .	10
Determining Multiple Roots	10
III. MULLER'S METHOD	12
Derivation of the Algorithm.	12
Procedure for Muller's Method.	19
Geometrical Interpretation of Muller's Method. . .	20
Determining Multiple Roots	20
IV. GREATEST COMMON DIVISOR METHOD.	22
Derivation of the Algorithm.	22
Determining Multiplicities	29
Procedure for the G.C.D. Method.	29
V. REPEATED GREATEST COMMON DIVISOR.	30
Derivation of the Algorithm.	30
Procedure for the Repeated G.C.D. Method	32
VI. CONCLUSION.	34
Polynomials With all Distinct Roots.	34
Polynomials With Multiple Roots.	35
Time Comparisons	40
Sample Results	42
REFERENCES	85
APPENDIX A - SPECIAL FEATURES OF NEWTON'S AND MULLER'S PROGRAMS.	86
Generating Approximations.	86
Altering Approximations.	88
Searching the Complex Plane.	89

Chapter	Page
Improving Zeros Found	91
Solving Quadratic Polynomial.	91
Missing Roots	92
Miscellaneous	92
APPENDIX B - NEWTON'S METHOD	100
Use of the Program.	100
Input Data for Newton's Method.	102
Variables Used in Newton's Method	107
Description of Program Output	107
Informative and Error Message	109
Flow Charts	120
Program Listing	126
APPENDIX C - MULLER'S METHOD	135
Use of the Program.	135
Input Data for Muller's Method.	137
Variables Used in Muller's Method	137
Description of Program Output	138
Informative and Error Messages.	139
Flow Charts	144
Program Listing	150
APPENDIX D - SPECIAL FEATURES OF THE G.C.D. AND THE REPEATED G.C.D. PROGRAMS.	163
Generating Approximations	163
Altering Approximations	165
Searching the Complex Plane	166
Improving Zeros Found	168
Solving Quadratic Polynomial.	168
Missing Roots	169
Miscellaneous	169
APPENDIX E - G.C.D. - NEWTON'S METHOD.	171
Use of the Program.	171
Input Data for G.C.D. - Newton's Method	173
Variables Used in G.C.D. - Newton's Method.	178
Description of Program Output	188
Informative and Error Messages.	189
Flow Charts	191
Program Listing	200

Chapter	Page
APPENDIX F - G.C.D. - MULLER'S METHOD	214
Use of the Program	214
Input Data for G.C.D. - Muller's Method	216
Variables Used in G.C.D. - Muller's Method	216
Description of Program Output	216
Informative and Error Messages	216
Flow Charts	222
Program Listing	234
APPENDIX G - REPEATED G.C.D. - NEWTON'S METHOD	253
Use of the Program	253
Input Data for Repeated G.C.D. - Newton's Method	255
Variables Used in Repeated G.C.D. - Newton's Method	255
Description of Program Output	255
Informative and Error Messages	256
Flow Charts	260
Program Listing	270
APPENDIX H - REPEATED G.C.D. - MULLER'S METHOD	286
Use of the Program	286
Input Data for Repeated G.C.D. - Muller's Method	287
Variables used in Repeated G.C.D. - Muller's Method	288
Description of Program Output	288
Informative and Error Messages	289
Flow Charts	290
Program Listing	301

CHAPTER I

INTRODUCTION

Frequently in scientific work it becomes necessary to find the zeros, real or complex, of the polynomial of degree N

$$P(X) = a_1 X^N + a_2 X^{N-1} + \dots + a_N X + a_{N+1}$$

where $a_1 \neq 0$ and the coefficients $a_1, a_2, \dots, a_{N+1}$ are complex numbers. Various classical methods calculate the exact roots of polynomials of degree 1, 2, 3, or 4. For polynomials of higher degree, no such methods exist. Thus, to solve for the zeros of such polynomials, numerical methods of iteration based on successive approximations must be employed. In the following material four such methods are given which are particularly suited for modern high speed computers.

Newton's method is an iterative procedure which generates a sequence of successive approximations of a zero of $P(X)$ by using the iteration formula

$$X_{n+1} = X_n - P(X_n)/P'(X_n).$$

An initial approximation to the zero is required to start the iterative process. Under certain conditions this sequence will converge quadratically to the desired root. It is, however, necessary to compute the value of the polynomial and its derivative for each step in the

iterative procedure. Once a zero of $P(X)$ has been found, it is divided out of $P(X)$, giving a deflated polynomial of lower degree. $P(X)$ is replaced by the deflated polynomial and the iterative process is applied to extract another zero of $P(X)$. This procedure is repeated until all zeros of $P(X)$ have been found. The zeros may then be rechecked and their accuracy possibly improved by using them as initial approximations with Newton's process applied to the full (undeflated) polynomial.

Muller's method is also an iterative procedure generating a sequence $X_1, X_2, \dots, X_n, \dots$ of successive approximations of a root of $P(X)$. This method converges almost quadratically near a zero and does not require the evaluation of the derivative of the polynomial. Muller's method requires three distinct approximations of a root to start the process of iteration. A quadratic equation is constructed through the three given points as an approximation of $P(X)$. The root of the quadratic closest to X_n is taken as X_{n+1} , the next approximation to the zero. This process is then repeated on the last three points of the sequence. After a root of $P(X)$ has been found, $P(X)$ is deflated, and replaced in the above procedure by the deflated polynomial. After all zeros of $P(X)$ are found from successive deflations, they are improved as in Newton's method.

The greatest common divisor method reduces the problem of finding all zeros (possibly multiple zeros) of $P(X)$ to one of extracting the zeros of a polynomial $P_1(X) = P(X)/D(X)$, all of whose zeros are simple. $D(X)$, the greatest common divisor of $P(X)$ and its derivative, $P'(X)$, is obtained by repeated application of the division algorithm. Once $P_1(X)$ is obtained, some suitable method such as Newton's or Muller's method

is used to find the zeros of $P_1(X)$. By finding all the zeros of $P_1(X)$, all the zeros of $P(X)$ are obtained. The multiplicity of each zero may then be determined.

The repeated greatest common divisor method repeatedly uses the greatest common divisor method to extract the zeros of $P(X)$ and their multiplicities at the same time. That is, the repeated greatest common divisor method reduces the problem of finding the zeros of $P(X)$, which possibly has multiple zeros, to one of finding the zeros of a polynomial which has only simple zeros and the zeros of this polynomial are all the zeros of $P(X)$ of a given multiplicity. The repeated greatest common divisor method must also use a supporting method such as Newton's method or Muller's method.

Chapters II-V contain the examinations of these methods. Each examination includes a development of the method together with the conditions necessary for convergence of the method. Chapter VI contains a comparison of the methods giving advantages and disadvantages of each method.

A complete set of documentations is given for six FORTRAN IV programs in Appendices A-H. Flow charts, program listings, definition of variables used in the program, and instructions for use of each program are included.

It should also be noted that the expressions "zero of a polynomial" and "root of a polynomial" and the words "zero" and "root" are used interchangeably in this material.

CHAPTER II

NEWTON'S METHOD

1. Derivation of the Algorithm

Newton's method is probably the most popular iterative procedure for finding the zeros of a polynomial. This fact is due to the excellent results obtained, the simplicity of the computational routine, and the fast rate of convergence obtained provided the initial approximation of a zero is close enough. Also, the method can be applied to the extraction of complex as well as real zeros.

Consider the polynomial

$$P(X) = a_1 X^N + a_2 X^{N-1} + \dots + a_N X + a_{N+1} \quad (2-1)$$

where $a_1 \neq 0$ and the coefficients $a_1, a_2, \dots, a_{N+1}$ are complex. The algorithm for Newton's method can be derived by approximating $P(X)$ by a Taylor series expansion about an approximation, X_0 , of a zero, α , of $P(X)$. Using only the first two terms of the expansion, the expression

$$P(X) \doteq P(X_0) + P'(X_0)(X - X_0)$$

is obtained. If this equation is solved for $P(X) = 0$, then

$$0 \doteq P(X_0) + P'(X_0)(X - X_0)$$

results. Rearranging terms produces

$$0 \doteq P(X_0) + P'(X_0) X - P'(X_0) X_0$$

followed by

$$P'(X_0) X_0 - P(X_0) \doteq P'(X_0) X$$

from which division by $P'(X_0)$ produces

$$X_0 - P(X_0)/P'(X_0) \doteq X$$

which is the basic formula for Newton's method. Thus, in general, we obtain the $(n+1)^{\text{th}}$ approximation, X_{n+1} , of α from the n^{th} approximation, X_n , by

$$X_{n+1} = X_n - P(X_n)/P'(X_n). \quad (2-2)$$

As a result of repeated use of this algorithm, we obtain the sequence

$$X_0, X_1, X_2, \dots, X_n, \dots \quad (2-3)$$

of successive approximations of the root, α . It should be noted that an initial approximation is necessary to start the iterative process for each new zero; that is, a polynomial of degree N may require N initial approximations.

In order to use equation (2-2), it is necessary to compute, for each X_n , the value of the polynomial, $P(X_n)$, and its derivative, $P'(X_n)$. The division algorithm states that if $P(X)$ and $G(X)$ are polynomials, then there exists polynomials $H(X)$ and $K(X)$ such that $P(X) = H(X) G(X) + K(X)$ where $K(X) = 0$ or $\deg. K(X) < \deg. G(X)$. From this expression of $P(X)$, the following remainder theorem is obtained:

Theorem 2.1. If $P(X)$ is a polynomial and c is a complex number, then the remainder obtained from dividing $P(X)$ by $(X - c)$ is $P(c)$.

The proof of Theorem 2.1 is given in [3, P. 102]. Thus, $P(X)$ can be written as $P(X) = (X - c) H(X) + R$ where $P(c) = R$. $P'(X)$ is then obtained by the following theorem, the proof of which can be found in [3, PP. 105-106].

Theorem 2.2. If $P(X)$ and $H(X)$ are polynomials and c is a complex number such that $P(X) = (X - c) H(X) + R$ where $P(c) = R$, then the remainder obtained from dividing $H(X)$ by $(X - c)$ is $P'(c)$.

From synthetic division, an algorithm known as Horner's Method is acquired for computing $P(X_n)$ and $P'(X_n)$.

Theorem 2.3. Let $P(X)$ be defined as in equation (2-1) and let d be a complex number. Define a sequence $b_1, b_2, \dots, b_{N+1}$ by

$$\begin{aligned} b_1 &= a_1 \\ b_i &= a_i + db_{i-1} \quad (i = 2, 3, \dots, N+1). \end{aligned}$$

Define another sequence $c_1, c_2, \dots, c_N$ by

$$\begin{aligned} c_1 &= b_1 \\ c_j &= b_j + dc_{j-1} \quad (j = 2, 3, \dots, N). \end{aligned}$$

Then $P(d) = b_{N+1}$ and $P'(d) = c_N$. The elements $b_1, b_2, \dots, b_N$ are the coefficients of the polynomial $H(X)$ in Theorem 2.2 when $P(X)$ is divided by $(X - d)$.

These formulas are derived in [3, PP. 106-107]. Thus with equation (2-2) and the iteration formulas of the previous theorem, Newton's method can now be applied to generate the sequence (2-3) which will converge to the root, α , if the convergence conditions given in Theorem 2.4 are satisfied.

A criterion is needed to determine when to terminate the sequence (2-3); that is, when has a zero been found? For convergence of the sequence, there must exist a term in the sequence beyond which the difference between any two successive terms is arbitrarily small. Therefore, it is desirable to make the quotient $|X_n/X_{n+1}|$ sufficiently near 1. From equation (2-2)

$$1 = \left| \frac{X_n}{X_{n+1}} - \frac{\frac{P(X_n)}{P'(X_n)}}{\frac{X_{n+1}}{X_{n+1}}} \right|$$

$$\geq \left| \frac{X_n}{X_{n+1}} \right| - \left| \frac{\frac{P(X_n)}{P'(X_n)}}{\frac{X_{n+1}}{X_{n+1}}} \right|.$$

Thus

$$1 + \left| \frac{\frac{P(X_n)}{P'(X_n)}}{\frac{X_{n+1}}{X_{n+1}}} \right| \geq \left| \frac{X_n}{X_{n+1}} \right|$$

where $P'(X_n)$ and $X_{n+1} \neq 0$. Thus, iterations are continued until an X_n is obtained such that $|P(X_n)/P'(X_n)|/|X_{n+1}|$ is as small as desired.

After a zero, α , of $P(X)$ has been found, the term $(X - \alpha)$ is synthetically divided out of $P(X)$ by deflation using Theorem 2.3 obtaining

a polynomial, $P_1(X)$, of degree $N-1$. The root finding process is then repeated to extract a zero, α_1 , of $P_1(X)$. $P(X)$ can be written as

$$P(X) = (X - \alpha) P_1(X) + R$$

where $R = P(\alpha)$. But $P(\alpha) = 0$. Therefore, substitution produces

$$P(X) = (X - \alpha) P_1(X).$$

Now $P_1(\alpha_1) = 0$ implies that $P(\alpha_1) = 0$. Hence, α_1 is a zero of $P(X)$.

By the process of root finding and successive deflations, zeros $\alpha_0, \alpha_1, \dots, \alpha_{N-1}$ of the deflated polynomials

$$P(X) = P_0(X), P_1(X), \dots, P_{N-1}(X),$$

respectively, are extracted. Each α_i ($i = 0, 1, 2, \dots, N-1$) is a zero of $P(X)$ since each α_i is a zero of $P_{i-1}(X), P_{i-2}(X), \dots, P_1(X), P(X)$.

After all zeros of $P(X)$ have been found, it may be possible to improve their accuracy by using them as initial approximations with Newton's method applied to the full (undeflated) polynomial, $P(X)$. This should correct any loss of accuracy which may have resulted from the successive deflations.

2. Convergence of Newton's Method

The following theorem from [2, PP. 79-81] gives sufficient conditions for the convergence of sequence (2-3).

Theorem 2.4. Let $P(X)$ be a polynomial and let the following conditions be satisfied on the closed interval $[a, b]$:

1. $P(a) P(b) < 0$
2. $P'(X) \neq 0, X \in [a, b]$
3. $P''(X)$ is either ≥ 0 or ≤ 0 for all $X \in [a, b]$
4. If c denotes the endpoint of $[a, b]$ at which $|P'(X)|$ is smaller, then $|P(c)/P'(c)| \leq b - a$.

Then Newton's method converges to the (only) solution, s , of $P(X) = 0$ for any choice of X_0 in $[a, b]$.

When convergence is obtained, it is quadratic; that is,

$$e_{i+1} = \frac{1}{2} F''(\eta_i) e_i^2$$

where $F(X_i) = X_i - P(X_i)/P'(X_i)$, η_i is between X_i and the zero, α , and e_i is the error in X_i . This means that the error obtained in the $(i+1)^{th}$ iteration of Newton's algorithm is proportional to the square of the error obtained in the i^{th} iteration. A proof of quadratic convergence can be found in [1, PP. 31-33].

3. Procedure for Newton's Method

The general procedure for applying Newton's method is enumerated sequentially as follows, starting with initial approximation X_0 :

1. Calculate a new approximation X_{n+1} by

$$X_{n+1} = X_n - P(X_n)/P'(X_n).$$

2. Test for convergence; that is, test

$$|P(X_n)/P'(X_n)| / |X_{n+1}| < \epsilon$$

for some ϵ chosen as small as desired.

3. If convergence is obtained, perform the following:

- a. Save X_{n+1} as the desired approximation to a zero of $P(X)$.
 - b. Deflate $P(X)$ using X_{n+1} .
 - c. Replace $P(X)$ by the deflated polynomial.
 - d. Return to step 1 with a new initial approximation.
4. If no convergence is obtained, increase n by 1 and return to step 1.

In order to prevent an unending iteration process in case the method does not produce convergence, a maximum number of iterations should be specified. If convergence is not obtained within this number of iterations, change the initial approximation and return to step 1 above.

4. Geometrical Interpretation of Newton's Method

A geometrical interpretation of Newton's method is given in Figure 2.1. X_1 is an approximation to the zero, α . $P'(X_1)$ is the slope of the line tangent to $P(X)$ at X_1 . X_{1+1} is the intersection of the tangent line with the x axis.

5. Determining Multiple Roots

If $P(X)$ has m distinct zeros, then $P(X)$ can be written as

$$P(X) = a_1 (X - \alpha_1)^{e_1} (X - \alpha_2)^{e_2} \dots (X - \alpha_m)^{e_m}, \quad (m \leq N)$$

where α_i is a zero of $P(X)$ and e_i is the multiplicity of α_i ($i = 1, 2, \dots, m$). Consider the root α_j . Dividing out the term

$(X - \alpha_j)$ by deflating $P(X)$ gives $P_1(X)$ of degree $N-1$ which can be written as

$$P_1(X) = (X - \alpha_1)^{e_1} (X - \alpha_2)^{e_2} \dots (X - \alpha_j)^{e_j-1} \dots (X - \alpha_m)^{e_m}.$$

Evaluating $P_1(X)$ at the zero, α_j , gives $P_1(\alpha_j) = 0$ if $e_j > 1$. Thus, after a zero, α , of $P(X)$ is determined by Newton's iterative process and the current polynomial is deflated giving $P_1(X)$, then $P_1(\alpha)$ is evaluated. If $P_1(\alpha) \leq \epsilon$ for some small number ϵ , α is a root of $P_1(X)$ and thus has multiplicity at least equal to two. $P_1(X)$ is then deflated giving $P_2(X)$. If $P_2(\alpha) \leq \epsilon$, α is of multiplicity at least three. This process is continued until a deflated polynomial $P_k(X)$ is encountered such that either $\deg. P_k(X) = 0$ or $P_k(\alpha) > \epsilon$. α is then a zero of multiplicity $k+1$.




Figure 2.1. Geometrical Interpretation of Newton's Method

CHAPTER III

MULLER'S METHOD

1. Derivation of the Algorithm

Muller's method in [4] is an iterative procedure designed to find any prescribed number of zeros, real or complex, of a polynomial. The method does not require the evaluation of the derivative and near a zero the convergence is almost quadratic.

Consider the polynomial

$$P(X) = a_1 X^N + a_2 X^{N-1} + \dots + a_N X + a_{N+1} \quad (3-1)$$

with complex coefficients such that $a_1 \neq 0$. Given three distinct approximations, X_{n-2}, X_{n-1}, X_n , to a root, α , of $P(X)$, the problem is to determine X_{n+1} in such a way as to generate a sequence

$$X_1, X_2, X_3, \dots, X_n, X_{n+1}, \dots \quad (3-2)$$

of approximations converging to α . The points $(X_{n-2}, P(X_{n-2}))$, $(X_{n-1}, P(X_{n-1}))$, and $(X_n, P(X_n))$ determine a unique quadratic polynomial, $Q(X)$, approximating $P(X)$ in the vicinity of X_{n-2}, X_{n-1}, X_n . A general proof of this can be found in [2, PP. 133-134]. Thus, the zeros of $Q(X)$ will be approximations of the zeros of $P(X)$ in this region of approximation. From the general representation in [2, P. 184] of the Lagrangian interpolating polynomial, the representation of $Q(X)$ is given by

$$\begin{aligned}
Q(X) &= \frac{(X - X_{n-1})(X - X_{n-2})}{(X_n - X_{n-1})(X_n - X_{n-2})} P(X_n) \\
&\quad + \frac{(X - X_n)(X - X_{n-2})}{(X_{n-1} - X_n)(X_{n-1} - X_{n-2})} P(X_{n-1}) \\
&\quad + \frac{(X - X_n)(X - X_{n-1})}{(X_{n-2} - X_n)(X_{n-2} - X_{n-1})} P(X_{n-2})
\end{aligned}$$

which can be rewritten as

$$\begin{aligned}
Q(X) &= Q(X - X_n + X_n) \\
&= \frac{(X - X_n + X_n - X_{n-1})(X - X_n + X_n - X_{n-1} + X_{n-1} - X_{n-2})}{(X_n - X_{n-1})(X_n - X_{n-1} + X_{n-1} - X_{n-2})} P(X_n) \\
&\quad - \frac{(X - X_n)(X - X_n + X_n - X_{n-1} + X_{n-1} - X_{n-2})}{(X_n - X_{n-1})(X_{n-1} - X_{n-2})} P(X_{n-1}) \\
&\quad + \frac{(X - X_n)(X - X_n + X_n - X_{n-1})}{(X_n - X_{n-1} + X_{n-1} - X_{n-2})(X_{n-1} - X_{n-2})} P(X_{n-2}).
\end{aligned}$$

In order to simplify this expression, introduce the quantities

$$h_n = X_n - X_{n-1}, \quad h = X - X_n.$$

Then

$$\begin{aligned}
Q(X) &= Q(X_n + h) \\
&= \frac{(h + h_n)(h + h_n + h_{n-1})}{h_n(h_n + h_{n-1})} P(X_n) \\
&\quad - \frac{h(h + h_n + h_{n-1})}{h_n h_{n-1}} P(X_{n-1})
\end{aligned}$$

$$\begin{aligned}
& + \frac{h(h + h_n)}{(h_n + h_{n-1})h_{n-1}} P(X_{n-2}) \\
& = \frac{h^2 + 2hh_n + hh_{n-1} + h_n^2 + h_n h_{n-1}}{h_n^2 + h_n h_{n-1}} P(X_n) \\
& \quad - \frac{h^2 + hh_n + hh_{n-1}}{h_n h_{n-1}} P(X_{n-1}) \\
& \quad + \frac{h^2 + hh_n}{h_n h_{n-1} + h_{n-1}^2} P(X_{n-2}).
\end{aligned}$$

Collecting terms containing like powers of h produces

$$Q(X) = Q(X_n + h)$$

$$\begin{aligned}
& = \left(\frac{P(X_n)}{h_n^2 + h_n h_{n-1}} - \frac{P(X_{n-1})}{h_n h_{n-1}} + \frac{P(X_{n-2})}{h_n h_{n-1} + h_{n-1}^2} \right) h^2 \\
& + \left(\frac{(2h_n + h_{n-1}) P(X_n)}{h_n^2 + h_n h_{n-1}} - \frac{(h_n + h_{n-1}) P(X_{n-1})}{h_n h_{n-1}} + \frac{h_n P(X_{n-2})}{h_n h_{n-1} + h_{n-1}^2} \right) h \\
& + \frac{h_n (h_n + h_{n-1}) P(X_n)}{h_n^2 + h_n h_{n-1}} \\
& = \left(\frac{P(X_n) h_{n-1}}{h_n^2 h_{n-1} + h_n h_{n-1}^2} - \frac{P(X_{n-1})}{h_n h_{n-1}} + \frac{P(X_{n-2})}{h_n h_{n-1} + h_{n-1}^2} \right) h^2 \\
& + \left(\frac{(2h_n h_{n-1} + h_{n-1}^2) P(X_n)}{h_n^2 h_{n-1} + h_n h_{n-1}^2} - \frac{(h_n + h_{n-1}) P(X_{n-1})}{h_n h_{n-1}} + \frac{h_n P(X_{n-2})}{h_n h_{n-1} + h_{n-1}^2} \right) h
\end{aligned}$$

$$+ \frac{(h_n^2 h_{n-1} + h_{n-1}^2 h_n) P(X_n)}{h_n^2 h_{n-1} + h_n h_{n-1}^2} .$$

Using the common denominator, $h_n^2 h_{n-1} + h_n h_{n-1}^2$, and combining terms yields

$$\begin{aligned} Q(X_n + h) = & \left(\frac{P(X_n) h_{n-1} - P(X_{n-1})(h_n + h_{n-1}) + P(X_{n-2}) h_n}{h_n^2 h_{n-1} + h_n h_{n-1}^2} \right) h^2 \\ & + \left(\frac{(2h_n h_{n-1} + h_{n-1}^2) P(X_n) - (h_n + h_{n-1})^2 P(X_{n-1}) + h_n^2 P(X_{n-2})}{h_n^2 h_{n-1} + h_n h_{n-1}^2} \right) h \\ & + \frac{(h_n^2 h_{n-1} + h_{n-1}^2 h_n) P(X_n)}{h_n^2 h_{n-1} + h_n h_{n-1}^2} . \end{aligned}$$

Multiplying by h_n/h_{n-1}^2 results in

$$\begin{aligned} Q(X_n + h) = & \left[\frac{P(X_n) \frac{h_n}{h_{n-1}} - P(X_{n-1}) \left(\left(\frac{h_n}{h_{n-1}} \right)^2 + \frac{h_n}{h_{n-1}} \right) + P(X_{n-2}) \left(\frac{h_n}{h_{n-1}} \right)^2}{\frac{h_n^3}{h_{n-1}} + h_n^2} \right] h^2 \\ & + \left[\frac{\left(\frac{h_n^2}{h_{n-1}} + h_n \right) P(X_n) - h_n \left[\left(\frac{h_n}{h_{n-1}} \right) + \left(\frac{h_{n-1}}{h_{n-1}} \right) \right]^2 P(X_{n-1}) + \frac{h_n^3}{h_{n-1}^2} P(X_{n-2})}{\frac{h_n^3}{h_{n-1}} + h_n^2} \right] h \end{aligned}$$

$$+ \left[\frac{\left(\frac{h_n^3}{h_{n-1}} + h_n^2 \right) P(X_n)}{\frac{h_n^3}{h_{n-1}} + h_n^2} \right].$$

Let $q_n = \frac{h_n}{h_{n-1}}$ and $q = \frac{h}{h_n}$. Then

$$\begin{aligned} Q(X_n + h) &= \left(\frac{P(X_n) q_n - P(X_{n-1}) (q_n^2 + q_n) + P(X_{n-2}) q_n^2}{q_n + 1} \right) q^2 \\ &\quad + \left(\frac{(2 q_n + 1) P(X_n) - (q_n + 1)^2 P(X_{n-1}) + q_n^2 P(X_{n-2})}{q_n + 1} \right) q \\ &\quad + \frac{(q_n + 1) P(X_n)}{q_n + 1}. \end{aligned}$$

Now let

$$\begin{aligned} A_n &= q_n P(X_n) - q_n (q_n + 1) P(X_{n-1}) + q_n^2 P(X_{n-2}) \\ B_n &= (2q_n + 1) P(X_n) - (q_n + 1)^2 P(X_{n-1}) + q_n^2 P(X_{n-2}) \\ C_n &= (q_n + 1) P(X_n). \end{aligned}$$

Then

$$Q(X_n + h) = Q(X_n + qh_n)$$

and

$$Q(X_n + qh_n) = \frac{A_n q^2 + B_n q + C_n}{q_n + 1}.$$

Solving the quadratic equation $Q(X_n + qh_n) = 0$ and denoting the result by q_{n+1} gives:

$$q_{n+1} = \frac{-B_n \pm \sqrt{B_n^2 - 4A_n C_n}}{2A_n}$$

and the new approximation is found as follows:

$$q_{n+1} = \frac{h_{n+1}}{h_n} = \frac{X_{n+1} - X_n}{h_n}.$$

Thus

$$X_{n+1} = X_n + h_n q_{n+1}$$

In order to avoid loss of accuracy, q_{n+1} can be written in a better form as follows:

$$\begin{aligned} q_{n+1} &= \frac{-B_n \pm \sqrt{B_n^2 - 4A_n C_n}}{2A_n} \cdot \frac{B_n \pm \sqrt{B_n^2 - 4A_n C_n}}{B_n \pm \sqrt{B_n^2 - 4A_n C_n}} \\ &= \frac{-B_n^2 + B_n^2 - 4A_n C_n}{2A_n (B_n \pm \sqrt{B_n^2 - 4A_n C_n})} \\ q_{n+1} &= \frac{-2C_n}{B_n \pm \sqrt{B_n^2 - 4A_n C_n}}. \end{aligned} \quad (3-3)$$

The sign in the denominator should be chosen such that the magnitude of the denominator is largest, thus causing $|q_{n+1}|$ to be smallest. This, in turn, will make X_{n+1} closest to X_n .

Note that each iteration of this process requires three approximations, X_{n-2}, X_{n-1}, X_n , in order to compute X_{n+1} . Thus, when X_{n+1} is found, X_{n-1}, X_n, X_{n+1} are used to compute X_{n+2} ; that is, the last three terms of the generated sequence are used to compute the next term.

Convergence of the sequence (3-2) to a zero is obtained when the elements X_k and X_{k+1} of the sequence are found such that

$$\frac{|X_{k+1} - X_k|}{|X_{k+1}|} < \epsilon, X_{k+1} \neq 0;$$

that is, the ratio of the change in the approximation to the approximation itself is as small as desired.

In order to use the iterative formulas, it is necessary to compute the value, $P(X_j)$, of the polynomial $P(X)$ at the approximation X_j . The procedure for doing this is discussed in Chapter II, § 1. The iteration formulas are given in Theorem 2.3 of Chapter II.

After a zero, α , of $P(X)$ has been found, $P(X)$ is deflated as described in Chapter II, § 1, and the process repeated to extract a zero, α_1 , of $P_1(X)$. By applying Muller's method to successively deflated polynomials, all the zeros of $P(X)$ are obtained. For more detailed discussion of this procedure see Chapter II, § 1, keeping in mind that Muller's instead of Newton's method is used.

Muller's method requires three initial approximations to a zero in order to start the iteration process. If three are not known, the values $X_1 = -1$, $X_2 = 1$, $X_3 = 0$ can be used.

Convergence of Muller's method is almost quadratic provided the three initial approximations are sufficiently close to a zero of $P(X)$. This is natural to expect since $P(X)$ is being approximated by a

quadratic polynomial. Quadratic convergence means that the error obtained in the $(n+1)^{\text{th}}$ step of the iterative process is proportional to the square of the error obtained in the n^{th} iteration. However, no general proof of convergence has been obtained for Muller's method. It has produced convergence in the majority of the cases tested.

In application of Muller's method, an alteration should be made to handle the case in which the denominator of equation (3-3) is zero (0). This occurs whenever $P(X_n) = P(X_{n-1}) = P(X_{n-2})$. If this happens, set $q_{n+1} = 1$.

Another alteration which should be made in actual practice is to compute the quantity $|P(X_{n+1})| / |P(X_n)|$ whenever the value $P(X_{n+1})$ is calculated. If the former quantity exceeds ten (10), q_{n+1} is halved and h_n , X_{n+1} , and $P(X_{n+1})$ are recomputed accordingly.

2. Procedure for Muller's Method

The basic steps performed by Muller's method are listed sequentially as follows, starting with initial approximations X_1 , X_2 , and X_3 .

1. Compute h_n , q_n , D_n , B_n , C_n , q_{n+1} as defined previously.
2. Compute the next approximation X_{n+1} by

$$X_{n+1} = X_n + h_n q_{n+1}.$$

3. Test for convergence; that is, test

$$|X_{n+1} - X_n| / |X_{n+1}| < \epsilon$$

for some suitably small number ϵ .

4. If the test fails, return to step 1 with the last three approximations X_{n+1} , X_n , X_{n-1} .

5. If the test passes, do the following:
 - a. Save X_{n+1} as the desired approximation to a zero.
 - b. Deflate the current polynomial using X_{n+1} .
 - c. Replace the current polynomial by the deflated polynomial.
 - d. Return to step 1 with a new set of initial approximations.

In order to avoid an unending iteration process in case the method does not produce convergence, a maximum number of iterations should be specified. If convergence is not obtained within this number of iterations, the initial approximations should be altered.

3. Geometrical Interpretation of Muller's Method

Figure 3.1. shows the geometrical interpretation of Muller's method for real roots of $P(X)$ and the quadratic $Q(X)$. The root of $Q(X)$ closest to X_1 is chosen as the next approximation X_{1+1} .

4. Determining Multiple Roots

For a discussion concerning multiple roots see Chapter II, § 5.




Figure 3.1. Geometrical Interpretation of Muller's Method

CHAPTER IV

GREATEST COMMON DIVISOR METHOD

1. Derivation of the Algorithm

The greatest common divisor (g.c.d.) method reduces the problem of finding all the zeros of a polynomial, possibly having multiple zeros, to one of solving for zeros of a polynomial all of whose zeros are simple.

Consider the N^{th} degree polynomial

$$P(X) = a_1 X^N + a_2 X^{N-1} + \dots + a_N X + a_{N+1}$$

where $a_1 \neq 0$ and $a_1, a_2, \dots, a_{N+1}$ are complex numbers. If $P(X)$ has m distinct zeros, $\alpha_1, \alpha_2, \dots, \alpha_m$, then $P(X)$ can be expressed in the form

$$P(X) = a_1 (X - \alpha_1)^{e_1} (X - \alpha_2)^{e_2} \dots (X - \alpha_m)^{e_m} \quad (4-1)$$

where e_i is the multiplicity of α_i , $i = 1, 2, \dots, m$. The derivative of $P(X)$ is

$$P'(X) = N a_1 X^{N-1} + (N-1) a_2 X^{N-2} + \dots + a_N$$

which can also be expressed as

$$P'(X) = a_1 (X - \alpha_1)^{e_1-1} (X - \alpha_2)^{e_2-1} \dots (X - \alpha_m)^{e_m-1} \sum_{i=1}^m e_i \prod_{\substack{j=1 \\ j \neq i}}^m (X - \alpha_j). \quad (4-2)$$

The greatest common divisor of $P(X)$ and $P'(X)$ is obtained from the following theorem.

Theorem 4.1. Let $P(X)$ be an N^{th} degree polynomial having m distinct zeros $\alpha_1, \alpha_2, \dots, \alpha_m$ of multiplicity $e_1, e_2, \dots, e_m$ respectively. Then the polynomial

$$D(X) = (X - \alpha_1)^{e_1-1} (X - \alpha_2)^{e_2-1} \dots (X - \alpha_m)^{e_m-1}$$

is the unique monic greatest common divisor of $P(X)$ and its derivative $P'(X)$.

Proof. Since the set of all polynomials over the complex number field is a unique factorization domain and since each factor $X - \alpha_i$ is irreducible, it follows from (4-1) and (4-2) that $D(X)$ is the unique monic greatest common divisor of $P(X)$ and $P'(X)$.

It follows from Theorem 4.1 that each zero of $D(X)$ is also a zero of $P(X)$ and $P'(X)$. Hence we have the following result.

Theorem 4.2. If $P(X)$ is a polynomial, then $P(X)$ and $P'(X)$ are relatively prime if and only if $P(X)$ has no multiple zeros.

Consider the polynomial $H(X)$ obtained by dividing $P(X)$ by its monic g.c.d., $D(X)$.

$$H(X) = P(X)/D(X)$$

$$= a_1 \prod_{i=1}^m (X - \alpha_i)^{e_i} \bigg/ \prod_{i=1}^m (X - \alpha_i)^{e_i-1}$$

$$= a_1 \prod_{i=1}^m (X - \alpha_i).$$

The zeros of $H(X)$ are all simple zeros and are also all the distinct zeros of $P(X)$. Use of the g.c.d. method involves computation of $H(X)$ when given $P(X)$.

In order to obtain $H(X)$, a computational algorithm is necessary to find the g.c.d. of $P(X)$ and $P'(X)$. The general method for computing the g.c.d. of two polynomials is as follows: Let $R_0(X)$ and $R_1(X)$ be two polynomials having degrees N_0 and N_1 respectively such that $N_1 \leq N_0$. The g.c.d. of $R_0(X)$ and $R_1(X)$ is desired. By the division algorithm, there exists polynomials $S_1(X)$ and $R_2(X)$ such that

$$R_0(X) = R_1(X) S_1(X) + R_2(X)$$

where either $R_2(X) = 0$ or $\deg. R_2(X) < \deg. R_1(X)$. Similarly if $R_2(X) \neq 0$, there exists polynomials $S_2(X)$ and $R_3(X)$ such that

$$R_1(X) = S_2(X) R_2(X) + R_3(X)$$

where either $R_3(X) = 0$ or $\deg. R_3(X) < \deg. R_2(X)$. Continuing in the above manner, suppose $R_i(X)$ and $R_{i+1}(X)$ have been found where $\deg. R_{i+1}(X) < \deg. R_i(X)$. Then there exists polynomials $R_{i+2}(X)$ and $S_{i+1}(X)$ such that

$$R_i(X) = R_{i+1}(X) S_{i+1}(X) + R_{i+2}(X)$$

where either $R_{i+2}(X) = 0$ or $\deg. R_{i+2}(X) < \deg. R_{i+1}(X)$. Then we obtain a sequence $R_0(X), R_1(X), \dots, R_K(X), R_{K+1}(X)$ such that $\deg. R_i(X) < \deg. R_{i-1}(X)$, $i = 1, 2, \dots, K+1$. Since a polynomial cannot have degree less than zero, the above process, in a finite number of steps (at most N_1), results in polynomials $R_{K-1}(X)$, $S_K(X)$ and $R_K(X)$ with $\deg. R_K(X) < \deg. R_{K-1}(X)$ such that

$$R_{K-1}(X) = R_K(X) S_K(X) + R_{K+1}(X)$$

and $R_{K+1}(X) = 0$.

Theorem 4.3. Let the sequence $R_0(X), R_1(X), \dots, R_K(X), R_{K+1}(X)$ be defined as above. Then $R_K(X)$ is the greatest common divisor of $R_0(X)$ and $R_1(X)$.

Proof. It is clear that $R_K(X)$ divides $R_{K-1}(X)$. If $R_K(X)$ divides $R_i(X)$ for $0 \leq j < i \leq k$, then $R_j(X) = R_{j+1}(X) S_{j+1}(X) + R_{j+2}(X)$. Thus, $R_K(X)$ divides $R_j(X)$ and it follows by induction that $R_K(X)$ divides both $R_0(X)$ and $R_1(X)$. By reversing the inductive argument given above, it is easy to see that if $L(X)$ divides $R_0(X)$ and $R_1(X)$, then $L(X)$ divides $R_i(X)$ for $i = 0, 1, \dots, K$. Therefore, $L(X)$ divides $R_K(X)$ which shows that $R_K(X)$ is the greatest common divisor of $R_0(X)$ and $R_1(X)$.

The above theorem tells how to obtain the greatest common divisor of two polynomials. A machine oriented method is now developed for computing the sequence of $R_j(X)$'s. Beginning the sequence with $R_0(X)$ and $R_1(X)$, the polynomial $R_{i+1}(X)$ of the sequence is derived from $R_i(X)$

and $R_{i-1}(X)$ as follows: Let $R_{i-1}(X)$ of degree N_{i-1} be given by

$$\begin{aligned} R_{i-1}(X) \\ = r_{i-1,1} X^{N_{i-1}} + r_{i-1,2} X^{N_{i-1}-1} + \dots + r_{i-1,N_{i-1}} X + r_{i-1,N_{i-1}+1} \end{aligned}$$

and $R_i(X)$ of degree N_i be given by

$$R_i(X) = r_{i,1} X^{N_i} + r_{i,2} X^{N_i-1} + \dots + r_{i,N_i} X + r_{i,N_i+1}$$

where $N_i \leq N_{i-1}$. Define $U_1(X)$ by

$$U_1(X) = (r_{i-1,1} / r_{i,1}) X^{N_{i-1}-N_i}.$$

Then define $T_1(X)$ by

$$\begin{aligned} T_1(X) &= R_{i-1}(X) - U_1(X) R_i(X) \\ &= [r_{i-1,1} - r_{i,1} (r_{i-1,1} / r_{i,1})] X^{N_{i-1}} \\ &\quad + [r_{i-1,2} - r_{i,2} (r_{i-1,1} / r_{i,1})] X^{N_{i-1}-1} \\ &\quad + \dots \\ &\quad + [r_{i-1,N_{i-1}+1} - r_{i,N_{i-1}+1} (r_{i-1,1} / r_{i,1})] \end{aligned}$$

where $r_{i,j} = 0$ for $j > N_i+1$.

We consider three cases.

- (1) If $T_1(X) = 0$, then $R_i(X) = R_K(X)$; that is, $R_i(X)$ is the g.c.d. of $R_0(X)$ and $R_1(X)$.
- (2) If $T_1(X) \neq 0$ and $\deg. T_1(X) < N_i$, then $R_{i+1}(X) = T_1(X)$.

- (3) If $T_1(X) \neq 0$ and $\deg. T_1(X) = M_1 \geq N_i$, then define $U_2(X)$ by

$$U_2(X) = (t_{1,1} / r_{i,1}) X^{M_1 - N_i}$$

where

$$T_1(X) = t_{1,1} X^{M_1} + t_{1,2} X^{M_1-1} + \dots + t_{1,M_1} X + t_{1,M_1+1}.$$

Define $T_2(X) = T_1(X) - U_2(X) R_i(X)$ which can be expressed by

$$\begin{aligned} T_2(X) &= [t_{1,1} - (t_{1,1} / r_{i,1}) r_{i,1}] X^{M_1} \\ &\quad + [t_{1,2} - (t_{1,1} / r_{i,1}) r_{i,2}] X^{M_1-1} \\ &\quad + \dots \\ &\quad + [t_{1,M_1+1} - (t_{1,1} / r_{i,1}) r_{i,M_1+1}] \end{aligned}$$

where $r_{i,j} = 0$ for $j > N_i+1$. We again consider the following three cases.

- (1) If $T_2(X) = 0$, then $R_i(X)$ is the g.c.d. of $R_0(X)$ and $R_1(X)$.
- (2) If $T_2(X) \neq 0$ and $\deg. T_2(X) < \deg. R_i(X)$, then $R_{i+1}(X) = T_2(X)$.
- (3) If $T_2(X) \neq 0$ and $\deg. T_2(X) = M_2 \geq N_i$, then define $U_3(X)$ by

$$U_3(X) = (t_{2,1} / r_{i,1}) X^{M_2 - N_i}$$

where

$$T_2(X) = t_{2,1} X^{M_2} + t_{2,2} X^{M_2-1} + \dots + t_{2,M_2} X + t_{2,M_2+1}.$$

Since $\deg. T_{i+1}(X) < \deg. T_i(X)$, then this process is finite (not to exceed N_{i-1}) ending, for some integer S , in $T_S(X)$ such that

- (1) $T_S(X) = 0$ and $R_i(X)$ is the g.c.d. of $R_0(X)$ and $R_1(X)$ or
- (2) $T_S(X) \neq 0$ but $\deg. T_S(X) < \deg. R_i(X)$, in which case
 $T_S(X) = R_{i+1}(X).$

Thus, using this algorithm and given $R_0(X)$ and $R_1(X)$, the sequence $R_0(X), R_1(X), R_2(X), \dots, R_i(X), R_{i+1}(X)$ can be generated such that either

- (1) $R_{i+1}(X) = 0$ and $R_i(X)$ is the g.c.d. of $R_0(X)$ and $R_1(X)$ or
- (2) $R_{i+1}(X) \neq 0$ and $N_{i+1} < N_i$. In a finite number of iterations, $R_K(X)$, the g.c.d. of $R_0(X)$ and $R_1(X)$, can be obtained.

Recall that we wanted to obtain the polynomial $H(X) = P(X)/D(X)$ where $D(X)$ is the g.c.d. of $P(X)$ and $P'(X)$. Thus, after obtaining $D(X)$ by the above algorithm, it is necessary to divide $P(X)$ by $D(X)$ obtaining $H(X)$ all whose zeros are simple.

Once $H(X)$ is obtained, an appropriate method such as Newton's method or Muller's method is applied to extract the zeros of $H(X)$. This gives all the zeros of $P(X)$.

As in Newton's or Muller's method, the zeros may be checked for accuracy and possibly improved by using them as initial approximations with the particular method applied to the full (undeflated) polynomial, $P(X)$.

2. Determining Multiplicities

After all zeros of $P(X)$ are found, the multiplicity of each zero can be determined by the process outlined in Chapter II, § 5.

3. Procedure for the G.C.D. Method

The basic steps performed by the greatest common divisor method are listed sequentially as follows:

1. Given a polynomial, $P(X)$, in the form

$$P(X) = a_1X^N + a_2X^{N-1} + \dots + a_NX + a_{N+1}.$$
2. Calculate the derivative, $P'(X)$, of $P(X)$ in the form

$$P'(X) = b_1X^{N-1} + b_2X^{N-2} + \dots + b_N$$
 where $b_1 = Na_1$,
 $b_2 = (N-1)a_2, \dots, b_N = a_N.$
3. Find $D(X)$, the g.c.d. of $P(X)$ and $P'(X)$, using the algorithms developed above.
4. Calculate $H(X) = P(X)/D(X)$, the polynomial having only simple zeros.
5. Use some appropriate method to extract the zeros of $H(X)$.
6. Determine the multiplicity of each of the zeros obtained in step 5.

CHAPTER V

REPEATED GREATEST COMMON DIVISOR METHOD

1. Derivation of the Algorithm

The repeated greatest common divisor (repeated g.c.d.) method makes repeated use of the g.c.d. method to extract the zeros and their multiplicities of a polynomial with complex coefficients. That is, the repeated g.c.d. method reduces the problem of finding the zeros of a polynomial, $P(X)$, which possibly has multiple zeros, to one of finding the zeros of a polynomial which has only simple zeros and the zeros of this polynomial are all the zeros of $P(X)$ of a given multiplicity.

Let

$$\begin{aligned} P(X) &= a_1 X^N + a_2 X^{N-1} + \dots + a_N X + a_{N+1} \\ &= a_1 (X - \alpha_1)^{e_1} (X - \alpha_2)^{e_2} \dots (X - \alpha_m)^{e_m} \end{aligned}$$

where $a_1 \neq 0$, each a_i is a complex number, and $\alpha_1, \alpha_2, \dots, \alpha_m$ are the distinct zeros of $P(X)$ having multiplicity $e_1, e_2, \dots, e_m$, respectively. If $D_1(X)$ is the monic greatest common divisor of $P(X)$ and $P'(X)$, then Theorem 4.1 shows that

$$D_1(X) = (X - \alpha_1)^{e_1-1} (X - \alpha_2)^{e_2-1} \dots (X - \alpha_m)^{e_m-1}$$

where we assume that if $e_j = 1$, then $X - \alpha_j$ does not appear in the

representation. Let $D_2(X)$ be the monic greatest common divisor of $D_1(X)$ and $D_1'(X)$. Then

$$D_2(X) = (X - \alpha_1)^{e_1-2} (X - \alpha_2)^{e_2-2} \dots (X - \alpha_m)^{e_m-2}$$

where we assume that if $e_j \leq 2$, then $X - \alpha_j$ does not appear in the representation. From the above it is clear that the zeros of $D_1(X)$ are just the multiple zeros of $P(X)$ to one lower power. The zeros of $D_2(X)$ are just the multiple zeros of $D_1(X)$ to one lower power. Thus, the zeros of $D_2(X)$ are just the zeros of $P(X)$ which have multiplicity greater than two, and their multiplicity in $D_2(X)$ is reduced by two. Therefore, it follows that

$$G_1(X) = [P(X)/D_1(X)]/[D_1(X)/D_2(X)] = P(X)D_2(X)/[D_1(X)]^2$$

has only simple zeros and they are just the simple zeros of $P(X)$. In general if $D_j(X)$ has been defined for $1 \leq j \leq i$ and if $D_{i+1}(X)$ is the monic greatest common divisor of $D_i(X)$ and $D_i'(X)$, then the zeros of $D_{i+1}(X)$ are the multiple zeros of $D_i(X)$ to one lower power. Thus, the zeros of $D_{i+1}(X)$ are just the zeros of $P(X)$ which have multiplicity greater than $i+1$ and their multiplicity in $D_{i+1}(X)$ is reduced by $i+1$. It follows that

$$\begin{aligned} G_i(X) &= [D_{i-1}(X)/D_i(X)]/[D_i(X)/D_{i+1}(X)] \\ &= D_{i-1}(X) D_{i+1}(X)/[D_i(X)]^2 \end{aligned}$$

has simple zeros and they are just the zeros of $P(X)$ that have multiplicity i . Thus, we have proven the following theorem.

Theorem 5.1. Let $P(X) = a_1X^N + a_2X^{N-1} + \dots + a_NX + a_{N+1}$ where $a_1 \neq 0$ and $a_1, a_2, \dots, a_{N+1}$ are complex numbers. If $D_0(X) = P(X)$ and if $D_{i+1}(X)$ is the monic greatest common divisor of $D_i(X)$ and $D_i'(X)$ for $i \geq 0$, then

$$G_i(X) = D_{i-1}(X) D_{i+1}(X) / [D_i(X)]^2$$

has only simple zeros and they are just the zeros of $P(X)$ that have multiplicity i .

Thus, by the above theorem we can generate a sequence of polynomials $G_1(X), G_2(X), \dots, G_K(X)$ where the set of zeros of $P(X)$ is the same as the set of zeros of this sequence and the multiplicity of each zero in $P(X)$ is given by the corresponding subscript on $G(X)$. Therefore, by using a method such as Newton's method or Muller's method to calculate the zeros of each $G_i(X)$, we will have the zeros of $P(X)$ along with their multiplicities.

2. Procedure for the Repeated G.C.D. Method

The basic steps performed by the greatest common divisor method are listed sequentially as follows:

1. Given a polynomial, $P(X)$, in the form

$$P(X) = a_1X^N + a_2X^{N-1} + \dots + a_NX + a_{N+1}.$$

2. Set $D_0(X) = P(X)$.

3. Calculate the derivative, $D_0'(X)$, of $D_0(X)$ in the form

$$D_0'(X) = b_1X^{M-1} + b_2X^{M-2} + \dots + b_M$$

where $\deg. D_0(X) = M$, $D_0(X) = d_1 X^M + \dots + d_{M+1}$,
 and $b_1 = M d_1$, $b_2 = (M-1) d_2$, $\dots$, $b_M = d_M$.

4. Find $D_1(X)$, the g.c.d. of $D_0(X)$ and $D_0'(X)$ using the algorithms developed in Chapter IV.
5. Similar to 3., calculate $D_1'(X)$.
6. Find $D_2(X)$, the g.c.d. of $D_1(X)$ and $D_1'(X)$ using the algorithms developed in Chapter IV.
7. Calculate $G(X) = D_0(X) D_2(X) / [D_1(X)]^2$.
8. Use some appropriate method to extract the zeros of $G(X)$ and assign these zeros the correct multiplicity as zeros of $P(X)$.
9. Set $D_0(X) = D_1(X)$, $D_0'(X) = D_1'(X)$, and $D_1(X) = D_2(X)$.
 Then repeat 5.-8. above until all the zeros of $P(X)$ are found.

CHAPTER VI

CONCLUSION

In order to compare Newton's, Muller's, the greatest common divisor, and the repeated greatest common divisor methods, we consider the polynomials as being divided into the following classes:

1. polynomials with all distinct zeros.
2. polynomials with multiple zeros.

The comparisons in the following material are results of tests made on the IBM 360/50 computer which has a 32 bit word. The programs were successfully run on the CDC 6600 and the UNIVAC 1108 which have a 60 bit word and a 36 bit word respectively. It was noted that the UNIVAC 1108 is about 15 times faster than the IBM 360/50. The CDC 6600 is faster than the UNIVAC 1108 but the difference is not as great as that between the UNIVAC 1108 and the IBM 360/50.

1. Polynomials With all Distinct Zeros

First we consider the class of polynomials having distinct zeros. Newton's method is particularly suited for this class of polynomials. Its quadratic convergence is very fast which can save time and money to the user. The accuracy obtained is excellent as shown in Exhibit 6.1 which presents the zeros of a 15th degree polynomial in double precision. In most cases, the method produces convergence for almost any initial approximation given.

Muller's method also produces good results on this class of polynomials. The rate of convergence is, however, somewhat slower than Newton's method. This fact is especially significant when working with polynomials of high degree. The accuracy obtained by Muller's method is comparable to, but does not exceed that of Newton's method. In most cases, the accuracy of the two methods does not differ by more than one or two decimal places. Exhibit 6.2 shows results of Muller's method for the polynomial of Exhibit 6.1. As in Newton's method, convergence is produced for almost any initial approximation given.

The g.c.d. method, whether used with Newton's or Muller's method as a supporting method on this class of polynomials, is no better than Newton's or Muller's method alone. The reason for this is that the greatest common divisor of the polynomial, $P(X)$, and its derivative is 1. Thus, $H(X) = P(X)/\text{g.c.d. } P(X) = P(X)$; that is, the polynomial solved by the supporting method is the same as the original polynomial. Thus, in this case the g.c.d. method will not produce better results than the supporting method used alone. The above comments also hold for the repeated g.c.d. method.

Thus, this class of polynomials presents no difficulty to any of these four methods. Newton's method, because of its speed, is therefore recommended.

2. Polynomials With Multiple Zeros

Next consider the class of polynomials containing multiple zeros. Exhibits 6.3 - 6.26 illustrate output from six different programs using the methods described in Chapters II - V. Four polynomials are used where the zeros of these polynomials are listed below. The number in

parentheses indicates the multiplicity of that zero.

<u>Polynomial #1</u>	<u>Polynomial #2</u>	<u>Polynomial #3</u>	<u>Polynomial #4</u>
2+2i (3)	-2.33 (1)	2+2i (3)	1+i (6)
1+2i (2)	.003 (2)	1+2i (2)	1-i (6)
-1+.5i (1)	i (2)	-1+.5i (3)	
	1.5i (2)		
	-1.5i (2)		
	3i (3)		
	-1-i (3)		

Note the relationship between polynomials #1 and #3.

This class presents considerable difficulty for Newton's method, especially those polynomials containing zeros of high multiplicity or containing a considerable number of multiple zeros. The iteration formula for Newton's method is

$$X_{n+1} = X_n - P(X_n)/P'(X_n).$$

If c is a multiple zero then $P(c) = P'(c) = 0$. Hence, as $X_n \rightarrow c$, $P(X_n) \rightarrow 0$ and $P'(X_n) \rightarrow 0$ and the iteration formula may be unstable, resulting in no convergence or bad accuracy. As the number of multiple zeros increases, the polynomial becomes more ill-conditioned, convergence becomes more difficult, and accuracy is lost. Thus, the possibility of convergence decreases. This also holds true if the multiplicities of the zeros are increased. The rate of convergence of Newton's method is much slower for multiple zeros than for distinct zeros. Exhibit 6.3 shows a polynomial (#1) containing two multiple zeros solved in double precision. Note the following from Exhibit 6.3.

1. Roots #2 and #3 are greatly improved by iterating on the original polynomial. Distinct roots are usually improved in this manner.

2. The time taken to solve this 6th degree equation with multiple roots is greater than the time taken by the same program to solve a 15th degree polynomial with all distinct roots (Exhibit 6.1).
3. Root #2 did not pass the convergence test after 200 iterations even though it was improved. This is probably due to the fact that the polynomial from which root 2 was extracted had only one multiple root but the original polynomial from which it was extracted the second time had two multiple roots; that is, the original polynomial is more ill-conditioned.
4. The accuracy of the roots before the attempt to improve accuracy is very poor. Root #2 is accurate to only three decimal places as compared to the 15 decimal places in Exhibit 6.1 for distinct roots. Root #3 is especially bad, the imaginary part being accurate to only one decimal place.

Exhibit 6.4 uses polynomial #2. Note the poor results obtained before the attempt to improve accuracy and the improvement afterward. Also note that after the attempt to improve accuracy, one of the zeros, namely 3i, is lost and an extra zero, namely 1.5, is included in the list. (See Appendix A, § 4.) A convergence requirement of 10^{-5} was used on this polynomial to get it to converge to all of the zeros in a maximum number of 200 iterations.

In many cases, Newton's method fails to converge altogether. Polynomial #3 could not be solved using Newton's method with a maximum

number of 200 iterations and a convergence requirement of 10^{-9} . Exhibit 6.5 illustrates the bad results for a convergence requirement of 10^{-5} which was needed in order to get convergence. In Exhibit 6.6 a convergence requirement of 10^{-3} was needed in order to get convergence to the zeros of polynomial #4.

Muller's method also encounters difficulty, although to a lesser degree than Newton's method, on this class of polynomials. In most cases, Muller's method produces convergence even when Newton's method completely fails. Newton's method completely failed for polynomials #3 and #4 with a convergence requirement of 10^{-9} but convergence was obtained using Muller's method as shown in Exhibits 6.9 and 6.10. The accuracy obtained by Muller's method is not good but usually better than Newton's method using the same convergence requirement. The rate of convergence of Muller's method is considerably slower for multiple zeros than for distinct zeros. However, for multiple zeros, Muller's method is as fast or faster than Newton's.

The g.c.d. method is perfectly suited for polynomials with multiple zeros. All multiple zeros are removed leaving only a polynomial of class 1 (all distinct roots) to be solved. This indicates that best results should be obtained by using Newton's method as the supporting method, since Newton's method enjoys the advantage of speed over Muller's method for distinct zeros. This has indeed proved to be true. The accuracy of the zeros obtained decreases, somewhat, when the number of multiple zeros is increased. This is due to accuracy lost in computing the g.c.d. and the quotient polynomial and not as a result of the supporting method. It is easy to see that the accuracy of the g.c.d. method is best when the degree of the greatest common divisor of

$P(X)$ and $P'(X)$ is maximum. This is due to the fact that the error in the greatest common divisor is minimized in this case. The accuracy obtained using Newton's method and Muller's method as supporting methods is about the same. This is verified by Exhibits 6.11 - 6.14 (g.c.d. method with Newton) and Exhibits 6.15 - 6.18 (g.c.d. method with Muller).

Multiplicities are determined with excellent accuracy. The g.c.d. method is not as sensitive to zeros of high multiplicity or polynomials containing a large number of multiple zeros as are both Newton's and Muller's methods. A quick comparison of Exhibits 6.11 - 6.14 and 6.15 - 6.18 with Exhibits 6.3 - 6.6 and 6.7 - 6.10 show that the g.c.d. method with either supporting method is much more accurate than either Newton's or Muller's method. For example, Exhibits 6.5 and 6.9 show polynomial #3 for which Newton's method and Muller's method both gave poor convergence. But Exhibits 6.13 and 6.17 show very accurate results for polynomial #3.

The repeated g.c.d. method is also suited very well for polynomials with multiple zeros. Exhibits 6.19 - 6.22 and Exhibits 6.23 - 6.26 are results of the repeated g.c.d. method with Newton's method and Muller's method as supporting methods, respectively. However, the results of the repeated g.c.d. method are not as good as those obtained from the g.c.d. method. Since the repeated g.c.d. method repeated uses the g.c.d. algorithm, the error tends to build up in this method when a polynomial has several zeros of different multiplicities. This can be observed by comparing Exhibits 6.20 and 6.24 with Exhibits 6.12 and 6.16 on polynomial #2 and by comparing Exhibits 6.21 and 6.25 with Exhibits 6.13 and 6.17 on polynomial #3. As was the case of the g.c.d.

method, there is little difference between the repeated g.c.d. method with Newton's method or Muller's method as a supporting method. This can be observed by comparing Exhibits 6.19 - 6.22 (Newton) with Exhibits 6.23 - 6.26 (Muller). Even though the results of the repeated g.c.d. method are not quite as good as the results of the g.c.d. method, they are far superior to the results of both Newton's method and Muller's method.

Table 6.I gives a comparison of the execution times of the six methods for polynomials #1 - #4.

TABLE 6.I

<u>METHOD</u>	<u>EXECUTION TIME*</u>
Newton	104.16 seconds
Muller	96.79 seconds
G.C.D. with Newton	7.51 seconds
G.C.D. with Muller	8.91 seconds
Repeated G.C.D. with Newton	7.71 seconds
Repeated G.C.D. with Muller	15.16 seconds

It is clear from Table 6.I that the g.c.d. and the repeated g.c.d. methods are much faster than both Newton's and Muller's method on

* These times are from execution runs on the IBM 360/50 WATFOR system.

polynomials with multiple zeros. Therefore, for polynomials with multiple zeros, the order in which the methods are recommended is as follows.

1. G.C.D. with Newton.
2. G.C.D. with Muller.
3. Repeated G.C.D. with Newton.
4. Repeated G.C.D. with Muller.
5. Muller.
6. Newton.

NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 7 OF DEGREE 15

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.3000000000000000 01 + 0.0000000000000000 00 I
P(2) = -0.1790000000000000 02 + -0.0000000000000000 00 I
P(3) = 0.2010000000000000 02 + -0.6576000000000000 02 I
P(4) = 0.1745000000000000 03 + 0.2843680000000000 03 I
P(5) = -0.7594200000000000 03 + 0.3118080000000000 03 I
P(6) = 0.8274360000000000 03 + -0.3069040000000000 04 I
P(7) = 0.1329084000000000 04 + 0.4710624000000000 04 I
P(8) = -0.5611860000000000 04 + -0.1674576000000000 04 I
P(9) = 0.7224756000000000 04 + -0.1548288000000000 04 I
P(10) = -0.2276992000000000 04 + 0.3046320000000000 04 I
P(11) = -0.1241472000000000 04 + -0.4097280000000000 04 I
P(12) = 0.5402800000000000 04 + 0.1263488000000000 04 I
P(13) = -0.6468336000000000 04 + 0.1236480000000000 04 I
P(14) = 0.1467456000000000 04 + 0.2272000000000000 02 I
P(15) = -0.1077120000000000 03 + -0.5675840000000000 03 I
P(16) = 0.3456000000000000 02 + 0.1267200000000000 03 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.3000000000000000 00 + 0.1609358115166531D-16 I	1	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.2000000000000000 00 + -0.2000000000000000 00 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = -0.2000000000000000 00 + 0.2000000000000000 00 I	1	0.3682284792654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = 0.2485026109655803D-17 + 0.1000000000000000 00 I	1	-0.5176382551966724D 00 + 0.1931851608368755D 01 I
ROOT(5) = -0.1000000000000000 01 + -0.5321953706459946D-16 I	1	-0.1767767147080701D 01 + 0.1767766758852015D 01 I
ROOT(6) = -0.1000000000000000 01 + -0.9999999999999999D 00 I	1	-0.2897777583074990D 01 + 0.7764567463987070D 00 I
ROOT(7) = -0.3100918759280349D-15 + -0.9999999999999999D 00 I	1	-0.3380740248331229D 01 + -0.9058671940816160D 00 I
ROOT(8) = -0.2000000000000000 01 + -0.3000000000000000 00 I	1	-0.2828426607107896D 01 + -0.2828427642384390D 01 I
ROOT(9) = 0.1000000000000000 16D 01 + -0.1003764996352100D-12 I	1	-0.1164684801399899D 01 + -0.434666459873368D 01 I
ROOT(10) = 0.2000000000000000 01 + -0.1000000000000032D 01 I	1	0.1294096345098645D 01 + -0.4829428831453027D 01 I
ROOT(11) = 0.1899999999999980D 01 + -0.1385507847222020D-12 I	1	0.3689089292979509D 01 + -0.3689086300072258D 01 I
ROOT(12) = 0.3000000000000000 01 + -0.1417549187449219D-13 I	1	0.579555539512303D 01 + -0.1552912644264674D 01 I
ROOT(13) = 0.3999999999999980D 01 + 0.400000000000003D 01 I	1	0.4278517377461250D 01 + 0.1662325708247752D 01 I
ROOT(14) = 0.9999999999999960D 00 + 0.1000000000000000D 00 I	1	SOLVED BY DIRECT METHOD
ROOT(15) = -0.333333333333334D 01 + -0.1650371707706594D-15 I	1	SOLVED BY DIRECT METHOD

Exhibit 6.1.

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF PIX) ARE

ROOTS OF PIX)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.3000000000000000 00 + 0.3328943537705913D-16 i	1	0.4829629115656279D 00 + 0.1294095286438187D 00 i
ROOT(2) = 0.2000000000000000 00 + 0.2000000000000000 00 i	1	0.7071067553044346D 00 + 0.7071068070664595D 00 i
ROOT(3) = -0.2000000000000000 00 + 0.2000000000000000 00 i	1	0.3882284792654056D 00 + 0.16888763117193D 01 i
ROOT(4) = -0.1126043043762753D-17 + 0.1000000000000000 01 i	1	-0.5176382551964724D 00 + 0.1931231608368755D 01 i
ROOT(5) = -0.1000000000000000 01 + 0.1882474030530796D-16 i	1	-0.174727167080761D 01 + 0.176169726852015D 01 i
ROOT(6) = -0.1000000000000000 01 + 0.1000000000000000 01 i	1	-0.2897771883074980D 01 + 0.7765517463987070D 00 i
ROOT(7) = 0.6131849319379140D-16 + 0.1000000000000000 01 i	1	-0.3380740248331229D 01 + 0.905882164298160D 00 i
ROOT(8) = 0.1000000000000000 01 + 0.3600000000000000 01 i	1	-0.2828426607107856D 01 + 0.282827164298160D 00 i
ROOT(9) = 0.2000000000000000 01 + 0.728776548089504D-16 i	1	-0.1184684801399890D 01 + 0.436666646873349D 01 i
ROOT(10) = 0.2000000000000000 01 + 0.999999999999999D 00 i	1	0.1294095634509845D 01 + 0.436666646873349D 01 i
ROOT(11) = 0.2000000000000000 01 + 0.3968039620811357D-14 i	1	0.3889088292979509D 01 + 0.436666646873349D 01 i
ROOT(12) = 0.2899999999999997D 01 + 0.2896889630437848D-14 i	1	0.5795555393512303D 01 + 0.1552912644268974D 01 i
ROOT(13) = 0.4000000000000000 01 + 0.4000000000000000 01 i	1	0.6278517357734125D 01 + 0.1552912644268974D 01 i
ROOT(14) = 0.999999999999999D 00 + 0.1000000000000000 01 i	1	SOLVED BY DIRECT METHOD
ROOT(15) = -0.3333333333333333D 01 + 0.8998063670517638D-16 i	1	SOLVED BY DIRECT METHOD

Exhibit 6.1, Roots Are: $-1 - i$, $1 + i$, $-2 - 3i$, $2 - i$, 3 , 2 , i , $-i$,
 $-10/3$, $.3$, -1 , 1 , $4 + 4i$, $-.2 + .2i$, $.2 - .2i$.

MULLERS METHOD FOR FINDING THE ZEROS OF A POLYNOMIAL
POLYNOMIAL NUMBER 7 OF DEGREE 15

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.3000000000000000 01 + 0.0000000000000000 00 I
P(2) = -0.1790000000000000 02 + -0.0000000000000000 00 I
P(3) = 0.2010000000000000 02 + -0.6576000000000000 02 I
P(4) = 0.1745000000000000 03 + 0.2843680000000000 03 I
P(5) = -0.7594200000000000 03 + 0.3118080000000000 03 I
P(6) = -0.8274360000000000 03 + -0.3069040000000000 04 I
P(7) = -0.1329084000000000 04 + 0.4710624000000000 04 I
P(8) = -0.5611860000000000 04 + -0.1674576000000000 04 I
P(9) = 0.7224756000000000 04 + -0.1548298000000000 04 I
P(10) = -0.7274992000000000 04 + -0.3046320000000000 04 I
P(11) = -0.1241472000000000 04 + -0.4097280000000000 04 I
P(12) = -0.5402800000000000 04 + 0.1263448000000000 04 I
P(13) = -0.6468336000000000 04 + 0.1236448000000000 04 I
P(14) = 0.1467456000000000 04 + 0.2272000000000000 02 I
P(15) = -0.1077120000000000 03 + -0.5475840000000000 03 I
P(16) = 0.3456000000000000 02 + 0.1267200000000000 03 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR MULTIPLICITIES. 0.10D-01
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

BEFORE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)

ROOT(1) = 0.3000000000000000 00 + 0.3383998751751856D-16 I
ROOT(2) = 0.2000000000000000 00 + -0.2000000000000000 00 I
ROOT(3) = 0.9999999999999999 00 + 0.9999999999999999 00 I
ROOT(4) = -0.2389082408313382D-16 + 0.1000000000000000 01 I
ROOT(5) = -0.1000000000000000 01 + 0.2531629277617640D-15 I
ROOT(6) = -0.3533333333333334 01 + -0.2144722669087737D-15 I
ROOT(7) = -0.1000000000000000 01 + -0.9999999999999999 01 I
ROOT(8) = -0.2000000000000000 01 + -0.3000000000000000 01 I
ROOT(9) = -0.1139816730936650D-12 + -0.9999999999999999 00 I
ROOT(10) = 0.1999999999999999 01 + -0.9999999999999999 00 I
ROOT(11) = 0.2999999999999999 01 + 0.192189120439124D-14 I
ROOT(12) = 0.200000000000122D 01 + -0.2254154879580170D-12 I
ROOT(13) = 0.3999999999999999 01 + 0.400000000000005D 01 I
ROOT(14) = 0.100000000000208D 01 + 0.2297444641937337D-12 I
ROOT(15) = -0.2000000000000366D 00 + 0.2000000000000595D 00 I

MULTIPLICITIES

1
1
1
1
1
1
1
1
1
1
1
1
1
1
1

INITIAL APPROXIMATION

0.4829629115636279D 00 + 0.1294095284438187D 00 I
0.7071067553046346D 00 + 0.7071068070684595D 00 I
0.3882284792654036D 00 + 0.1448888763117193D 01 I
-0.5176382551966724D 00 + 0.1931851408368755D 01 I
-0.1767761437080701D 01 + 0.176766758852015D 01 I
-0.38977758304990D 01 + 0.7784587463967070D 00 I
-0.3380740248331229D 01 + -0.9058671940916160D 00 I
-0.282842860707896D 01 + -0.2828427642384390D 01 I
-0.1164684801399899D 01 + -0.4346666459873368D 01 I
0.1294096345098645D 01 + -0.4829628831453027D 01 I
0.3889088297979509D 01 + -0.3889086300072258D 01 I
0.5795555393512303D 01 + -0.1552912644268974D 01 I
0.6278517357734125D 01 + 0.1682325708247752D 01 I
SOLVED BY DIRECT METHOD
SOLVED BY DIRECT METHOD

Exhibit 6.2.

AFTER THE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
R00T(1) = 0.30000000000000000000 00 + 0.33767394620632440-16 i	1	0.48296291156562790 00 + 0.12940952844381870 00 i
R00T(2) = 0.20200000000000000000 00 + -0.20000000000000000000 00 i	1	0.70710675530463460 00 + 0.70710680706845950 00 i
R00T(3) = 0.99999999999999999999 00 + 0.10000000000000000000 01 i	1	0.38822847926540560 00 + 0.14488887631171930 01 i
R00T(4) = 0.1153775751536680-16 + 0.10000000000000000000 01 i	1	-0.51763825519667240 00 + 0.19318516083687550 01 i
R00T(5) = -0.10000000000000000000 01 + -0.99356260300168480-17 i	1	-0.17677671470807010 01 + 0.1767667588520150 01 i
R00T(6) = -0.33333333333333340 01 + -0.19191253971536640-15 i	1	-0.2897775830749900 01 + 0.77645674639870700 00 i
R00T(7) = -0.10000000000000000000 01 + -0.99999999999999999999 00 i	1	-0.33837407483117290 01 + -0.90586719408161600 00 i
R00T(8) = -0.20000000000000000000 01 + -0.30000000000000000000 01 i	1	-0.2828426607078960 01 + -0.28284276423843900 01 i
R00T(9) = 0.55291352338277240-16 + -0.10000000000000000000 01 i	1	-0.11646848013998990 01 + -0.43466664598733680 01 i
R00T(10) = 0.20000000000000000000 01 + -0.99999999999999999999 00 i	1	0.12940963450986450 01 + -0.48296288314530270 01 i
R00T(11) = 0.29999999999999999999 01 + 0.24394773729111440-14 i	1	0.38890882929795090 01 + -0.38890863000722540 01 i
R00T(12) = 0.20000000000000000000 01 + -0.41020504900166570-14 i	1	0.57955553935123030 01 + -0.15529126442689740 01 i
R00T(13) = 0.40000000000000000000 01 + 0.40000000000000000000 01 i	1	0.62785173577341259 01 + 0.16823257082477520 01 i
R00T(14) = 0.10000000000000000000 01 + 0.76824382838570700-16 i	1	SOLVED BY DIRECT METHOD
R00T(15) = -0.20000000000000000000 00 + 0.20000000000000000000 00 i	1	SOLVED BY DIRECT METHOD

Exhibit 6.2. Roots Are: $-1 - i, 1 + i, -2 - 3i, 2 - i, -i, -10/3, .3, -1, 1, 4 + 4i, -.2 + .2i, .2 - .2i$.

NEWTON'S METHOD TO FIND ZEROS OF POLYNOMIALS POLYNOMIAL NUMBER 1 OF DEGREE 6

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(2) = -0.7000000000000000 01 + -0.1050000000000000 02 I
P(3) = -0.2800000000000000 02 + 0.5800000000000000 02 I
P(4) = 0.1710000000000000 03 + 0.1500000000000000 01 I
P(5) = -0.7300000000000000 02 + -0.2510000000000000 03 I
P(6) = -0.2280000000000000 03 + 0.1040000000000000 03 I
P(7) = 0.7200000000000000 02 + 0.1040000000000000 03 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.9999998886125019D 00 + 0.2000000052138284D 01 I	2	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.1996737810257486D 01 + 0.1995253821143684D 01 I	3	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = -0.9902131979974624D 00 + 0.5142384322923812D 00 I	1	SOLVED BY DIRECT METHOD

IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(2) = 0.1996737810257486D 01 + 0.1995253821143684D 01 I DID NOT CONVERGE.
THE PRESENT APPROXIMATION AFTER 200 ITERATIONS IS PRINTED BELOW.

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.9999998886125019D 00 + 0.2000000052138284D 01 I	2	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.19999992907503309D 01 + 0.1999959474001489D 01 I	3	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = -0.999999999999999D 00 + 0.500000000000000D 00 I	1	SOLVED BY DIRECT METHOD

Exhibit 6.3. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$

NEWTON'S METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 2 OF DEGREE 15

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.4800000000000000 02 + 0.0000000000000000 00 I
P(2) = 0.2557120000000000 03 + -0.3840000000000000 03 I
P(3) = -0.7353568000000000 02 + -0.2189696000000000 04 I
P(4) = -0.3855565960000000 04 + -0.6946851456000000 04 I
P(5) = -0.1733386464800000 05 + -0.1420625972800000 05 I
P(6) = -0.4967989270400000 05 + -0.1765857464000000 05 I
P(7) = -0.1022394522130000 06 + -0.6030664232000000 04 I
P(8) = -0.1642742200560000 06 + -0.4137366230400000 05 I
P(9) = -0.2036625888420000 06 + -0.1093999227670000 06 I
P(10) = -0.1871255780010000 06 + -0.1929865440330000 06 I
P(11) = -0.1274997298590000 06 + -0.2171341227420000 06 I
P(12) = -0.2814692716800000 05 + -0.1928489727960000 06 I
P(13) = -0.1329434348000000 05 + -0.1038130226550000 06 I
P(14) = -0.3052900774700000 05 + -0.2998989141300000 05 I
P(15) = -0.1835889020000000 03 + -0.1827432160000000 03 I
P(16) = 0.2755620000000000 00 + 0.2755620000000000 00 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-05
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.300000007563981848-02 + 0.2609394904552981D-08 I	2	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.8626623124099139D-04 + 0.100008017322988D 01 I	2	0.7071067553048346D 00 + 0.7071066070664595D 00 I
ROOT(3) = 0.5768342464927569D-02 + 0.1508145174316570D 01 I	1	0.3882286792654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = -0.592663447105876D-02 + 0.1491722999922961D 01 I	1	-0.517638251966724D 00 + 0.1931851608368755D 01 I
ROOT(5) = -0.1014904754401601D 01 + -0.1028420931384370D 01 I	1	-0.1767767147080701D 01 + 0.1767766798852015D 01 I
ROOT(6) = -0.2333333617834863D 01 + 0.2168624494945200D-05 I	1	-0.269777583074990D 01 + 0.774567463987070D 00 I
ROOT(7) = -0.1018508435678558D 01 + -0.972859560275189D 00 I	1	-0.3380740248331229D 01 + -0.905867194081616D 00 I
ROOT(8) = -0.9645441989282835D 00 + -0.998774789527914D 00 I	1	-0.2828426607107896D 01 + -0.2828427642384390D 01 I
ROOT(9) = -0.2481483709937032D-02 + -0.1497879862710794D 01 I	2	-0.1164484801399899D 01 + -0.4346666459873368D 01 I
ROOT(10) = 0.1560514417537318D 00 + 0.2495207685212091D 01 I	1	0.1294096343098645D 01 + -0.4829628831453027D 01 I
ROOT(11) = 0.3325675653760710D 00 + 0.3365993627402638D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(12) = -0.4837126081291009D 00 + 0.3134585687890842D 01 I	1	SOLVED BY DIRECT METHOD

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

Exhibit 6.4.

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.3000000368372262D-02 + 0.1304067616398651D-08 i	2	0.4829629115656279D 00 + 0.1294095284438187D 00 i
ROOT(2) = 0.6742035082463408D-06 + 0.1000000627692775D 01 i	2	0.7071067553046346D 00 + 0.7071068070684595D 00 i
ROOT(3) = 0.7624724932810605D-06 + 0.15000001030008833D 01 i	1	0.3882284792654056D 00 + 0.144888763117193D 01 i
ROOT(4) = -0.6671875662730541D-06 + 0.1499999011165093D 01 i	1	-0.5176382551966724D 00 + 0.1931851608368755D 01 i
ROOT(5) = -0.9999960111018027D 00 + -0.1000007493843641D 01 i	1	-0.1767767147080701D 01 + 0.1767766758852015D 01 i
ROOT(6) = -0.2333333333311710D 01 + 0.6499059058277495D-11 i	1	-0.2897777583074990D 01 + 0.7764567463987070D 00 i
ROOT(7) = -0.1000007692970472D 01 + -0.9999983698619370D 00 i	1	-0.3380740248331229D 01 + -0.9058671940816160D 00 i
ROOT(8) = -0.9999951740158410D 00 + -0.9999950263677619D 00 i	1	-0.2828426607107896D 01 + -0.2828427642384390D 01 i
ROOT(9) = -0.5924878530578477D-06 + -0.1499999487187738D 01 i	2	-0.1164684801399899D 01 + -0.4346666459873368D 01 i
ROOT(10) = -0.6093654451589193D-06 + 0.1500001089824981D 01 i	1	0.1294096345098645D 01 + -0.4829628831453027D 01 i
ROOT(11) = 0.2619599103083468D-04 + 0.3000007838671785D 01 i	1	SOLVED BY DIRECT METHOD
ROOT(12) = -0.9819291180928265D-05 + 0.2999974121589575D 01 i	1	SOLVED BY DIRECT METHOD

Exhibit 6.4. Roots Are: $-2.33, .003(2), i(2),$
 $1.5i(2), -1.5i(2) 3i(3), -1-i(3)$

NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 3 OF DEGREE 8

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(2) = -0.5000000000000000 01 + -0.1150000000000000 02 I
P(3) = -0.5175000000000000 02 + 0.4300000000000000 02 I
P(4) = 0.1572500000000000 03 + 0.1446250000000000 03 I
P(5) = 0.3075000000000000 03 + -0.3475000000000000 03 I
P(6) = -0.4952500000000000 03 + -0.4948750000000000 03 I
P(7) = -0.5857500000000000 03 + 0.4247500000000000 03 I
P(8) = 0.1810000000000000 03 + 0.4420000000000000 03 I
P(9) = 0.1580000000000000 03 + 0.5000000000000000 01 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-05
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.9999992498047420 00 + 0.1999998467904090D 01 I	2	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = -0.9921687016763280 01 + 0.5027384405227564D 00 I	2	0.7071067553046346D 00 + 0.7071068070484595D 00 I
ROOT(3) = 0.1916659864272422D 01 + 0.1966959787044516D 01 I	1	0.3882284732654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = 0.2013726806167612D 01 + 0.2086055012105922D 01 I	1	-0.5176382551966724D 00 + 0.1931851608366755D 01 I
ROOT(5) = 0.2069406641003821D 01 + 0.1946950095885711D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(6) = -0.1015484227697537D 01 + 0.4945588881101592D 00 I	1	SOLVED BY DIRECT METHOD

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.9999996239397971D 00 + 0.1999999233348969D 01 I	2	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = -0.9999426539148827D 00 + 0.499597557955272D 00 I	2	0.7071067553046346D 00 + 0.7071068070484595D 00 I
ROOT(3) = 0.199996221578001D 01 + 0.2000020836036100D 01 I	1	0.3882284732654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = 0.2000005629322798D 01 + 0.1999952499501282D 01 I	1	-0.5176382551966724D 00 + 0.1931851608366755D 01 I
ROOT(5) = 0.2000000873850570D 01 + 0.1999950053877640D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(6) = -0.10000005373961587D 01 + 0.49999950920121687D 00 I	1	SOLVED BY DIRECT METHOD

Exhibit 6.5. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$ (3)

2 NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS POLYNOMIAL NUMBER 4 OF DEGREE 12

THE COEFFICIENTS OF P(X) ARE

```
P(1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(2) = -0.1200000000000000 02 + -0.0000000000000000 00 I
P(3) = 0.7200000000000000 02 + 0.0000000000000000 00 I
P(4) = -0.2800000000000000 03 + -0.0000000000000000 00 I
P(5) = 0.7800000000000000 03 + 0.0000000000000000 00 I
P(6) = -0.1632000000000000 04 + -0.0000000000000000 00 I
P(7) = 0.2624000000000000 04 + 0.0000000000000000 00 I
P(8) = -0.3264000000000000 04 + -0.0000000000000000 00 I
P(9) = 0.3120000000000000 04 + 0.0000000000000000 00 I
P(10) = -0.2240000000000000 04 + -0.0000000000000000 00 I
P(11) = 0.1152000000000000 04 + 0.0000000000000000 00 I
P(12) = -0.3840000000000000 03 + -0.0000000000000000 00 I
P(13) = 0.6400000000000000 02 + 0.0000000000000000 00 I
```

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-03
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)

```
ROOT( 1) = 0.9994078033691520 00 + 0.9951391489643941D 00 I
ROOT( 2) = 0.10200968856097200 01 + -0.5420646130387889D 00 I
ROOT( 3) = 0.10027211066816580 01 + 0.1023310661526634D 01 I
ROOT( 4) = 0.6189968959361400 00 + -0.7949702013736144D 00 I
ROOT( 5) = 0.9851422047423414D 00 + -0.1395697998133038D 01 I
ROOT( 6) = 0.6515828215165421D 00 + -0.1206987666454936D 01 I
ROOT( 7) = 0.1393799984400508D 01 + -0.827192431176851D 00 I
ROOT( 8) = 0.1330620199428514D 01 + -0.1232093496230541D 01 I
```

IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(2) = 0.10200968856097200 01 + -0.5420646130387889D 00 I DID NOT CONVERGE.
THE PRESENT APPROXIMATION AFTER 200 ITERATIONS IS PRINTED BELOW.

IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(5) = 0.9851422047423414D 00 + -0.1395697998133038D 01 I DID NOT CONVERGE.
THE PRESENT APPROXIMATION AFTER 200 ITERATIONS IS PRINTED BELOW.

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

MULTIPLICITIES INITIAL APPROXIMATION

```
5 0.4829629115656279D 00 + 0.1294095284438187D 00 I
1 0.7071067553046346D 00 + 0.7071068070684595D 00 I
1 0.38822847925654056D 00 + 0.1448888763117193D 01 I
1 -0.5176382551966724D 00 + 0.1931851608368755D 01 I
1 -0.1767767147080701D 01 + 0.1767766758852015D 01 I
1 -0.289777563074990D 01 + 0.7764567463987070D 00 I
1 SOLVED BY DIRECT METHOD
1 SOLVED BY DIRECT METHOD
```

Exhibit 6.6.

```

ROOTS OF P(X)
MULTIPLICITIES
INITIAL APPROXIMATION
ROOT( 1) = 0.95949113795146260 00 + 0.99560175860617460 00 I
ROOT( 2) = 0.9592312317176770 00 + -0.10017819876754680 01 I
ROOT( 3) = 0.9585780717884340 00 + 0.10031671344932110 01 I
ROOT( 4) = 0.9999156996089460 00 + -0.99503023164852320 00 I
ROOT( 5) = 0.99991042189066210 00 + -0.99411359917024930 00 I
ROOT( 6) = 0.9960538588568690 00 + -0.10027599236645720 01 I
ROOT( 7) = 0.10037198385540710 01 + -0.99684656745825120 00 I
ROOT( 8) = 0.10001682256999940 01 + -0.99537592890632930 00 I
0.482962911156562790 00 + 0.12940952844381870 00 I
0.70710675530463460 00 + 0.73710680706845950 00 I
0.38822847926540560 00 + 0.14488887631171930 01 I
-0.51763825519667240 00 + 0.19318516083687550 01 I
-0.17677671470807010 01 + 0.117677667588520150 01 I
-0.2897775830749900 01 + 0.77645674639870700 00 I
SOLVED BY DIRECT METHOD
SOLVED BY DIRECT METHOD

```

Exhibit 6.6. Roots Are: $1+i(6)$, $1-i(6)$

MULLERS METHOD FOR FINDING THE ZEROS OF A POLYNOMIAL
POLYNOMIAL NUMBER 1 OF DEGREE 6

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.100000000000000001 + 0.000000000000000000 00 I
P(2) = -0.700000000000000010 + -0.105000000000000000 02 I
P(3) = -0.280000000000000000 + 0.580000000000000010 02 I
P(4) = 0.171000000000000000 + 0.150000000000000000 01 I
P(5) = -0.730000000000000000 + -0.251000000000000000 03 I
P(6) = -0.228000000000000000 + 0.104000000000000000 03 I
P(7) = 0.720000000000000010 + 0.104000000000000000 03 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.199995488749810D 01 + 0.2000017733363912D 01 I	3	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.1013502627672869D 01 + 0.1996163083160882D 01 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = 0.9866348122507864D 00 + 0.2003804882325521D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(4) = -0.1000002106173084D 01 + 0.4999768344218605D 00 I	1	SOLVED BY DIRECT METHOD

AFTER THE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.1999954835419787D 01 + 0.2000017627898503D 01 I	3	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.1000000159779278D 01 + 0.199999979600045D 01 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = 0.9999998306711213D 00 + 0.2000000071685089D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(4) = -0.999999999999998D 00 + 0.500000000000000D 00 I	1	SOLVED BY DIRECT METHOD

Exhibit 6.7. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$

MULLERS METHOD FOR FINDING THE ZEROS OF A POLYNOMIAL
POLYNOMIAL NUMBER 2 OF DEGREE 15

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.4800000000000000 02 + 0.0000000000000000 00 I
P(2) = 0.2557120000000000 03 + -0.3840000000000000 03 I
P(3) = -0.7353536000000000 02 + -0.2189696000000000 04 I
P(4) = -0.3855656960000000 04 + -0.6946851458000000 05 I
P(5) = -0.1733866468000000 05 + -0.1420625972800000 05 I
P(6) = -0.4967989270400000 05 + -0.1765857464000000 05 I
P(7) = -0.1022394522130000 06 + -0.6030664232000000 04 I
P(8) = -0.1642742200560000 06 + 0.4137366230400000 05 I
P(9) = -0.2036625888420000 06 + 0.1093899227670000 06 I
P(10) = -0.1871255780010000 06 + 0.1929865440330000 06 I
P(11) = -0.1274997298590000 06 + 0.2171341227420000 06 I
P(12) = -0.2814692716800000 05 + 0.1928489727960000 06 I
P(13) = 0.1329434434800000 05 + 0.1038130226500000 06 I
P(14) = 0.3053900774700000 05 + 0.2998989141300000 05 I
P(15) = -0.1835899020000000 03 + -0.1827632160000000 03 I
P(16) = 0.2755620000000000 00 + 0.2755620000000000 00 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

ROOT(1) = 0.3000000056259372D-02 + 0.8277931832119355D-12 I 2
ROOT(2) = 0.1686472961057889D-04 + 0.1000001835704009D 01 I 1
ROOT(3) = 0.1280227161896980D-04 + 0.1499998729963919D 01 I 1
ROOT(4) = 0.6019678782146761D-04 + 0.299999366616581900D 01 I 1
ROOT(5) = -0.1686485393117943D-04 + 0.99999816357761600D 01 I 1
ROOT(6) = -0.23333333333333649D 01 + 0.3249409455271494D-12 I 1
ROOT(7) = -0.1000163769022359D 01 + -0.100017371459914D 01 I 2
ROOT(8) = -0.334905054308976D-03 + -0.1500133733984800D 01 I 1
ROOT(9) = -0.9996724783532904D 00 + -0.9996521399325563D 00 I 1
ROOT(10) = 0.334811914008498D-03 + -0.149986916248616D 01 I 1
ROOT(11) = -0.1715220973482156D-03 + 0.2999990216209916D 01 I 1
ROOT(12) = 0.1113136898246577D-03 + 0.3000073149878609D 01 I 1
ROOT(13) = -0.1276492775835444D-04 + 0.1500001215505238D 01 I 1

AFTER THE ATTEMPT TO IMPROVE ACCURACY

Exhibit 6.8.

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.300000056258119D-02 +	2	0.4829629115656279D 00 + 0.1296095254438187D 00 I
ROOT(2) = 0.3001086172156215D-07 +	1	0.7071087553048346D 00 + 0.7071086070684595D 00 I
ROOT(3) = 0.6189053790319132D-07 +	1	0.3882284792654056D 00 + 0.1446888763117193D 01 I
ROOT(4) = 0.2397327410073823D-04 +	1	-0.5176382551966724D 00 + 0.1931851608368750D 01 I
ROOT(5) = 0.3973984540428445D-07 +	1	-0.1767767147080701D 01 + 0.1767766758852015D 01 I
ROOT(6) = -0.2333333333333334D 01 +	1	-0.289777583074990D 01 + 0.7764567463987070D 00 I
ROOT(7) = -0.1000006990404978D 01 +	2	-0.3380740248331229D 01 + -0.9058671940816160D 00 I
ROOT(8) = 0.9284422544336306D-08 +	1	-0.2828426607107896D 01 + -0.2828427642384390D 01 I
ROOT(9) = -0.1000006251006585D 01 +	1	-0.1164684801399899D 01 + -0.4346666459873368D 01 I
ROOT(10) = 0.1671714279306380D-07 +	1	0.1294096345098645D 01 + -0.4829628831453027D 01 I
ROOT(11) = -0.2558768623037301D-04 +	1	0.3889088292979509D 01 + -0.3889086300072258D 01 I
ROOT(12) = 0.2110926108982832D-04 +	1	SOLVED BY DIRECT METHOD
ROOT(13) = -0.54537993864692274D-07 +	1	SOLVED BY DIRECT METHOD

Exhibit 6.8. Roots Are: $-2.33, .003(2), i(2), 1.5i(2),$
 $-1.5i(2) 3i(3), -1-i(3)$

MULLERS METHOD FOR FINDING THE ZEROS OF A POLYNOMIAL
POLYNOMIAL NUMBER 3 OF DEGREE 8

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.100000000000000000 01 + 0.000000000000000000 00 I
P(2) = -0.500000000000000001 01 + -0.115000000000000000 02 I
P(3) = -0.517500000000000002 02 + 0.430000000000000000 02 I
P(4) = 0.157250000000000003 03 + 0.144625000000000000 03 I
P(5) = 0.307500000000000003 03 + -0.347500000000000000 03 I
P(6) = -0.495250000000000003 03 + -0.494875000000000000 03 I
P(7) = -0.585750000000000003 03 + 0.424750000000000000 03 I
P(8) = 0.181000000000000003 03 + 0.442000000000000000 03 I
P(9) = 0.158000000000000003 03 + 0.600000000000000000 01 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.2000042202873018D 01 + 0.2000028743998294D 01 I	3	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.1000563806908988D 01 + 0.1980638952568216D 01 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = -0.1063528418749844D 01 + 0.4942372379199120D 00 I	1	0.3882284792654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = -0.9722122516869824D 00 + 0.560379457973125D 00 I	1	-0.5176382551966724D 00 + 0.1931851608368755D 01 I
ROOT(5) = 0.9994370680506410D 00 + 0.2019094286923692D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(6) = -0.96438868131418565D 00 + 0.4459563547509665D 00 I	1	SOLVED BY DIRECT METHOD

AFTER THE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.2000042175889591D 01 + 0.20000287507113895D 01 I	3	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.100000023165255D 01 + 0.199999980577700D 01 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = -0.1000006594122314D 01 + 0.4999952821126052D 00 I	1	0.3882284792654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = -0.1000000972571865D 01 + 0.5000084175726651D 00 I	1	-0.5176382551966724D 00 + 0.1931851608368755D 01 I
ROOT(5) = 0.9999999168409400D 00 + 0.2000000100132659D 01 I	1	SOLVED BY DIRECT METHOD
ROOT(6) = -0.9999925738129557D 00 + 0.4999962818337703D 00 I	1	SOLVED BY DIRECT METHOD

Exhibit 6.9. Roots Are: 2+2i (3), 1+2i (2), -1+.5i (3)

MULLERS METHOD FOR FINDING THE ZEROS OF A POLYNOMIAL
POLYNOMIAL NUMBER 4 OF DEGREE 12

THE COEFFICIENTS OF P(X) ARE

P(1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(2) = -0.1200000000000000 02 + -0.0000000000000000 00 I
P(3) = 0.7200000000000000 03 + 0.0000000000000000 00 I
P(4) = -0.2800000000000000 04 + -0.0000000000000000 00 I
P(5) = 0.7800000000000000 05 + 0.0000000000000000 00 I
P(6) = -0.1632000000000000 06 + 0.0000000000000000 00 I
P(7) = 0.2624000000000000 07 + 0.0000000000000000 00 I
P(8) = -0.3264000000000000 08 + 0.0000000000000000 00 I
P(9) = 0.3120000000000000 09 + 0.0000000000000000 00 I
P(10) = -0.2240000000000000 04 + 0.0000000000000000 00 I
P(11) = 0.1152000000000000 03 + 0.0000000000000000 00 I
P(12) = -0.3840000000000000 04 + 0.0000000000000000 00 I
P(13) = 0.6400000000000000 02 + 0.0000000000000000 00 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
RADIUS TO START SEARCH. 0.100-01
RADIUS TO END SEARCH. 0.000 00
0.000 00

BEFORE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.1004468192948739D 01 + 0.1002186117532751D 01 I	4	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.9858978428495002D 00 + 0.1006804518996020D 01 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = 0.984226248473947D 00 + 0.9844382337895168D 00 I	1	0.3882284792654056D 00 + 0.1448888763117193D 01 I
ROOT(4) = 0.1007069490520364D 01 + -0.8278425247752856D 00 I	2	-0.5176382551966724D 00 + 0.1931851408368755D 01 I
ROOT(5) = 0.7584518107423498D 00 + -0.9410320205052345D 00 I	1	-0.1767767147080701D 01 + 0.1767766758852015D 01 I
ROOT(6) = 0.8435901394416417D 00 + -0.12151582772720118D 01 I	1	-0.289777583074990D 01 + 0.7764567463987070D 00 I
ROOT(7) = 0.1245479586151531D 01 + -0.9609751096770448D 00 I	1	SOLVED BY DIRECT METHOD
ROOT(8) = 0.11388031428157103D 01 + -0.1227138567895571D 01 I	1	SOLVED BY DIRECT METHOD

AFTER THE ATTEMPT TO IMPROVE ACCURACY

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.1003576740125457D 01 + 0.1001742634790606D 01 I	4	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) = 0.9990524786861438D 00 + 0.1004512114384271D 01 I	1	0.7071067553046346D 00 + 0.7071068070684595D 00 I
ROOT(3) = 0.1003596793567675D 01 + 0.9965600317041306D 00 I	1	0.3882284792654056D 00 + 0.1448888763117193D 01 I

Exhibit 6.10.

ROOT(4) = 0.10007324697602040 01 + -0.99544295721310670 00 I
ROOT(5) = 0.10047654833334880 01 + -0.100069571831889740 01 I
ROOT(6) = 0.99547825662743310 00 + -0.99607660789158700 00 I
ROOT(7) = 0.99541663095228950 00 + -0.1000189829218090 01 I
ROOT(8) = 0.10000743905941600 01 + -0.99512853288497680 00 I

2
1
1
1
1
1
1
1
SOLVED BY DIRECT METHOD
SOLVED BY DIRECT METHOD

-0.5176392579567240 00 + 0.19318516083687550 01 I
-0.17677671470807010 01 + 0.1767767588520150 01 I
-0.2897775830769500 01 + 0.77645074639870700 00 I

Exhibit 6.10. Roots Are: 1+i (6), 1-i (6)

GREATEST COMMON DIVISOR METHOD USED WITH NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 1

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.10D-02
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.10D-19
TEST FOR MULTIPLICITIES. 0.10D-01
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

THE DEGREE OF P(X) IS 6 THE COEFFICIENTS ARE

P(7) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(6) = -0.7000000000000000 01 + -0.1050000000000000 02 I
P(5) = -0.2800000000000000 02 + 0.5800000000000000 02 I
P(4) = 0.1710000000000000 03 + 0.1500000000000000 01 I
P(3) = -0.7300000000000000 02 + -0.2510000000000000 03 I
P(2) = -0.2280000000000000 03 + 0.1040000000000000 03 I
P(1) = 0.7200000000000000 02 + 0.1040000000000000 03 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 3 THE COEFFICIENTS ARE

Q(4) = 0.1000000000000000 01 + 0.0000000000000000 00 I
Q(3) = -0.2000000000000150 01 + -0.4500000000000180 01 I
Q(2) = -0.7000000000000420 01 + 0.3500000000000060 01 I
Q(1) = 0.9999999999997762 00 + 0.6999999999999812 01 I

ROOTS OF Q(X)

ROOT(1) = 0.9999999999997565 00 + 0.1999999999999574 01 I
ROOT(2) = 0.2000000000000347 01 + 0.2000000000000529 01 I
ROOT(3) = -0.9999999999995390 00 + 0.5000000000000149 00 I

INITIAL APPROXIMATION

0.4829629115656279D 00 + 0.1294095284438187D 00 I
RESULTS OF SUBROUTINE QUAD
RESULTS OF SUBROUTINE QUAD

ROOTS OF P(X)

ROOT(1) = 0.9999999999997565 00 + 0.1999999999999574 01 I
ROOT(2) = 0.2000000000000347 01 + 0.2000000000000530 01 I
ROOT(3) = -0.9999999999995390 00 + 0.5000000000000147 00 I

MULTIPLICITIES

2
3
1

INITIAL APPROXIMATION

0.4829629115656279D 00 + 0.1294095284438187D 00 I
RESULTS OF SUBROUTINE QUAD
RESULTS OF SUBROUTINE QUAD

Exhibit 6.11. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$

GREATEST COMMON DIVISOR METHOD USED WITH NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 2

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.10D-02
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.10D-19
TEST FOR MULTIPLICITIES. 0.10D-01
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

THE DEGREE OF P(X) IS 15 THE COEFFICIENTS ARE

P(16) = 0.480000000000000000 02 + 0.000000000000000000 00 I
P(15) = 0.255712000000000000 03 + -0.384000000000000000 03 I
P(14) = -0.735355680000000000 02 + -0.218969600000000000 04 I
P(13) = -0.385556696000000000 04 + -0.694685145600000010 04 I
P(12) = -0.173338646480000000 05 + -0.142062597280000000 05 I
P(11) = -0.496798927040000010 05 + -0.176585746400000000 05 I
P(10) = -0.102239452213000000 06 + -0.603066423200000010 04 I
P(9) = -0.164276220056000000 06 + -0.413736623040000000 05 I
P(8) = -0.203662368842000000 06 + 0.109389922767000000 06 I
P(7) = -0.187125578001000000 06 + 0.192986544093000000 06 I
P(6) = -0.127499729890000000 06 + 0.217134122742000000 06 I
P(5) = -0.281469271680000000 05 + 0.192848972796000000 06 I
P(4) = 0.132943443480000000 05 + 0.103813022655000000 06 I
P(3) = 0.305390077470000000 05 + 0.299898914130000000 05 I
P(2) = -0.183589902000000000 03 + -0.1827632160000000 03 I
P(1) = 0.275562000000000000 00 + 0.275562000000000000 00 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 7 THE COEFFICIENTS ARE

Q(8) = 0.480000000000000000 02 + 0.000000000000000000 00 I
Q(7) = 0.1598559999590830 03 + -0.14400000000184630 03 I
Q(6) = 0.2675199998682870 03 + -0.527568000000682380 03 I
Q(5) = 0.32719599997819040 03 + -0.9144416000000759420 03 I
Q(4) = 0.230159999960445530 02 + -0.152125200000037680 04 I
Q(3) = -0.72072000047996880 02 + -0.13274279999966590 04 I
Q(2) = -0.75578400005415630 03 + -0.75200400003332210 03 I
Q(1) = 0.22680001335502310 01 + 0.22679999258286330 01 I

ROOTS OF Q(X)

ROOT(1) = 0.3000000038645997D-02 + -0.1370543176638413D-09 I
ROOT(2) = 0.1466635543509054D-09 + 0.1000000000302556D 01 I
ROOT(3) = -0.162620608809517D-09 + 0.1499999999807067D 01 I
ROOT(4) = -0.1167767202194363D-10 + -0.15000000000006763D 01 I

INITIAL APPROXIMATION

0.4829629115656279D 00 + 0.1294095284438187D 00 I
0.7071067553046346D 00 + 0.7071068070684595D 00 I
0.3882284792654056D 00 + 0.144888763117193D 01 I
-0.5176382551966724D 00 + 0.1931851608368755D 01 I

Exhibit 6.12.

GREATEST COMMON DIVISOR METHOD USED WITH NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 3

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 8 THE COEFFICIENTS ARE

P(9) = 0.1800000000000000 01 + 0.0000000000000000 00 I
P(8) = -0.5000000000000000 01 + -0.1150000000000000 02 I
P(7) = -0.5175000000000000 02 + 0.4300000000000000 02 I
P(6) = 0.1572500000000000 03 + 0.1446250000000000 03 I
P(5) = 0.3075000000000000 03 + -0.3475000000000000 03 I
P(4) = -0.4952500000000000 03 + -0.4948750000000000 03 I
P(3) = -0.5857500000000000 03 + 0.4247500000000000 03 I
P(2) = 0.1810000000000000 03 + 0.4420000000000000 03 I
P(1) = 0.1580000000000000 03 + 0.6000000000000000 01 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 3 THE COEFFICIENTS ARE

Q(4) = 0.1000000000000000 01 + 0.0000000000000000 00 I
Q(3) = -0.2000000000000273D 01 + -0.4500000000000275D 01 I
Q(2) = -0.7000000000000935D 01 + 0.35000000000000469D 01 I
Q(1) = 0.9999999999994173D 00 + 0.70000000000000498D 01 I

ROOTS OF Q(X)

ROOT(1) = 0.9999999999995483D 00 + 0.1999999999999755D 01 I
ROOT(2) = 0.2000000000000665D 01 + 0.2000000000000539D 01 I
ROOT(3) = -0.999999999999408D 00 + 0.4999999999999814D 00 I

ROOTS OF P(X)

ROOT(1) = 0.9999999999995483D 00 + 0.1999999999999755D 01 I
ROOT(2) = 0.2000000000000665D 01 + 0.2000000000000539D 01 I
ROOT(3) = -0.999999999999408D 00 + 0.4999999999999812D 00 I

INITIAL APPROXIMATION

0.4829629115656279D 00 + 0.1294095284438187D 00 I
RESULTS OF SUBROUTINE QUAD
RESULTS OF SUBROUTINE QUAD

MULTIPLICITIES

INITIAL APPROXIMATION

2 0.4829629115656279D 00 + 0.1294095284438187D 00 I
3 RESULTS OF SUBROUTINE QUAD
3 RESULTS OF SUBROUTINE QUAD

Exhibit 6.13. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$ (3)

GREATEST COMMON DIVISOR METHOD USED WITH NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 4

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.10D-02
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.10D-19
TEST FOR MULTIPLICITIES. 0.10D-01
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

THE DEGREE OF P(X) IS 12 THE COEFFICIENTS ARE

P(13) = 0.100000000000000000 01 + 0.000000000000000000 00 I
P(12) = -0.1200000000000000 02 + -0.000000000000000000 00 I
P(11) = 0.7200000000000000 02 + 0.000000000000000000 00 I
P(10) = -0.2800000000000000 03 + -0.000000000000000000 00 I
P(9) = 0.7800000000000000 03 + 0.000000000000000000 00 I
P(8) = -0.1632000000000000 04 + -0.000000000000000000 00 I
P(7) = 0.2624000000000000 04 + 0.000000000000000000 00 I
P(6) = -0.3264000000000000 04 + -0.000000000000000000 00 I
P(5) = 0.3120000000000000 04 + 0.000000000000000000 00 I
P(4) = -0.2240000000000000 04 + -0.000000000000000000 00 I
P(3) = 0.1152000000000000 04 + 0.000000000000000000 00 I
P(2) = -0.3840000000000000 03 + -0.000000000000000000 00 I
P(1) = 0.6400000000000000 02 + 0.000000000000000000 00 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 2 THE COEFFICIENTS ARE

Q(3) = 0.100000000000000000 01 + 0.000000000000000000 00 I
Q(2) = -0.2000000000000000 01 + -0.000000000000000000 00 I
Q(1) = 0.199999999999998300 01 + 0.000000000000000000 00 I

ROOTS OF P(X)

MULTIPLICITIES

ROOT(1) = 0.100000000000000000 01 + 0.99999999999990740 00 I
ROOT(2) = 0.100000000000000000 01 + -0.99999999999990740 00 I

RESULTS OF SUBROUTINE QUAD
RESULTS OF SUBROUTINE QUAD

Exhibit 6.14. Roots Are: 1+i (6), 1-i (6)

GREATEST COMMON DIVISOR METHOD USED WITH MULLERS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 1

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 6 THE COEFFICIENTS ARE

P(7) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(6) = -0.7000000000000010 01 + -0.1050000000000000 02 I
P(5) = -0.2800000000000000 02 + 0.5800000000000000 02 I
P(4) = 0.1710000000000000 03 + 0.1500000000000000 01 I
P(3) = -0.7300000000000000 02 + -0.2510000000000000 03 I
P(2) = -0.2280000000000000 03 + 0.1040000000000000 03 I
P(1) = 0.7200000000000000 02 + 0.1040000000000000 03 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 3 THE COEFFICIENTS ARE

Q(4) = 0.1000000000000000 01 + 0.0000000000000000 00 I
Q(3) = -0.2000000000001500 01 + -0.4500000000000118 01 I
Q(2) = -0.7000000000000420 01 + 0.3500000000000060 01 I
Q(1) = 0.9999999999997762 00 + 0.6999999999999812 01 I

ROOTS OF Q(X)

INITIAL APPROXIMATION

0.4829629115656279D 00 + 0.1294095284438187D 00 I
SOLVED BY DIRECT METHOD
SOLVED BY DIRECT METHOD

ROOT(1) = 0.99999999999997565D 00 + 0.19999999999999574D 01 I
ROOT(2) = 0.20000000000000347D 01 + 0.20000000000000529D 01 I
ROOT(3) = -0.9999999999999539D 00 + 0.50000000000000149D 00 I

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

ROOT(1) = 0.99999999999997568D 00 + 0.19999999999999574D 01 I 2
ROOT(2) = 0.20000000000000347D 01 + 0.20000000000000530D 01 I 3
ROOT(3) = -0.9999999999999541D 00 + 0.50000000000000148D 00 I 1
RESULTS OF SUBROUTINE QUAD
RESULTS OF SUBROUTINE QUAD

Exhibit 6.15. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$

GREATEST COMMON DIVISOR METHOD USED WITH MULLERS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 2

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 15 THE COEFFICIENTS ARE

P(16) = 0.4800000000000000 02 + 0.0000000000000000 00 I
P(15) = 0.2557120000000000 03 + -0.3840000000000000 03 I
P(14) = -0.7353556800000000 02 + -0.2189696000000000 04 I
P(13) = -0.3855565600000000 04 + -0.6948651456000000 04 I
P(12) = -0.1733386464000000 05 + -0.1420625972800000 05 I
P(11) = -0.4967989270400000 05 + -0.1765857464000000 05 I
P(10) = -0.1022394522130000 06 + -0.6030664232000000 04 I
P(9) = -0.1642742200560000 06 + 0.4137366230400000 05 I
P(8) = -0.2036625888420000 06 + 0.1093899227670000 06 I
P(7) = -0.1871255780010000 06 + 0.1929865440330000 06 I
P(6) = -0.1274997298590000 06 + 0.2171341227420000 06 I
P(5) = -0.2814692716800000 05 + 0.1928489727960000 06 I
P(4) = 0.1329434434800000 05 + 0.1038130226550000 06 I
P(3) = 0.3053900774700000 05 + 0.2998989141300000 05 I
P(2) = -0.1835899020000000 03 + -0.1827632160000000 03 I
P(1) = 0.2755620000000000 00 + 0.2755620000000000 00 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 7 THE COEFFICIENTS ARE

Q(8) = 0.4800000000000000 02 + 0.0000000000000000 00 I
Q(7) = 0.1598559999590830 03 + -0.14400000000184530 03 I
Q(6) = 0.26751999998662870 03 + -0.52756800000682380 03 I
Q(5) = 0.32719599997819040 03 + -0.91441600000759420 03 I
Q(4) = 0.230159999960445530 02 + -0.15212520000037680 04 I
Q(3) = -0.72072000047996880 02 + -0.13274279999966590 04 I
Q(2) = -0.75578400005415630 03 + -0.75200400003332210 03 I
Q(1) = 0.22680001335502310 01 + 0.22679999258286330 01 I

ROOTS OF Q(X)

ROOT(1) = 0.30000000386459970-02 + -0.13705431778101480-09 I
ROOT(2) = 0.1468683724911731D-09 + 0.1000000003025560 01 I
ROOT(3) = -0.1626201046532216D-09 + 0.14999999998070670 01 I
ROOT(4) = 0.1140454591987512D-09 + 0.30000000000000000 01 I

INITIAL APPROXIMATION

0.48296291156562790 00 + 0.12940952844381870 00 I
0.7071067553043460 00 + 0.70710680706845950 00 I
0.38822847926540560 00 + 0.14488887631171950 01 I
-0.51763825519667240 00 + 0.19318516089687550 01 I

Exhibit 6.16.

-0.1767767147080701D 01 + 0.1767766758852015D 01 I
SOLVED BY DIRECT METHOD
SOLVED BY DIRECT METHOD

[illegible]ROOTS OF $P(X)$

MULTIPLICITIES

INITIAL APPROXIMATION

RRDRT(1)	=	0.300000000386659970	-02	+	-0.137054317781874509	-09	1	2	0.482962911565642790	00	+	0.12940952584438187D	00	0
RRDRT(2)	=	0.1662636090771709D	-09	+	0.100000000003225560	01	1	2	0.070104807068845950	00	+	0.70104807068845950	00	1
RRDRT(3)	=	-0.16262091734869740D	-09	+	0.149999999998070660	01	1	2	0.388225847628634056D	00	+	0.1448868763117930	01	1
RRDRT(4)	=	0.11404556580943080D	-09	+	0.3000000000053940	01	1	3	-0.51763825519686724D	00	+	0.1931816083687550	01	1
RRDRT(5)	=	-0.23333333340885D	-11	+	0.4291402714880148	-11	1	1	-0.1767767147080701D	01	+	0.176776758852015D	01	1
RRDRT(6)	=	-0.11677375717066360D	-10	+	0.1500000000006763D	00	1	2	RESULTS OF SUBROUTINE QUAD					
RRDRT(7)	=	-0.1000000000322600D	-10	+	0.9999999999955779D	00	1	3	RESULTS OF SUBROUTINE QUAD					

Exhibit 6.16. Roots Are: $-2.33, .003(2), i(2), 1.5i(2), -1.5i(2), 3i(3), -1-i(3)$

GREATEST COMMON DIVISOR METHOD USED WITH MULLERS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 3

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.10D-02
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.10D-19
TEST FOR MULTIPLICITIES. 0.10D-01
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

THE DEGREE OF P(X) IS 8 THE COEFFICIENTS ARE

P(9) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(8) = -0.5000000000000000 01 + -0.1150000000000000 02 I
P(7) = -0.5175000000000000 02 + 0.4300000000000000 02 I
P(6) = 0.1572500000000000 03 + 0.1446250000000000 03 I
P(5) = 0.3075000000000000 03 + -0.3475000000000000 03 I
P(4) = -0.4952500000000000 03 + -0.4948750000000000 03 I
P(3) = -0.5857500000000000 03 + 0.4247500000000000 03 I
P(2) = 0.1810000000000000 03 + 0.4420000000000000 03 I
P(1) = 0.1580000000000000 03 + 0.6000000000000000 01 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 3 THE COEFFICIENTS ARE

Q(4) = 0.1000000000000000 01 + 0.0000000000000000 00 I
Q(3) = -0.2000000000000000 01 + -0.4500000000000000 01 I
Q(2) = -0.7000000000000000 01 + 0.3500000000000000 01 I
Q(1) = 0.9999999999999999 173D 00 + 0.7000000000000000 01 I

ROOTS OF Q(X)

ROOT(1) =	0.9999999999999999 5483D 00 + 0.19999999999999755D 01 I	INITIAL APPROXIMATION
ROOT(2) =	0.2000000000000000 0665D 01 + 0.20000000000000539D 01 I	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(3) =	-0.9999999999999999 408D 00 + 0.49999999999999814D 00 I	SOLVED BY DIRECT METHOD SOLVED BY DIRECT METHOD

ROOTS OF P(X)

		MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) =	0.9999999999999999 5483D 00 + 0.19999999999999755D 01 I	2	0.4829629115656279D 00 + 0.1294095284438187D 00 I
ROOT(2) =	0.2000000000000000 0665D 01 + 0.20000000000000539D 01 I	3	RESULTS OF SUBROUTINE QUAD
ROOT(3) =	-0.9999999999999999 410D 00 + 0.49999999999999812D 00 I	3	RESULTS OF SUBROUTINE QUAD

Exhibit 6.17. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$ (3)

GREATEST COMMON DIVISOR METHOD USED WITH MULLERS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 4

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 12 THE COEFFICIENTS ARE

P(13) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(12) = -0.1200000000000000 02 + -0.0000000000000000 00 I
P(11) = 0.7200000000000010 02 + 0.0000000000000000 00 I
P(10) = -0.2800000000000000 03 + -0.0000000000000000 00 I
P(9) = 0.7800000000000000 03 + 0.0000000000000000 00 I
P(8) = -0.1632000000000000 04 + -0.0000000000000000 00 I
P(7) = 0.2624000000000000 04 + 0.0000000000000000 00 I
P(6) = -0.3264000000000000 04 + -0.0000000000000000 00 I
P(5) = 0.3120000000000000 04 + 0.0000000000000000 00 I
P(4) = -0.2240000000000000 04 + -0.0000000000000000 00 I
P(3) = 0.1152000000000000 04 + 0.0000000000000000 00 I
P(2) = -0.3840000000000000 03 + -0.0000000000000000 00 I
P(1) = 0.6400000000000010 02 + 0.0000000000000000 00 I

Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X).
THE DEGREE OF Q(X) IS 2 THE COEFFICIENTS ARE

Q(3) = 0.1000000000000000 01 + 0.0000000000000000 00 I
Q(2) = -0.2000000000000150 01 + -0.0000000000000000 00 I
Q(1) = 0.1999999999999850 01 + 0.0000000000000000 00 I

ROOTS OF P(X)

MULTIPLICITIES

	RESULTS OF SUBROUTINE QUAD	RESULTS OF SUBROUTINE QUAD
RDDT(1) =	0.1000000000000007D 01 + 0.9999999999999074D 00 I	6
RDDT(2) =	0.1000000000000007D 01 + -0.9999999999999074D 00 I	6

Exhibit 6.18. Roots Are: $1+i(6)$, $1-i(6)$

REPEATED USE OF THE GREATEST COMMON DIVISOR AND NEWTONS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS. POLYNOMIAL NUMBER 1

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 6 THE COEFFICIENTS ARE

P(7) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(6) = -0.7000000000000010 01 + -0.1050000000000000 02 I
P(5) = -0.2800000000000000 02 + 0.5800000000000010 02 I
P(4) = 0.1710000000000000 03 + 0.1500000000000000 01 I
P(3) = -0.7300000000000000 02 + -0.2510000000000000 03 I
P(2) = -0.2280000000000000 03 + 0.1040000000000000 03 I
P(1) = 0.7200000000000010 02 + 0.1040000000000000 03 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 1

G(2) = 0.1000000000000000 01 + 0.0000000000000000 00 I
G(1) = 0.9999999999973500 00 + -0.5000000000175420 00 I

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

ROOT(1) = -0.9999999999973500 00 + 0.500000000000175420 00 I 1 NO INITIAL APPROXIMATIONS

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 2

G(2) = 0.1000000000000000 01 + 0.0000000000000000 00 I
G(1) = -0.9999999999991790 00 + -0.1999999999968450 01 I

Exhibit 6.19.

ROOTS OF P(X) MULTIPLICITIES INITIAL APPROXIMATION

ROOT(1) = 0.999999999999179D 00 + 0.1999999999996845D 01 I 2 NO INITIAL APPROXIMATIONS

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 3

2) = 0.1000000000000000D 01 + 0.0000000000000000D 00 I
 1) = -0.1999999999999967D 01 + -0.20000000000001519D 01 I

ROOTS OF P(X) MULTIPLICITIES INITIAL APPROXIMATION

ROOT(3) = 0.1999999999999967D 01 + 0.20000000000001519D 01 I 3 NO INITIAL APPROXIMATIONS

Exhibit 6.19. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$

REPEATED USE OF THE GREATEST COMMON DIVISOR AND NEWTONS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS
POLYNOMIAL NUMBER 2

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF PIX IS 15 THE COEFFICIENTS ARE

P(16) = 0.4800000000000000 02 + 0.0000000000000000 00 I
P(15) = 0.2557120000000000 03 + -0.3840000000000000 03 I
P(14) = -0.7355568000000000 02 + -0.2189696000000000 04 I
P(13) = -0.3855656960000000 04 + -0.6946851456000000 04 I
P(12) = -0.1733386464000000 05 + -0.1420625972800000 05 I
P(11) = -0.4947989270400000 05 + -0.1765857644000000 05 I
P(10) = -0.1022394522130000 06 + -0.6030864232000000 04 I
P(9) = -0.1642742200560000 06 + 0.4137366236400000 05 I
P(8) = -0.2038625888420000 06 + 0.1093899227670000 06 I
P(7) = -0.1871255780010000 06 + 0.1929865440330000 06 I
P(6) = -0.1274997298590000 06 + 0.2171341227420000 06 I
P(5) = -0.2814692716800000 05 + 0.1528489727960000 06 I
P(4) = 0.1329434434800000 05 + 0.1038130226550000 06 I
P(3) = 0.3053900774700000 05 + 0.2998989141300000 05 I
P(2) = -0.1835899020000000 03 + -0.1827632160000000 03 I
P(1) = 0.2755620000000000 00 + 0.2755620000000000 00 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF PIX WHICH HAVE MULTIPLICITY 1

G(2) = 0.4800000000000000 02 + 0.0000000000000000 00 I
G(1) = 0.1119999992896031D 03 + -0.8274988658740768D-06 I

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = -0.233333331853397D 01 + 0.1723955970570993D-07 I	1	NO INITIAL APPROXIMATIONS

Exhibit 6.20.

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 2

G15) = 0.1000000000000000 01 + 0.0000000000000000 00 I
 G14) = -0.2999970655863438D-02 + -0.9999999456362725D 00 I
 G13) = 0.2249999931178885D 01 + 0.3000058598117361D-02 I
 G12) = -0.6750109902460278D-02 + -0.2250000129067665D 01 I
 G11) = 0.4643479465382683D-06 + 0.6749603982170172D-02 I

ROOTS OF G(X)

INITIAL APPROXIMATION

ROOT(1) = 0.2999823203921417D-02 + -0.2056988264688649D-06 I
 ROOT(2) = 0.4767581845696651D-06 + 0.100000055850184D 01 I
 ROOT(3) = 0.2331721616933697D-07 + -0.149999972431725D 01 I
 ROOT(4) = -0.3526234587180479D-06 + 0.1499999587916640D 01 I

ROOTS OF P(X)

INITIAL APPROXIMATION

ROOT(1) = 0.2999823203921417D-02 + -0.2056988264688649D-06 I 2
 ROOT(2) = 0.4767581845696651D-06 + 0.100000055850184D 01 I 2
 ROOT(3) = 0.2331721616933697D-07 + -0.149999972431725D 01 I 2
 ROOT(4) = -0.3526234587180479D-06 + 0.1499999587916640D 01 I 2

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 3

G13) = 0.1000000000000000 01 + 0.0000000000000000 00 I
 G12) = 0.9999999853705541D 00 + -0.2000000017162632D 01 I
 G11) = 0.300000000253335D 01 + -0.3000000031791514D 01 I

ROOTS OF P(X)

INITIAL APPROXIMATION

ROOT(1) = 0.146293042349907D-07 + 0.3000000017162633D 01 I 3
 ROOT(2) = -0.9999999999998584D 00 + -0.100000000000002D 01 I 3

Exhibit 6.20. Roots Are: -2.33, .003 (2), i (2), 1.5i (2),
 -1.5i (2), 3i (3), -1-i (3)

REPEATED USE OF THE GREATEST COMMON DIVISOR AND NEWTONS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS
POLYNOMIAL NUMBER 3

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.10D-02
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.10D-19
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

THE DEGREE OF P(X) IS 8 THE COEFFICIENTS ARE

P(9) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(8) = -0.5000000000000000 01 + -0.1150000000000000 02 I
P(7) = -0.5175000000000000 02 + 0.4300000000000000 02 I
P(6) = 0.1572500000000000 03 + 0.1446250000000000 03 I
P(5) = 0.3075000000000000 03 + -0.3475000000000000 03 I
P(4) = -0.4952500000000000 03 + -0.4948750000000000 03 I
P(3) = -0.5857500000000000 03 + 0.4247500000000000 03 I
P(2) = 0.1810000000000000 03 + 0.4420000000000000 03 I
P(1) = 0.1580000000000000 03 + 0.6000000000000000 01 I

NO ROOTS OF MULTIPLICITY 1

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 2

G(2) = 0.1000000000000000 01 + 0.0000000000000000 00 I
G(1) = -0.9999999999999999 00 + -0.1999999999999999 01 I

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.9999999999999999 00 + 0.1999999999999999 01 I	2	NO INITIAL APPROXIMATIONS

Exhibit 6.21.

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 3

G(3) = 0.100000000000000 01 + 0.000000000000000 00 I
G(2) = -0.100000000000000 01 + -0.250000000000000 01 I
G(1) = -0.500000000000000 01 + -0.100000000000000 01 I

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.200000000000000 01 + 0.200000000000000 01 I	3	NO INITIAL APPROXIMATIONS
ROOT(2) = -0.999999999999999 00 + 0.499999999999999 00 I	3	NO INITIAL APPROXIMATIONS

Exhibit 6.21. Roots Are: 2+2i (3), 1+2i (2), -1+.5i (3)

REPEATED USE OF THE GREATEST COMMON DIVISOR AND NEWTONS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS
POLYNOMIAL NUMBER 4

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 12 THE COEFFICIENTS ARE

P(13) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(12) = -0.1200000000000000 02 + -0.0000000000000000 00 I
P(11) = 0.1200000000000000 03 + 0.0000000000000000 00 I
P(10) = -0.2800000000000000 03 + -0.0000000000000000 00 I
P(9) = 0.7800000000000000 03 + 0.0000000000000000 00 I
P(8) = -0.1632000000000000 04 + -0.0000000000000000 00 I
P(7) = 0.2624000000000000 04 + 0.0000000000000000 00 I
P(6) = -0.3264000000000000 04 + -0.0000000000000000 00 I
P(5) = 0.3120000000000000 04 + 0.0000000000000000 00 I
P(4) = -0.2240000000000000 04 + -0.0000000000000000 00 I
P(3) = 0.1152000000000000 04 + 0.0000000000000000 00 I
P(2) = -0.3840000000000000 03 + -0.0000000000000000 00 I
P(1) = 0.6400000000000000 02 + 0.0000000000000000 00 I

NO ROOTS OF MULTIPLICITY 1

NO ROOTS OF MULTIPLICITY 2

Exhibit 6.22.

NO ROOTS OF MULTIPLICITY 3

NO ROOTS OF MULTIPLICITY 4

NO ROOTS OF MULTIPLICITY 5

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 6

G(3) = 0.1000000000000000 01 + 0.0000000000000000 00 I
 G(2) = -0.2000000000000000 01 + -0.0000000000000000 00 I
 G(1) = 0.2000000000000000 01 + 0.0000000000000000 00 I

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = 0.1000000000000000 330 01 + 0.999999999997060 00 I	6	NO INITIAL APPROXIMATIONS
ROOT(2) = 0.1000000000000000 330 01 + -0.999999999997060 00 I	6	NO INITIAL APPROXIMATIONS

Exhibit 6.22. Roots Are: $1+i(6)$, $1-i(6)$

REPEATED USE OF THE GREATEST COMMON DIVISOR AND MULLERS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS POLYNOMIAL NUMBER 1

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 6 THE COEFFICIENTS ARE

P(7) = 0.100000000000000 01 + 0.000000000000000 00 I
P(6) = -0.700000000000010 01 + -0.105000000000000 02 I
P(5) = -0.280000000000000 02 + 0.580000000000000 02 I
P(4) = 0.170000000000000 03 + 0.150000000000000 01 I
P(3) = -0.730000000000000 02 + -0.251000000000000 03 I
P(2) = -0.228000000000000 03 + 0.104000000000000 03 I
P(1) = 0.720000000000000 02 + 0.104000000000000 03 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 1

G(2) = 0.100000000000000 01 + 0.000000000000000 00 I
G(1) = 0.99999999999973500 00 + -0.50000000000175420 00 I

ROOTS OF G(X)

ROOT(1) = -0.99999999999973500 00 + 0.500000000000175410 00 I 0.48296291156562790 00 + 0.12940952844381870 00 I

ROOTS OF P(X)

MULTIPLICITIES INITIAL APPROXIMATION

ROOT(1) = -0.99999999999973500 00 + 0.500000000000175410 00 I 1 0.48296291156562790 00 + 0.12940952844381870 00 I

Exhibit 6.23.

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 2

G(2) = 0.1000000000000000 01 + 0.0000000000000000 00 I
G(1) = -0.9999999999999999 79D 00 + -0.1999999999999999 6845D 01 I

ROOTS OF G(X)

INITIAL APPROXIMATION

ROOT(1) = 0.9999999999999999 179D 00 + 0.1999999999999999 6845D 01 I 0.4829629115656279D 00 + 0.1294095284438187D 00 I

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

ROOT(1) = 0.9999999999999999 179D 00 + 0.1999999999999999 6845D 01 I 2 0.4829629115656279D 00 + 0.1294095284438187D 00 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 3

2) = 0.1000000000000000 01 + 0.0000000000000000 00 I
1) = -0.1999999999999999 67D 01 + -0.2000000000001519D 01 I

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

ROOT(3) = 0.1999999999999999 67D 01 + 0.2000000000001519D 01 I 3 NO INITIAL APPROXIMATIONS

Exhibit 6.23. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+.5i$

REPEATED USE OF THE GREATEST COMMON DIVISOR AND MULLERS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS
POLYNOMIAL NUMBER 2

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF P(X) IS 15 THE COEFFICIENTS ARE

P(16) = 0.4800000000000000 02 + 0.0000000000000000 00 I
P(15) = 0.2557120000000000 03 + -0.3840000000000000 03 I
P(14) = -0.7355358600000000 02 + -0.2189600000000000 04 I
P(13) = -0.3859565600000000 04 + -0.6946851456000000 04 I
P(12) = -0.1733386464000000 05 + -0.1420625972800000 05 I
P(11) = -0.4967989270400000 05 + -0.1765857464000000 05 I
P(10) = -0.1022394522130000 06 + -0.6030664232000000 04 I
P(9) = -0.1642742200560000 06 + 0.4137366230400000 05 I
P(8) = -0.2036625884200000 06 + 0.1093899227600000 06 I
P(7) = -0.1871255780010000 06 + 0.1929865440330000 06 I
P(6) = -0.1274997298590000 06 + 0.2171341227420000 06 I
P(5) = -0.2814692716800000 05 + 0.1928489727960000 06 I
P(4) = 0.1329434434800000 05 + 0.1038130226550000 06 I
P(3) = -0.3059300774700000 05 + -0.2998989141300000 05 I
P(2) = -0.1835899020000000 03 + -0.1827632160000000 03 I
P(1) = 0.2755620000000000 00 + 0.2755620000000000 00 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 1

G(2) = 0.4800000000000000 02 + 0.0000000000000000 00 I
G(1) = 0.1119999928960310 03 + -0.8274988658740768D-06 I

ROOTS OF G(X)

INITIAL APPROXIMATION

ROOT(1) = -0.23333333318533397D 01 + 0.1723955970570993D-07 I 0.4829629115656279D 00 + 0.1294095284438187D 00 I

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

Exhibit 6.24.

ROOT(1) = -0.23333333331853397D 01 + 0.1723955970570993D-07 I 1 0.4829629115656279D 00 + 0.1294095284438187D 00 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 2

G(5) = 0.1000000000000000D 01 + 0.0000000000000000D 00 I
 G(4) = -0.2999970655863438D-02 + -0.9999999656362725D 00 I
 G(3) = 0.224999993117885D 01 + 0.3000058598117361D-02 I
 G(2) = -0.6750109902460278D-02 + -0.225000129067665D 01 I
 G(1) = 0.4643479465382683D-06 + 0.6749603982170172D-02 I

ROOTS OF G(X)

INITIAL APPROXIMATION

ROOT(1) = 0.2999823203921417D-02 + -0.2056988264688640D-06 I
 ROOT(2) = 0.4767581845683165D-06 + 0.100000055850184D 01 I
 ROOT(3) = 0.233172161693712D-07 + -0.149999972431725D 01 I
 ROOT(4) = -0.3526234587180127D-06 + 0.149999958791664D 01 I

0.4829629115656279D 00 + 0.1294095284438187D 00 I
 0.7071067553046346D 00 + 0.7071068070684595D 00 I
 SOLVED BY DIRECT METHOD
 SOLVED BY DIRECT METHOD

ROOTS OF P(X)

INITIAL APPROXIMATION

ROOT(1) = 0.2999823203921417D-02 + -0.2056988264688650D-06 I 2
 ROOT(2) = 0.4767581845683165D-06 + 0.100000055850184D 01 I 2
 ROOT(3) = 0.2340917191041074D-07 + -0.149999972431725D 01 I 2
 ROOT(4) = -0.3527154144586705D-06 + 0.149999958791664D 01 I 2

0.4829629115656279D 00 + 0.1294095284438187D 00 I
 0.7071067553046346D 00 + 0.7071068070684595D 00 I
 NO INITIAL APPROXIMATIONS
 NO INITIAL APPROXIMATIONS

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 3

G(3) = 0.1000000000000000D 01 + 0.0000000000000000D 00 I
 G(2) = 0.9999999853705541D 00 + -0.2600000017162632D 01 I
 G(1) = 0.300000002533335D 01 + -0.3000000031791514D 01 I

ROOTS OF G(X)

INITIAL APPROXIMATION

Exhibit 6.24.

ROOT(1) = -0.9999999999998587D 08 + -0.10000000000000000002D 01 I
ROOT(2) = 0.14629304614932610-07 + 0.30000000017162634D 01 I
0.4829629115656279D 00 + 0.1294095284436187D 00 I
SOLVED BY DIRECT METHOD

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT(1) = -0.9999999999998586D 00 + -0.10000000000000000002D 01 I	3	0.4829629115656279D 00 + 0.1294095284436187D 00 I
ROOT(2) = 0.1462930456963442D-07 + 0.30000000017162634D 01 I	3	NO INITIAL APPROXIMATIONS

Exhibit 6.24. Roots Are: -2.33 , $.003(2)$, $i(2)$, $1.5i(2)$,
 $-1.5i(2)$, $3i(3)$, $-1-i(3)$

REPEATED USE OF THE GREATEST COMMON DIVISOR AND MULLERS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS
POLYNOMIAL NUMBER 3

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.100-02
TEST FOR CONVERGENCE. 0.100-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.100-19
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

THE DEGREE OF PIX) IS 8 THE COEFFICIENTS ARE

P(9) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(8) = -0.5000000000000000 01 + -0.1150000000000000 02 I
P(7) = -0.5175000000000000 02 + 0.4300000000000000 02 I
P(6) = 0.1572500000000000 03 + 0.1446250000000000 03 I
P(5) = 0.3075000000000000 03 + -0.3475000000000000 03 I
P(4) = -0.4952500000000000 03 + -0.4948750000000000 03 I
P(3) = -0.5857500000000000 03 + 0.4247500000000000 03 I
P(2) = 0.1810000000000000 03 + 0.4420000000000000 03 I
P(1) = 0.1580000000000000 03 + 0.6000000000000000 01 I

NO ROOTS OF MULTIPLICITY 1

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF PIX) WHICH HAVE MULTIPLICITY 2

G(2) = 0.1000000000000000 01 + 0.0000000000000000 00 I
G(1) = -0.9999999999999999 00 + -0.1999999999999999 01 I

ROOTS OF G(X)

INITIAL APPROXIMATION

ROOT(1) = 0.9999999999999999 00 + 0.1999999999999999 01 I
0.4829629115656279D 00 + 0.1294095284438187D 00 I

Exhibit 6.25.

ROOTS OF P(X) MULTIPLICITIES INITIAL APPROXIMATION

ROOT(1) = 0.99999999999946498D 00 + 0.19999999999996468D 01 I 2 0.4829629115656279D 00 + 0.1294095284438187D 00 I

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 3

G(3) = 0.1000000000000000D 01 + 0.0000000000000000D 00 I
 G(2) = -0.10000000000002539D 01 + -0.25000000000001629D 01 I
 G(1) = -0.30000000000003296D 01 + -0.10000000000000421D 01 I

ROOTS OF G(X) INITIAL APPROXIMATION

ROOT(1) = -0.9999999999999928D 00 + 0.4999999999999759D 00 I 0.4829629115656279D 00 + 0.1294095284438187D 00 I
 ROOT(2) = 0.20000000000002532D 01 + 0.20000000000001653D 01 I SOLVED BY DIRECT METHOD

IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(2) = 0.20000000000002532D 01 + 0.20000000000001653D 01 I
 DID NOT CONVERGE AFTER 200 ITERATIONS
 THE PRESENT APPROXIMATION IS 0.2001999974253328D 01 + 0.2001999974252448D 01 I

ROOTS OF P(X) MULTIPLICITIES INITIAL APPROXIMATION

ROOT(1) = -0.9999999999999930D 00 + 0.4999999999999759D 00 I 3 0.4829629115656279D 00 + 0.1294095284438187D 00 I

NOT ALL ROOTS OF THE ABOVE POLYNOMIAL, G, WERE FOUND

Exhibit 6.25. Roots Are: $2+2i$ (3), $1+2i$ (2), $-1+5i$ (3)

REPEATED USE OF THE GREATEST COMMON DIVISOR AND MULLERS METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIALS
POLYNOMIAL NUMBER 4

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR ZERO IN SUBROUTINE GCD. 0.10D-02
TEST FOR CONVERGENCE. 0.10D-09
TEST FOR ZERO IN SUBROUTINE QUAD. 0.10D-19
RADIUS TO START SEARCH. 0.00D 00
RADIUS TO END SEARCH. 0.00D 00

THE DEGREE OF PIX IS 12 THE COEFFICIENTS ARE

P(13) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P(12) = -0.1200000000000000 02 + -0.0000000000000000 00 I
P(11) = 0.7200000000000000 02 + 0.0000000000000000 00 I
P(10) = -0.2500000000000000 03 + -0.0000000000000000 00 I
P(9) = 0.7800000000000000 03 + 0.0000000000000000 00 I
P(8) = -0.1832000000000000 04 + -0.0000000000000000 00 I
P(7) = 0.2824000000000000 04 + 0.0000000000000000 00 I
P(6) = -0.3264000000000000 04 + -0.0000000000000000 00 I
P(5) = 0.3120000000000000 04 + 0.0000000000000000 00 I
P(4) = -0.2240000000000000 04 + -0.0000000000000000 00 I
P(3) = 0.1152000000000000 04 + 0.0000000000000000 00 I
P(2) = -0.3840000000000000 03 + -0.0000000000000000 00 I
P(1) = 0.6400000000000000 02 + 0.0000000000000000 00 I

NO ROOTS OF MULTIPLICITY 1

NO ROOTS OF MULTIPLICITY 2

Exhibit 6.26.

NO ROOTS OF MULTIPLICITY 3

NO ROOTS OF MULTIPLICITY 4

NO ROOTS OF MULTIPLICITY 5

THE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE ROOTS OF P(X) WHICH HAVE MULTIPLICITY 6

G13) = 0.1000000000000000 01 + 0.0000000000000000 00 I
G12) = -0.2000000000000000 01 + -0.0000000000000000 00 I
G11) = 0.2000000000000000 01 + 0.0000000000000000 00 I

ROOTS OF G(X)

INITIAL APPROXIMATION

R00T(1) = 0.1000000000000000 33D 01 + 0.9999999999999707D 00 I 0.4829629115656279D 00 + 0.1294095284438187D 00 I
R00T(2) = 0.1000000000000000 33D 01 + -0.9999999999999707D 00 I SOLVED BY DIRECT METHOD

ROOTS OF P(X)

MULTIPLICITIES

INITIAL APPROXIMATION

R00T(1) = 0.1000000000000000 33D 01 + 0.9999999999999707D 00 I 6 0.4829629115656279D 00 + 0.1294095284438187D 00 I
R00T(2) = 0.1000000000000000 33D 01 + -0.9999999999999707D 00 I 6 NO INITIAL APPROXIMATIONS

Exhibit 6.26. Roots Are: $1+i(6)$, $1-i(6)$

REFERENCES

1. S. D. Conte, Elementary Numerical Analysis, McGraw Hill, New York, 1965.
2. Peter Henrici, Elements of Numerical Analysis, John Wiley and Sons, Inc., New York, 1964.
3. Thomas R. McCalla, Introduction to Numerical Methods and FORTRAN Programming, John Wiley and Sons, Inc., New York, 1967.
4. David E. Muller, A method for solving algebraic equations using an automatic computer, Math. Tables and Aids to Comp., 10 (1956), 208-215.

APPENDIX A

SPECIAL FEATURES OF NEWTON'S AND MULLER'S PROGRAMS

Several special features have been provided in each program as an aid to the user and to improve accuracy of the results. These are explained and illustrated below.*

1. Generating Approximations

If the user does not have initial approximations available, subroutine GENAPP can systematically generate, for an N^{th} degree polynomial, N initial approximations of increasing magnitude, beginning with the magnitude specified by XSTART. If XSTART is 0., XSTART is automatically initialized to 0.5 to avoid the approximation $0. + 0.i$. The approximations are generated according to the formula:

$$X_K = (XSTART + 0.5K) (\cos \beta + i \sin \beta)$$

where

$$\beta = \frac{\pi}{12} + K \frac{\pi}{6}, K = 0, 1, 2, \dots$$

To accomplish this, the user defined the number of initial approximations to be read (NIAP) on the control card to be zero (0) or these

*These illustrations are representative of Newton's method in double precision. The control cards for Muller's method are similarly prepared.

columns (7-8) may be left blank. If XSTART is left blank, it is interpreted as 0.

For example, a portion of a control card which generates initial approximations beginning at the origin for a seventh degree polynomial is shown in Example A.1.

Variable Name

Card Columns

1	2		4	5		7	8		6		7		7		7		8
									4		0		2		8		0
N	O	P		N		N	I										
O						A											
P						P											
O																	
L																	
Y																	
1				7													

Example

Example A.1

The approximations are generated in a spiral configuration as illustrated in Figure A.1. Exhibit 6.1 is an example of output resulting from generated approximations.

Example A.2 shows a portion of a control card which generated initial approximations beginning at a magnitude of 25.0 for a sixth degree polynomial.

Each altered approximation is then taken as a starting approximation. Each initial or altered approximation which does not produce convergence is printed as in Exhibit A.1. If none of the six starting approximations produce convergence, the next initial approximation is taken, and the process repeated. The six approximations are spaced 60 degrees apart on a circle of radius $|X_0|$ centered at the origin as illustrated in Figure A.2.

3. Searching the Complex Plane

By use of initial approximations and the altering technique, any region of the complex plane in the form of an annulus centered at the origin can be searched for roots. This procedure can be accomplished in two ways.

The first way is more versatile but requires more effort on the part of the user. Specifically selected initial approximations can be used to define particular regions to be searched. For example, if the roots of a particular polynomial are known to have magnitudes between 20 and 40, an annulus of inner radius 20 and outer radius 40 could be searched by using the initial approximations $20. + i$, $23. + i$, $26. + i$, $29. + i$, $32. + i$, $35. + i$, $38. + i$, $40. + i$.

By generating initial approximations internally, the program can search an annulus centered at the origin of inner radius XSTART and outer radius XEND. Values for XSTART and XEND are supplied on the control card by the user. Example A.3 shows a portion of a control card to search the above annulus of inner radius 20.0 and outer radius 40.0.

4. Improving Zeros Found

After the zeros of a polynomial are found, they are printed under the heading "Before the Attempt to Improve Accuracy." They are then used as initial approximations with Newton's (Muller's) method applied each time to the full (undeflated) polynomial. In most cases, zeros that have lost accuracy due to roundoff error in the deflation process are improved. The improved zeros are then printed under the heading "After the Attempt to Improve Accuracy." Since each root is used as an approximation to the original (undeflated) polynomial, it is possible that the root may converge to an entirely different root. This is especially true where several zeros are close together. Therefore, the user should check both lists of zeros to determine whether or not this has occurred. See Exhibit 6.4.

5. Solving Quadratic Polynomial

After $N-2$ roots of an N^{th} degree polynomial have been extracted, the remaining quadratic, $aX^2 + bX + c$, is solved using the quadratic formula

$$X = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

for the two remaining roots. These are indicated by the words "Solved By Direct Method" in the initial approximation column. If only a polynomial of degree 1 is to be solved, the solution is found directly as $(X - C) = 0$ implies $X = C$.

6. Missing Roots

If not all N roots of an N^{th} degree polynomial are found, the coefficients of the remaining deflated polynomial are printed under the heading "Coefficients of Deflated Polynomial For Which No Zeros Were Found." The user may then work with this polynomial in an attempt to find the remaining roots. The coefficient of the highest degree term will be printed first (Exhibit A.2).

7. Miscellaneous

By using various combinations of values for NIAP, XSTART, and XEND, the user has several options available as illustrated below.

Example A.5 shows the control card for a seventh degree polynomial. Three initial approximations are supplied by the user. At most three distinct roots will be found and the remaining deflated polynomial will be printed (Exhibit A.2).

1	2		4	5		7	8		6	7		7	7	8
									4	0		2	8	0
N						N								
O						I								
P						A								
O						P								
L														
Y														
1				7		3								

Example A.5

Note that if several roots are known to the user, they may be "divided out" of the original polynomial by using this procedure.

Example A.6 indicates that 2 initial approximations are supplied by the user to a 7th degree polynomial. After these approximations are used the circle of radius 15 will be searched for the remaining roots.

1	2		4	5		7	8		6	7	7	7	8
									4	0	2	8	0
N			N			N			XSTART		XEND		
P						I							
O						A							
L						P							
Y													
1			7			2					1.5D+01		

Example A.6

By defining XSTART between 0. and 15. an annulus instead of the circle will be searched (Exhibit A.3).




Figure A.1. Generating Initial Approximations




Figure A.2. Altering Approximations

Figure A.3. Distribution of Approximations

NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 2 OF DEGREE 3

THE COEFFICIENTS OF P(X) ARE

P1 1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
P1 2) = 0.2000000000000000 01 + 0.0000000000000000 00 I
P1 3) = -0.1000000000000000 01 + -0.0000000000000000 00 I
P1 4) = -0.2000000000000000 01 + -0.0000000000000000 00 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 0
MAXIMUM NUMBER OF ITERATIONS. 3
TEST FOR CONVERGENCE. 0.100-03
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

NO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AFTER 3 ITERATIONS.

0.48296291156562790 00 + 0.12940952844381870 00 I INITIAL APPROXIMATION
-0.48296291156562790 00 + -0.12940952844381870 00 I ALTERED APPROXIMATION
0.12940949308846860 00 + 0.48296292103906440 00 I ALTERED APPROXIMATION
-0.12940949308846860 00 + -0.48296292103906440 00 I ALTERED APPROXIMATION
-0.35355342941614020 00 + 0.35355335177040300 00 I ALTERED APPROXIMATION
0.35355342941614020 00 + -0.35355335177040300 00 I ALTERED APPROXIMATION
0.70710675530463460 00 + 0.70710680706845950 00 I INITIAL APPROXIMATION
-0.70710675530463460 00 + -0.70710680706845950 00 I ALTERED APPROXIMATION
-0.25881912759833590 00 + 0.96592580418437740 00 I ALTERED APPROXIMATION
0.25881912759833590 00 + -0.96592580418437740 00 I ALTERED APPROXIMATION
-0.96592586102499680 00 + 0.25881891546623570 00 I ALTERED APPROXIMATION
0.96592586102499680 00 + -0.25881891546623570 00 I ALTERED APPROXIMATION
0.38822847926540560 00 + 0.14488887631171930 01 I INITIAL APPROXIMATION
-0.38822847926540560 00 + -0.14488887631171930 01 I ALTERED APPROXIMATION
-0.10606602882484210 01 + 0.10606600553112090 01 I ALTERED APPROXIMATION
0.10606602882484210 01 + -0.10606600553112090 01 I ALTERED APPROXIMATION
-0.14488886778562400 01 + -0.38822879746355020 00 I ALTERED APPROXIMATION
0.14488886778562400 01 + 0.38822879746355020 00 I ALTERED APPROXIMATION

COEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO ZEROS WERE FOUND

D1 1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
D1 2) = 0.2000000000000000 01 + 0.0000000000000000 00 I
D1 3) = -0.1000000000000000 01 + -0.0000000000000000 00 I
D1 4) = -0.2000000000000000 01 + -0.0000000000000000 00 I

Exhibit A.1.

NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS
POLYNOMIAL NUMBER 1 OF DEGREE 7

THE COEFFICIENTS OF P(X) ARE

P1 1 = 0.1000000000000000 01 + 0.0000000000000000 00 I
P1 2 = -0.1000000000000000 01 + 0.1100000000000000 02 I
P1 3 = -0.5900000000000000 02 + -0.2900000000000000 02 I
P1 4 = 0.1950000000000000 03 + -0.1890000000000000 03 I
P1 5 = 0.7000000000000000 02 + 0.7230000000000000 03 I
P1 6 = -0.1624000000000000 04 + -0.6960000000000000 03 I
P1 7 = 0.1922000000000000 04 + -0.1832000000000000 04 I
P1 8 = 0.1596000000000000 04 + 0.1640000000000000 04 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 3
MAXIMUM NUMBER OF ITERATIONS. 200
TEST FOR CONVERGENCE. 0.100-09
TEST FOR MULTIPLICITIES. 0.100-01
RADIUS TO START SEARCH. 0.000 00
RADIUS TO END SEARCH. 0.000 00

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT1 1) = -0.2999999999999970 01 + -0.3000000000000020 01 I	1	-0.3500000000000000 01 + -0.3500000000000000 01 I
ROOT1 2) = 0.2000000000000000 01 + 0.2000000000000000 01 I	1	0.2500000000000000 01 + 0.2500000000000000 01 I
ROOT1 3) = -0.9999999999999962 00 + -0.3999999999999940 01 I	1	-0.1500000000000000 01 + -0.4500000000000000 01 I

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT1 1) = -0.2999999999999960 01 + -0.3000000000000010 01 I	1	-0.3500000000000000 01 + -0.3500000000000000 01 I
ROOT1 2) = 0.2000000000000000 01 + 0.2000000000000000 01 I	1	0.2500000000000000 01 + 0.2500000000000000 01 I
ROOT1 3) = -0.9999999999999970 00 + -0.3999999999999960 01 I	1	-0.1500000000000000 01 + -0.4500000000000000 01 I

COEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO ZEROS WERE FOUND

D1 1) = 0.1000000000000000 01 + 0.0000000000000000 00 I
D1 2) = -0.2999999999999990 01 + 0.4000000000000000 01 I
D1 3) = -0.2000000000000000 02 + -0.1899999999999990 02 I
D1 4) = 0.4100000000000000 02 + -0.2200000000000000 02 I
D1 5) = 0.2300000000000000 02 + 0.4100000000000000 02 I

Exhibit A.2. Roots Are: $-1 - 4i, -2 - 3i, -3 - 3i, -1 - i, 2 + 2i, 4 - i, 2 - i$.

NEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS POLYNOMIAL NUMBER 1 OF DEGREE 7

THE COEFFICIENTS OF P(X) ARE

P1 1 = 0.1000000000000000 01 + 0.0000000000000000 00 I
P1 2 = -0.1000000000000000 01 + 0.1100000000000000 02 I
P1 3 = -0.5900000000000000 02 + -0.2900000000000000 03 I
P1 4 = 0.1950000000000000 03 + -0.1690000000000000 04 I
P1 5 = 0.7000000000000000 04 + 0.7230000000000000 05 I
P1 6 = -0.1624000000000000 05 + -0.6960000000000000 06 I
P1 7 = 0.1922000000000000 06 + -0.1832000000000000 07 I
P1 8 = 0.1596000000000000 07 + 0.1692000000000000 08 I

NUMBER OF INITIAL APPROXIMATIONS GIVEN. 2

MAXIMUM NUMBER OF ITERATIONS. 200

TEST FOR CONVERGENCE. 0.100-09

TEST FOR MULTIPLICITIES. 0.100-01

RADIUS TO START SEARCH. 0.700 01

RADIUS TO END SEARCH. 0.150 02

BEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)

	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT1 1) = -0.2999999999999970 01 + -0.3000000000000000 01 I	1	-0.3500000000000000 01 + -0.3500000000000000 01 I
ROOT1 2) = 0.2000000000000000 01 + 0.2000000000000000 01 I	1	0.2500000000000000 01 + 0.2500000000000000 01 I
ROOT1 3) = 0.4000000000000000 01 + -0.1000000000000000 01 I	1	0.6761480761918791 01 + 0.1811733982134620 01 I
ROOT1 4) = 0.1999999999999970 01 + -0.9999999999999970 00 I	1	0.5303300664784760 01 + 0.5303301053013447 01 I
ROOT1 5) = -0.9999999999999980 00 + -0.1000000000000000 01 I	1	0.2070551889415497 01 + 0.7727406736625032 01 I
ROOT1 6) = -0.9999999999999970 00 + -0.3999999999999950 01 I	1	SOLVED BY DIRECT METHOD
ROOT1 7) = -0.2000000000000000 01 + -0.3000000000000000 01 I	1	SOLVED BY DIRECT METHOD

AFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS OF P(X) ARE

ROOTS OF P(X)

	MULTIPLICITIES	INITIAL APPROXIMATION
ROOT1 1) = -0.2999999999999960 01 + -0.3000000000000000 01 I	1	-0.3500000000000000 01 + -0.3500000000000000 01 I
ROOT1 2) = 0.2000000000000000 01 + 0.2000000000000000 01 I	1	0.2500000000000000 01 + 0.2500000000000000 01 I
ROOT1 3) = 0.4000000000000000 01 + -0.1000000000000000 01 I	1	0.6761480761918791 01 + 0.1811733982134620 01 I
ROOT1 4) = 0.2000000000000000 01 + -0.9999999999999980 00 I	1	0.5303300664784760 01 + 0.5303301053013447 01 I
ROOT1 5) = -0.9999999999999970 00 + -0.1000000000000000 01 I	1	0.2070551889415497 01 + 0.7727406736625032 01 I
ROOT1 6) = -0.9999999999999980 00 + -0.4000000000000000 01 I	1	SOLVED BY DIRECT METHOD
ROOT1 7) = -0.2000000000000000 01 + -0.3000000000000000 01 I	1	SOLVED BY DIRECT METHOD

Exhibit A.3. Roots Are: $-1 - 4i$, $-2 - 3i$, $-3 - 3i$, $-1 - i$, $2 + 2i$, $4 - i$, $2 - i$.

APPENDIX B

NEWTON'S METHOD

1. Use of the Program

A double precision FORTRAN IV program using Newton's method is presented here. Flow charts for this program are given in Figure B.6 while Table B.VIII gives a FORTRAN IV listing of this program. Single precision variables are listed in some of the tables. The single precision variables are used in the flow charts and the corresponding double precision variables can be obtained from the appropriate tables.

The program is designed to solve polynomials of degree 25 or less. Both the coefficient of the highest degree term and the constant coefficient should be non-zero. In order to solve polynomials of degree N , where $N > 25$, certain array dimensions must be changed. These are listed in Table B.I for the main program and subprograms in double precision.

TABLE B.I
PROGRAM CHANGES FOR SOLVING POLYNOMIALS
OF DEGREE GREATER THAN 25
BY NEWTON'S METHOD

Double Precision

Main Program

RA(N+1), VA(N+1)
RB(N+1), VB(N+1)
RC(N+1), VC(N+1)
RD(N+1), VD(N+1)
RCOEF(N+1), VCOEF(N+1)
MULT(N)
RXZERO(N), VXZERO(N)
RX(N), VX(N)
RXINIT(N), VXINIT(N)

Subroutine HORNER

RA(N+1), VA(N+1)
RB(N+1), VB(N+1)
RC(N+1), VC(N+1)

Subroutine BETTER

RXZERO(N), VXZERO(N)
RX(N), VX(N)
RA(N+1), VA(N+1)
RCOEF(N+1), VCOEF(N+1)
RC(N+1), VC(N+1)
RB(N+1), VB(N+1)

Subroutine GENAPP

APPR(N), APPI(N)

Subroutine QUAD

UA(N+1), VA(N+1)
UROOT(N), VROOT(N)
MULTI(N)

Table B.II lists the system functions used in the program of Newton's method. In the table "d" denotes a double precision variable name.

TABLE B.II
SYSTEM FUNCTIONS USED IN NEWTON'S METHOD

Double Precision

DABS(d)	- obtain absolute value
DCOS(d)	- obtain cosine of angle
DSIN(d)	- obtain sine of angle
DATAN2(d ₁ ,d ₂)	- arctangent of d ₁ /d ₂
DSQRT(d)	- square root

2. Input Data for Newton's Method

The input data for Newton's method is grouped into polynomial data sets. Each polynomial data set consists of the data for one and only one polynomial. As many polynomials as the user desires may be solved by placing the polynomial data sets one behind the other. Each polynomial data set consists of three kinds of information placed in the following order:

1. Control information.
2. Coefficients of the polynomial.
3. Initial approximations. These may be omitted as described in Appendix A, § 1.

An end card follows the entire collection of data sets. It indicates that there is no more data to follow and terminates execution of the program. This information is displayed in Figure B.1 and described below. For the double precision data, the D-type specification should

be used. All data should be right justified. The recommendations given in Table B.III are those found to give best results on the IBM 360/50 computer which has a 32 bit word.

Control Information

The control card is the first card of the polynomial data set and contains the information given in Table B.III. See Figure B.2.

TABLE B.III
CONTROL DATA FOR NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
NOPOLY	c.c. 1-2	Number of the polynomial. Integer. Right justified.
N	c.c. 4-5	Degree of the polynomial. Integer. Right justified.
NIAP	c.c. 7-8	Number of initial approxi- mations to be read. Integer. If no approximations are given, this should be left blank.
MAX	c.c. 19-21	Maximum number of iterations. Integer. Right justified. 200 is recommended.
EPSCNV	c.c. 30-35	Convergence requirement. Double precision. 1.D-10 is recommended.

TABLE B.III (Continued)

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
EPSQ	c.c. 37-42	Tolerance check for zero (0) in subroutine QUAD. Double precision. Right justify. 1.D-20 is recommended.
EPSMUL	c.c. 44-49	Multiplicity requirement. Double precision. Right justify. 1.D-02 is recommended.
XSTART	c.c. 64-70	Magnitude at which to begin generating initial approximations. Double precision. Right justify. This is a special feature of the program and may be omitted.
XEND	c.c. 72-78	Magnitude at which to end the generating of initial approximations. Double precision. Right justify. This is a special feature of the program and may be omitted.
KCHECK	c.c. 80	This should be left blank.

Coefficients of the Polynomial

The coefficient cards follow the control card. For an N^{th} degree polynomial, $N+1$ coefficients must be entered one per card. The coefficient of the highest degree term is entered first. For example, if the polynomial $X^5 + 3X^4 + 2X + 5$ were to be solved, the order in which the coefficients would be entered is: 1, 3, 0, 0, 2, 5. Each

coefficient is entered, one per card, as described in Table B.IV and illustrated in Figure B.3.

TABLE B.IV
COEFFICIENT DATA FOR NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
RA (A in single precision)	c.c. 1-30	Real part of complex coefficient. Double precision. Right justify. If none, leave blank or enter 0.0D00.
VA (A in single precision)	c.c. 31-60	Imaginary part of complex coefficient. Double precision. Right justify. If none, leave blank or enter 0.0D00.

Initial Approximations

The initial approximation cards follow the set of coefficient cards. The number of initial approximations read must be the number specified on the control card and are entered, one per card, as given in Table B.V and illustrated in Figure B.4.

TABLE B.V
INITIAL APPROXIMATION DATA FOR NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
RXZERO (XZERO in single precision)	c.c. 1-30	Real part of complex number. Double precision. Right justify. If none, leave blank or enter 0.0D00.
VXZERO (XZERO in single precision)	c.c. 31-60	Imaginary part of complex number. Double precision. Right justify. If none, leave blank or enter 0.0D00.

End Card

The end card is the last card of the input data to the program. It indicates that there is no more data to be read. When this card is read, program execution is terminated. This card is described in Table B.VI and illustrated in Figure B.5.

TABLE B.VI
DATA TO END EXECUTION OF NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
KCHECK	c.c. 80	Must contain the number 1. Integer.

3. Variables Used in Newton's Method

The definitions of the major variables used in Newton's method are given in Table B.VII. The symbols used to indicate type are:

R - real variable
I - integer variable
C - complex variable
D - double precision
L - logical variable
A - alphanumeric variable

When two variables are listed, the one on the left is the real part of the corresponding single precision complex variable; the one on the right is the imaginary part. The symbols used to indicate disposition are:

E - entered
R - returned
ECR - entered, changed, and returned
C - variable in common

4. Description of Program Output

The output from Newton's method programs consist of the following information.

The number and degree of the polynomial are printed in the heading (Exhibit 6.1).

The coefficients are printed under the heading "THE COEFFICIENTS OF P(X) ARE." The coefficient of the highest degree term is listed first (Exhibit 6.1).

As an aid to ensure the control information is correct, the number of initial approximations given, maximum number of iterations, test for convergence, test for multiplicities, radius to start search, and radius to end search are printed as read from the control card (Exhibit 6.1).

The zeros found before and after the attempt to improve accuracy are printed. See Appendix A, § 4 for further explanation (Exhibit 6.1).

If not all zeros of the polynomial are found, the coefficients of the remaining unsolved polynomial will be printed, with coefficient of highest degree term first, under the heading "COEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO ZEROS WERE FOUND." See Appendix A, § 6. This is illustrated in Exhibit A.2.

The multiplicity of each zero is given under the title "MULTIPLICITIES" (Exhibit 6.1).

The initial approximation producing convergence to a root is printed to the right of the corresponding root and headed by "INITIAL APPROXIMATION." The initial approximations may be those supplied by the user, or generated by the program, or a combination of both (Exhibit A.3). See Appendix A, § 1 and § 2 for discussion of approximations. The message "SOLVED BY DIRECT METHOD" indicates that the corresponding root or roots was obtained by Subroutine QUAD. See Appendix A, § 5.

If an approximation does not produce convergence within the maximum number of iterations, it is printed under the heading "NO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AFTER XXX ITERATIONS." XXX is replaced by the maximum number of iterations. The type of the approximation, that is, initial approximation or altered approximations is given (Exhibit A.1). See Appendix A, § 1 and § 2 for discussion of approximations.

5. Informative and Error Messages

The output may contain informative or error messages. These are intended as an aid to the user and are described as follows:

"IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(X) = YYY DID NOT CONVERGE THE PRESENT APPROXIMATION AFTER ZZZ ITERATIONS IS PRINTED BELOW." X is the number of the zero, YYY is the value of the zero before the attempt to improve accuracy, ZZZ is the maximum number of iterations. This message indicates that a zero found before attempting to improve accuracy did not converge sufficiently when being used as an initial approximation on the full (undeflated) polynomial. The current approximation is printed in the list of improved zeros. In many cases, this failure to converge is a result of an ill-conditioned polynomial and this current approximation of the root may be better than its approximation before the attempt to improve accuracy. In most cases, the polynomial from which this root was first extracted had fewer multiple roots, due to deflations, than the original polynomial.

"THE VALUE OF THE DERIVATIVE AT X0 = XXX IS ZERO."

This message is printed as a result of the value of the derivative of the original polynomial at an approximation, XXX, being zero (0). It occurred in the attempt to improve the accuracy of a zero. The previous message is then printed.




Figure B.1. Sequence of Input Data for Newton's Method

[illegible]

Figure B.2. Control Card for Newton's Method

TABLE B. VII

VARIABLES USED IN NEWTON'S METHOD

<u>Single Precision</u> <u>Variable</u>	<u>Type</u>	<u>Double Precision</u> <u>Variable</u>	<u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
Main Program					
NOPOLY	I	NOPOLY	I		Number of the polynomial
N	I	N	I		Degree of the polynomial
NIAP	I	NIAP	I		Number of initial approximations to be read
MAX	I	MAX	I		Maximum number of iterations to be performed
EPSCNV	R	EPSCNV	D		Tolerance check for convergence
EPSMUL	R	EPSMUL	D		Tolerance check for multiplicities
EPSQ	R	EPSQ	D		Tolerance check for zero in subroutine QUAD
XSTART	R	XSTART	D		Magnitude from which to begin the search for zeros
XEND	R	XEND	D		Magnitude to end the search for zeros
KCHECK	I	KCHECK	I		Program Control. When KCHECK = 1, program will terminate execution.
NA	I	NA	I		Number of coefficients or original polynomial
A	C	RA,VA	D		Array containing the coefficients of original polynomial P(X)
NDEF	I	NDEF	I		Degree of current deflated polynomial
L	I	L	I		Counter for number of initial approximations used
ITER	I	ITER	I		Counter for number of iterations
NROOT	I	NROOT	I		Counter for number of roots found (counting multiplicities)
IALTER	I	IALTER	I		Counter for number of alterations of each initial approximation
ITIME	I	ITIME	I		Program control
K	I	K	I		Counter for number of distinct roots found
ND	I	ND	I		Program control & number of coefficient of deflated polynomial for which no zeros were found

TABLE B. VII (Continued)

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition of Argument</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>		
XO	C	RXO, VXO	D		Current approximation (X_n) to root
COEF	C	RCOEF, VCOEF	D		Working array containing coefficients of current deflated polynomial
DPX	C	RDPX, VDPX	D		Derivative of $P(X)$ at some value X
PX	C	RPX, VPX	D		Value of $P(X)$ at some point X
XZERO	C	RXZERO, VZXERO	D		Array containing the initial approximations
XNEW	C	RXNEW, VXNEW	D		New approximation (X_{n+1}) obtained from old approximation (X_n) by Newton's Algorithm
KANS	I	KANS	I		KANS = 1 implies convergence, KANS = 0 implies no convergence
MULT	I	MULT	I		Array containing the number of multiplicities of each root
X	C	RX, VX	D		Array containing the zeros of $P(X)$
XINIT	C	RXINIT, VXINIT	D		Array containing the initial or altered approximations which produced convergence to each root
NUM	I	NUM	I		Number of coefficients of current deflated polynomial
B	C	RB, VB	D		Array containing the coefficients of newly deflated polynomial
IROOT	I	IROOT	I		Number of distinct roots found by Newton's method, i.e. not solved for directly by subroutine QUAD
D	C	RD, VD	D		Array containing the coefficients of deflated polynomial for which no zeros were found
I01	I	I01	I		Unit number of input device
I02	I	I02	I		Unit number of output device
C	C	RC, VC	D		Array containing sequence of values leading to the derivative
EPSCHK	R	EPSCHK	D		Current tolerance for checking convergence or multiplicity

TABLE B. VII (Continued)

Single Precision		Double Precision		Disposition of Argument	Description
Variable	Type	Variable	Type		
Subroutine HORNER					
A	C	RA, VA	D	E	Array of coefficients of polynomial
B	C	RB, VB	D	R	Array of coefficients of deflated polynomial
NDEF	I	NDEF	I	E	Degree of polynomial
NUM	I	NUM	I		Number of coefficients of polynomial
XO	C	RXO, VXO	D	E	Point (X_n) at which to evaluate the polynomial and its derivative. Also current approximation (X_{n+1}) used to deflate the polynomial
PX	C	RPX, VPX	D	R	Value of polynomial at X_n
DPX	C	RDPX, VDPX	D	R	Value of the derivative of polynomial at X_n
C	C	RC, VC	D	R	Array of containing sequence of values leading to the derivative
Subroutine NEWTON					
PX	C	RPX, VPX	D	E	Value of polynomial at X_n
DPX	C	RDPX, VDPX	D	E	Derivative of polynomial at X_n
XO	C	RXO, VXO	D	E	Current approximation (X_n) to root
XNEW	C	RXNEW, VXNEW	D	R	New approximation (X_{n+1}) to root
Subroutine CHECK					
EPSLON	R	EPS	D	C	Tolerance for convergence or multiplicity check
PX	C	RPX, VPX	D	E	Value of P(X) at X_n
DPX	C	RDPX, VDPX	D	E	Derivative of P(X) at X_n
XO	C	RXO, VXO	D	E	Current approximations (X_{n+1}) to root
I02	I	I02	I	C	Unit number of output device
KANS	I	KANS	I	R	KANS = 1 implies convergence, KANS = 0 implies no convergence

TABLE B. VII (Continued)


<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition of Argument</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>		
Subroutine BETTER					
I02	I	I02	I	C	Unit number of output device
XZERO	C	RXZERO, VXZERO	D	E	Array of approximations
X	C	RX,VX	D	ECR	Array of roots
A	C	RA,VA	D	E	Coefficients of original (undeflated) polynomial, P(X)
COEF	C	RCOEF,VCOEF	D	E	Working array for coefficients of polynomial
NA	I	NA	I	E	Number of coefficients of original polynomial
X0	C	RX0,VX0	D		Current approximation (X _n) to root
DPX	C	RDPX,VDPX	D		Derivative of P(X) at X _n
PX	C	RPX,VPX	D		Value of P(X) at X _n
KANS	I	KANS	I		KANS = 1 implies convergence; KANS = 0 implies no convergence
ITER	I	ITER	I		Counter for number of iterations
XNEW	C	RXNEW,VXNEW	D		New approximation (X _{n+1}) to root
NN	I	NN	I		Degree of polynomial
C	C	RC,VC	D	E	Array containing the sequence of values leading to the derivative
K	I	K	I	E	Number of distinct roots of P(X) found
N	I	N	I	E	Degree of polynomial P(X)
B	C	RB,VB	D	E	Array of coefficients of deflated polynomial
MAX	I	MAX	I	C	Maximum number of iterations permitted
EPSCHK	R	EPS	D	C	Tolerance for checking convergence
Subroutine GENAPP					
APP	C	APPR,APPI	D	R	Array containing initial approximations
NAPP	I	NAPP	I	E	Number of initial approximations to be generated

TABLE B. VII (Continued)

Single Precision Variable	Double Precision Variable	Precision Type	Disposition of Argument	Description
XSTART	XSTART	R	D	ECR
BETA	BETA	R	D	Magnitude at which to begin generating approximations; also magnitude of the approximation being generated
U	APPR(I)	R	D	Argument of the complex approximation being generated
V	APPI(I)	R	D	Real part of complex approximation
				Imaginary part. of complex approximation
Subroutine ALTER				
XOLD	XOLDR,XOLDI	C	D	ECR
NALTER	NALTER	I	I	ECR
ITIME	ITIME	I	I	E
MAX	MAX	I	I	C
Y	XOLDI	R	D	Maximum number of iterations permitted Imaginary part of original initial approximation (unaltered)
X	XOLDR	R	D	Real part of original unaltered initial approximation
R	R	R	D	Magnitude of original unaltered initial approximation
BETA	BETA	R	D	Argument of new approximation
XOLDR	XOLDR	R	D	Real part of new approximation
XOLDI	XOLDI	R	D	Imaginary part of new approximation
I02	I02	I	I	C
				Unit number of output device
Subroutine QUAD				
A	UA,VA	C	D	E
NA	NA	I	I	E
ROOT	UROOT,VROOT	C	D	ECR
NROOT	NROOT	I	I	ECR
				Coefficients of polynomial to be solved Degree of polynomial Array of roots of P(X) (original polynomial) Number of distinct roots of P(X) (the original polynomial)

TABLE B. VII (Continued)

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>	<u>of Argument</u>	
MULTI	I	MULTI	I	ECR	Array containing multiplicities of each root
EPST	R	EPST	D	E	Tolerance check for the number zero
DISC	C	UDISC,VDISC	D		Value of the discriminant ($b^2 - 4ac$) of Quadratic
Subroutine COMSQT					
		UX,VX	D	E	Complex number for which the square root is desired
		UY,VY	D	R	Square root of the complex number






Figure B.6. (Continued)

BETTER

CHECK




Figure B.6. (Continued)

HORNER

NEWTON





Figure B.6. (Continued)

GENAPP



ALTER





Figure B.6. (Continued)

QUAD



COMSQT




Figure B.6. (Continued)

TABLE B. VIII

PROGRAM FOR NEWTON'S METHOD

```

C *****
C *
C * DOUBLE PRECISION PROGRAM FOR NEWTON'S METHOD
C *
C *
C * NEWTONS METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C * POLYNOMIAL OF MAXIMUM DEGREE 25 BY COMPUTING A SEQUENCE OF APPROX-
C * IMATIONS CONVERGING TO A ZERO OF THE POLYNOMIAL USING THE ITERATION
C * FORMULA
C *
C *       $X(N+1) = X(N) - P(X(N)) / P'(X(N))$ .
C *
C *****
0001  DOUBLE PRECISION RA,VA,RXZERO,VXZERO,RB,VB,RCOEF,VCOEF,RX,VX,RXINI
      IT,VXINIT,RC,VC,RO,VD,RPX,VPX,ROPX,ROPX,RXNEW,VXNEW,RXD,VXD,EPSCNK,
0002  2EPSCNV,EPSCQ,EPSCMUL,XSTART,XEND,ABPX,ABDPX
      DIMENSION RA(26),VA(26),RB(26),VB(26),RC(26),VC(26),RO(26),VD(26),R
      1RCOEF(26),VCOEF(26),MULT(25),RXZERO(25),VXZERO(25),RX(25),VX(25),R
      2XINIT(25),VXINIT(25)
0003  COMMON EPSCNK,MAX,IO2
0004  IO1=5
0005  IO2=6
0006  1 READ(IO1,1000) NOPOLY,N,NIAP,MAX,EPSCNV,EPSCQ,EPSCMUL,XSTART,XEND,KC
      1HECK
0007  IF(KCHECK.EQ.1) STOP
0008  NA=N+1
0009  READ(IO1,1010) (RA(I),VA(I),I=1,NA)
0010  WRITE(IO2,1030) NOPOLY,N
0011  WRITE(IO2,1040) (I,RA(I),VA(I),I=1,NA)
0012  WRITE(IO2,2060)
0013  WRITE(IO2,2000) NIAP
0014  WRITE(IO2,2010) MAX
0015  WRITE(IO2,2020) EPSCNV
0016  WRITE(IO2,2030) EPSCMUL
0017  WRITE(IO2,2040) XSTART
0018  WRITE(IO2,2050) XEND
0019  IF(NIAP.NE.0) GO TO 3
0020  NIAP=N
0021  CALL GENAPP(RXZERO,VXZERO,NIAP,XSTART)
0022  GO TO 4
0023  3 READ(IO1,1020) (RXZERO(I),VXZERO(I),I=1,NIAP)
0024  4 NDEF=N
0025  L=1
0026  ITER=0
0027  NROOT=0
0028  IROOT=0
0029  ITIME=0
0030  NO=0
0031  IALTER=0
0032  K=0
0033  RXO=RXZERO(L)
0034  VXO=VXZERO(L)
0035  DO 5 I=1,NA
0036  RCOEF(I)=RA(I)
0037  5 VCOEF(I)=VA(I)
0038  10 CALL HORNER(RCOEF,VCOEF,RXO,VXO,NDEF,RB,VB,RC,VC,RPX,VPX,ROPX,ROPX
      1)
0039  ABPX=DSQRT(RPX*RPX+VPX*VPX)
0040  ABDPX=DSQRT(ROPX*ROPX+VDPX*VDPX)

```

TABLE B. VIII (Continued)

0041	IF(ABDPX.NE.0.0) GO TO 20	
0042	IF(ABPX.EQ.0.0) GO TO 70	
0043	GO TO 110	
0044	20 CALL NEWTON(RPX,VPX,RPX,VPX,RXQ,VXQ,RXNEW,VXNEW)	
0045	ITER=ITER+1	200
0046	RXQ=RXNEW	204
0047	VXQ=VXNEW	205
0048	EPSCHK=EPSCHV	
0049	CALL CHECK(RPX,VPX,RPX,VPX,RXQ,VXQ,KANS)	
0050	IF(KANS.EQ.1) GO TO 70	194
0051	IF(ITER.GE.MAX) GO TO 40	
0052	GO TO 10	208
0053	40 CALL ALTER(RXZERO(L),VXZERO(L),IALTER,ITIME)	
0054	IF(IALTER.GT.5) GO TO 110	
0055	RXQ=RXZERO(L)	
0056	VXQ=VXZERO(L)	
0057	ITER=0	244
0058	GO TO 10	248
0059	60 ND=NDEF+1	
0060	DO 65 J=1,ND	
0061	RD(J)=RCOEF(J)	
0062	65 VD(J)=VCOEF(J)	
0063	GO TO 140	
0064	70 NROOT=NROOT+1	268
0065	K=K+1	272
0066	MULT(K)=1	276
0067	RX(K)=RXQ	280
0068	VX(K)=VXQ	281
0069	RXINIT(K)=RXZERO(L)	288
0070	VXINIT(K)=VXZERO(L)	289
0071	CALL HORNER(RCOEF,VCOEF,RXQ,VXQ,NDEF,RB,VB,RC,VC,RPX,VPX,RPX,VPX)	
0072	1)	
0073	80 IF(NROOT.GE.N) GO TO 147	
0074	NDEF=NDEF-1	
0075	NUM=NDEF+1	294
0076	DO 105 I=1,NUM	296
0077	RCOEF(I)=RB(I)	297
0078	105 VCOEF(I)=VB(I)	
0079	CALL HORNER(RCOEF,VCOEF,RXQ,VXQ,NDEF,RB,VB,RC,VC,RPX,VPX,RPX,VPX)	
0080	1)	
0081	ABPX=DSQRT(RPX*RPX+VPX*VPX)	
0082	ABDPX=DSQRT(RDPX*RDVX+VDPX*VDPX)	
0083	IF(ABDPX.NE.0.0) GO TO 107	
0084	IF(ABPX.EQ.0.0) GO TO 130	
0085	GO TO 110	
0086	107 CONTINUE	
0087	EPSCHK=EPSMUL	
0088	CALL CHECK(RPX,VPX,RPX,VPX,RXQ,VXQ,KANS)	
0089	IF(KANS.EQ.1) GO TO 130	300
0090	110 IF(NDEF.GT.2) GO TO 113	
0091	IROOT=K	
0092	CALL QUAD(RCOEF,VCOEF,NDEF,RX,VX,K,MULT,EPSQ)	
0093	GO TO 150	
0094	113 IF(L.LT.NIAP) GO TO 115	
0095	IF(XEND.EQ.0.0) GO TO 60	
0096	IF(XSTART.GT.XEND) GO TO 60	
0097	NIAP=N	
0098	CALL GENAPP(RXZERO,VXZERO,NIAP,XSTART)	

TABLE B. VIII (Continued)

0097	L=0	
0098	115 L=L+1	
0099	RXD=RXZERO(L)	312
0100	VXD=VXZERO(L)	313
0101	ITER=0	316
0102	IALTER=0	320
0103	GO TO 10	324
0104	130 MULT(K)=MULT(K)+1	328
0105	NROOT=NROOT+1	332
0106	GO TO 80	336
0107	140 IF(K.EQ.0) GO TO 160	338
0108	147 IROOT=K	
0109	150 WRITE(102,1025)	
0110	WRITE(102,1050)	
0111	WRITE(102,1060) (I,RX(I),VX(I),MULT(I),RXINIT(I),VXINIT(I),I=1,IRO 10T)	
0112	KKK=IROOT+1	
0113	IF(IROOT.LT.K) WRITE(102,1062) (I,RX(I),VX(I),MULT(I),I=KKK,K)	
0114	EPSCHE=EPSCNV	
0115	CALL BETTER(K,RXZERO,VXZERO,RX,VX,NA,RA,VA,RCOEF,VCOEF,N,RC,VC,RB, 1VB)	
0116	160 IF(K.EQ.0) GO TO 170	
0117	WRITE(102,1065)	
0118	WRITE(102,1050)	
0119	WRITE(102,1060) (I,RX(I),VX(I),MULT(I),RXINIT(I),VXINIT(I),I=1,IRO 10T)	
0120	KKK=IROOT+1	
0121	IF(IROOT.LT.K) WRITE(102,1062) (I,RX(I),VX(I),MULT(I),I=KKK,K)	
0122	170 IF(ND.EQ.0) GO TO 1	
0123	WRITE(102,1070)	396
0124	WRITE(102,1075) (J,RD(J),VD(J),J=1,ND)	
0125	GO TO 1	444
0126	1000 FORMAT(3(I2,1X),9X,I3,8X,3(D6.0,1X),13X,2(D7.0,1X),11)	
0127	1010 FORMAT(2D30.0)	
0128	1030 FORMAT(1H1,8X,43HNEWTONS METHOD TO FIND ZEROS OF POLYNOMIALS/9X,18 1HPOLYNOMIAL NUMBER ,I2,11H OF DEGREE ,I2,///1X,28HTHE COEFFICIENT 2S OF P(X) ARE/)	
0129	1040 FORMAT(3X,2HP(,I2,4H) = ,D23.16,3H + ,D23.16,2H I)	
0130	1020 FORMAT(2D30.0)	
0131	1025 FORMAT(///1X,61HBEFORE THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS 1OF P(X) ARE)	
0132	1050 FORMAT(///2X,13HROOTS OF P(X),52X,14HMULTIPLICITIES,17X,21HINITIAL 1 APPROXIMATION/)	
0133	1060 FORMAT(3X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,7X,D23. 116,3H + ,D23.16,2H I)	
0134	1062 FORMAT(3X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,8X,23HS 1OLVED BY DIRECT METHOD)	
0135	1065 FORMAT(///1X,61HAFTER THE ATTEMPT TO IMPROVE ACCURACY, THE ZEROS 1OF P(X) ARE)	
0136	1070 FORMAT(///1X,65HCOEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO Z 1EROS WERE FOUND/)	
0137	1075 FORMAT(3X,2HD(,I2,4H) = ,D23.16,3H + ,D23.16,2H I)	
0138	2000 FORMAT(1X,41HNUMBER OF INITIAL APPROXIMATIONS GIVEN. ,I2)	
0139	2010 FORMAT(1X,29HMAXIMUM NUMBER OF ITERATIONS.,11X,I3)	
0140	2020 FORMAT(1X,21HTEST FOR CONVERGENCE.,13X,D9.2)	
0141	2030 FORMAT(1X,24HTEST FOR MULTIPLICITIES.,10X,D9.2)	
0142	2040 FORMAT(1X,23HRAIUS TO START SEARCH.,11X,D9.2)	
0143	2050 FORMAT(1X,21HRAIUS TO END SEARCH.,13X,D9.2)	
0144	2060 FORMAT(///1X)	
0145	END	450

TABLE B. VIII (Continued)

```

0001      SUBROUTINE GENAPP(APPR,APPI,NAPP,XSTART)
C      *****
C      *
C      * SUBROUTINE GENAPP GENERATES N INITIAL APPROXIMATIONS, WHERE N IS THE
C      * DEGREE OF THE ORIGINAL POLYNOMIAL.
C      *
C      *****
0002      DOUBLE PRECISION APPR,APPI,XSTART,DUMMY,BETA
0003      DIMENSION APPR(25),APPI(25)
0004      COMMON DUMMY,MAX,I02
0005      IF(XSTART.EQ.0.0) XSTART=0.5
0006      BETA=0.2617994
0007      DO 10 I=1,NAPP
0008      APPR(I)=XSTART*DCOS(BETA)
0009      APPI(I)=XSTART*DSIN(BETA)
0010      BETA=BETA+0.5235988
0011      10 XSTART=XSTART+0.5
0012      RETURN
0013      END

0001      SUBROUTINE ALTER(XOLDR,XOLDI,NALTER,ITIME)
C      *****
C      *
C      * SUBROUTINE ALTER ALTERS THE INITIAL APPROXIMATIONS WHICH PRODUCE NO
C      * CONVERGENCE TO A ZERO. THIS IS DONE A MAXIMUM OF 5 TIMES FOR EACH ROOT.
C      *
C      *****
0002      DOUBLE PRECISION XOLDR,XOLDI,DUMMY,ABXOLD,BETA
0003      COMMON DUMMY,MAX,I02
0004      IF(ITIME.NE.0) GO TO 5
0005      ITIME =1
0006      WRITE(I02,1010) MAX
0007      5 IF(NALTER.EQ.0) GO TO 10
0008      WRITE(I02,1000) XOLDR,XOLDI
0009      GO TO 20
0010      10 ABXOLD=DSQRT(XOLDR*XOLDR+XOLDI*XOLDI)
0011      BETA=DATAN2(XOLDI,XOLDR)
0012      WRITE(I02,1020) XOLDR,XOLDI
0013      20 NALTER=NALTER+1
0014      IF(NALTER.GT.5) RETURN
0015      GO TO (30,40,30,40,30),NALTER
0016      30 XOLDR=-XOLDR
0017      XOLDI=-XOLDI
0018      GO TO 50
0019      40 BETA=BETA+1.0471976
0020      XOLDR=ABXOLD*DCOS(BETA)
0021      XOLDI=ABXOLD*DSIN(BETA)
0022      50 RETURN
0023      1000 FORMAT(1X,D23.16,3H + ,D23.16,2H I,10X,21HALTERED APPROXIMATION)
0024      1010 FORMAT(///1X,54HNO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AF
0025      1020 FORMAT(/1X,D23.16,3H + ,D23.16,2H I,10X,21HINITIAL APPROXIMATION)
0026      1TER ,13,12H ITERATIONS.//)
0026      END

```


TABLE B. VIII (Continued)

```

0001      SUBROUTINE QUAD(UA,VA,NA,UROOT,VROOT,NROOT,MULTI,EPST)
C      *****
C      *
C      * SUBROUTINE QUAD SOLVES DIRECTLY FOR THE ZEROS AND THEIR MULTIPLICITIES *
C      * OF EITHER A QUADRATIC POLYNOMIAL OR A LINEAR FACTOR. SOLUTION OF THE *
C      * QUADRATIC IS DONE USING THE QUADRATIC FORMULA. *
C      *
C      *****
0002      DOUBLE PRECISION UA,VA,UROOT,VROOT,888,UAAA,VAAA,UDISC,VDISC,UDUMM
0003      1Y,VDUMMY,RDUMMY,SDUMMY,EPST,UBBB,VBBB
0004      DIMENSION UA(26),VA(26),UROOT(25),VROOT(25),MULTI(25)
0005      IF(NA.EQ.2) GO TO 7
0006      IF(NA.EQ.1) GO TO 5
0007      UROOT(NROOT+1)=0.0
0008      VROOT(NROOT+1)=0.0
0009      MULTI(NROOT+1)=1
0010      NROOT=NROOT+1
0011      GO TO 50
0012      5 BBB=UA(1)*UA(1)+VA(1)*VA(1)
0013      UROOT(NROOT+1)=(-UA(2)*UA(1)-VA(2)*VA(1))/BBB
0014      VROOT(NROOT+1)=(-VA(2)*UA(1)+UA(2)*VA(1))/BBB
0015      MULTI(NROOT+1)=1
0016      NROOT=NROOT+1
0017      GO TO 50
0018      7 UDISC=(UA(2)*UA(2)-VA(2)*VA(2))-(4.0*(UA(1)*UA(3)-VA(1)*VA(3)))
0019      VDISC=(VA(2)*UA(2)+UA(2)*VA(2))-(4.0*(VA(1)*UA(3)+UA(1)*VA(3)))
0020      BBB=DSQRT(UDISC*UDISC+VDISC*VDISC)
0021      IF(BBB.LT.EPST) GO TO 10
0022      CALL COMSQT(UDISC,VDISC,UDUMMY,VDUMMY)
0023      UBBB=-UA(2)+UDUMMY
0024      VBBB=-VA(2)+VDUMMY
0025      RDUMMY=-UA(2)-UDUMMY
0026      SDUMMY=-VA(2)-VDUMMY
0027      UAAA=2.0*UA(1)
0028      VAAA=2.0*VA(1)
0029      BBB=UAAA*UAAA+VAAA*VAAA
0030      UROOT(NROOT+1)=(UBBB*UAAA+VBBB*VAAA)/BBB
0031      VROOT(NROOT+1)=(VBBB*UAAA-UBBB*VAAA)/BBB
0032      UROOT(NROOT+2)=(RDUMMY*UAAA+SDUMMY*VAAA)/BBB
0033      VROOT(NROOT+2)=(SDUMMY*UAAA-RDUMMY*VAAA)/BBB
0034      MULTI(NROOT+1)=1
0035      MULTI(NROOT+2)=1
0036      NROOT=NROOT+2
0037      GO TO 50
0038      10 UAAA=2.0*UA(1)
0039      VAAA=2.0*VA(1)
0040      BBB=UAAA*UAAA+VAAA*VAAA
0041      UROOT(NROOT+1)=(-UA(2)*UAAA-VA(2)*VAAA)/BBB
0042      VROOT(NROOT+1)=(-VA(2)*UAAA+UA(2)*VAAA)/BBB
0043      MULTI(NROOT+1)=2
0044      NROOT=NROOT+1
0045      50 RETURN
      END

```

TABLE B. VIII (Continued)

```

0001      SUBROUTINE COMSQT(UX,VX,UY,VY)
C      *****
C      *
C      * THIS SUBROUTINE COMPUTES THE SQUARE ROOT OF A COMPLEX NUMBER.
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UY,VY,DUMMY,R,AAA,BBB
0003      R=DSQRT(UX*UX+VX*VX)
0004      AAA=DSQRT(DABS((R+UX)/2.0))
0005      BBB=DSQRT(DABS((R-UX)/2.0))
0006      IF(VX) 10,20,30
0007      10 UY=AAA
0008      VY=-1.0*BBB
0009      GO TO 100
0010      20 IF(UX) 40,50,60
0011      30 UY=AAA
0012      VY=BBB
0013      GO TO 100
0014      40 DUMMY=DABS(UX)
0015      UY=0.0
0016      VY=DSQRT(DUMMY)
0017      GO TO 100
0018      50 UY=0.0
0019      VY=0.0
0020      GO TO 100
0021      60 DUMMY=DABS(UX)
0022      UY=DSQRT(DUMMY)
0023      VY=0.0
0024      100 RETURN
0025      END

```

TABLE B. VIII (Continued)

```

0001 SUBROUTINE HORNER(RA,VA,RXO,VXO,NDEF,RB,VB,RC,VC,RPX,VPX,ROPX,VOPX
C *****
C *
C * HORNER'S METHOD COMPUTES THE VALUE OF A POLYNOMIAL P(X) AT A POINT D AND *
C * ITS DERIVATIVE AT D. SYNTHETIC DIVISION IS USED TO DEFLATE THE *
C * POLYNOMIAL BY DIVIDING OUT THE FACTOR (X-D). *
C *
C *****
0002 1)
0003 DOUBLE PRECISION VDPX,RXO,VXO,RB,VB,RC,VC,RPX,VPX,ROPX,RA,VA
0004 DIMENSION RA(26),VA(26),RB(26),VB(26),RC(26),VC(26)
0005 RB(1)=RA(1) 516
0006 VB(1)=VA(1) 517
0007 NUM=NDEF+1 520
0008 DO 10 I=2,NUM 524
0009 10 RB(I)=RA(I)+(RB(I-1)*RXO-VB(I-1)*VXO)
0010 RPX=RB(NUM) 532
0011 VPX=VB(NUM) 533
0012 RC(1)=RB(1) 540
0013 VC(1)=VB(1) 541
0014 IF(NDEF.LT.2) GO TO 25
0015 DO 20 J=2,NDEF 544
0016 RC(J)=RB(J)+(RC(J-1)*RXO-VC(J-1)*VXO)
0017 20 VC(J)=VB(J)+(VC(J-1)*RXO+RC(J-1)*VXO)
0018 25 ROPX=RC(NDEF)
0019 VOPX=VC(NDEF) 553
0020 RETURN 572
0021 END 580

```

```

0001      SUBROUTINE NEWTON(RPX,VPX,RPX,VDPX,RXD,VXD,RXNEW,VXNEW)      600
C      *****
C      *
C      * THIS SUBROUTINE CALCULATES A NEW APPROXIMATION FROM THE OLD APPROX-
C      * IMATION BY USING THE ITERATION FORMULA
C      *  $X(N+1) = X(N) - P(X(N)) / P'(X(N))$ .
C      *
C      *****
0002      DOUBLE PRECISION RPX,VPX,RPX,VDPX,RXD,VXD,RXNEW,VXNEW,ARG
0003      DOUBLE PRECISION DDD
0004      ARG=RPX*RPX+VDPX*VDPX
0005      DDD=DSQRT(ARG)
0006      IF(DDD.EQ.0.0) RETURN
0007      RXNEW=RXD-((RPX*RPX+VPX*VDPX)/ARG)
0008      VXNEW=VXD-((VPX*RPX-RPX*VDPX)/ARG)
0009      RETURN
0010      END

```

TABLE B. VIII (Continued)

```

0001      SUBROUTINE CHECK(RPX,VPX,RPX,VPX,RPX,VPX,RXO,VXO,KANS)
C      *****
C      *
C      * THIS SUBROUTINE CHECKS FOR CONVERGENCE OF THE SEQUENCE OF APPROX-
C      * IMATIONS BY TESTING THE EXPRESSION
C      * ABSOLUTE VALUE OF (P(X(N))/P'(X(N)))/ABSOLUTE VALUE OF X(N+1).
C      * WHEN IT IS AS SMALL AS DESIRED, CONVERGENCE IS OBTAINED.
C      *
C      *****
0002      DOUBLE PRECISION RPX,VPX,RPX,VPX,RPX,VPX,RXO,VXO,ABSXO,ABSQUO,RDUMMY,VDU    749
      IMY, EPS    750
0003      DOUBLE PRECISION ARG
0004      DOUBLE PRECISION DDD
0005      COMMON EPS,MAX,IO2
0006      ABSXO=DSQRT(RXO*RXO+VXO*VXO)
0007      IF(ABSXO.EQ.0.) GO TO 25
0008      ARG=RPX*RPX+VPX*VPX
0009      DDD=DSQRT(ARG)
0010      IF(DDD.EQ.0.) GO TO 25
0011      RDUMMY=(RPX*RPX+VPX*VPX)/ARG
0012      VDUMMY=(VPX*RPX-RPX*VPX)/ARG
0013      ABSQUO=DSQRT(RDUMMY*RDUMMY+VDUMMY*VDUMMY)
0014      IF(ABSQUO/ABSXO.LT.EPS) GO TO 10
0015      KANS=0    760
0016      RETURN    764
0017      10 KANS=1    768
0018      RETURN    772
0019      25 KANS=0
0020      RETURN
0021      END    780

```

TABLE B. VIII (Continued)

```

0001      SUBROUTINE BETTER(K,RXZERO,VXZERO,RX,VX,NA,RA,VA,RCOEF,VCOEF,N,RC,
          1VC,RB,VB)
C      *****
C      *
C      * SUBROUTINE BETTER ATTEMPTS TO IMPROVE THE ACCURACY OF THE ZEROS FOUND
C      * BY USING THEM AS INITIAL APPROXIMATIONS WITH NEWTON'S METHOD APPLIED TO
C      * THE FULL, UNDEFLATED POLYNOMIAL.
C      *
C      *****
0002      DOUBLE PRECISION RXZERO,VXZERO,RX,VX,RA,VA,RCOEF,VCOEF,RC,VC,RB,VB      805
0003      1,RXO,VXO,RPX,VPX, RDPX, VDPX, RXNEW, VXNEW, EPS
          DIMENSION RXZERO(25),VXZERO(25),RX(25),VX(25),RA(26),VA(26),RCOEF(
          126),VCOEF(26),RC(26),VC(26),RB(26),VB(26)
0004      DOUBLE PRECISION ABPX,ABDPX
0005      COMMON EPS,MAX,IO2
0006      DO 10 I=1,K
0007          RXZERO(I)=RX(I)
0008          VXZERO(I)=VX(I)
0009      DO 20 I=1,NA
0010          RCOEF(I)=RA(I)
0011          VCOEF(I)=VA(I)
0012      DO 50 J=1,K
0013          RXO=RXZERO(J)
0014          VXO=VXZERO(J)
0015          NN=N
0016          ITER=0
0017      30 CALL HORNER(RCOEF,VCOEF,RXO,VXO,NN,RB,VB,RC,VC,RPX,VPX, RDPX, VDPX)
0018          ABPX=DSQRT(RPX*RPX+VPX*VPX)
0019          ABDPX=DSQRT(RDPX*RDPX+VDPX*VDPX)
0020          IF(ABDPX.NE.0.0) GO TO 33
0021          IF(ABPX.EQ.0.0) GO TO 40
0022          GO TO 34
0023      33 CALL NEWTON(RPX,VPX, RDPX, VDPX, RXO, VXO, RXNEW, VXNEW)
0024          ITER=ITER+1
0025          RXO=RXNEW
0026          VXO=VXNEW
0027          CALL CHECK(RPX,VPX, RDPX, VDPX, RXO, VXO, KANS)
0028          IF(KANS.EQ.1) GO TO 40
0029          IF(ITER.GE.MAX) GO TO 35
0030          GO TO 30
0031      34 WRITE(IO2,1112) RXO,VXO
0032      35 WRITE(IO2,100) J,RXZERO(J),VXZERO(J)
0033          WRITE(IO2,200) MAX
0034      40 RX(J)=RXO
0035          VX(J)=VXO
0036      50 CONTINUE
0037          RETURN
0038      1112 FORMAT(1H0,36HTHE VALUE OF THE DERIVATIVE AT XO = ,D23.16,3H + ,D2
          13.16,2H I,10H IS ZERO.)
0039      100 FORMAT(42H0IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(,I2,4H) = ,D23
          1.16,3H + ,D23.16,2H I,18H DID NOT CONVERGE.)
0040      200 FORMAT(33H THE PRESENT APPROXIMATION AFTER ,I3,29H ITERATIONS IS P
          RINTED BELOW.)
0041      END

```

APPENDIX C

MULLER'S METHOD

2. Use of the Program

A double precision FORTRAN IV program using Muller's method is presented in this appendix. Flow charts for this program are given in Figure C.1 while Table C.V gives a FORTRAN IV listing of this program. Single precision variables are listed in some of the tables. The single precision variables are used in the flow charts and the corresponding double precision variables can be obtained from the appropriate tables.

The program is designed to solve polynomials of degree 25 or less. Both the coefficient of the highest degree term and the constant coefficient should be non-zero. In order to solve polynomials of degree N , where $N > 25$, certain array dimensions must be changed. These are listed in Table C.I for the main program and subprograms in double precision.

TABLE C.I
 PROGRAM CHANGES FOR SOLVING POLYNOMIALS OF
 DEGREE GREATER THAN 25
 BY MULLER'S METHOD

Double Precision

Main Program

UROOT(N),VROOT(N)
 MULT(N)
 UAPP(N,3),VAPP(N,3)
 UWORK(N+1),VWORK(N+1)
 UB(N+1),VB(N+1)
 UA(N+1),VA(N+1)
 URAPP(N,3),VRAPP(N,3)

Subroutine BETTER

UROOT(N),VROOT(N)
 UA(N+1),VA(N+1)
 UBAPP(N,3),VBAPP(N,3)
 UB(N+1),VB(N+1)
 UROOTS(N),VROOTS(N)
 URAPP(N,3),VRAPP(N,3)
 MULT(N)

Subroutine GENAPP

APPR(N,3),APPI(N,3)

Subroutine HORNER

UA(N+1),VA(N+1)
 UB(N+1),VB(N+1)

Subroutine QUAD

UA(N+1),VA(N+1)
 UROOT(N),VROOT(N)
 MULTI(N)

Table C.II lists the system functions used in the program of Muller's method. In the table "d" denotes a double precision variable name.

TABLE C.II
SYSTEM FUNCTIONS USED IN MULLER'S METHOD

Double Precision

DABS(d)	- obtain absolute value
DATAN2(d ₁ ,d ₂)	- arctangent of d ₁ /d ₂
DSQRT(d)	- square root
DCOS(d)	- cosine of angle
DSIN(d)	- sine of angle
DSQRT(d)	- square root

2. Input Data for Muller's Method

The input data for Muller's method is identical to the input data for Newton's method as described in Appendix B, § 2 except for the variable names. The correspondence of input variable names is given in Table C.III. Only one (not three) initial approximation, X_0 , is given for each root. The other two required by Muller's method are constructed within the program and are $.9X_0$ and $1.1X_0$.

3. Variables Used in Muller's Method

The definitions of the major variables used in Muller's method are given in Table C.IV. For definitions of variables not listed in this table see the definitions of variables for the corresponding subroutine in Table B.VII. The notation and symbols used here are the same as for Table B.VII and are described in Appendix B, § 3.

TABLE C.III
CORRESPONDENCE OF NEWTON'S AND MULLER'S
INPUT DATA VARIABLES

<u>Newton's Method</u>	<u>Muller's Method</u>
Control Card	
NOPOLY	NOPOLY
N	NP
NIAP	NAPP
MAX	MAX
EPSCNV	EPS
EPSQ	EPSQ
EPSMUL	EPSM
XSTART	XSTART
XEND	XEND
KCHECK	KCHECK
Coefficient Card	
A (RA)	A (UA)
A (VA)	A (VA)
Initial Approximation Card	
XZERO (RXZERO)	APP (UAPP)
XZERO (VXZERO)	APP (VAPP)
End Card	
KCHECK	KCHECK

4. Description of Program Output

The output from Muller's method is the same as that for Newton's method as described in Appendix B, § 4. Only one initial approximation, Z, (not three) is printed for each root. It is either that supplied by the user or generated by the program. The other two approximations used were 0.9Z and 1.1Z.

5. Informative and Error Messages

The output may contain informative messages printed as an aid to the user. These are:

"NO ZEROS WERE FOUND FOR POLYNOMIAL NUMBER XX."

XX is the number of the polynomial. This message is printed if no roots of the polynomial were found.

"IN THE ATTEMPT TO IMPROVE ACCURACY, $\text{ROOT}(X) = YYY$

DID NOT CONVERGE AFTER ZZZ ITERATIONS

THE PRESENT APPROXIMATION IS AAA"

X is the number of the root before the attempt to improve accuracy, YYY is the value of the root before attempt to improve accuracy, ZZZ is the maximum number of iterations, and AAA is the current approximation after the maximum number of iterations. This message has the same meaning as the corresponding message in Appendix B, § 5.

TABLE C. IV
VARIABLES USED IN MULLER'S METHOD

<u>Single Precision</u> <u>Variable</u>	<u>Precision</u> <u>Type</u>	<u>Double Precision</u> <u>Variable</u>	<u>Precision</u> <u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
Main Program					
NP	I	NP	I		Degree of polynomial $P(X)$
NROOT	I	NROOT	I		Number of distinct roots found
NOMULT	I	NOMULT	I		Number of roots (counting multiplicities)
ROOT	C	UROOT, VROOT	D		Array containing the roots
NAPP	I	NAPP	I		Number of initial approximations to be read in
APP	C	UAPP, VAPP	D		Array of initial approximations
WORK	C	UWORK, VWORK	D		Working array containing coefficients of current polynomial
B	C	UB, VB	D		Array containing coefficients of deflated polynomial
A	C	UA, VA	D		Array containing coefficients of original polynomial, $P(X)$
RAPP	C	URAPP, VRAPP	D		Array of initial or altered approximations for which convergence was obtained
X1	C	UX1, VX1	D		One of three current approximations to a root
X2	C	UX2, VX2	D		One of three current approximations to a root
X3	C	UX3, VX3	D		One of three current approximations to a root
PX1	C	UPX1, VPX1	D		Value of polynomial $P(X)$ at X_1
PX2	C	UPX2, VPX2	D		Value of polynomial $P(X)$ at X_2
PX3	C	UPX3, VPX3	D		Value of polynomial $P(X)$ at X_3
X4	C	UX4, VX4	D		Newest approximation (X_{n+1}) to root
PX4	C	UPX4, VPX4	D		Value of polynomial $P(X)$ at X_4
MULT	I	MULT	I		Array containing the multiplicities of each root found
ITER	I	ITER	I		Counter for iterations
IO1	I	IO1	I		Unit number of input device
IO2	I	IO2	I		Unit number of output device

TABLE C.IV. (Continued)

<u>Single Precision</u> <u>Variable</u>	<u>Type</u>	<u>Double Precision</u> <u>Variable</u>	<u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
EPSRT	R	EPSRT	D		Number used in subroutine BETTER to generate two approximations from the one given
NOPLY	I	NOPLY	I		Number of the polynomial
MAX	I	MAX	I		Maximum number of iterations
EPS	R	EPS	D		Tolerance check for convergence
EPSO	R	EPSO	D		Tolerance check for zero (0)
EPSM	R	EPSM	D		Tolerance check for multiplicities
KCHECK	I	KCHECK	I		Program control, KCHECK = 1 stops execution of program
XSTART	R	XSTART	D		Magnitude at which to start generating initial approximations
XEND	R	XEND	D		Magnitude at which to end generating initial approximations
NWORK	I	NWORK	I		Degree of current deflated polynomial whose coefficients are in WORK
ITIME	I	ITIME	I		Program control
NALTER	I	NALTER	I		Number of alterations which have been performed on an initial approximation
IAPP	I	IAPP	I		Counter for number of initial approximations used
CONV	L	CONV	L		When CONV is true, convergence has been obtained
IROOT	I	IROOT	I		Number of distinct roots solved by Muller's method, i.e. not solved directly by subroutine QUAD
Subroutine HORNER					
A	C	UA, VA	D	E	Array of current polynomial coefficients (to be deflated or evaluated)
NA	I	NA	I	E	Degree of polynomial to be deflated or evaluated
X	C	UX, VX	D	E	Approximation at which to evaluate or deflate the polynomial

TABLE C. IV (Continued)

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition of Argument</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>		
B	C	UB, VB	D	R	Array containing the coefficients of the deflated polynomial
PX	C	UPX, VPX	D	R	Value of the polynomial at X
NUM	I	NUM	I		Number of coefficients of polynomial to be deflated
Subroutine TEST					
X3	C	UX3, VX3	D	E	Approximation to Root (old) (X_n)
X4	C	UX4, VX4	D	E	New approximation to root (X_{n+1})
CONV	L	CONV	L	R	CONV = true implies convergence has been obtained
EPS	R	EPS	D	C	Tolerance for convergence test
EPSO	R	EPSO	D	C	Tolerance check for zero (0)
DENOM	R	DENOM	D		Magnitude of new approximation, (X_{n+1})
Subroutine BETTER					
MULT	I	MULT	I	ECR	Array of multiplicities of each root
A	C	UA, VA	D	E	Array of coefficients of original undeflated polynomial
NP	I	NP	I	E	Degree of original polynomial
ROOT	C	UROOT, VROOT	D	ECR	Array of roots
NROOT	I	NROOT	I	ECR	Number of roots stored in root
BAPP	C	UBAPP, VBAPP	D	E	Array of initial approximations (old roots)
IROOT	I	IROOT	I	ECR	Number of roots solved by the iterative process (Not QUAD)
ROOTS	C	UROOTS, VROOTS	D		Temporary storage for new (better) roots
L	I	L	I		Number of roots found by BETTER
EPSRT	R	EPSRT	D	C	A small number used to generate two of the three approximations when given one
ITER	I	ITER	I	C	Counter for number of iterations

TABLE C.IV (Continued)

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition of Argument</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>		
B	C	UB,VB	D		Array containing coefficients of deflated polynomial
X1	C	UX1,VX1	D		One of three approximations to the root
X2	C	UX2,VX2	D		One of three approximations to the root
X3	C	UX3,VX3	D		One of three approximations to the root
PX1	C	UPX1,VPX1	D		Value of polynomial (P(X)) at X1
PX2	C	UPX2,VPX2	D		Value of polynomial (P(X)) at X2
PX3	C	UPX3,VPX3	D		Value of polynomial (P(X)) at X3
CONV	L	CONV	L		CONV = true implies convergence has been obtained
X4	C	UX4,VX4	D		Newest approximation to root
J	I	J	I		Program control - counts the number of roots used as initial approximations
MAX	I	MAX	I	C	Maximum number of iterations permitted
I02	I	I02	I	C	Unit number of output device
Subroutine ALTER					
X1	C	X1R,X1I	D	ECR	One of the three approximations to be altered
X2	C	X2R,X2I	D	ECR	One of the three approximations to be altered
X3	C	X3R,X3I	D	ECR	One of the three approximations to be altered
X2R	R	X2R	D		Real part of complex approximation
X2I	R	X2I	D		Imaginary part of complex approximation
Subroutine QUAD					
EPST	R	EPST	D	E	Tolerance check for zero (0)
Subroutine CALC					

These variables are dummy variables used for temporary storage and thus, are not defined.

MAIN PROGRAM




Figure C.1. Flow Charts for Muller's Method





Figure C.1. (Continued)

CALC



TEST





Figure C.1. (Continued)

GENAPP



ALTER





Figure C.1. (Continued)

QUAD



COMSQT




Figure C.1. (Continued)

TABLE C.V.

PROGRAM FOR MULLER'S METHOD

```

C *****
C *
C * DOUBLE PRECISION PROGRAM FOR MULLER'S METHOD
C *
C *
C * MULLER'S METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C * POLYNOMIAL OF MAXIMUM DEGREE 25. THROUGH THREE GIVEN POINTS THE
C * POLYNOMIAL IS APPROXIMATED BY A QUADRATIC. THE ZERO OF THE QUADRATIC
C * CLOSEST TO THE OLD APPROXIMATION IS TAKEN AS THE NEW APPROXIMATION.
C * IN THIS MANNER A SEQUENCE IS OBTAINED CONVERGING TO A ZERO.
C *****
0001  DOUBLE PRECISION UPX3,VPX3,UPX2,VPX2,UROOT,VROOT,UX1,VX1,UAPP,VAPP
      1,UX2,VX2,UWORK,VWORK,UX3,VX3,UB,VB,UX4,VX4,UA,VA,UPX1,VPX1,URAPP,V
      2RAPP,UPX4,VPX4,EPSRT,EPSO,EPS,CCC,EPSM,UH3,VH3,UQ4,VQ4,ABPX4,ABPX3
      3,QQQ,XSTART,XEND
0002  DIMENSION UROOT(25),VROOT(25),MULT(25),UAPP(25,3),VAPP(25,3),UWORK
      1(26),VWORK(26),UB(26),VB(26),UA(26),VA(26),URAPP(25,3),VRAPP(25,3)
0003  DATA PNAME,DNAME/2HP1,2HD1/
0004  LOGICAL CONV
0005  COMMON EPSRT,EPSO,EPS,I02,MAX
0006  I01=5
0007  I02=6
0008  EPSRT=0.999
0009  10 NROOT=0
0010  IROOT=0
0011  IPATH=1
0012  NOMULT=0
0013  NALTER=0
0014  ITIME=0
0015  IAPP=1
0016  ITER=1
0017  READ(I01,1000) NOPOLY,NP,NAPP,MAX,EPS,EPSO,EPSM,XSTART,XEND,KCHECK
0018  IF(KCHECK.EQ.1) STOP
0019  KKK=NP+1
0020  READ(I01,1010) (UA(I),VA(I),I=1,KKK)
0021  WRITE(I02,1020) NOPOLY,NP
0022  WRITE(I02,1035) (PNAME,I,UA(I),VA(I),I=1,KKK)
0023  WRITE(I02,2060)
0024  WRITE(I02,2000) NAPP
0025  WRITE(I02,2010) MAX
0026  WRITE(I02,2020) EPS
0027  WRITE(I02,2030) EPSM
0028  WRITE(I02,2040) XSTART
0029  WRITE(I02,2050) XEND
0030  IF(NP.GT.2) GO TO 15
0031  CALL QUAD(UA,VA,NP,UROOT,VROOT,NROOT,MULT,EPSO)
0032  WRITE(I02,1037)
0033  WRITE(I02,1086) (I,UROOT(I),VROOT(I),MULT(I),I=1,NROOT)
0034  GO TO 10
0035  15 IF(NAPP.NE.0) GO TO 20
0036  NAPP=NP
0037  CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0038  GO TO 27
0039  20 READ(I01,1030) (UAPP(I,2),VAPP(I,2),I=1,NAPP)
0040  DO 25 I=1,NAPP
0041  UAPP(I,1)=0.9*UAPP(I,2)
0042  VAPP(I,1)=0.9*VAPP(I,2)

```

TABLE C.V (Continued)

```

0043      UAPP(I,3)=1.1*UAPP(I,2)
0044 25 VAPP(I,3)=1.1*VAPP(I,2)
0045 27 KKK=NP+1
0046      DO 30 I=1, KKK
0047      UWORK(I)=UA(I)
0048 30 VWORK(I)=VA(I)
0049      NWORK=NP
0050 40 UX1=UAPP(IAPP,1)
0051      VX1=VAPP(IAPP,1)
0052      UX2=UAPP(IAPP,2)
0053      VX2=VAPP(IAPP,2)
0054      UX3=UAPP(IAPP,3)
0055      VX3=VAPP(IAPP,3)
0056      CALL HORNER(NWORK,UWORK,VWORK,UX1,VX1,UB,VB,UPX1,VPX1)
0057      CALL HORNER(NWORK,UWORK,VWORK,UX2,VX2,UB,VB,UPX2,VPX2)
0058      CALL HORNER(NWORK,UWORK,VWORK,UX3,VX3,UB,VB,UPX3,VPX3)
0059 50 CALL CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UX
      14,VX4,UQ4,VQ4,UH3,VH3)
0060 60 CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0061      ABPX4=DSQRT(UPX4*UPX4+VPX4*VPX4)
0062      ABPX3=DSQRT(UPX3*UPX3+VPX3*VPX3)
0063      IF(ABPX3.EQ.0.0) GO TO 70
0064      QQQ=ABPX4/ABPX3
0065      IF(QQQ.LE.10.) GO TO 70
0066      UQ4=0.5*UQ4
0067      VQ4=0.5*VQ4
0068      UX4=UX3+(UH3*UQ4-VH3*VQ4)
0069      VX4=VX3+(VH3*UQ4+UH3*VQ4)
0070      GO TO 60
0071 70 CALL TEST(UX3,VX3,UX4,VX4,CONV)
0072      IF(CONV) GO TO 120
0073      IF(ITER.LT.MAX) GO TO 110
0074      CALL ALTER(UAPP(IAPP,1),VAPP(IAPP,1),UAPP(IAPP,2),VAPP(IAPP,2),UAP
      1P(IAPP,3),VAPP(IAPP,3),NALTER,ITIME)
0075      IF(NALTER.GT.5) GO TO 75
0076      ITER=1
0077      GO TO 40
0078 75 IF(IAPP.LT.NAPP) GO TO 100
0079      IF(XEND.EQ.0.0) GO TO 77
0080      IF(XSTART.GT.XEND) GO TO 77
0081      NAPP=NP
0082      CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0083      IAPP=0
0084      GO TO 100
0085 77 WRITE(102,1090)
0086      KKK=NWORK+1
0087      WRITE(102,1035) (DNAME,J,UWORK(J),VWORK(J),J=1,KKK)
0088 80 IF(NROOT.EQ.0) GO TO 90
0089      WRITE(102,1060)
0090      IF(IPATH.EQ.1) GO TO 82
0091 81 IPATH=2
0092      CALL BETTER(UA,VA,NP,UROOT,VROOT,NROOT,URAPP,VRAPP,IROOT,MULT)
0093      WRITE(102,1200)
0094 82 IF(NROOT.EQ.0) GO TO 90
0095      IF(IROOT.EQ.0) GO TO 85
0096      WRITE(102,1080)
0097      DO 55 I=1, IROOT
0098 55 WRITE(102,1085) I,UROOT(I),VROOT(I),MULT(I),URAPP(I,2),VRAPP(I,2)

```

TABLE C.V. (Continued)

```

0099      IF(IROOT.LT.NROOT) GO TO 85
0100      GO TO 87
0101      85 KKK=IROOT+1
0102      WRITE(102,1086) (I,UROOT(I),VROOT(I),MULT(I),I=KKK,NROOT)
0103      87 IF(IPATH.EQ.1) GO TO 81
0104      GO TO 10
0105      90 WRITE(102,1070) NOPOLY
0106      GO TO 10
0107      100 IAPP=IAPP+1
0108      ITER=1
0109      NALTER=0
0110      GO TO 40
0111      120 NROOT=NROOT+1
0112      IROOT=NROOT
0113      MULT(NROOT)=1
0114      NOMULT=NOMULT+1
0115      UROOT(NROOT)=UX4
0116      VROOT(NROOT)=VX4
0117      URAPP(NROOT,1)=UAPP(IAPP,1)
0118      VRAPP(NROOT,1)=VAPP(IAPP,1)
0119      URAPP(NROOT,2)=UAPP(IAPP,2)
0120      VRAPP(NROOT,2)=VAPP(IAPP,2)
0121      URAPP(NROOT,3)=UAPP(IAPP,3)
0122      VRAPP(NROOT,3)=VAPP(IAPP,3)
0123      125 IF(NOMULT.LT.NP) GO TO 130
0124      GO TO 80
0125      130 CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0126      NWORK=NWORK-1
0127      KKK=NWORK+1
0128      DO 140 I=1,KKK
0129      UWORK(I)=UB(I)
0130      VWORK(I)=VB(I)
0131      CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0132      CCC=DSQRT(UPX4*UPX4+VPX4*VPX4)
0133      IF(CCC.LT.EPSM) GO TO 150
0134      IF(NWORK.GT.2) GO TO 75
0135      IROOT=NROOT
0136      CALL QUAD(UWORK,VWORK,NWORK,UROOT,VROOT,NROOT,MULT,EPSO)
0137      GO TO 80
0138      150 MULT(NROOT)=MULT(NROOT)+1
0139      NOMULT=NOMULT+1
0140      GO TO 125
0141      110 UX1=UX2
0142      VX1=VX2
0143      UX2=UX3
0144      VX2=VX3
0145      UX3=UX4
0146      VX3=VX4
0147      UPX1=UPX2
0148      VPX1=VPX2
0149      UPX2=UPX3
0150      VPX2=VPX3
0151      UPX3=UPX4
0152      VPX3=VPX4
0153      ITER=ITER+1
0154      GO TO 50
0155      1010 FORMAT(2D30.0)
0156      1020 FORMAT(1H1,1X,52HMULLERS METHOD FOR FINDING THE ZEROS OF A POLYNOM

```

TABLE C.V (Continued)

```

11AL/1H ,1X,18HPOLYNOMIAL NUMBER ,I2,11H OF DEGREE ,I2///1H ,1X,28H
2THE COEFFICIENTS OF P(X) ARE//)
0157 1030 FORMAT(2D30.0)
0158 1090 FORMAT(///,1X,65HCOEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO
1ZEROS WERE FOUND//)
0159 1080 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES,17X,21HINITIAL
1 APPROXIMATION//)
0160 1070 FORMAT(///,43H NO ZEROS WERE FOUND FOR POLYNOMIAL NUMBER ,I2)
0161 1086 FORMAT(2X,5HROOT(I,I2,4H) = ,D23.16,3H + ,D23.16,2H 1,8X,12,9X,23HS
1OLVED BY DIRECT METHOD)
0162 1037 FORMAT(///,1X,13HZEROS OF P(X),51X,14HMULTIPLICITIES//)
0163 1035 FORMAT(3X,A2,I2,4H) = ,D23.16,3H + ,D23.16,2H 1)
0164 1085 FORMAT(2X,5HROOT(I,I2,4H) = ,D23.16,3H + ,D23.16,2H 1,8X,12,8X,D23.
116,3H + ,D23.16,2H 1)
0165 1000 FORMAT(3(I2,1X),9X,I3,8X,3(D6.0,1X),13X,2(D7.0,1X),11)
0166 1060 FORMAT(///35H BEFORE ATTEMPT TO IMPROVE ACCURACY)
0167 1200 FORMAT(///1X,37HAFTER THE ATTEMPT TO IMPROVE ACCURACY)
0168 2000 FORMAT(1X,41HNUMBER OF INITIAL APPROXIMATIONS GIVEN. ,I2)
0169 2010 FORMAT(1X,29HMAXIMUM NUMBER OF ITERATIONS.,11X,I3)
0170 2020 FORMAT(1X,21HTEST FOR CONVERGENCE.,13X,D9.2)
0171 2030 FORMAT(1X,24HTEST FOR MULTIPLICITIES.,10X,D9.2)
0172 2040 FORMAT(1X,23HRADIUS TO START SEARCH.,11X,D9.2)
0173 2050 FORMAT(1X,21HRADIUS TO END SEARCH.,13X,D9.2)
0174 2060 FORMAT(//1X)
0175 END

```


TABLE C.V (Continued)

```

0001      SUBROUTINE ALTER(X1R,X1I,X2R,X2I,X3R,X3I,NALTER,ITIME)
C      *****
C      *
C      * SUBROUTINE ALTER ALTERS THE INITIAL APPROXIMATIONS WHICH PRODUCE NO
C      * CONVERGENCE TO A ZERO. THIS IS DONE A MAXIMUM OF 5 TIMES FOR EACH ROOT.
C      *
C      *****
0002      DOUBLE PRECISION X1R,X1I,X2R,X2I,X3R,X3I,EPS1,EPS2,EPS3,R,BETA
0003      COMMON EPS1,EPS2,EPS3,IO2,MAX
0004      IF(ITIME.NE.0) GO TO 5
0005      ITIME=1
0006      WRITE(IO2,1010) MAX
0007      5 IF(NALTER.EQ.0) GO TO 10
0008      WRITE(IO2,1000) X1R,X1I,X2R,X2I,X3R,X3I
0009      GO TO 20
0010      10 R=DSQRT(X2R*X2R+X2I*X2I)
0011      BETA=DATAN2(X2I,X2R)
0012      WRITE(IO2,1020) X1R,X1I,X2R,X2I,X3R,X3I
0013      20 NALTER=NALTER+1
0014      IF(NALTER.GT.5) RETURN
0015      GO TO (30,40,30,40,30),NALTER
0016      30 X2R=-X2R
0017      X2I=-X2I
0018      GO TO 50
0019      40 BETA=BETA+1.0471976
0020      X2R=R*DCOS(BETA)
0021      X2I=R*DSIN(BETA)
0022      50 X1R=0.9*X2R
0023      X1I=0.9*X2I
0024      X3R=1.1*X2R
0025      X3I=1.1*X2I
0026      RETURN
0027      1000 FORMAT(1X,5HX1 = ,D23.16,3H + ,D23.16,2H I,10X,22HALTERED APPROXIM
1ATIONS/1X,5HX2 = ,D23.16,3H + ,D23.16,2H I/1X,5HX3 = ,D23.16,3H +
2,D23.16,2H I/)
0028      1020 FORMAT(1H0,5HX1 = ,D23.16,3H + ,D23.16,2H I,10X,22INITIAL APPROXI
1MATIONS/1X,5HX2 = ,D23.16,3H + ,D23.16,2H I/1X,5HX3 = ,D23.16,3H +
2 ,D23.16,2H I/)
0029      1010 FORMAT(///1X,54HNO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AF
ITER ,13,12H ITERATIONS.//)
0030      END

```

TABLE C.V (Continued)

```

0001      SUBROUTINE GENAPP(APPR,APPI,NAPP,XSTART)
C      *****
C      *
C      * SUBROUTINE GENAPP GENERATES N INITIAL APPROXIMATIONS, WHERE N IS THE *
C      * DEGREE OF THE ORIGINAL POLYNOMIAL. *
C      *
C      *****
0002      DOUBLE PRECISION APPR,APPI,XSTART,EPS1,EPS2,EPS3,BETA
0003      DIMENSION APPR(25,3),APPI(25,3)
0004      COMMON EPS1,EPS2,EPS3,IQ2,MAX
0005      IF(XSTART.EQ.0.0) XSTART=0.5
0006      BETA=0.2617994
0007      DO 10 I=1,NAPP
0008      APPR(I,2)=XSTART*OCOS(BETA)
0009      APPI(I,2)=XSTART*DSIN(BETA)
0010      BETA=BETA+0.5235988
0011  10 XSTART=XSTART+0.5
0012      DO 20 I=1,NAPP
0013      APPR(I,1)=0.9*APPR(I,2)
0014      APPI(I,1)=0.9*APPI(I,2)
0015      APPR(I,3)=1.1*APPR(I,2)
0016  20 APPI(I,3)=1.1*APPI(I,2)
0017      RETURN
0018      END

```

TABLE C.V (Continued)

```

0001      SUBROUTINE BETTER(UA,VA,NP,UROOT,VROOT,NROOT,URAPP,VRAPP,IROOT,MUL
          1T)
C      *****
C      *
C      * SUBROUTINE BETTER ATTEMPTS TO IMPROVE THE ACCURACY OF THE ZEROS FOUND *
C      * BY USING THEM AS INITIAL APPROXIMATIONS WITH MULLER'S METHOD APPLIED TO *
C      * THE FULL, UNDEFLATED POLYNOMIAL. *
C      *
C      *****
0002      DOUBLE PRECISION UROOT,VROOT,UA,VA,UBAPP,VBAPP,UX1,VX1,UX2,VX2,UX3
          1,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UB,VB,UROOTS,VROOTS,EPSRT,UX4,V
          2X4,URAPP,VRAPP,EPSO,EPS,UQ4,VQ4,UH3,VH3
0003      LOGICAL CONV
0004      DIMENSION UROOT(25),VROOT(25),UA(26),VA(26),UBAPP(25,3),VBAPP(25,3
          1),UB(26),VB(26),UROOTS(25),VROOTS(25),URAPP(25,3),VRAPP(25,3),MULT
          3(25)
0005      COMMON EPSRT,EPSO,EPS,IQ2,MAX
0006      IF(NROOT.LE.1) RETURN
0007      L=0
0008      DO 10 I=1,NROOT
0009      UBAPP(I,1)=UROOT(I)*EPSRT
0010      VBAPP(I,1)=VROOT(I)*EPSRT
0011      UBAPP(I,2)=UROOT(I)
0012      VBAPP(I,2)=VROOT(I)
0013      UBAPP(I,3)=UROOT(I)*(2.0-EPSRT)
0014      10 VBAPP(I,3)=VROOT(I)*(2.0-EPSRT)
0015      DO 100 J=1,NROOT
0016      UX1=UBAPP(J,1)
0017      VX1=VBAPP(J,1)
0018      UX2=UBAPP(J,2)
0019      VX2=VBAPP(J,2)
0020      UX3=UBAPP(J,3)
0021      VX3=VBAPP(J,3)
0022      ITER=1
0023      CALL HORNER(NP,UA,VA,UX1,VX1,UB,VB,UPX1,VPX1)
0024      CALL HORNER(NP,UA,VA,UX2,VX2,UB,VB,UPX2,VPX2)
0025      20 CALL HORNER(NP,UA,VA,UX3,VX3,UB,VB,UPX3,VPX3)
0026      CALL CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UX
          14,VX4,UQ4,VQ4,UH3,VH3)
0027      30 CALL TEST(UX3,VX3,UX4,VX4,CONV)
0028      IF(CONV) GO TO 50
0029      IF(ITER.LT.MAX) GO TO 40
0030      WRITE(102,1000) J,UROOT(J),VROOT(J),MAX
0031      WRITE(102,1010) UX4,VX4
0032      IF(J.LT.IROOT) GO TO 33
0033      IF(J.EQ.IROOT) GO TO 35
0034      GO TO 100
0035      33 KKK=IROOT-1
0036      DO 34 K=J,KKK
0037      URAPP(K,1)=URAPP(K+1,1)
0038      VRAPP(K,1)=VRAPP(K+1,1)
0039      URAPP(K,2)=URAPP(K+1,2)
0040      VRAPP(K,2)=VRAPP(K+1,2)
0041      URAPP(K,3)=URAPP(K+1,3)
0042      VRAPP(K,3)=VRAPP(K+1,3)
0043      34 VRAPP(K,3)=VRAPP(K+1,3)
0044      35 IROOT=IROOT-1
0045      GO TO 100
          40 UX1=UX2

```

TABLE C.V (Continued)

```

0046      VX1=VX2
0047      UX2=UX3
0048      VX2=VX3
0049      UX3=UX4
0050      VX3=VX4
0051      UPX1=VPX2
0052      VPX1=VPX2
0053      UPX2=UPX3
0054      VPX2=VPX3
0055      ITER=ITER+1
0056      GO TO 20
0057      50 L=L+1
0058          UROOTS(L)=UX4
0059          VROOTS(L)=VX4
0060          MULT(L)=MULT(J)
0061      100 CONTINUE
0062          IF(L.EQ.0) GO TO 120
0063          DO 110 I=1,L
0064              UROOT(I)=UROOTS(I)
0065          110 VROOT(I)=VROOTS(I)
0066              NROOT=L
0067              RETURN
0068      120 NROOT=0
0069              RETURN
0070      1000 FORMAT(///42H IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(,I2,4H) = ,
1023.16,3H + ,D23.16,2H I/24H DID NOT CONVERGE AFTER ,I3,11H ITERAT
210NS)
0071      1010 FORMAT(30H THE PRESENT APPROXIMATION IS ,D23.16,3H + ,D23.16,2H I/
1/)
0072      END

```

TABLE C.V (Continued)

```

0001      SUBROUTINE CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,V
          1PX3,UX4,VX4,UQ4,VQ4,UH3,VH3)
C      *****
C      *
C      * GIVEN THREE APPROXIMATIONS X(N-2), X(N-1), AND X(N), SUBROUTINE CALC
C      * APPROXIMATES THE POLYNOMIAL BY A QUADRATIC AND SOLVES FOR THE ZERO OF
C      * THE QUADRATIC CLOSEST TO X(N). THIS ZERO IS THE NEW APPROXIMATION
C      * X(N+1) TO THE ZERO OF THE POLYNOMIAL.
C      *
C      *****
0002      DOUBLE PRECISION ARG1,ARG2
0003      DOUBLE PRECISION UPX3,VPX3,UPX2,VPX2,UX1,VX1,UX2,VX2,UX3,VX3,UPX1,
          1VPX1,UH3,VH3,UH2,VH2,UQ3,VQ3,UD,VD,U8,V8,UC,VC,UDISC,VDISC,UCCC,VC
          2CC,UDEN1,VDEN1,UDEN2,VDEN2,UQ4,VQ4,UX4,VX4,EPSRT,EPS0,EPS,UDDD,VDD
          3D,AAA,BBB,RAD,UAAA,VAAA,U BBB,VBBB
0004      DOUBLE PRECISION THETA,ANGLE,UTEST,VTEST
0005      COMMON EPSRT,EPS0,EPS,IO2,MAX
0006      UH3=UX3-UX2
0007      VH3=VX3-VX2
0008      UH2=UX2-UX1
0009      VH2=VX2-VX1
0010      BBB=UH2*UH2+VH2*VH2
0011      UQ3=(UH3*UH2+VH3*VH2)/BBB
0012      VQ3=(VH3*UH2-UH3*VH2)/BBB
0013      UDDD=1.0+UQ3
0014      VDDD=VQ3
0015      UD=(UPX3-(UDDD*UPX2-VDDD*VPX2))+{UQ3*UPX1-VQ3*VPX1}
0016      VD=(VPX3-(VDDD*UPX2+UDDD*VPX2))+{VQ3*UPX1+UQ3*VPX1}
0017      UAAA=2.0*UQ3
0018      VAAA=2.0*VQ3
0019      UAAA=UAAA+1.0
0020      UBBB=UDDD*UDDD-VDDD*VDDD
0021      VBBB=VDDD*UDDD+UDDD*VDDD
0022      UCCC=UQ3*UQ3-VQ3*VQ3
0023      VCCC=VQ3*UQ3+UQ3*VQ3
0024      UB=(UAAA*UPX3-VAAA*VPX3)-(UBBB*UPX2-VBBB*VPX2)+(UCCC*UPX1-VCCC*V
          1PX1)
0025      VB=(VAAA*UPX3+UAAA*VPX3)-(VBBB*UPX2+UBBB*VPX2)+(VCCC*UPX1+UCCC*V
          1PX1)
0026      UC=UDDD*UPX3-VDDD*VPX3
0027      VC=VDDD*UPX3+UDDD*VPX3
0028      UDISC=(UB*UB-VB*VB)-(4.0*(UD*UC-VD*VC))
0029      VDISC=(2.0*(VB*UB))-(4.0*(VD*UC+UD*VC))
0030      AAA=DSQRT(UDISC*UDISC+VDISC*VDISC)
0031      IF(AAA.EQ.0.0) GO TO 5
0032      GO TO 7
0033      5 THETA=0.0
0034      GO TO 9
0035      7 THETA=DATAN2(VDISC,UDISC)
0036      9 RAD=DSQRT(AAA)
0037      ANGLE=THETA/2.0
0038      UTEST=RAD*DCOS(ANGLE)
0039      VTEST=RAD*DSIN(ANGLE)
0040      UDEN1=UB+UTEST
0041      VDEN1=VB+VTEST
0042      UDEN2=UB-UTEST
0043      VDEN2=VB-VTEST
0044      ARG1=UDEN1*UDEN1+VDEN1*VDEN1

```

TABLE C.V (Continued)

```

0045      ARG2=UDEN2*UDEN2+VDEN2*VDEN2
0046      AAA=DSQRT(ARG1)
0047      BBB=DSQRT(ARG2)
0048      IF(AAA.LT.BBB) GO TO 10
0049      IF(AAA.EQ.0.0) GO TO 60
0050      UAAA=-2.0*UC
0051      VAAA=-2.0*VC
0052      UQ4=(UAAA*UDEN1+VAAA*VDEN1)/ARG1
0053      VQ4=(VAAA*UDEN1-UAAA*VDEN1)/ARG1
0054      GO TO 50
0055  10 IF(BBB.EQ.0.0) GO TO 60
0056      UAAA=-2.0*UC
0057      VAAA=-2.0*VC
0058      UQ4=(UAAA*UDEN2+VAAA*VDEN2)/ARG2
0059      VQ4=(VAAA*UDEN2-UAAA*VDEN2)/ARG2
0060      GO TO 50
0061  50 UX4=UX3+(UH3*UQ4-VH3*VQ4)
0062      VX4=VX3+(VH3*UQ4+UH3*VQ4)
0063      RETURN
0064  60 UQ4=1.0
0065      VQ4=0.0
0066      GO TO 50
0067      END

```

TABLE C.V (Continued)

```

0001      SUBROUTINE TEST(UX3,VX3,UX4,VX4,CONV)
C      *****
C      *
C      * SUBROUTINE TEST CHECKS FOR CONVERGENCE OF THE SEQUENCE OF APPROX-
C      * IMATIONS BY TESTING THE EXPRESSION
C      * ABSOLUTE VALUE OF (X(N+1)-X(N))/ABSOLUTE VALUE OF X(N+1).
C      * WHEN IT IS AS SMALL AS DESIRED, CONVERGENCE IS OBTAINED.
C      *
C      *****
0002      DOUBLE PRECISION UX3,VX3,UX4,VX4,EPSRT,EPSO,EPS,AAA,UDUMMY,VDUMMY,
0003      1DENOM
0004      LOGICAL CONV
0005      COMMON EPSRT,EPSO,EPS,102,MAX
0006      UDUMMY=UX4-UX3
0007      VDUMMY=VX4-VX3
0008      AAA=DSQRT(UDUMMY*UDUMMY+VDUMMY*VDUMMY)
0009      DENOM=DSQRT(UX4*UX4+VX4*VX4)
0010      IF(DENOM.LT.EPSO) GO TO 20
0011      IF(AAA/DENOM.LT.EPS) GO TO 10
0012      5 CONV=.FALSE.
0013      GO TO 100
0014      10 CONV=.TRUE.
0015      GO TO 100
0016      20 IF(AAA.LT.EPSO) GO TO 10
0017      GO TO 5
0018      100 RETURN
      END

0001      SUBROUTINE HORNER(NA,UA,VA,UX,VX,UB,VB,UPX,VPX)
C      *****
C      *
C      * HORNER'S METHOD COMPUTES THE VALUE OF THE POLYNOMIAL P(X) AT A POINT D.
C      * SYNTHETIC DIVISION IS USED TO DEFLATE THE POLYNOMIAL BY DIVIDING OUT THE
C      * FACTOR (X-D).
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UPX,VPX,UB,VB,UA,VA
0003      DIMENSION UA(26),VA(26),UB(26),VB(26)
0004      UB(1)=UA(1)
0005      VB(1)=VA(1)
0006      NUM=NA+1
0007      DO 10 I=2,NUM
0008      UB(I)=UA(I)+(UB(I-1)*UX-VB(I-1)*VX)
0009      10 VB(I)=VA(I)+(VB(I-1)*UX+UB(I-1)*VX)
0010      UPX=UB(NUM)
0011      VPX=VB(NUM)
0012      RETURN
0013      END

```

TABLE C.V (Continued)

```

0001      SUBROUTINE QUAD(UA,VA,NA,UROOT,VROOT,NROOT,MULTI,EPST)
C      *
C      * SUBROUTINE QUAD SOLVES DIRECTLY FOR THE ZEROS AND THEIR MULTIPLICITIES *
C      * OF EITHER A QUADRATIC POLYNOMIAL OR A LINEAR FACTOR. SOLUTION OF THE *
C      * QUADRATIC IS DONE USING THE QUADRATIC FORMULA. *
C      *
C      *
0002      DOUBLE PRECISION UA,VA,UROOT,VROOT,BBB,UAAA,VAAA,UDISC,VDISC,UDUMM
1Y,VDUMMY,RDUMMY,SDUMMY,EPST,U888,V888
0003      DIMENSION UA(26),VA(26),UROOT(25),VROOT(25),MULTI(25)
0004      IF(NA.EQ.2) GO TO 7
0005      IF(NA.EQ.1) GO TO 5
0006      UROOT(NROOT+1)=0.0
0007      VROOT(NROOT+1)=0.0
0008      MULTI(NROOT+1)=1
0009      NROOT=NROOT+1
0010      GO TO 50
0011      5 BBB=UA(1)*UA(1)+VA(1)*VA(1)
0012      UROOT(NROOT+1)=(-UA(2)*UA(1)-VA(2)*VA(1))/BBB
0013      VROOT(NROOT+1)=(-VA(2)*UA(1)+UA(2)*VA(1))/BBB
0014      MULTI(NROOT+1)=1
0015      NROOT=NROOT+1
0016      GO TO 50
0017      7 UDISC=(UA(2)*UA(2)-VA(2)*VA(2))-(4.0*(UA(1)*UA(3)-VA(1)*VA(3)))
0018      VDISC=(VA(2)*UA(2)+UA(2)*VA(2))-(4.0*(VA(1)*UA(3)+UA(1)*VA(3)))
0019      BBB=DSQRT(UDISC*UDISC+VDISC*VDISC)
0020      IF(BBB.LT.EPST) GO TO 10
0021      CALL COMSQT(UDISC,VDISC,UDUMMY,VDUMMY)
0022      U888=-UA(2)+UDUMMY
0023      V888=-VA(2)+VDUMMY
0024      RDUMMY=-UA(2)-UDUMMY
0025      SDUMMY=-VA(2)-VDUMMY
0026      UAAA=2.0*UA(1)
0027      VAAA=2.0*VA(1)
0028      BBB=UAAA*UAAA+VAAA*VAAA
0029      UROOT(NROOT+1)=(U888*UAAA+V888*VAAA)/BBB
0030      VROOT(NROOT+1)=(V888*UAAA-U888*VAAA)/BBB
0031      UROOT(NROOT+2)=(RDUMMY*UAAA+SDUMMY*VAAA)/BBB
0032      VROOT(NROOT+2)=(SDUMMY*UAAA-RDUMMY*VAAA)/BBB
0033      MULTI(NROOT+1)=1
0034      MULTI(NROOT+2)=1
0035      NROOT=NROOT+2
0036      GO TO 50
0037      10 UAAA=2.0*UA(1)
0038      VAAA=2.0*VA(1)
0039      BBB=UAAA*UAAA+VAAA*VAAA
0040      UROOT(NROOT+1)=(-UA(2)*UAAA-VA(2)*VAAA)/BBB
0041      VROOT(NROOT+1)=(-VA(2)*UAAA+UA(2)*VAAA)/BBB
0042      MULTI(NROOT+1)=2
0043      NROOT=NROOT+1
0044      50 RETURN
0045      END

```


TABLE C.V (Continued)

```

0001      SUBROUTINE COMSQT(UX,VX,UY,VY)
C      *****
C      *
C      * THIS SUBROUTINE COMPUTES THE SQUARE ROOT OF A COMPLEX NUMBER.
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UY,VY,DUMMY,R,AAA,BBB
0003      R=DSQRT(UX*UX+VX*VX)
0004      AAA=DSQRT(DABS((R+UX)/2.0))
0005      BBB=DSQRT(DABS((R-UX)/2.0))
0006      IF(VX) 10,20,30
0007      10 UY=AAA
0008      VY=-1.0*BBB
0009      GO TO 100
0010      20 IF(UX) 40,50,60
0011      30 UY=AAA
0012      VY=BBB
0013      GO TO 100
0014      40 DUMMY=DABS(UX)
0015      UY=0.0
0016      VY=DSQRT(DUMMY)
0017      GO TO 100
0018      50 UY=0.0
0019      VY=0.0
0020      GO TO 100
0021      60 DUMMY=DABS(UX)
0022      UY=DSQRT(DUMMY)
0023      VY=0.0
0024      100 RETURN
0025      END

```

APPENDIX D

SPECIAL FEATURES OF THE G.C.D. AND THE REPEATED G.C.D. PROGRAMS

Several special features have been provided in each program as an aid to the user and to improve accuracy of the results. These are explained and illustrated below.*

1. Generating Approximations

If the user does not have initial approximations available, subroutine GENAPP can systematically generate, for an N^{th} degree polynomial, N initial approximations of increasing magnitude, beginning with the magnitude specified by XSTART. If XSTART is 0., XSTART is automatically initialized to 0.5 to avoid the approximation $0. + 0.i$. The approximations are generated according to the formula:

$$X_K = (XSTART + 0.5K) (\cos \beta + i \sin \beta)$$

where

$$\beta = \frac{\pi}{12} + K \frac{\pi}{6}, K = 0, 1, 2, \dots$$

To accomplish this, the user defines the number of initial approximations to be read (NAPP) on the control card to be zero (0) or these columns

*These illustrations are representative of G.C.D.-Newton's method in double precision. Control cards for other methods should be prepared accordingly.

(7-8) may be left blank. If XSTART is left blank, it is interpreted as 0.

For example, a portion of a control card which generates initial approximations beginning at the origin for a seventh degree polynomial is shown in Example D.1.

Variable Name

Card Columns

1	2		4	5		7	8		6		7		7		7		8
									4		0		2		8		0
N	O					N											
P						A											
O						P											
L																	
Y																	
1																	

Example

Example D.1

The approximations are generated in a spiral configuration as illustrated in Figure A.1.

Example D.2 shows a portion of a control card which generates initial approximations beginning at a magnitude of 25.0 for a sixth degree polynomial.

1 2	4 5	7 8		6 7 4 0	7 7 2 8	8 0
N O P O L Y	N P	N A P P		XSTART		
2	6			2.5D+01		

Example D.2

Note that if the approximations are generated beginning at the origin, the order in which the roots are found will probably be of increasing magnitude. Roots obtained in this way are usually more accurate.

2. Altering Approximations

If an initial approximation, X_0 , does not produce convergence to a root within the maximum number of iterations, it is systematically altered a maximum of five times until convergence is possibly obtained according to the following formulas:

If the number of the alteration is odd: ($j = 1, 3$)

$$X_{j+1} = |X_0| (\cos \beta + i \sin \beta) \text{ where}$$

$$\beta = \tan^{-1} \frac{\operatorname{Im} X_0}{\operatorname{Re} X_0} + K \frac{\pi}{3}; \quad K = 1 \text{ if } j = 1 \\ K = 2 \text{ if } j = 3.$$

If the number of the alteration is even: ($j = 0, 2, 4$)

$$X_{j+1} = -X_j.$$

Each altered approximation is then taken as a starting approximation. If none of the six starting approximations produce convergence, the next initial approximation is taken, and the process repeated. The six approximations are spaced 60 degrees apart on a circle of radius $|X_0|$ centered at the origin as illustrated in Figure A.2.

3. Searching the Complex Plane

By use of initial approximations and the altering technique, any region of the complex plane in the form of an annulus centered at the origin can be searched for roots. This procedure can be accomplished in two ways.

The first way is more versatile but requires more effort on the part of the user. Specifically selected initial approximation can be used to define particular regions to be searched. For example, if the roots of a particular polynomial are known to have magnitudes between 20 and 40 an annulus of inner radius 20 and outer radius 40 could be searched by using the initial approximations $20. + i$, $23. + i$, $26. + i$, $29. + i$, $32. + i$, $35. + i$, $38. + i$, $40. + i$.

By generating initial approximations internally, the program can search an annulus centered at the origin of inner radius XSTART and outer radius XEND. Values for XSTART and XEND are supplied on the control card by the user. Example D.3 shows a portion of a control card to search the above annulus of inner radius 20.0 and outer radius 40.0.

1	2		4	5		7	8		6	7		7	7	8
									4	0		2	8	0
N			N			N								
O			P			A								
P						P								
O														
L														
Y														
1			7											
									2.0D+01			4.0D+01		

Example D.3

Note that since not less than N initial approximations can be generated at one time, the outer radius of the annulus actually searched may be greater than XEND but not greater than $XEND + .5N$.

Example D.4 shows a control card to search a circle of radius 15.

1	2		4	5		7	8		6	7		7	7	8
									4	0		2	8	0
N			N			N								
O			P			A								
P						P								
O														
L														
Y														
1			7									1.5D+01		

Example D.4

Figure A.3 shows the distribution of initial and altered approximations for an annulus of width 2 and inner radius a.

4. Improving Zeros Found

After the zeros of a polynomial are found, they are printed under the heading "Roots of Q(X)." They are then used as initial approximations with Newton's (Muller's) method applied each time to the full (undeflated) polynomial Q(X), which contains only distinct roots. In most cases, zeros that have lost accuracy due to roundoff error in the deflation process are improved. The improved zeros are then printed under the heading "Roots of P(X)." Since each root is used as an approximation to the original (undeflated) polynomial Q(X), it is possible that the root may converge to an entirely different root. This is especially true where several zeros are close together. Therefore, the user should check both lists of zeros to determine whether or not this has occurred.

5. Solving Quadratic Polynomial

After N-2 roots of an Nth degree polynomial have been extracted, the remaining quadratic, $aX^2 + bX + c$, is solved using the quadratic formula

$$X = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

for the two remaining roots. These are indicated by the words "Results of Subroutine QUAD" in the initial approximation column. If only a polynomial of degree 1 is to be solved, the solution is found directly as $(X - C) = 0$ implies $X = C$.

6. Missing Roots

If not all N roots of an N^{th} degree polynomial are found, the coefficients of the remaining deflated polynomial are printed under the heading "Coefficients of Deflated Polynomial For Which No Zeros Were Found." The user may then work with this polynomial in an attempt to find the remaining roots. The leading coefficient (coefficient of the highest degree term) will be printed first (Exhibit 6.11)

7. Miscellaneous

By using various combinations of values for NAPP, XSTART, and XEND, the user has several options available as illustrated below.

Example D.5 shows the control card for a seventh degree polynomial. Three initial approximations are supplied by the user. At most three roots will be found and the coefficients of the remaining deflated polynomial will be printed.

1	2		4	5		7	8		6		7		7		7		8
									4		0		2		8		0
N						N											
O						A											
P						P											
O																	
L																	
Y																	
1			7			3											

Example D.5

Note that if several roots are known to the user, they may be "divided out" of the original polynomial by using this procedure.

Example D.6 indicates that 2 initial approximations are supplied by the user to a 7th degree polynomial. After these approximations are used the circle of radius 15 will be searched for the remaining roots.

1	2		4	5		7	8		6	7		7	7		8
									4	0		2	8		0
N						N									
O						A									
P						P									
O															
L															
Y															
1			7			2							1.5D+01		

Example D.6

By defining XSTART between 0. and 15. an annulus instead of the circle will be searched.

APPENDIX E

G.C.D. - NEWTON'S METHOD

1. Use of the Program

A double precision FORTRAN IV program using the G.C.D. method with Newton's method as a supporting method is presented here. Flow charts for this program are given in Figure E.6 while Table E.VII gives a FORTRAN IV listing of this program. Single precision variables are listed in some of the tables. The simple precision variables are used in the flow charts and the corresponding double precision variables can be obtained from the appropriate tables.

This program is designed to solve polynomials having degree less than or equal to 25. In order to solve polynomials of degree N where $N > 25$, the data statement and array dimensions given in Table E.I must be changed.

In this program both the leading coefficient and the constant coefficient are assumed to be non-zero.

TABLE E.I

PROGRAM CHANGES NECESSARY TO SOLVE POLYNOMIALS OF DEGREE
GREATER THAN 25 BY G.C.D. - NEWTON'S METHOD

Main Program

Data Entry/1H1,1H2,...,1H9,2H10,2H11,...,2HXX/where XX = N+1

UP(N+1), VP(N+1)
UAPP(N), VAPP(N)
UROOT(N), VROOT(N)
MULT(N)
UDP(N+1), VDP(N+1)
UD(N+1), VD(N+1)
UQ(N+1), VQ(N+1)
UQQ(N+1), VQQ(N+1)
UAP(N), VAP(N)
UQD(N+1), VQD(N+1)
ENTRY(N+1)
UROOTS(N), VROOTS(N)

Subroutine GENAPP

APPR(N), APPI(N)

Subroutine GCD

UR(N+1), VR(N+1)
US(N+1), VS(N+1)
USS(N+1), VSS(N+1)
URR(N+1), VRR(N+1)
UT(N+1), VT(N+1)

Subroutine QUAD

UA(N+1), VA(N+1)
UROOT(N), VROOT(N)
MULT(N)

Subroutine NEWTON

UP(N+1), VP(N+1)
UB(N+1), VB(N+1)

Subroutine DIVIDE

UP(N+1), VP(N+1)
UD(N+1), VD(N+1)
UQ(N+1), VQ(N+1)

TABLE E.I (Continued)

Subroutine HORNER

UP(N+1), VP(N+1)

UB(N+1), VB(N+1)

Subroutine DERIV

UP(N+1), VP(N+1)

UA(N+1), VA(N+1)

Subroutine MULTI

UP(N+1), VP(N+1)

UROOT(N), VROOT(N)

UA(N+1), VA(N+1)

UB(N+1), VB(N+1)

MULT(N)

2. Input Data for G.C.D. - Newton's Method

The input data for G.C.D. - Newton's method is grouped into polynomial data sets. Each polynomial data set consists of the data for one and only one polynomial. As many polynomials as the user desires may be solved by placing the polynomial data sets one behind the other. Each polynomial data set consists of three kinds of information placed in the following order:

1. Control information.
2. Coefficients of the polynomial.
3. Initial approximations. These may be omitted as described in Appendix D, § 1.

An end card follows the entire collection of data sets. It indicates that there is no more data to follow and terminates execution of the

program. This information is displayed in Figure E.1 and described below. All data should be right justified and the D-type specification should be used. The recommendations given in Table E.II are those found to give best results on the IBM 360/50 computer which has a 32 bit word.

Control Information

The control card is the first card of the polynomial data set and contains the information given in Table E.II. See Figure E.2.

TABLE E.II
CONTROL DATA FOR G.C.D. - NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
NOPOLY	c.c. 1-2	Number of the polynomial. Integer. Right justified.
NP	c.c. 4-5	Degree of the polynomial. Integer. Right justified.
NAPP	c.c. 7-8	Number of initial approximations to be read. Integer. Right justified. If no initial approximations are given, leave blank.
MAX	c.c. 19-21	Maximum number of iterations. Integer. Right justified. 200 is recommended.
EPS1	c.c. 23-28	Test for zero in subroutine GCD. Double precision. Right justify. 1.D-03 is recommended.

TABLE E.II (Continued)

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
EPS2	c.c. 30-35	Convergence requirement. Double precision. Right justify. 1.D-10 is recommended.
EPS3	c.c. 37-42	Test for zero in subroutine QUAD. Double precision. Right justify. 1.D-20 is recommended.
EPS4	c.c. 44-49	Multiplicity requirement. Double precision. Right justify. 1.D-02 is recommended.
XSTART	c.c. 64-70	Magnitude at which to begin generating initial approximations. Double precision. Right justify. This is a special feature of the program and may be omitted.
XEND	c.c. 72-78	Magnitude at which to end the generating of initial approximations. Double precision. Right justify. This is a special feature of the program and may be omitted.
KCHECK	c.c. 80	This should be left blank.

Coefficients of the Polynomial

The coefficient cards follow the control card. For an N^{th} degree polynomial, $N+1$ coefficients must be entered one per card. The coefficient of the highest degree term is entered first; that is, the leading coefficient is entered first. For example, if the polynomial $x^5 + 3x^4 + 2x + 5$ were to be solved for its zeros, the order in which

the coefficients would be entered is: 1, 3, 0, 0, 2, 5. Each real or complex coefficient is entered, one per card, as described in Table E.III and illustrated in Figure E.3.

TABLE E.III
COEFFICIENT DATA FOR G.C.D. - NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
UP (P in single precision)	c.c. 1-30	Real part of complex coefficient. Double precision. Right justify. If none, leave blank or enter 0.0D00.
VP (P in single precision)	c.c. 31-60	Imaginary part of complex coefficient. Double precision. Right justify. If none, leave blank or enter 0.0D00.

Initial Approximations

The initial approximation cards follow the set of coefficient cards. The number of initial approximations read must be the number specified on the control card and are entered, one per card, as given in Table E.IV and illustrated in Figure E.4.

TABLE E.IV
INITIAL APPROXIMATION DATA FOR G.C.D. - NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Columns</u>	<u>Description</u>
UAPP (APP in single precision)	c.c. 1-30	Real part of complex number. Double precision. Right justify. If none, leave blank or enter 0.0D00.
VAPP (APP in single precision)	c.c. 31-60	Imaginary part of complex number. Double precision. Right justify. If none, leave blank or enter 0.0D00.

End Card

The end card is the last card of the input data to the program. It indicates that there is no more data to be read. When this card is read, program execution is terminated. This card is described in Table E.V and illustrated in Figure E.5.

TABLE E.V
DATA TO END EXECUTION OF G.C.D. - NEWTON'S METHOD

<u>Variable Name</u>	<u>Card Column</u>	<u>Description</u>
KCHECK	c.c. 80	Must contain the number 1. Integer.

3. Variables Used in G.C.D. - Newton's Method

The definitions of the major variables used in G.C.D. - Newton's method are given in Table E.VI. The symbols used to indicate type are:

R - real variable
I - integer variable
D - double precision
C - complex variable
L - logical variable
A - alphanumeric variable

When two variables are listed, the one on the left is the real part of the corresponding single precision complex variable; the one on the right is the imaginary part. The symbols used to indicate disposition are:

E - entered
R - returned
ECR - entered, changed, and returned
C - variable in common




Figure E.1. Sequence of Input Data for G.C.D.-Newton's Method

Variable Name									
Card Columns									
00000000	11111111	11111111	11111111	22222222	22222222	33333333	33333333	44444444	55555555
1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890
N	O	P	N	A	P	P			
MAX									
EPS1									
EPS2									
EPS3									
EPS4									
XSTART									
XEND									
1	7	7	200	1.D-03	1.D-10	1.D-20	1.D-02	1.0D+01	5.0D+02
Example									

Figure E.2. Control Card for G.C.D. - Newton's Method

Variable Name									
Card Columns									
00000000	11111111	11111111	11111111	22222222	22222222	33333333	33333333	44444444	55555555
1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890	1234567890
UP									
VP									
+0.125768D+01									
-0.37225D+02									
Example									

Figure E.3. Coefficient Card for G.C.D. - Newton's Method




Figure E.4. Initial Approximation Card for G.C.D. - Newton's Method




Figure E.5. End Card for G.C.D. - Newton's Method

TABLE E.VI
VARIABLES USED IN G.C.D. - NEWTON'S METHOD

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>	<u>of Argument</u>	
Main Program					
J	I	J	I		Number of distinct roots found
ITIME	I	ITIME	I		Program control
NOPOLY	I	NOPOLY	I		Number of the polynomial
NP	I	NP	I		Degree of the original polynomial
P	C	UP,VP	D		Array of coefficients of original polynomial, P(X)
NAPP	I	NAPP	I		Number of initial approximation to be read
EPS1	R	EPS1	D		Tolerance check for zero (0) in Subroutine GCD
EPS2	R	EPS2	D		Tolerance check for convergence
EPS3	R	EPS3	D		Tolerance check for zero (0) in Subroutine QUAD
MAX	I	MAX	I		Maximum number of iterations permitted
IO1	I	IO1	I		Unit number of input device
IO2	I	IO2	I		Unit number of output device
KCHECK	I	KCHECK	I		Program control, KCHECK = 1 implies stop execution
APP	C	UAPP,VAPP	D		Array of initial approximations
XSTART	R	XSTART	D		Magnitude at which to start search for roots
XEND	R	XEND	D		Magnitude at which to end search for roots
ANAME	A	ANAME	A		Contains name of method used "NEWTONS"
ROOT	C	UROOT,VROOT	D		Array of roots found
MULT	I	MULT	I		Array of multiplicities
DP	C	UDP,VDP	D		Array containing coefficients of the derivative, (P'(X)), of P(X)
NDP	I	NDP	I		Degree of the derivative of original polynomial
D	C	UD,VD	D		Array of coefficients of the greatest common divisor of P(X) and P'(X)
ND	I	ND	I		Degree of g.c.d. of P(X) and P'(X)
Q	C	UQ,VQ	D		Array of coefficients of quotient polynomial P(X)/g.c.d.

TABLE E.VI (Continued)

Single Precision		Double Precision		Disposition of Argument	Description
Variable	Type	Variable	Type		
NQ	I	NQ	I		Degree of quotient polynomial Q(X)
ZRO	C	UZRO,VZRO	D		Value at which to evaluate or deflate polynomial
DUMMY	C	UDUMMY,VDUMMY	D		Dummy variable
QQ	C	UQQ,VQQ	D		Working array of coefficients of current polynomial
NQQ	I	NQQ	I		Degree of current polynomial, QQ(X)
IALTER	I	IALTER	I		Number of alterations of an initial approximation
CONV	L	CONV	L		CONV = TRUE implies convergence to a root
EPS4	R	EPS4	D		Tolerance for checking multiplicities
AP	C	UAP,VAP	D		Array of approximations (initial or altered) producing convergence
QD	C	UQD,VQD	D		Array of coefficients of newly deflated polynomial
JAP	I	JAP	I		Number of distinct roots found by iterative process i.e. not as a result of Subroutine QUAD
J1	I	J1	I		Number of distinct roots found in the attempt to improve roots
ROOTS	C	UROOTS,VROOTS	D		Array of improved roots
NEWT	L	NEWT	L		Program control. NEWT = TRUE implies that Newton's method was used instead of Subroutine QUAD
Subroutine NEWTON					
X	C	UX,VX	D	E	Starting approximation (initial or altered)
N	I	N	I	E	Degree of current polynomial
P	C	UP,VP	D	E	Array of coefficients of current polynomial
MAX	I	MAX	I	C	Maximum number of iterations
EPSLON	R	EPSLON	D	C	Tolerance for checking convergence
X0	C	UX0,VX0	D	R	Current approximation to root
B	C	UB,VB	D		Array of coefficients of newly deflated polynomial
DPX0	C	UDPX0,VDPX0	D		Derivative of the polynomial at X0

TABLE E.VI (Continued)

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition of Argument</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>		
DIFF	C	UDIFF,VDIFF	D		PX0/DPX0
PX0	C	UPX0,VPX0	D		Value of polynomial at X0
CONV	L	CONV	L	R	CONV = TRUE implies convergence to root
Subroutine HORNER					
X	C	UX,VX	D	E	Value at which to evaluate or deflate polynomial
N	I	N	I	E	Degree of polynomial
P	C	UP,VP	D		Array of coefficients of polynomial
C	C	UC,VC	D	R	Updated at each iteration to yield derivative of polynomial at X
B	C	UB,VB	D		Array of coefficients of newly deflated polynomial
Subroutine QUAD					
N	I	N	I	E	Degree of polynomial to be solved
A	C	UA,VA	D	E	Array of coefficients of polynomial to be solved
J	I	J	I	ECR	Number of distinct roots found of original polynomial (J = -1 implies original polynomial is of degree 2 or 1)
ROOT	C	UROOT,VROOT	D	ECR	Array of roots found
MULT	I	MULT	I	ECR	Array of multiplicities
DISC	C	UDISC,VDISC	D		Discriminate of quadratic
TEMP	C	UTEMP,VTEMP	D		$\sqrt{\text{DISC}}$
EPSLON	R	EPSLON	D	C	Tolerance for zero (0)
D	C	UD,VD	D		Twice leading coefficient of quadratic

TABLE E.VI (Continued)

Single Precision		Double Precision		Disposition of Argument	Description
Variable	Type	Variable	Type		
Subroutine GCD					
R	C	UR,VR	D	E	Array of coefficients of original polynomial, P(X)
S	C	US,VS	D	E	Array of coefficients of derivative polynomial, P'(X)
N	I	N	I	E	Degree of original polynomial, P(X)
M	I	M	I	E	Degree of derivative polynomial, P'(X)
RR	C	URR,VRR	D		Array of coefficients of dividend polynomial
SS	C	USS,VSS	D	R	Array of coefficients of divisor polynomial also array of coefficients of g.c.d. of P(X) and P'(X) when returned
N1	I	N1	I		Degree of dividend polynomial, RR(X)
M1	I	M1	I	R	Degree of divisor polynomial, SS(X), also degree of g.c.d. of P(X) and P'(X) when returned
D	C	UD,VD	D		Quotient RR_{N1+1}/SS_{M1+1}
T	C	UT,VT	D		Array of coefficients of difference polynomial (RR - D(SS))
K	I	K	I		Degree of difference polynomial T(X)
EPSLON	R	EPSLON	D	C	Tolerance check for zero (0)
Subroutine MULTI					
N	I	N	I	E	Degree of original polynomial, P(X)
P	C	UP,VP	D	E	Array of coefficients of original polynomial, P(X)
J	I	J	I	E	Number of distinct roots of P(X)
ROOT	C	UROOT,VROOT	D	E	Array of distinct roots of P(X)
A	C	UA,VA	D		Working array of coefficients of current polynomial
M	I	M	I		Degree of current polynomial, A(X)
MULT	I	MULT	I	R	Array of multiplicities of the roots
IO2	I	IO2	I	C	Unit number of output device
B	C	UB,VB	D		Array of coefficients of newly deflated polynomial
C	C	UC,VC	D		Derivative of polynomial at ROOT _i
EPSLON	R	EPSLON	D	C	Tolerance for checking multiplicities

TABLE E.VI (Continued)

<u>Single Precision</u> <u>Variable</u>	<u>Double Precision</u> <u>Variable</u>	<u>Precision</u> <u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
Subroutine DERIV				
N	N	I	E	Degree of polynomial, P(X)
P	UP, VP	D	E	Array of coefficients of polynomial, P(X)
A	UA, VA	D	R	Array of coefficients of derivative, P'(X)
M	M	I	R	Degree of derivative polynomial, P'(X)
Subroutine DIVIDE				
P	UP, VP	D	E	Array of coefficients of dividend polynomial
N	N	I	E	Degree of dividend polynomial
D	UD, VD	D	E	Array of coefficients of divisor polynomial
M	M	I	E	Degree of divisor polynomial
Q	UQ, VQ	D	R	Array of coefficients of quotient polynomial P(X)/D(X)
K	K	I	R	Degree of quotient polynomial, Q(X)
J	J	I		Counter
TERM	UTERM, VTERM	D		Dummy variable used for temporary storage of products
KK	KK	I		Number of coefficients of quotient polynomial, Q(X)
Subroutine GENAPP				
APP	APPR, APPI	D	R	Array containing initial approximations
NAPP	NAPP	I	E	Number of initial approximations to be generated
XSTART	XSTART	D	ECR	Magnitude at which to begin generating approximations; also magnitude of the approximation being generated
BETA	BETA	D		Argument of complex approximation being generated
U	APPR(I)	D		Real part of complex approximation
V	APPI(I)	D		Imaginary part of complex approximation

TABLE E.VI (Continued)

<u>Single Precision Variable</u>	<u>Precision Type</u>	<u>Double Precision Variable</u>	<u>Precision Type</u>	<u>Disposition of Argument</u>	<u>Description</u>
Subroutine ALTER					
XOLD	C	XOLDR, XOLDI	D	ECR	Old approximation to be altered to new approximation
NALTER	I	NALTER	I	ECR	Number of alterations performed on an initial approximation
ITIME	I	ITIME	I	E	Program control
MAX	I	MAX	I	C	Maximum number of iterations permitted
Y	R	XOLDI	D		Imaginary part of original initial approximation (unaltered)
X	R	XOLDR	D		Real part of original, unaltered initial approximation
R	R	ABXOLD	D		Magnitude of original unaltered initial approximation
BETA	R	BETA	D		Argument of new approximation
XOLDR	R	XOLDR	D		Real part of new approximation
XOLDI	R	XOLDI	D		Imaginary part of new approximation
I02	I	I02	I	C	Unit number of output device
Subroutine COMSQT					
		UX, VX	D	E	Complex number for which the square root is desired
		UY, VY	D	R	Square root of the complex number

4. Description of Program Output

The output from G.C.D. - Newton's method consists of the following information.

The heading is "GREATEST COMMON DIVISOR METHOD USED WITH NEWTON'S METHOD TO FIND ZEROS OF POLYNOMIALS NUMBER XX." XX represents the number of the polynomial.

As an aid to ensure that the control information is correct, the number of initial approximations given, maximum number of iterations, test for zero in subroutine GCD, test for convergence, test for zero in subroutine QUAD, test for multiplicities, radius to start search, and radius to end search are printed as read from the control card.

The coefficients of the polynomial are printed under the heading "THE DEGREE OF P(X) IS XX THE COEFFICIENTS ARE." XX represents the degree of the polynomial. The coefficient of the highest degree term is printed first.

The polynomial obtained after dividing the original polynomial, $P(X)$, by the greatest common divisor of $P(X)$ and its derivative, $P'(X)$, is printed under the heading "Q(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE DISTINCT ROOTS OF P(X). THE DEGREE OF Q(X) IS XX THE COEFFICIENTS ARE." XX represents the degree of this polynomial. This polynomial contains all distinct roots and is solved by Newton's method. The coefficient of the highest degree term is printed first; that is, the leading coefficient is printed first.

The zeros found before the attempt to improve accuracy are printed under the heading "ROOTS OF Q(X)."

The initial approximation producing convergence to a root is

printed to the right of the corresponding root and headed by "INITIAL APPROXIMATION." The initial approximations may be those supplied by the user, or generated by the program or a combination of both. The message "RESULTS OF SUBROUTINE QUAD" indicates that the corresponding root was obtained by subroutine QUAD. See Appendix D, § 5.

The zeros found after the attempt to improve accuracy are printed under the heading "ROOTS OF P(X)." The corresponding initial approximation producing convergence is printed as described above.

The multiplicity of each zero is given under the title "MULTIPLICITIES."

5. Informative Messages and Error Messages

The output may contain informative or error messages. These are intended as an aid to the user and are described as follows.

If not all roots of a polynomial were found before the attempt to improve accuracy, the remaining unsolved polynomial will be printed, with the leading coefficient first, under the heading "COEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO ZEROS WERE FOUND." See Appendix D, § 6.

"NO ROOTS FOR INITIAL APPROXIMATION ROOT XX = YYY." This message is printed if a root fails to produce convergence when trying to improve accuracy. XX represents the number of the root and YYY represents the value of the root before the attempt to improve accuracy.

"NO ROOTS FOR THE POLYNOMIAL Q(X) OF DEGREE XX WITH GENERATED INITIAL APPROXIMATIONS." XX represents the degree of the polynomial Q(X). This message is printed if none of the roots produce convergence in the attempt to improve accuracy.

"THE EPSILON (XXX) CHECK IN SUBROUTINE MULTI INDICATES THAT ROOT YY = ZZZ IS NOT CLOSE ENOUGH TO BE A TRUE ROOT. IT IS PRINTED BELOW WITH MULTIPLICITY 0." XXX represents the multiplicity requirement (EPS4 on the control card), YY represents the number of the root, and ZZZ represents the value of the root after the attempt to improve accuracy. The message indicates that this root does not meet the requirement for multiplicities. It is, however, usually a good approximation to the true root since convergence was obtained both before and after the attempt to improve accuracy.

MAIN PROGRAM




Figure E.6. Flow Charts for G.C.D.-Newton's Method





Figure E.6. (Continued)

QUAD



NEWTON






Figure E.6. (Continued)

DIVIDE



HORNER



DERIV




Figure E.6. (Continued)

MULTI




Figure E.6. (Continued)

COMSQT





Figure E.6. (Continued)

GENAPP



ALTER




Figure E.6. (Continued)

TABLE E.VII

PROGRAM FOR G.C.D.-NEWTON'S METHOD

```

C *****
C *
C * DOUBLE PRECISION PROGRAM FOR G.C.D. - NEWTON'S METHOD
C *
C *
C * THE G.C.D. METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C * POLYNOMIAL OF MAXIMUM DEGREE 25. ALL MULTIPLE ROOTS ARE REMOVED BY
C * DIVIDING THE POLYNOMIAL BY THE GREATEST COMMON DIVISOR OF THE POLYNOMIAL
C * AND ITS DERIVATIVE. THE ZEROS OF THE RESULTING POLYNOMIAL ARE EXTRACTED
C * AND THEIR MULTIPLICITIES DETERMINED.
C *
C *****
0001  DOUBLE PRECISION UP,VP,UAPP,VAPP,UROOT,VROOT,UDP,VDP,UD,VD,UZRO,VZ
      1RO,UQ,VQ,UDUMMY,VDUMMY,UQQ,VQQ,UAP,VAP,UQD,VQD,UROOTS,VROOTS,EPS1,
      2EPS2,EPS3,EPS4
0002  DOUBLE PRECISION XSTART
0003  DOUBLE PRECISION XEND
0004  DIMENSION UP(26),VP(26),UAPP(25),VAPP(25),UROOT(25),VROOT(25),MULT
      1(25),UDP(26),VDP(26),UD(26),VD(26),UQ(26),VQ(26),UQQ(26),VQQ(26),U
      2AP(25),VAP(25),UQD(26),VQD(26),ANAME(2),ENTRY(26),UROOTS(25),VROOT
      3S(25)
0005  COMMON EPS1,EPS2,EPS3,EPS4,IO2,MAX
0006  LOGICAL NEWT,CONV
0007  DATA PNAME,QNAME,QQNAME/2HP(,2HQ(,3HQQ(/
0008  DATA ENTRY/1H1,1H2,1H3,1H4,1H5,1H6,1H7,1H8,1H9,2H10,2H11,2H12,2H13
      1,2H14,2H15,2H16,2H17,2H18,2H19,2H20,2H21,2H22,2H23,2H24,2H25,2H26/
0009  DATA ANAME(1),ANAME(2)/4HNEWT,4HONS /
0010  IO1=5
0011  IO2=6
0012  10 J=0
0013  ITIME=0
0014  READ(IO1,1000) NOPOLY,NP,NAPP,MAX,EPS1,EPS2,EPS3,EPS4,XSTART,XEND,
      1KCHECK
0015  IF(KCHECK.EQ.1) STOP
0016  WRITE(IO2,1020) ANAME(1),ANAME(2),NOPOLY
0017  WRITE(IO2,2000) NAPP
0018  WRITE(IO2,2010) MAX
0019  WRITE(IO2,2070) EPS1
0020  WRITE(IO2,2020) EPS2
0021  WRITE(IO2,2080) EPS3
0022  WRITE(IO2,2030) EPS4
0023  WRITE(IO2,2040) XSTART
0024  WRITE(IO2,2050) XEND
0025  WRITE(IO2,2060)
0026  KKK=NP+1
0027  NNN=KKK+1
0028  DO 20 I=1,KKK
0029  JJJ=NNN-I
0030  20 READ(IO1,1010) UP(JJJ),VP(JJJ)
0031  IF(NAPP.NE.0) GO TO 22
0032  NAPP=NP
0033  CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0034  GO TO 23
0035  22 READ(IO1,1015) (UAPP(I),VAPP(I),I=1,NAPP)
0036  23 WRITE(IO2,1030) NP
0037  KKK=NP+1
0038  NNN=KKK+1
0039  DO 25 I=1,KKK

```

TABLE E.VII (Continued)

```

0040      JJJ=NNN-I
0041 25 WRITE(102,1040) PNAME,ENTRY(JJJ),UP(JJJ),VP(JJJ)
0042    IF(NP.GE.3) GO TO 30
0043      J=-1
0044      CALL QUAD(NP,UP,VP,J,UROOT,VROOT,MULT)
0045      WRITE(102,1070)
0046      WRITE(102,1165) (I,UROOT(I),VROOT(I),MULT(I),I=1,J)
0047      GO TO 10
0048 30 CALL DERIV(NP,UP,VP,NDP,UDP,VDP)
0049      CALL GCD(NP,UP,VP,NDP,UDP,VDP,ND,UD,VD)
0050      IF(ND.GT.1) GO TO 70
0051      IF(ND.EQ.0) GO TO 65
0052      UDUMMY=(UD(2)*UD(2))+(VD(2)*VD(2))
0053      UZRO=(-(UD(1)*UD(2))-(VD(1)*VD(2)))/UDUMMY
0054      VZRO=(-(UD(2)*VD(1))+(UD(1)*VD(2)))/UDUMMY
0055      CALL HORNER(UZRO,VZRO,NP,UP,VP,UQ,VQ,UDUMMY,VDUMMY)
0056      NQ=NP-1
0057      DO 60 I=1,NP
0058        UQ(I)=UQ(I+1)
0059 60 VQ(I)=VQ(I+1)
0060      GO TO 80
0061 65 KKK=NP+1
0062      DO 66 I=1,KKK
0063        UQ(I)=UP(I)
0064 66 VQ(I)=VP(I)
0065      NQ=NP
0066      GO TO 80
0067 70 CALL DIVIDE(NP,UP,VP,ND,UD,VD,NQ,UQ,VQ)
0068 80 WRITE(102,1120) NQ
0069      KKK=NQ+1
0070      NNN=KKK+1
0071      DO 83 I=1,KKK
0072        JJJ=NNN-I
0073 83 WRITE(102,1040) QNAME,ENTRY(JJJ),UQ(JJJ),VQ(JJJ)
0074      IF(NQ.GE.3) GO TO 85
0075      GO TO 110
0076 85 KKK=NQ+1
0077      DO 90 I=1,KKK
0078        UQQ(I)=UQ(I)
0079 90 VQQ(I)=VQ(I)
0080      NQQ=NQ
0081      GO TO 120
0082 110 CALL QUAD(NQ,UQ,VQ,J,UROOT,VROOT,MULT)
0083      NEWT=.FALSE.
0084      GO TO 310
0085 120 DO 200 I=1,NAPP
0086      IALTER=0
0087 130 CALL NEWTON(UAPP(I),VAPP(I),NQQ,UQQ,VQQ,UZRO,VZRO,CONV)
0088      IF(CONV) GO TO 160
0089      CALL ALTER(UAPP(I),VAPP(I),IALTER,ITIME)
0090      IF(IALTER.GT.5) GO TO 200
0091      GO TO 130
0092 160 J=J+1
0093      UROOT(J)=UZRO
0094      VROOT(J)=VZRO
0095      UAP(J)=UAPP(I)
0096      VAP(J)=VAPP(I)
0097      CALL HORNER(UZRO,VZRO,NQQ,UQQ,VQQ,UQD,VQD,UDUMMY,VDUMMY)

```


TABLE E.VII (Continued)

```

0098      DO 180 I1=1,NQQ
0099      UQQ(I1)=UQD(I1+1)
0100 180 VQQ(I1)=VQD(I1+1)
0101      NQQ=NQQ-1
0102      IF(NQQ.LT.3) GO TO 220
0103 200 CONTINUE
0104      IF(J.GE.NQ) GO TO 205
0105      IF(XEND.EQ.0.0) GO TO 205
0106      IF(XSTART.GT.XEND) GO TO 205
0107      NAPP=NQ
0108      CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0109      GO TO 120
0110 205 IF(NQQ.LE.2) GO TO 210
0111      WRITE(102,1200)
0112      KKK=NQQ+1
0113      NNN=KKK+1
0114      DO 157 L=1,KKK
0115      JJJ=NNN-L
0116 157 WRITE(102,1100) QQNAME,ENTRY(JJJ),UQQ(JJJ),VQQ(JJJ)
0117 210 IF(J.EQ.0) GO TO 10
0118      JAP=J
0119      GO TO 230
0120 220 JAP=J
0121      CALL QUAD(NQQ,UQQ,VQQ,J,UROOT,VROOT,MULT)
0122 230 WRITE(102,1132)
0123      WRITE(102,1133) (I,UROOT(I),VROOT(I),UAP(I),VAP(I),I=1,JAP)
0124      IF(JAP.LT.J) GO TO 235
0125      GO TO 240
0126 235 KKK=JAP+1
0127      WRITE(102,1134) (I,UROOT(I),VROOT(I),I=KKK,J)
0128 240 J1=0
0129      DO 300 I=1,J
0130      CALL NEWTON(UROOT(I),VROOT(I),NQ,UQ,VQ,UZRO,VZRO,CONV)
0131      IF(CONV) GO TO 280
0132      WRITE(102,1140) I,UROOT(I),VROOT(I)
0133      IF(I.LT.JAP) GO TO 241
0134      IF(I.EQ.JAP) GO TO 250
0135      GO TO 300
0136 241 KKK=JAP-1
0137      DO 245 I1=I,KKK
0138      UAP(I1)=UAP(I1+1)
0139 245 VAP(I1)=VAP(I1+1)
0140 250 JAP=JAP-1
0141      GO TO 300
0142 280 J1=J1+1
0143      UROOTS(J1)=UZRO
0144      VROOTS(J1)=VZRO
0145 300 CONTINUE
0146      IF(J1.EQ.0) GO TO 305
0147      J=J1
0148      DO 303 I=1,J
0149      UROOT(I)=UROOTS(I)
0150 303 VROOT(I)=VROOTS(I)
0151      GO TO 307
0152 305 WRITE(102,1150) NQ
0153      KKK=NQ+1
0154      NNN=KKK+1
0155      DO 306 L=1,KKK

```

TABLE E.VII (Continued)

```

0156      JJJ=NNN-L
0157      306 WRITE(IO2,1040) QNAME,ENTRY(JJJ),UQ(JJJ),VQ(JJJ)
0158      GO TO 10
0159      307 NEWT=.TRUE.
0160      310 CALL MULTI(NP,UP,VP,J,UROOT,VROOT,MULT)
0161      IF(NEWT) GO TO 330
0162      WRITE(IO2,1070)
0163      WRITE(IO2,1165) (L,UROOT(L),VROOT(L),MULT(L),L=1,J)
0164      GO TO 10
0165      330 WRITE(IO2,1180)
0166      WRITE(IO2,1190) (L,UROOT(L),VROOT(L),MULT(L),UAP(L),VAP(L),L=1,JAP
0167      )
0168      KKK=JAP+1
0169      IF(JAP.LT.J) WRITE(IO2,1165) (L,UROOT(L),VROOT(L),MULT(L),L=KKK,J)
0170      GO TO 10
0171      1000 FORMAT(3(I2,1X),9X,I3,1X,4(D6.0,1X),13X,2(D7.0,1X),I1)
0172      1010 FORMAT(2D30.0)
0173      1015 FORMAT(2D30.0)
0174      1020 FORMAT(1H1,10X,41HGREATEST COMMON DIVISOR METHOD USED WITH ,2(A4),
0175      135HMETHOD TO FIND ZEROS OF POLYNOMIALS/1X,18HPOLYNOMIAL NUMBER ,I
0176      22//)
0177      1030 FORMAT(1X,22HTHE DEGREE OF P(X) IS ,I2,22H THE COEFFICIENTS ARE//
0178      1)
0179      1040 FORMAT(2X,A2,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0180      1070 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES//)
0181      1080 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,10X,I2)
0182      1100 FORMAT(2X,A3,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0183      1120 FORMAT(///1X,73HQ(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE
0184      1DISTINCT ROOTS OF P(X)./1X,22HTHE DEGREE OF Q(X) IS ,I2,22H THE C
0185      20COEFFICIENTS ARE//)
0186      1200 FORMAT(///1X,70HCOEFFICIENTS OF THE DEFLATED POLYNOMIAL FOR WHICH
0187      1NO ZEROS WERE FOUND.//)
0188      1132 FORMAT(///1X,13HROOTS OF Q(X),84X,21HINITIAL APPROXIMATION//)
0189      1133 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,17X,D23.16,3H
0190      1 + ,D23.16,2H I)
0191      1134 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,20X,26HRESULT
0192      1S OF SUBROUTINE QUAD)
0193      1140 FORMAT(///,1X,40HNO ROOTS FOR INITIAL APPROXIMATION ROOT(,I2,4H) =
0194      1 ,D23.16,3H + ,D23.16,2H I)
0195      1150 FORMAT(///,1X,45HNO ROOTS FOR THE POLYNOMIAL Q(X) OF DEGREE = ,I2,
0196      138H WITH GENERATED INITIAL APPROXIMATIONS//)
0197      1165 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,10X,26H
0198      1RESULTS OF SUBROUTINE QUAD)
0199      1180 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES,17X,21HINITIAL
0200      1 APPROXIMATION//)
0201      1190 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,7X,D23.
0202      116,3H + ,D23.16,2H I)
0203      2000 FORMAT(1X,41HNUMBER OF INITIAL APPROXIMATIONS GIVEN. ,I2)
0204      2010 FORMAT(1X,29HMAXIMUM NUMBER OF ITERATIONS.,11X,I3)
0205      2020 FORMAT(1X,21HTEST FOR CONVERGENCE.,13X,D9.2)
0206      2030 FORMAT(1X,24HTEST FOR MULTIPLICITIES.,10X,D9.2)
0207      2040 FORMAT(1X,23HRADIUS TO START SEARCH.,11X,D9.2)
0208      2050 FORMAT(1X,21HRADIUS TO END SEARCH.,13X,D9.2)
0209      2060 FORMAT(//1X)
0210      2070 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE GCD. ,D9.2)
0211      2080 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE QUAD. ,D9.2)
0212      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE GENAPP(APPR,APPI,NAPP,XSTART)
C      *****
C      *
C      * SUBROUTINE GENAPP GENERATES N INITIAL APPROXIMATIONS, WHERE N IS THE
C      * DEGREE OF THE ORIGINAL POLYNOMIAL.
C      *
C      *****
0002      DOUBLE PRECISION APPR,APPI,XSTART,BETA, EPS1,EPS2,EPS3,EPS4
0003      DIMENSION APPR(25),APPI(25)
0004      COMMON EPS1,EPS2,EPS3,EPS4,IO2,MAX
0005      IF(XSTART.EQ.0.0) XSTART=0.5
0006      BETA=0.2617994
0007      DO 10 I=1,NAPP
0008      APPR(I)=XSTART*DCOS(BETA)
0009      APPI(I)=XSTART*DSIN(BETA)
0010      BETA=BETA+0.5235988
0011 10 XSTART=XSTART+0.5
0012      RETURN
0013      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE ALTER(XOLDR,XOLDI,NALTER,ITIME)
C      *****
C      *
C      * SUBROUTINE ALTER ALTERS THE INITIAL APPROXIMATIONS WHICH PRODUCE NO
C      * CONVERGENCE TO A ZERO. THIS IS DONE A MAXIMUM OF 5 TIMES FOR EACH ROOT.
C      *
C      *****
0002      DOUBLE PRECISION XOLDR,XOLDI,ABXOLD,BETA,EPS1,EPS2,EPS3,EPS4
0003      COMMON EPS1,EPS2,EPS3,EPS4,IO2,MAX
0004      IF(ITIME.NE.0) GO TO 5
0005      ITIME =1
0006      WRITE(IO2,1010) MAX
0007      5 IF(NALTER.EQ.0) GO TO 10
0008      WRITE(IO2,1000) XOLDR,XOLDI
0009      GO TO 20
0010      10 ABXOLD=DSQRT((XOLDR*XOLDR)+(XOLDI*XOLDI))
0011      BETA=DATAN2(XOLDI,XOLDR)
0012      WRITE(IO2,1020) XOLDR,XOLDI
0013      20 NALTER=NALTER+1
0014      IF(NALTER.GT.5) RETURN
0015      GO TO (30,40,30,40,30),NALTER
0016      30 XOLDR=-XOLDR
0017      XOLDI=-XOLDI
0018      GO TO 50
0019      40 BETA=BETA+1.0471976
0020      XOLDR=ABXOLD*DCOS(BETA)
0021      XOLDI=ABXOLD*DSIN(BETA)
0022      50 RETURN
0023      1000 FORMAT(1X,D23.16,3H + ,D23.16,2H I,10X,21HALTERED APPROXIMATION)
0024      1010 FORMAT(//1X,54HNO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AF
          1TER ,I3,12H ITERATIONS.//)
0025      1020 FORMAT(1X,D23.16,3H + ,D23.16,2H I,10X,21HINITIAL APPROXIMATION)
0026      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE GCD(N,UR,VR,M,US,VS,M1,USS,VSS)
C      *****
C      *
C      * GIVEN POLYNOMIALS P(X) AND DP(X) WHERE DEG. DP(X) IS LESS THAN DEG.
C      * P(X), SUBROUTINE GCD COMPUTES THE GREATEST COMMON DIVISOR OF P(X) AND
C      * DP(X).
C      *
C      *****
0002      DOUBLE PRECISION USSSSS,VSSSSS
0003      DOUBLE PRECISION UR,VR,US,VS,USS,VSS,URR,VRR,UD,VD,UT,VT,EPSLON,EP
0004      IS2,EPS3,EPS4,BBB
0005      DIMENSION UR(26),VR(26),US(26),VS(26),USS(26),VSS(26),URR(26),VRR(
0006      26),UT(26),VT(26)
0007      COMMON EPSLON,EPS2,EPS3,EPS4,IO2,MAX
0008      N1=N
0009      M1=M
0010      KKK=N+1
0011      DO 20 I=1,KKK
0012      URR(I)=UR(I)
0013      VRR(I)=VR(I)
0014      KKK=M+1
0015      DO 25 I=1,KKK
0016      USS(I)=US(I)
0017      VSS(I)=VS(I)
0018      30 BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0019      UD=(URR(N1+1)*USS(M1+1)+VRR(N1+1)*VSS(M1+1))/BBB
0020      VD=(USS(M1+1)*VRR(N1+1)-URR(N1+1)*VSS(M1+1))/BBB
0021      KKK=N1+1-M1
0022      DO 40 I=KKK,N1
0023      UT(I)=URR(I)-(UD*USS(I-N1+M1)-VD*VSS(I-N1+M1))
0024      VT(I)=VRR(I)-(UD*VSS(I-N1+M1)+VD*USS(I-N1+M1))
0025      IF(M1.EQ.N1) GO TO 70
0026      KKK=N1-M1
0027      DO 60 I=1,KKK
0028      UT(I)=URR(I)
0029      VT(I)=VRR(I)
0030      70 DO 90 I=1,N1
0031      BBB=DSQRT(UT(N1+1-I)*UT(N1+1-I)+VT(N1+1-I)*VT(N1+1-I))
0032      IF(BBB.GT.EPSLON) GO TO 100
0033      90 CONTINUE
0034      DO 95 I=1,M1
0035      BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0036      USSSSS=(USS(I)*USS(M1+1)+VSS(I)*VSS(M1+1))/BBB
0037      VSSSSS=(VSS(I)*USS(M1+1)-USS(I)*VSS(M1+1))/BBB
0038      USS(I)=USSSSS
0039      VSS(I)=VSSSSS
0040      95 VSS(I)=VSSSSS
0041      USS(M1+1)=1.0
0042      VSS(M1+1)=0.0
0043      GO TO 200
0044      100 K=N1-I
0045      IF(K.EQ.0) GO TO 170
0046      IF(K.LT.M1) GO TO 140
0047      KKK=K+1
0048      DO 130 J=1,KKK
0049      URR(J)=UT(J)
0050      VRR(J)=VT(J)
0051      N1=K
0052      GO TO 30

```

TABLE E.VII (Continued)

```

0050      140 KKK=K+1
0051          DO 150 J=1, KKK
0052              URR(J)=USS(J)
0053              VRR(J)=VSS(J)
0054              USS(J)=UT(J)
0055      150 VSS(J)=VT(J)
0056          KKK=K+2
0057          NNN=M1+1
0058          DO 160 J=KKK, NNN
0059              URR(J)=USS(J)
0060      160 VRR(J)=VSS(J)
0061          N1=M1
0062          M1=K
0063          GO TO 30
0064      170 USS(1)=1.0
0065          VSS(1)=0.0
0066          M1=0
0067      200 RETURN
0068          END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE QUAD(N,UA,VA,J,URROOT,VROOT,MULT)
C      *****
C      *
C      * SUBROUTINE QUAD SOLVES DIRECTLY FOR THE ZEROS AND THEIR MULTIPLICITIES *
C      * OF EITHER A QUADRATIC POLYNOMIAL OR A LINEAR FACTOR. SOLUTION OF THE *
C      * QUADRATIC IS DONE USING THE QUADRATIC FORMULA. *
C      *
C      *****
0002      DOUBLE PRECISION UA,VA,URROOT,VROOT,UDISC,VDISC,UTEMP,VTEMP,UD,VD,E
1PS1,EPS2,EPS4,EPSLON,BBB
0003      DIMENSION UA(26),VA(26),URROOT(25),VROOT(25),MULT(25)
0004      COMMON EPS1,EPS2,EPSLON,EPS4,IQ2,MAX
0005      IF(N.GT.1) GO TO 60
0006      IF(J.LT.0) GO TO 40
0007      J=J+1
0008      GO TO 50
0009      40 MULT(1)=1
0010      J=1
0011      50 BBB=UA(2)*UA(2)+VA(2)*VA(2)
0012      URROOT(J)=- (UA(1)*UA(2)+VA(1)*VA(2))/BBB
0013      VROOT(J)=- (VA(1)*UA(2)-UA(1)*VA(2))/BBB
0014      GO TO 200
0015      60 UDISC=(UA(2)*UA(2)-VA(2)*VA(2))- (4.0*(UA(3)*UA(1)-VA(3)*VA(1)))
0016      VDISC=(2.0*UA(2)*VA(2))- (4.0*(UA(3)*VA(1)+VA(3)*UA(1)))
0017      BBB=DSQRT(UDISC*UDISC+VDISC*VDISC)
0018      IF(BBB.LE.EPSLON) GO TO 100
0019      IF(J.GE.0) GO TO 80
0020      MULT(1)=1
0021      MULT(2)=1
0022      J=0
0023      80 CALL COMSQT(UDISC,VDISC,UTEMP,VTEMP)
0024      UD=2.0*UA(3)
0025      VD=2.0*VA(3)
0026      BBB=UD*UD+VD*VD
0027      URROOT(J+1)=((-UA(2)+UTEMP)*UD+(-VA(2)+VTEMP)*VD)/BBB
0028      VROOT(J+1)=((-VA(2)+VTEMP)*UD-(-UA(2)+UTEMP)*VD)/BBB
0029      URROOT(J+2)=((-UA(2)-UTEMP)*UD+(-VA(2)-VTEMP)*VD)/BBB
0030      VROOT(J+2)=((-VA(2)-VTEMP)*UD-(-UA(2)-UTEMP)*VD)/BBB
0031      J=J+2
0032      GO TO 200
0033      100 IF(J.LT.0) GO TO 110
0034      J=J+1
0035      GO TO 130
0036      110 MULT(1)=2
0037      J=1
0038      130 UD=2.0*UA(3)
0039      VD=2.0*VA(3)
0040      BBB=UD*UD+VD*VD
0041      URROOT(J)=(-UA(2)*UD-VA(2)*VD)/BBB
0042      VROOT(J)=(-VA(2)*UD+UA(2)*VD)/BBB
0043      200 RETURN
0044      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE NEWTON(UX,VX,N,UP,VP,UXO,VXO,CONV)
C      *****
C      *
C      * THIS SUBROUTINE CALCULATES A NEW APPROXIMATION FROM THE OLD APPROX-
C      * IMATION BY USING THE ITERATION FORMULA
C      *
C      *       $X(N+1) = X(N) - P(X(N))/P'(X(N))$ .
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UP,VP,UXO,VXO,UB,VB,UDPXO,VDPXO,UPXO,VPXO,U
1DIFF,VDIFF,EPS1,EPSLON,EPS3,EPS4,AAA,BBB
0003      DOUBLE PRECISION DDD
0004      DOUBLE PRECISION ABPXO
0005      DIMENSION UP(26),VP(26),UB(26),VB(26)
0006      COMMON EPS1,EPSLON,EPS3,EPS4,IO2,MAX
0007      LOGICAL CONV
0008      UXO=UX
0009      VXO=VX
0010      DO 10 I=1,MAX
0011      CALL HORNER(UXO,VXO,N,UP,VP,UB,VB,UDPXO,VDPXO)
0012      UPXO=UB(1)
0013      VPXO=VB(1)
0014      DDD=DSQRT(UDPXO*UDPXO+VDPXO*VDPXO)
0015      IF(DDD.NE.0.0) GO TO 5
0016      ABPXO=DSQRT(UPXO*UPXO+VPXO*VPXO)
0017      IF(ABPXO.EQ.0.0) GO TO 20
0018      GO TO 15
0019      5 BBB=UDPXO*UDPXO+VDPXO*VDPXO
0020      UDIFF=(UPXO*UDPXO+VPXO*VDPXO)/BBB
0021      VDIFF=(VPXO*UDPXO-UPXO*VDPXO)/BBB
0022      UXO=UXO-UDIFF
0023      VXO=VXO-VDIFF
0024      AAA=DSQRT(UDIFF*UDIFF+VDIFF*VDIFF)
0025      BBB=DSQRT(UXO*UXO+VXO*VXO)
0026      IF(BBB.EQ.0.0) GO TO 10
0027      IF(AAA/BBB.LT.EPSLON) GO TO 20
0028      10 CONTINUE
0029      15 CONV=.FALSE.
0030      RETURN
0031      20 CONV=.TRUE.
0032      RETURN
0033      END

```


TABLE E.VII (Continued)

```

0001      SUBROUTINE DIVIDE(N,UP,VP,M,UD,VD,K,UQ,VQ)
C      *****
C      *
C      * GIVEN TWO POLYNOMIALS F(X) AND G(X), SUBROUTINE DIVIDE COMPUTES THE
C      * QUOTIENT POLYNOMIAL H(X) = F(X)/G(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UD,VD,UQ,VQ,UTERM,VTERM,UDUMMY
0003      DIMENSION UP(26),VP(26),UD(26),VD(26),UQ(26),VQ(26)
0004      K=N-M
0005      UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0006      UQ(K+1)=(UP(N+1)*UD(M+1)+VP(N+1)*VD(M+1))/UDUMMY
0007      VQ(K+1)=(VP(N+1)*UD(M+1)-UP(N+1)*VD(M+1))/UDUMMY
0008      IF(K.EQ.0) GO TO 100
0009      J=-1
0010      DO 50 I=1,K
0011      J=J+1
0012      UTERM=UP(N-J)
0013      VTERM=VP(N-J)
0014      KK=K+1
0015      NNN=M-J
0016      DO 40 M=NNN,M
0017      IF(KK.GT.1) GO TO 10
0018      GO TO 45
0019      10 IF(M1.GE.1) GO TO 20
0020      GO TO 40
0021      20 UTERM=UTERM-(UQ(KK)*UD(M1)-VQ(KK)*VD(M1))
0022      VTERM=VTERM-(UQ(KK)*VD(M1)+VQ(KK)*UD(M1))
0023      40 KK=KK-1
0024      45 UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0025      UQ(K+1-I)=(UTERM*UD(M+1)+VTERM*VD(M+1))/UDUMMY
0026      50 VQ(K+1-I)=(VTERM*UD(M+1)-UTERM*VD(M+1))/UDUMMY
0027      100 RETURN
0028      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE HORNER(UX,VX,N,UP,VP,UB,VB,UC,VC)
C      *****
C      *
C      * HORNER'S METHOD COMPUTES THE VALUE OF THE POLYNOMIAL P(X) AT A
C      * POINT D AND ITS DERIVATIVE AT D. SYNTHETIC DIVISION IS USED TO
C      * DEFLATE THE POLYNOMIAL BY DIVIDING OUT THE FACTOR (X - D).
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UP,VP,UB,VB,UC,VC
0003      DOUBLE PRECISION UDUMMY,VDUMMY
0004      DIMENSION UP(26),VP(26),UB(26),VB(26)
0005      UB(N+1)=UP(N+1)
0006      VB(N+1)=VP(N+1)
0007      UB(N)=(UX*UB(N+1)-VX*VB(N+1))+UP(N)
0008      VB(N)=(UX*VB(N+1)+VX*UB(N+1))+VP(N)
0009      UC=UB(N+1)
0010      VC=VB(N+1)
0011      KKK=N-1
0012      DO 10 I=1,KKK
0013          UB(KKK+1-I)=(UX*UB(KKK+2-I)-VX*VB(KKK+2-I))+UP(KKK+1-I)
0014          VB(KKK+1-I)=(UX*VB(KKK+2-I)+VX*UB(KKK+2-I))+VP(KKK+1-I)
0015          UDUMMY=UX*UC-VX*VC
0016          VDUMMY=UX*VC+VX*UC
0017          UC=UDUMMY+UB(KKK+2-I)
0018      10 VC=VDUMMY+VB(KKK+2-I)
0019      RETURN
0020      END

0001      SUBROUTINE DERIV(N,UP,VP,M,UA,VA)
C      *****
C      *
C      * GIVEN A POLYNOMIAL P(X), SUBROUTINE DERIV COMPUTES THE COEFFICIENTS OF
C      * ITS DERIVATIVE P'(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UA,VA,AAA
0003      DIMENSION UP(26),VP(26),UA(26),VA(26)
0004      KKK=N+1
0005      DO 10 I=2,KKK
0006          AAA=I-1
0007          UA(I-1)=AAA*UP(I)
0008      10 VA(I-1)=AAA*VP(I)
0009      M=N-1
0010      RETURN
0011      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE MULTI(N,UP,VP,J,UROOT,VROOT,MULT)
C      *****
C      *
C      * GIVEN N ZEROS OF A POLYNOMIAL, SUBROUTINE MULTI COMPUTES THEIR
C      * MULTIPLICITIES.
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UROOT,VROOT,UA,VA,UB,VB,UC,VC,EPS1,EPS2,EPS
0003      1LON,EPS3,BBB
0004      DIMENSION UP(26),VP(26),UROOT(25),VROOT(25),UA(26),VA(26),UB(26),V
0005      1B(26),MULT(25)
0006      COMMON EPS1,EPS2,EPS3,EPSLON,IO2,MAX
0007      DO 100 I=1,J
0008      KKK=N+1
0009      DO 10 K=1,KKK
0010      UA(K)=UP(K)
0011      VA(K)=VP(K)
0012      M=N
0013      MULT(I)=0
0014      20 CALL HORNER(UROOT(I),VROOT(I),M,UA,VA,UB,VB,UC,VC)
0015      BBB=DSQRT(UB(1)*UB(1)+VB(1)*VB(1))
0016      IF(BBB.LT.EPSLON) GO TO 50
0017      IF(MULT(I).EQ.0) GO TO 40
0018      GO TO 100
0019      40 WRITE(IO2,1000) EPSLON,I,UROOT(I),VROOT(I)
0020      GO TO 100
0021      50 MULT(I)=MULT(I)+1
0022      IF(M.GT.1) GO TO 60
0023      GO TO 100
0024      60 DO 70 K=1,M
0025      UA(K)=UB(K+1)
0026      VA(K)=VB(K+1)
0027      M=M-1
0028      GO TO 20
0029      70 CONTINUE
0030      RETURN
0031      1000 FORMAT(///15H THE EPSILON (,D10.3,48H) CHECK IN SUBROUTINE MULTI
0032      1INDICATES THAT ROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,/80H IS NO
0033      2T CLOSE ENOUGH TO BE A TRUE ROOT. IT IS PRINTED BELOW WITH MULTIP
0034      3LICITY 0//)
0035      END

```

TABLE E.VII (Continued)

```

0001      SUBROUTINE CONSQT(UX,VX,UY,VY)
C      *****
C      *
C      * THIS SUBROUTINE COMPUTES THE SQUARE ROOT OF A COMPLEX NUMBER.
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UY,VY,DUMMY,R,AAA,BBB
0003      R=DSQRT(UX*UX+VX*VX)
0004      AAA=DSQRT(DABS((R+UX)/2.0))
0005      BBB=DSQRT(DABS((R-UX)/2.0))
0006      IF(VX) 10,20,30
0007      10 UY=AAA
0008      VY=-1.0*BBB
0009      GO TO 100
0010      20 IF(UX) 40,50,60
0011      30 UY=AAA
0012      VY=BBB
0013      GO TO 100
0014      40 DUMMY=DABS(UX)
0015      UY=0.0
0016      VY=DSQRT(DUMMY)
0017      GO TO 100
0018      50 UY=0.0
0019      VY=0.0
0020      GO TO 100
0021      60 DUMMY=DABS(UX)
0022      UY=DSQRT(DUMMY)
0023      VY=0.0
0024      100 RETURN
0025      END

```

APPENDIX F

G.C.D. - MULLER'S METHOD

1. Use of the Program

A double precision FORTRAN IV program using the G.C.D. method with Muller's method as a supporting method is presented here. Flow charts for this program are given in Figure F.1 while Table F.III gives a FORTRAN IV listing of this program. Single precision variables are listed in Table F.II. The single precision variables are used in the flow charts and the corresponding double precision variables can be obtained from Table F.II.

This program is designed to solve polynomials having degree less than or equal to 25. In order to solve polynomials of degree N where $N > 25$, the data statement and array dimensions given in Table F.I must be changed.

In this program both the leading coefficient and the constant coefficient are assumed to be non-zero.

TABLE F.I

PROGRAM CHANGES NECESSARY TO SOLVE POLYNOMIALS OF DEGREE
GREATER THAN 25 BY G.C.D. - MULLER'S METHOD

Main Program

Data Entry/1H1,1H2,...,1H9,2H10,2H11,...,2HXX/where XX = N+1

URAPP(N,3), VRAPP(N,3)
UAPP(N,3), VAPP(N,3)
UP(N+1), VP(N+1)
UROOT(N), VROOT(N)
MULT(N)
UDP(N+1), VDP(N+1)
UD(N+1), VD(N+1)
UQ(N+1), VQ(N+1)
UQQ(N+1), VQQ(N+1)
UB(N+1), VB(N+1)
ENTRY(N+1)

Subroutines MULTI, DIVIDE, DERIV, GCD, and QUAD

See corresponding subroutines in Table E.I.

Subroutine MULLER

UROOT(N), VROOT(N)
MULT(N)
UAPP(N,3), VAPP(N,3)
UWORK(N+1), VWORK(N+1)
UB(N+1), VB(N+1)
UA(N+1), VA(N+1)
URAPP(N,3), VRAPP(N,3)

Subroutine BETTER

UROOT(N), VROOT(N)
UA(N+1), VA(N+1)
UBAPP(N,3), VBAPP(N,3)
UB(N+1), VB(N+1)
UROOTS(N), VROOTS(N)
URAPP(N,3), VRAPP(N,3)
MULT(N)

Subroutine GENAPP

APPR(N,3) APPI(N,3)

Subroutine HORNER

UA(N+1), VA(N+1)
UB(N+1), VB(N+1)

2. Input Data for G.C.D. - Muller's Method

The input data for G.C.D. - Muller's method is prepared exactly as described in Appendix E, § 2 for G.C.D. - Newton's method.

3. Variables Used in G.C.D. - Muller's Method

The main variables used in G.C.D. - Muller's method are given in Table F.II. The symbols used to indicate type and disposition are described in Appendix E, § 3. For variables not listed in Table F.II, see the main program or corresponding subprogram of Table E.VI.

4. Description of Program Output

The output from G.C.D. - Muller's method is identical to that for G.C.D. - Newton's method as described in Appendix E, § 4, keeping in mind that Muller's instead of Newton's method is used. The expression "SOLVED BY DIRECT METHOD" is equivalent to "RESULTS OF SUBROUTINE QUAD." Only one initial approximation, X_0 , (not three) is printed. The other two required by Muller's method were $.9X_0$ and $1.1X_0$.

5. Informative Messages and Error Messages

The informative messages and error messages in this program are described as follows. For other messages not listed here, see Appendix E, § 5.

"THE EPSILON (XXX) CHECK IN SUBROUTINE MULTI INDICATES THAT ROOT YY = ZZZ IS NOT CLOSE ENOUGH TO BE A TRUE ROOT. IT IS PRINTED BELOW WITH MULTIPLICITY 0." This message is described in Appendix E, § 5.

"COEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO ZEROS WERE FOUND." This message is described in Appendix E, § 5.

"NO ZEROS WERE FOUND FOR POLYNOMIAL NUMBER XX." XX represents the number of the polynomial for which no zeros were extracted.

"IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT XX = YYY DID NOT CONVERGE AFTER ZZZ ITERATIONS." This message indicates that a root did not produce convergence during the attempt to improve accuracy. XX represents the number of the root before the attempt to improve accuracy, YYY represents its value, and ZZZ represents the maximum number of iterations. The following message then follows. "THE PRESENT APPROXIMATION IS AAA." AAA represents the present approximation to the root after the maximum number of iterations.

TABLE F.II

VARIABLES USED IN G.C.D. - MULLER'S METHOD

<u>Single Precision</u> <u>Variable</u>	<u>Type</u>	<u>Double Precision</u> <u>Variable</u>	<u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
Subroutine MULLER					
NP	I	NP	I	E	Degree of polynomial P(X)
NROOT	I	NROOT	I	R	Number of distinct roots found
NOMULT	I	NOMULT	I		Number of roots (counting multiplicities)
ROOT	C	UROOT, VROOT	D	R	Array containing the roots
NAPP	I	NAPP	I	E	Number of initial approximations
APP	C	UAPP, VAPP	D	E	Array of initial approximations
WORK	C	UWORK, VWORK	D		Working array containing coefficients of current polynomial
B	C	UB, VB	D		Array containing coefficients of deflated polynomial
A	C	UA, VA	D	E	Array containing coefficients of original polynomial, P(X)
RAPP	C	URAPP, VRAPP	D	R	Array of initial or altered approximation for which convergence was obtained
X1	C	UX1, VX1	D		One of three current approximations to a root
X2	C	UX2, VX2	D		One of three current approximations to a root
X3	C	UX3, VX3	D		One of three current approximations to a root
PX1	C	UPX1, VPX1	D		Value of polynomial P(X) at X1
PX2	C	UPX2, VPX2	D		Value of polynomial P(X) at X2
PX3	C	UPX3, VPX3	D		Value of polynomial P(X) at X3
X4	C	UX4, VX4	D		Newest approximation (X _{n+1}) to root
PX4	C	UPX4, VPX4	D		Value of polynomial P(X) at X4
MULT	I	MULT	I		Array containing the multiplicities of each root found
ITER	I	ITER	I		Counter for iterations
I01	I	I01	I		Unit number of input device
I02	I	I02	I	C	Unit number of output device
EPSRT	R	EPSRT	D	C	Number used in subroutine BETTER to generate two approximations from the one given
NOPOLY	I	NOPOLY	I	E	Number of the polynomial

TABLE F.II (Continued)

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition of Argument</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>		
MAX	I	MAX	I	C	Maximum number of iterations
EPS	R	EPS	D	C	Tolerance check for convergence
EPSO	R	EPSO	D	C	Tolerance check for zero (0)
EPSM	R	EPSM	D	C	Tolerance check for multiplicities
KCHECK	I	KCHECK	I		Program control, KCHECK = 1 stops execution of program
XSTART	R	XSTART	D	E	Magnitude at which to start generating initial approximations
XEND	R	XEND	D	E	Magnitude at which to end generating initial approximations
NWORK	I	NWORK	I		Degree of current deflated polynomial whose coefficients are in WORK
ITIME	I	ITIME	I		Program control
NALTER	I	NALTER	I		Number of alterations which have been performed on an initial approximation
IAPP	I	IAPP	I		Counter for number of initial approximations used
CONV	L	CONV	L		When CONV is true, convergence has been obtained
IROOT	I	IROOT	I	R	Number of distinct roots solved by Muller's method, i.e. not solved directly by subroutine QUAD
Subroutine HORNER					
A	C	UA, VA	D	E	Array of current polynomial coefficients (to be deflated or evaluated)
NA	I	NA	I	E	Degree of polynomial to be deflated or evaluated
X	C	UX, VX	D	E	Approximation at which to evaluate or deflate the polynomial
B	C	UB, VB	D	R	Array containing the coefficients of the deflated polynomial
PX	C	UPX, VPX	D	R	Value of the polynomial at X
NUM	I	NUM	I		Number of coefficients of polynomial to be deflated

TABLE F.II (Continued)

Single Precision Variable	Double Precision Variable	Disposition of Argument	Description
Subroutine TEST			
X3	UX3, VX3	D	Approximation to root (old) (X_n)
X4	UX4, VX4	D	New approximation to root (X_{n+1})
CONV	CONV	L	CONV = True implies convergence has been obtained
EPS	EPS	D	Tolerance for convergence test
EPSO	EPSO	D	Tolerance check for zero (0)
DENOM	DENOM	D	Magnitude of new approximation, (X_{n+1})
Subroutine BETTER			
MULT	MULT	I	Array of multiplicities of each root
A	UA, VA	D	Array of coefficients of original undeflated polynomial
NP	NP	I	Degree of original polynomial
ROOT	UROOT, VROOT	D	Array of ROOTS
NROOT	NROOT	I	Number of roots stored in ROOT
BAPP	UBAPP, VBAPP	D	Array of initial approximations (old roots)
IROOT	IROOT	I	Number of roots solved by the iterative process (Not QUAD)
ROOTS	UROOTS, VROOTS	D	Temporary storage for new (better) roots
L	L	I	Number of roots found by BETTER
EPSRT	EPSRT	D	A small number used to generate two of the three approximations when given one
ITER	ITER	I	Counter for number of iterations
B	UB, VB	D	Array containing coefficients of deflated polynomial
X1	UX1, VX1	D	One of three approximations to the root
X2	UX2, VX2	D	One of three approximations to the root
X3	UX3, VX3	D	One of three approximations to the root
PX1	UPX1, VPX1	D	Value of polynomial (P(X)) at X1
PX2	UPX2, VPX2	D	Value of polynomial (P(X)) at X2
PX3	UPX3, VPX3	D	Value of polynomial (P(X)) at X3

TABLE F.II (Continued)

<u>Single Precision</u> <u>Variable</u>	<u>Type</u>	<u>Double Precision</u> <u>Variable</u>	<u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
CONV	L	CONV	L		CONV = true implies convergence has been obtained
X4	C	UX4, VX4	D		Newest approximation to root
J	I	J	I		Program control - counts the number of roots used as initial approximations
MAX	I	MAX	I	C	Maximum number of iterations permitted
I02	I	I02	I	C	Unit number of output device
Subroutine ALTER					
X1	C	X1R, X1I	D	ECR	One of the three approximations to be altered
X2	C	X2R, X2I	D	ECR	One of the three approximations to be altered
X3	C	X3R, X3I	D	ECR	One of the three approximations to be altered
X2R	R	X2R	D		Real part of complex approximation
X2I	R	X2I	D		Imaginary part of complex approximation
Subroutine CALC					

These variables are dummy variables used for temporary storage and thus, are not defined.

MAIN PROGRAM




Figure F.1. Flow Charts for G.C.D.-Muller's Method




Figure F.1. (Continued)

MULLER




Figure F.1. (Continued)





Figure F.1. (Continued)

DIVIDE



DERIV




Figure F.1. (Continued)




Figure F.1. (Continued)

QUAD




Figure F.1. (Continued)

BETTER

HORNER





Figure F.1. (Continued)

GENAPP



ALTER





Figure F.1. (Continued)

CALC



TEST




Figure F.1. (Continued)

COMSQT




Figure F.1. (Continued)

TABLE F.III

PROGRAM FOR G.C.D.-MULLER'S METHOD

```

C *****
C *
C * DOUBLE PRECISION PROGRAM FOR G.C.D. - MULLER'S METHOD
C *
C *
C * THE G.C.D. METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C * POLYNOMIAL OF MAXIMUM DEGREE 25. ALL MULTIPLE ROOTS ARE REMOVED BY
C * DIVIDING THE POLYNOMIAL BY THE GREATEST COMMON DIVISOR OF THE POLYNOMIAL
C * AND ITS DERIVATIVE. THE ZEROS OF THE RESULTING POLYNOMIAL ARE EXTRACTED
C * AND THEIR MULTIPLICITIES DETERMINED.
C *
C *****
0001 DOUBLE PRECISION URAPP,VRAPP
0002 DOUBLE PRECISION UP,VP,UAPP,VAPP,UROOT,VROOT,UDP,VDP,UD,VD,UZRO,VZ
0003 1RO,UQ,VQ,UDUMMY,VDDUMMY,UQQ,VQQ,UB,VB,EPS1,EPS2,EPS3,EPS4
0004 DIMENSION URAPP(25,3),VRAPP(25,3),UAPP(25,3),VAPP(25,3)
      DIMENSION UP(26),VP(26),UROOT(25),VROOT(25),MULT(25),UDP(26),VDP(2
16),UD(26),VD(26),UQ(26),VQ(26),UQQ(26),VQQ(26),UB(26),VB(26),ANAME
2(2),ENTRY(26)
0005 DOUBLE PRECISION XSTART
0006 DOUBLE PRECISION XEND
0007 DOUBLE PRECISION EPSRT
0008 COMMON EPSRT,EPS1,EPS2,EPS3,EPS4,IO2,MAX
0009 DATA PNAME,QNAME,QQNAME/2HP(,2HQ(,3HQQ(/
0010 DATA ENTRY/1H1,1H2,1H3,1H4,1H5,1H6,1H7,1H8,1H9,2H10,2H11,2H12,2H13
1,2H14,2H15,2H16,2H17,2H18,2H19,2H20,2H21,2H22,2H23,2H24,2H25,2H26/
0011 DATA ANAME(1),ANAME(2)/4HMULL,4HERS /
0012 LOGICAL NEWT
0013 IO1=5
0014 IO2=6
0015 10 J=0
0016 ITIME=0
0017 READ(IO1,1000) NOPOLY,NP,NAPP,MAX,EPS1,EPS2,EPS3,EPS4,XSTART,XEND,
1KCHECK
0018 IF(KCHECK.EQ.1) STOP
0019 WRITE(IO2,1020) ANAME(1),ANAME(2),NOPOLY
0020 WRITE(IO2,2000) NAPP
0021 WRITE(IO2,2010) MAX
0022 WRITE(IO2,2070) EPS1
0023 WRITE(IO2,2020) EPS2
0024 WRITE(IO2,2080) EPS3
0025 WRITE(IO2,2030) EPS4
0026 WRITE(IO2,2040) XSTART
0027 WRITE(IO2,2050) XEND
0028 WRITE(IO2,2060)
0029 KKK=NP+1
0030 NNN=KKK+1
0031 DO 20 I=1,KKK
0032 JJJ=NNN-I
0033 20 READ(IO1,1010) UP(JJJ),VP(JJJ)
0034 IF(NAPP.NE.0) GO TO 22
0035 NAPP=NP
0036 CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0037 GO TO 23
0038 22 READ(IO1,1015) (UAPP(I,2),VAPP(I,2),I=1,NAPP)
0039 23 WRITE(IO2,1030) NP
0040 KKK=NP+1
0041 NNN=KKK+1

```

TABLE F.III (Continued)

```

0042      DO 25 I=1,KKK
0043      JJJ=NNN-1
0044      25 WRITE(102,1040) PNAME,ENTRY(JJJ),UP(JJJ),VP(JJJ)
0045      IF(NP.GE.3) GO TO 30
0046      J=-1
0047      CALL QUAD(NP,UP,VP,J,UROOT,VROOT,MULT)
0048      WRITE(102,1070)
0049      WRITE(102,1165) (I,UROOT(I),VROOT(I),MULT(I),I=1,J)
0050      GO TO 10
0051      30 CALL DERIV(NP,UP,VP,NDP,UDP,VDP)
0052      CALL GCD(NP,UP,VP,NDP,UDP,VDP,ND,UD,VD)
0053      IF(ND.GT.1) GO TO 70
0054      IF(ND.EQ.0) GO TO 65
0055      UDUMMY=UD(2)*UD(2)+VD(2)*VD(2)
0056      UZRO=-(UD(1)*UD(2)+VD(1)*VD(2))/UDUMMY
0057      VZRO=-(UD(2)*VD(1)-UD(1)*VD(2))/UDUMMY
0058      KKK=NP+1
0059      DO 55 I=1,KKK
0060      UQQ(I)=UP(KKK+1-I)
0061      55 VQQ(I)=VP(KKK+1-I)
0062      NQQ=NP
0063      CALL HORNER(NQQ,UQQ,VQQ,UZRO,VZRO,UB,VB,UDUMMY,VDUMMY)
0064      NQ=NP-1
0065      DO 60 I=1,NP
0066      UQ(I)=UB(NP+1-I)
0067      60 VQ(I)=VB(NP+1-I)
0068      GO TO 80
0069      65 KKK=NP+1
0070      DO 66 I=1,KKK
0071      UQ(I)=UP(I)
0072      66 VQ(I)=VP(I)
0073      NQ=NP
0074      GO TO 80
0075      70 CALL DIVIDE(NP,UP,VP,ND,UD,VD,NQ,UQ,VQ)
0076      80 WRITE(102,1120) NQ
0077      KKK=NQ+1
0078      NNN=KKK+1
0079      DO 83 I=1,KKK
0080      JJJ=NNN-I
0081      83 WRITE(102,1040) QNAME,ENTRY(JJJ),UQ(JJJ),VQ(JJJ)
0082      IF(NQ.GE.3) GO TO 85
0083      GO TO 110
0084      85 KKK=NQ+1
0085      DO 90 I=1,KKK
0086      UQQ(I)=UQ(KKK+1-I)
0087      90 VQQ(I)=VQ(KKK+1-I)
0088      NQQ=NQ
0089      GO TO 120
0090      110 CALL QUAD(NQ,UQ,VQ,J,UROOT,VROOT,MULT)
0091      NEWT=.FALSE.
0092      GO TO 310
0093      120 CALL MULLER(UQQ,VQQ,NQQ,UAPP,VAPP,NAPP,XSTART,XEND,UROOT,VROOT,J,J
0094      IAP,URAPP,VRAPP,NOPOLY)
0095      NEWT=.TRUE.
0096      310 CALL MULTI(NP,UP,VP,J,UROOT,VROOT,MULT)
0097      IF(NEWT) GO TO 330
0098      WRITE(102,1070)
0099      WRITE(102,1165) (L,UROOT(L),VROOT(L),MULT(L),L=1,J)

```

TABLE F.III (Continued)

```

0099      GO TO 10
0100      330 WRITE(102,1180)
0101      DO 350 L=1,JAP
0102      350 WRITE(102,1190) L,URROOT(L),VROOT(L),MULT(L),URAPP(L,2),VRAPP(L,2)
0103      KKK=JAP+1
0104      IF(JAP.LT.J) WRITE(102,1165) (L,URROOT(L),VROOT(L),MULT(L),L=KKK,J)
0105      GO TO 10
0106      1000 FORMAT(3(I2,1X),9X,I3,1X,4(D6.0,1X),13X,2(D7.0,1X),I1)
0107      1010 FORMAT(2D30.0)
0108      1015 FORMAT(2D30.0)
0109      1020 FORMAT(1H1,10X,41HGREATEST COMMON DIVISOR METHOD USED WITH ,2(A4),
        135HMETHOD TO FIND ZEROS OF POLYNOMIALS/11X,18HPOLYNOMIAL NUMBER ,I
        22//)
0110      1030 FORMAT(1X,22HTHE DEGREE OF P(X) IS ,I2,22H THE COEFFICIENTS ARE//
        1)
0111      1040 FORMAT(2X,A2,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0112      1070 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES//)
0113      1080 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,10X,I2)
0114      1100 FORMAT(2X,A3,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0115      1120 FORMAT(///1X,73HQ(X) IS THE POLYNOMIAL WHICH HAS AS ITS ROOTS THE
        1DISTINCT ROOTS OF P(X)./1X,22HTHE DEGREE OF Q(X) IS ,I2,22H THE C
        20EFFICIENTS ARE//)
0116      1165 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,10X,26H
        1RESULTS OF SUBROUTINE QUAD)
0117      1180 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES,17X,21HINITIAL
        1 APPROXIMATION//)
0118      1190 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,9X,D23.
        116,3H + ,D23.16,2H I)
0119      2000 FORMAT(1X,41HNUMBER OF INITIAL APPROXIMATIONS GIVEN. ,I2)
0120      2010 FORMAT(1X,29HMAXIMUM NUMBER OF ITERATIONS.,11X,I3)
0121      2020 FORMAT(1X,21HTEST FOR CONVERGENCE.,13X,D9.2)
0122      2030 FORMAT(1X,24HTEST FOR MULTIPLICITIES.,10X,D9.2)
0123      2040 FORMAT(1X,23HRAIUS TO START SEARCH.,11X,D9.2)
0124      2050 FORMAT(1X,21HRAIUS TO END SEARCH.,13X,D9.2)
0125      2060 FORMAT(///1X)
0126      2070 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE GCD. ,D9.2)
0127      2080 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE QUAD. ,D9.2)
0128      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE MULTI(N,UP,VP,J,UROOT,VROOT,MULT)
C      *
C      *
C      * GIVEN N ZEROS OF A POLYNOMIAL, SUBROUTINE MULTI COMPUTES THEIR
C      * MULTIPLICITIES.
C      *
C      *
0002      DOUBLE PRECISION UP,VP,UROOT,VROOT,UA,VA,UB,VB,UC,VC,EPS1,EPS2,EPS
      ILON,EPS3,BBB
0003      DIMENSION UP(26),VP(26),UROOT(25),VROOT(25),UA(26),VA(26),UB(26),V
      IB(26),MULT(25)
0004      DOUBLE PRECISION EPSRT
0005      COMMON EPSRT,EPS1,EPS2,EPS3,EPSLON,IO2,MAX
0006      DO 100 I=1,J
0007          KKK=N+1
0008          DO 10 K=1,KKK
0009              UA(K)=UP(KKK+1-K)
0010          10 VA(K)=VP(KKK+1-K)
0011              M=N
0012              MULT(I)=0
0013          20 CALL HORNER(M,UA,VA,UROOT(I),VROOT(I),UB,VB,UC,VC)
0014              BBB=DSQRT(UC*UC+VC*VC)
0015              IF(BBB.LT.EPSLON) GO TO 50
0016              IF(MULT(I).EQ.0) GO TO 40
0017              GO TO 100
0018          40 WRITE(IO2,1000) EPSLON,I,UROOT(I),VROOT(I)
0019              GO TO 100
0020          50 MULT(I)=MULT(I)+1
0021              IF(M.GT.1) GO TO 60
0022              GO TO 100
0023          60 DO 70 K=1,M
0024              UA(K)=UB(K)
0025          70 VA(K)=VB(K)
0026              M=M-1
0027              GO TO 20
0028          100 CONTINUE
0029          RETURN
0030          1000 FORMAT(///15H THE EPSILON (,D10.3,48H) CHECK IN SUBROUTINE MULTI
      1INDICATES THAT ROOT(I,I2,4H) = ,D23.16,3H + ,D23.16,2H I,/80H IS NO
      2T CLOSE ENOUGH TO BE A TRUE ROOT. IT IS PRINTED BELOW WITH MULTIP
      3LICITY 0//)
0031      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE DIVIDE(N,UP,VP,M,UD,VD,K,UQ,VQ)
C      *****
C      *
C      * GIVEN TWO POLYNOMIALS F(X) AND G(X), SUBROUTINE DIVIDE COMPUTES THE
C      * QUOTIENT POLYNOMIAL H(X) = F(X)/G(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UD,VD,UQ,VQ,UTERM,VTERM,UDUMMY
0003      DIMENSION UP(26),VP(26),UD(26),VD(26),UQ(26),VQ(26)
0004      K=N-M
0005      UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0006      UQ(K+1)=(UP(N+1)*UD(M+1)+VP(N+1)*VD(M+1))/UDUMMY
0007      VQ(K+1)=(VP(N+1)*UD(M+1)-UP(N+1)*VD(M+1))/UDUMMY
0008      IF(K.EQ.0) GO TO 100
0009      J=-1
0010      DO 50 I=1,K
0011      J=J+1
0012      UTERM=UP(N-J)
0013      VTERM=VP(N-J)
0014      KK=K+1
0015      NNN=M-J
0016      DO 40 M1=NNN,M
0017      IF(KK.GT.1) GO TO 10
0018      GO TO 45
0019      10 IF(M1.GE.1) GO TO 20
0020      GO TO 40
0021      20 UTERM=UTERM-(UQ(KK)*UD(M1)-VQ(KK)*VD(M1))
0022      VTERM=VTERM-(UQ(KK)*VD(M1)+VQ(KK)*UD(M1))
0023      40 KK=KK-1
0024      UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0025      UQ(K+1-I)=(UTERM*UD(M+1)+VTERM*VD(M+1))/UDUMMY
0026      50 VQ(K+1-I)=(VTERM*UD(M+1)-UTERM*VD(M+1))/UDUMMY
0027      100 RETURN
0028      END

0001      SUBROUTINE DERIV(N,UP,VP,M,UA,VA)
C      *****
C      *
C      * GIVEN A POLYNOMIAL P(X), SUBROUTINE DERIV COMPUTES THE COEFFICIENTS OF
C      * ITS DERIVATIVE P'(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UA,VA,AAA
0003      DIMENSION UP(26),VP(26),UA(26),VA(26)
0004      KKK=N+1
0005      DO 10 I=2,KKK
0006      AAA=I-1
0007      UA(I-1)=AAA*UP(I)
0008      10 VA(I-1)=AAA*VP(I)
0009      M=N-1
0010      RETURN
0011      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE GCD(N,UR,VR,M,US,VS,M1,USS,VSS)
C      *****
C      *
C      * GIVEN POLYNOMIALS P(X) AND DP(X) WHERE DEG. DP(X) IS LESS THAN DEG.
C      * P(X), SUBROUTINE GCD COMPUTES THE GREATEST COMMON DIVISOR OF P(X) AND
C      * DP(X).
C      *
C      *****
0002      DOUBLE PRECISION EPSRT
0003      DOUBLE PRECISION USSSSS,VSSSSS
0004      DOUBLE PRECISION UR,VR,US,VS,USS,VSS,URR,VRR,UD,VD,UT,VT,EPSLON,EP
0005      IS2,EPS3,EPS4,BBB
0006      DIMENSION UR(26),VR(26),US(26),VS(26),USS(26),VSS(26),URR(26),VRR(
0007      26),UT(26),VT(26)
0008      COMMON EPSRT,EPSLON,EPS2,EPS3,EPS4,IO2,MAX
0009      N1=N
0010      M1=M
0011      KKK=N+1
0012      DO 20 I=1,KKK
0013      URR(I)=UR(I)
0014      VRR(I)=VR(I)
0015      KKK=M+1
0016      DO 25 I=1,KKK
0017      USS(I)=US(I)
0018      VSS(I)=VS(I)
0019      25 BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0020      UD=(URR(N1+1)*USS(M1+1)+VRR(N1+1)*VSS(M1+1))/BBB
0021      VD=(USS(M1+1)*VRR(N1+1)-URR(N1+1)*VSS(M1+1))/BBB
0022      KKK=N1+1-M1
0023      DO 40 I=KKK,N1
0024      UT(I)=URR(I)-(UD*USS(I-N1+M1)-VD*VSS(I-N1+M1))
0025      VT(I)=VRR(I)-(UD*VSS(I-N1+M1)+VD*USS(I-N1+M1))
0026      IF(M1.EQ.N1) GO TO 70
0027      KKK=N1-M1
0028      DO 60 I=1,KKK
0029      UT(I)=URR(I)
0030      VT(I)=VRR(I)
0031      DO 70 I=1,N1
0032      BBB=DSQRT(UT(N1+1-I)*UT(N1+1-I)+VT(N1+1-I)*VT(N1+1-I))
0033      IF(BBB.GT.EPSLON) GO TO 100
0034      90 CONTINUE
0035      DO 95 I=1,M1
0036      BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0037      USSSSS=(USS(I)*USS(M1+1)+VSS(I)*VSS(M1+1))/BBB
0038      VSSSSS=(VSS(I)*USS(M1+1)-USS(I)*VSS(M1+1))/BBB
0039      USS(I)=USSSSS
0040      VSS(I)=VSSSSS
0041      USS(M1+1)=1.0
0042      VSS(M1+1)=0.0
0043      GO TO 200
0044      100 K=N1-I
0045      IF(K.EQ.0) GO TO 170
0046      IF(K.LT.M1) GO TO 140
0047      KKK=K+1
0048      DO 130 J=1,KKK
0049      URR(J)=UT(J)
0050      VRR(J)=VT(J)
0051      N1=K

```

TABLE F.III (Continued)

```

0050      GO TO 30
0051 140 KKK=K+1
0052      DO 150 J=1, KKK
0053          URR(J)=USS(J)
0054          VRR(J)=VSS(J)
0055          USS(J)=UT(J)
0056 150 VSS(J)=VT(J)
0057      KKK=K+2
0058      NNN=M1+1
0059      DO 160 J=KKK, NNN
0060          URR(J)=USS(J)
0061 160 VRR(J)=VSS(J)
0062      M1=M1
0063      M1=K
0064      GO TO 30
0065 170 USS(1)=1.0
0066      VSS(1)=0.0
0067      M1=0
0068 200 RETURN
0069      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE QUAD(N,UA,VA,J,UROOT,VROOT,MULT)
C      *****
C      *
C      * SUBROUTINE QUAD SOLVES DIRECTLY FOR THE ZEROS AND THEIR MULTIPLICITIES *
C      * OF EITHER A QUADRATIC POLYNOMIAL OR A LINEAR FACTOR. SOLUTION OF THE *
C      * QUADRATIC IS DONE USING THE QUADRATIC FORMULA. *
C      *
C      *****
0002      DOUBLE PRECISION EPSRT
0003      DOUBLE PRECISION UA,VA,UROOT,VROOT,UDISC,VDISC,UTEMP,VTEMP,UD,VD,E
0004      EPS1,EPS2,EPS4,EPSLON,BBB
0005      DIMENSION UA(26),VA(26),UROOT(25),VROOT(25),MULT(25)
0006      COMMON EPSRT,EPS1,EPS2,EPSLON,EPS4,I02,MAX
0007      IF(N.GT.1) GO TO 60
0008      IF(J.LT.0) GO TO 40
0009      J=J+1
0010      GO TO 50
0011      40 MULT(1)=1
0012      J=1
0013      50 BBB=UA(2)*UA(2)+VA(2)*VA(2)
0014      UROOT(J)=-(UA(1)*UA(2)+VA(1)*VA(2))/BBB
0015      VROOT(J)=-(VA(1)*UA(2)-UA(1)*VA(2))/BBB
0016      GO TO 200
0017      60 UDISC=(UA(2)*UA(2)-VA(2)*VA(2))-(4.0*(UA(3)*UA(1)-VA(3)*VA(1)))
0018      VDISC=(2.0*UA(2)*VA(2))-(4.0*(UA(3)*VA(1)+VA(3)*UA(1)))
0019      BBB=DSQRT(UDISC*UDISC+VDISC*VDISC)
0020      IF(BBB.LE.EPSLON) GO TO 100
0021      IF(J.GE.0) GO TO 80
0022      MULT(1)=1
0023      MULT(2)=1
0024      J=0
0025      80 CALL COMSQT(UDISC,VDISC,UTEMP,VTEMP)
0026      UD=2.0*UA(3)
0027      VD=2.0*VA(3)
0028      BBB=UD*UD+VD*VD
0029      UROOT(J+1)=((-UA(2)+UTEMP)*UD+(-VA(2)+VTEMP)*VD)/BBB
0030      VROOT(J+1)=((-VA(2)+VTEMP)*UD-(-UA(2)+UTEMP)*VD)/BBB
0031      UROOT(J+2)=((-UA(2)-UTEMP)*UD+(-VA(2)-VTEMP)*VD)/BBB
0032      VROOT(J+2)=((-VA(2)-VTEMP)*UD-(-UA(2)-UTEMP)*VD)/BBB
0033      J=J+2
0034      GO TO 200
0035      100 IF(J.LT.0) GO TO 110
0036      J=J+1
0037      GO TO 130
0038      110 MULT(1)=2
0039      J=1
0040      130 UD=2.0*UA(3)
0041      VD=2.0*VA(3)
0042      BBB=UD*UD+VD*VD
0043      UROOT(J)=(-UA(2)*UD-VA(2)*VD)/BBB
0044      VROOT(J)=(-VA(2)*UD+UA(2)*VD)/BBB
0045      200 RETURN
      END

```


TABLE F.III (Continued)

```

0001      SUBROUTINE MULLER(UA,VA,NP,UAPP,VAPP,NAPP,XSTART,XEND,UROOT,VROOT,
          INROOT,IROOT,URAPP,VAPP,NOPOLY)
C      *
C      * *****
C      * MULLER'S METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C      * POLYNOMIAL OF MAXIMUM DEGREE 25. THROUGH THREE GIVEN POINTS THE
C      * POLYNOMIAL IS APPROXIMATED BY A QUADRATIC. THE ZERO OF THE QUADRATIC
C      * CLOSEST TO THE OLD APPROXIMATION IS TAKEN AS THE NEW APPROXIMATION.
C      * IN THIS MANNER A SEQUENCE IS OBTAINED CONVERGING TO A ZERO.
C      *
C      * *****
0002      DOUBLE PRECISION UPX3,VPX3,UPX2,VPX2,UROOT,VROOT,UX1,VX1,UAPP,VAPP
          1,UX2,VX2,UWORK,VWORK,UX3,VX3,UB,VB,UX4,VX4,UA,VA,UPX1,VPX1,URAPP,V
          2RAPP,UPX4,VPX4,EPSRT,EPS0,EPS,CCC,EPSP,UMH3,VH3,UQ4,VQ4,ABPX4,ABPX3
          3,QQQ,XSTART,XEND
0003      DIMENSION UROOT(25),VROOT(25),MULT(25),UAPP(25,3),VAPP(25,3),UWORK
          1(26),VWORK(26),UB(26),VB(26),UA(26),VA(26),URAPP(25,3),VRAPP(25,3)
0004      LOGICAL CONV
0005      DOUBLE PRECISION EPS1
0006      COMMON EPSRT,EPS1,EPS,EPSP,EPSP,IO2,MAX
0007      DATA PNAME,DNAME/2HP/,2HD/
0008      EPSRT=0.999
0009      NROOT=0
0010      IROOT=0
0011      IPATH=1
0012      NMULT=0
0013      NALTER=0
0014      ITIME=0
0015      IAPP=1
0016      ITER=1
0017      IF(NAPP.NE.0) GO TO 18
0018      NAPP=NP
0019      CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0020      GO TO 27
0021      18 DO 25 I=1,NAPP
0022          UAPP(I,1)=0.9*UAPP(I,2)
0023          VAPP(I,1)=0.9*VAPP(I,2)
0024          UAPP(I,3)=1.1*UAPP(I,2)
0025          VAPP(I,3)=1.1*VAPP(I,2)
0026      27 KKK=NP+1
0027          DO 30 I=1,KKK
0028              UWORK(I)=UA(I)
0029              VWORK(I)=VA(I)
0030          NWORK=NP
0031      40 UX1=UAPP(IAPP,1)
0032          VX1=VAPP(IAPP,1)
0033          UX2=UAPP(IAPP,2)
0034          VX2=VAPP(IAPP,2)
0035          UX3=UAPP(IAPP,3)
0036          VX3=VAPP(IAPP,3)
0037          CALL HORNER(NWORK,UWORK,VWORK,UX1,VX1,UB,VB,UPX1,VPX1)
0038          CALL HORNER(NWORK,UWORK,VWORK,UX2,VX2,UB,VB,UPX2,VPX2)
0039          CALL HORNER(NWORK,UWORK,VWORK,UX3,VX3,UB,VB,UPX3,VPX3)
0040      50 CALL CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UX
          14,VX4,UQ4,VQ4,UMH3,VH3)
0041      60 CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0042          ABPX4=DSQRT(UPX4*UPX4+VPX4*VPX4)
0043          ABPX3=DSQRT(UPX3*UPX3+VPX3*VPX3)

```

TABLE F.III (Continued)

```

0044      IF(ABPX3.EQ.0.0) GO TO 70
0045      QQQ=ABPX4/ABPX3
0046      IF(QQQ.LE.10.) GO TO 70
0047      UQ4=0.5*UQ4
0048      VQ4=0.5*VQ4
0049      UX4=UX3+(UH3*UQ4-VH3*VQ4)
0050      VX4=VX3+(VH3*UQ4+UH3*VQ4)
0051      GO TO 60
0052 70 CALL TEST(UX3,VX3,UX4,VX4,CONV)
0053      IF(CONV) GO TO 120
0054      IF(ITER.LT.MAX) GO TO 110
0055      CALL ALTER(UAPP(IAPP,1),VAPP(IAPP,1),UAPP(IAPP,2),VAPP(IAPP,2),UAP
1P(IAPP,3),VAPP(IAPP,3),NALTER,ITIME)
0056      IF(NALTER.GT.5) GO TO 75
0057      ITER=1
0058      GO TO 40
0059 75 IF(IAPP.LT.NAPP) GO TO 100
0060      IF(XEND.EQ.0.0) GO TO 77
0061      IF(XSTART.GT.XEND) GO TO 77
0062      NAPP=NP
0063      CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0064      IAPP=0
0065      GO TO 100
0066 77 WRITE(IO2,1090)
0067      KKK=NWORK+1
0068      WRITE(IO2,1035) (DNAME,J,UWORK(J),VWORK(J),J=1,KKK)
0069 80 IF(NROOT.EQ.0) GO TO 90
0070      IF(IPATH.EQ.1) GO TO 82
0071 81 IPATH=2
0072      CALL BETTER(UA,VA,NP,UROOT,VROOT,NROOT,URAPP,VRAPP,IROOT,MULT)
0073      RETURN
0074 82 IF(NROOT.EQ.0) GO TO 90
0075      IF(IROOT.EQ.0) GO TO 85
0076      WRITE(IO2,1080)
0077      DO 55 I=1,IROOT
0078 55 WRITE(IO2,1085) I,UROOT(I),VROOT(I),URAPP(I,2),VRAPP(I,2)
0079      IF(IROOT.LT.NROOT) GO TO 85
0080      GO TO 87
0081 85 KKK=IROOT+1
0082      WRITE(IO2,1086) (I,UROOT(I),VROOT(I),I=KKK,NROOT)
0083 87 IF(IPATH.EQ.1) GO TO 81
0084      RETURN
0085 90 WRITE(IO2,1070) NOPOLY
0086      RETURN
0087 100 IAPP=IAPP+1
0088      ITER=1
0089      NALTER=0
0090      GO TO 40
0091 120 NROOT=NROOT+1
0092      IROOT=NROOT
0093      MULT(NROOT)=1
0094      NOMULT=NOMULT+1
0095      UROOT(NROOT)=UX4
0096      VROOT(NROOT)=VX4
0097      URAPP(NROOT,1)=UAPP(IAPP,1)
0098      VRAPP(NROOT,1)=VAPP(IAPP,1)
0099      URAPP(NROOT,2)=UAPP(IAPP,2)
0100      VRAPP(NROOT,2)=VAPP(IAPP,2)

```

TABLE F.III (Continued)

```

0101      URAPP(NROOT,3)=UAPP(IAPP,3)
0102      VRAPP(NROOT,3)=VAPP(IAPP,3)
0103      125 IF(NOMULT.LT.NP) GO TO 130
0104      GO TO 80
0105      130 CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0106      NWORK=NWORK-1
0107      KKK=NWORK+1
0108      DO 140 I=1,KKK
0109      UWORK(I)=UB(I)
0110      140 VWORK(I)=VB(I)
0111      CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0112      CCC=DSQRT(UPX4*UPX4+VPX4*VPX4)
0113      IF(CCC.LT.EPSM) GO TO 150
0114      IF(NWORK.GT.2) GO TO 75
0115      IROOT=NROOT
0116      KKK=NWORK+1
0117      DO 145 I=1,KKK
0118      UB(I)=UWORK(KKK+1-I)
0119      145 VB(I)=VWORK(KKK+1-I)
0120      CALL QUAD(NWORK,UB,VB,NROOT,UROOT,VROOT,MULT)
0121      GO TO 80
0122      150 MULT(NROOT)=MULT(NROOT)+1
0123      NOMULT=NOMULT+1
0124      GO TO 125
0125      110 UX1=UX2
0126      VX1=VX2
0127      UX2=UX3
0128      VX2=VX3
0129      UX3=UX4
0130      VX3=VX4
0131      UPX1=UPX2
0132      VPX1=VPX2
0133      UPX2=UPX3
0134      VPX2=VPX3
0135      UPX3=UPX4
0136      VPX3=VPX4
0137      ITER=ITER+1
0138      GO TO 50
0139      1090 FORMAT(///,1X,65HCOEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO
        1ZEROS WERE FOUND//)
0140      1080 FORMAT(///1X,13HROOTS OF Q(X),83X,21HINITIAL APPROXIMATION//)
0141      1070 FORMAT(//,43H NO ZEROS WERE FOUND FOR POLYNOMIAL NUMBER ,I2)
0142      1086 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,19X,23HSOLVED
        1 BY DIRECT METHOD)
0143      1035 FORMAT(3X,A2,I2,4H) = ,D23.16,3H + ,D23.16,2H I)
0144      1050 FORMAT(82X,D23.16,3H + ,D23.16,2H I/82X,D23.16,3H + ,D23.16,2H I/)
0145      1085 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,18X,D23.16,3H
        1 + ,D23.16,2H I)
0146      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE BETTER(UA,VA,NP,UROOT,VROOT,NROOT,URAPP,VRAPP,IROOT,MUL
          IT)
C      *****
C      *
C      * SUBROUTINE BETTER ATTEMPTS TO IMPROVE THE ACCURACY OF THE ZEROS FOUND *
C      * BY USING THEM AS INITIAL APPROXIMATIONS WITH MULLER'S METHOD APPLIED TO *
C      * THE FULL, UNDEFLATED POLYNOMIAL. *
C      *
C      *****
0002      DOUBLE PRECISION UROOT,VROOT,UA,VA,UBAPP,VBAPP,UX1,VX1,UX2,VX2,UX3
          1,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UB,VB,UROOTS,VROOTS,EPSRT,UX4,V
          2X4,URAPP,VRAPP,EPSO,EPS,UQ4,VQ4,UH3,VH3
0003      LOGICAL CONV
0004      DIMENSION UROOT(25),VROOT(25),UA(26),VA(26),UBAPP(25,3),VBAPP(25,3
          1),UB(26),VB(26),UROOTS(25),VROOTS(25),URAPP(25,3),VRAPP(25,3),MULT
          3(25)
0005      DOUBLE PRECISION EPS1,EPSM
0006      COMMON EPSRT,EPS1,EPS,EPSO,EPSM,I02,MAX
0007      IF(NROOT.LE.1) RETURN
0008      L=0
0009      DO 10 I=1,NROOT
0010          UBAPP(I,1)=UROOT(I)*EPSRT
0011          VBAPP(I,1)=VROOT(I)*EPSRT
0012          UBAPP(I,2)=UROOT(I)
0013          VBAPP(I,2)=VROOT(I)
0014          UBAPP(I,3)=UROOT(I)*(2.0-EPSRT)
0015      10 VBAPP(I,3)=VROOT(I)*(2.0-EPSRT)
0016      DO 100 J=1,NROOT
0017          UX1=UBAPP(J,1)
0018          VX1=VBAPP(J,1)
0019          UX2=UBAPP(J,2)
0020          VX2=VBAPP(J,2)
0021          UX3=UBAPP(J,3)
0022          VX3=VBAPP(J,3)
0023          ITER=1
0024          CALL HORNER(NP,UA,VA,UX1,VX1,UB,VB,UPX1,VPX1)
0025          CALL HORNER(NP,UA,VA,UX2,VX2,UB,VB,UPX2,VPX2)
0026      20 CALL HORNER(NP,UA,VA,UX3,VX3,UB,VB,UPX3,VPX3)
0027          CALL CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UX
          14,VX4,UQ4,VQ4,UH3,VH3)
0028      30 CALL TEST(UX3,VX3,UX4,VX4,CONV)
0029          IF(CONV) GO TO 50
0030          IF(ITER.LT.MAX) GO TO 40
0031          WRITE(I02,1000) J,UROOT(J),VROOT(J),MAX
0032          WRITE(I02,1010) UX4,VX4
0033          IF(J.LT.IROOT) GO TO 33
0034          IF(J.EQ.IROOT) GO TO 35
0035          GO TO 100
0036      33 KKK=IROOT-1
0037          DO 34 K=J,KKK
0038              URAPP(K,1)=URAPP(K+1,1)
0039              VRAPP(K,1)=VRAPP(K+1,1)
0040              URAPP(K,2)=URAPP(K+1,2)
0041              VRAPP(K,2)=VRAPP(K+1,2)
0042              URAPP(K,3)=URAPP(K+1,3)
0043      34 VRAPP(K,3)=VRAPP(K+1,3)
0044      35 IROOT=IROOT-1
0045          GO TO 100

```

TABLE F.III (Continued)

```

0046      40 UX1=UX2
0047      VX1=VX2
0048      UX2=UX3
0049      VX2=VX3
0050      UX3=UX4
0051      VX3=VX4
0052      UPX1=UPX2
0053      VPX1=VPX2
0054      UPX2=UPX3
0055      VPX2=VPX3
0056      ITER=ITER+1
0057      GO TO 20
0058
0059      50 L=L+1
0060      UROOTS(L)=UX4
0061      VROOTS(L)=VX4
0062      100 CONTINUE
0063      IF(L.EQ.0) GO TO 120
0064      DO 110 I=1,L
0065      UROOT(I)=UROOTS(I)
0066      VROOT(I)=VROOTS(I)
0067      NROOT=L
0068      RETURN
0069      120 NROOT=0
0070      RETURN
0071      1000 FORMAT(///42H IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(I,12,4H) = ,
0072      1D23.16,3H + ,D23.16,2H I/24H DID NOT CONVERGE AFTER ,I3,11H ITERAT
0073      2IONS)
0074      1010 FORMAT(30H THE PRESENT APPROXIMATION IS ,D23.16,3H + ,D23.16,2H I/
0075      1/)
0076      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE ALTER(X1R,X1I,X2R,X2I,X3R,X3I,NALTER,ITIME)
C      *****
C      *
C      * SUBROUTINE ALTER ALTERS THE INITIAL APPROXIMATIONS WHICH PRODUCE NO
C      * CONVERGENCE TO A ZERO. THIS IS DONE A MAXIMUM OF 5 TIMES FOR EACH ROOT.
C      *
C      *****
0002      DOUBLE PRECISION X1R,X1I,X2R,X2I,X3R,X3I,EPS1,EPS2,EPS3,R,BETA
0003      DOUBLE PRECISION EPS4,EPS5
0004      COMMON EPS1,EPS2,EPS3,EPS4,EPS5,IO2,MAX
0005      IF(ITIME.NE.0) GO TO 5
0006      ITIME=1
0007      WRITE(IO2,1010) MAX
0008      5 IF(NALTER.EQ.0) GO TO 10
0009      WRITE(IO2,1000) X1R,X1I,X2R,X2I,X3R,X3I
0010      GO TO 20
0011      10 R=DSQRT(X2R*X2R+X2I*X2I)
0012      BETA=DATAN2(X2I,X2R)
0013      WRITE(IO2,1020) X1R,X1I,X2R,X2I,X3R,X3I
0014      20 NALTER=NALTER+1
0015      IF(NALTER.GT.5) RETURN
0016      GO TO (30,40,30,40,30),NALTER
0017      30 X2R=-X2R
0018      X2I=-X2I
0019      GO TO 50
0020      40 BETA=BETA+1.0471976
0021      X2R=R*DCOS(BETA)
0022      X2I=R*DSIN(BETA)
0023      50 X1R=0.9*X2R
0024      X1I=0.9*X2I
0025      X3R=1.1*X2R
0026      X3I=1.1*X2I
0027      RETURN
0028      1000 FORMAT(1X,5HX1 = ,D23.16,3H + ,D23.16,2H I,10X,22HALTERED APPROXIM
1ATIONS/1X,5HX2 = ,D23.16,3H + ,D23.16,2H I/1X,5HX3 = ,D23.16,3H +
2,D23.16,2H I/)
0029      1020 FORMAT(1H0,5HX1 = ,D23.16,3H + ,D23.16,2H I,10X,22INITIAL APPROXI
1MATIONS/1X,5HX2 = ,D23.16,3H + ,D23.16,2H I/1X,5HX3 = ,D23.16,3H +
2 ,D23.16,2H I/)
0030      1010 FORMAT(///1X,54HNO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AF
1TER ,I3,12H ITERATIONS.//)
0031      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE GENAPP(APPR,APPI,NAPP,XSTART)
C      *****
C      *
C      * SUBROUTINE GENAPP GENERATES N INITIAL APPROXIMATIONS, WHERE N IS THE *
C      * DEGREE OF THE ORIGINAL POLYNOMIAL. *
C      *
C      *****
0002      DOUBLE PRECISION APPR,APPI,XSTART,EPS1,EPS2,EPS3,BETA
0003      DOUBLE PRECISION EPSRT,EPS4
0004      DIMENSION APPR(25,3),APPI(25,3)
0005      COMMON EPSRT,EPS1,EPS2,EPS3,EPS4,IQ2,MAX
0006      IF(XSTART.EQ.0.0) XSTART=0.5
0007      BETA=0.2617994
0008      DO 10 I=1,NAPP
0009      APPR(I,2)=XSTART*DCOS(BETA)
0010      APPI(I,2)=XSTART*DSIN(BETA)
0011      BETA=BETA+0.5235988
0012      10 XSTART=XSTART+0.5
0013      DO 20 I=1,NAPP
0014      APPR(I,1)=0.9*APPR(I,2)
0015      APPI(I,1)=0.9*APPI(I,2)
0016      APPR(I,3)=1.1*APPR(I,2)
0017      20 APPI(I,3)=1.1*APPI(I,2)
0018      RETURN
0019      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE TEST(UX3,VX3,UX4,VX4,CONV)
C      *****
C      *
C      * SUBROUTINE TEST CHECKS FOR CONVERGENCE OF THE SEQUENCE OF APPROX-
C      * IMATIONS BY TESTING THE EXPRESSION
C      * ABSOLUTE VALUE OF (X(N+1)-X(N))/ABSOLUTE VALUE OF X(N+1).
C      * WHEN IT IS AS SMALL AS DESIRED, CONVERGENCE IS OBTAINED.
C      *
C      *****
0002      DOUBLE PRECISION UX3,VX3,UX4,VX4,EPSRT,EPS0,EPS,AAA,UDUMMY,VDUMMY,
0003      IDENOM
0004      LOGICAL CONV
0005      DOUBLE PRECISION EPS1,EPSM
0006      COMMON EPSRT,EPS1,EPS,EPS0,EPSM,IO2,MAX
0007      UDUMMY=UX4-UX3
0008      VDUMMY=VX4-VX3
0009      AAA=DSQRT(UDUMMY*UDUMMY+VDUMMY*VDUMMY)
0010      DENOM=DSQRT(UX4*UX4+VX4*VX4)
0011      IF(DENOM.LT.EPS0) GO TO 20
0012      IF(AAA/DENOM.LT.EPS) GO TO 10
0013      5 CONV=.FALSE.
0014      GO TO 100
0015      10 CONV=.TRUE.
0016      GO TO 100
0017      20 IF(AAA.LT.EPS0) GO TO 10
0018      GO TO 5
0019      100 RETURN
      END

0001      SUBROUTINE HORNER(NA,UA,VA,UX,VX,UB,VB,UPX,VPX)
C      *****
C      *
C      * HORNER'S METHOD COMPUTES THE VALUE OF THE POLYNOMIAL P(X) AT A POINT D.
C      * SYNTHETIC DIVISION IS USED TO DEFLATE THE POLYNOMIAL BY DIVIDING OUT THE
C      * FACTOR (X-D).
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UPX,VPX,UB,VB,UA,VA
0003      DIMENSION UA(26),VA(26),UB(26),VB(26)
0004      UB(1)=UA(1)
0005      VB(1)=VA(1)
0006      NUM=NA+1
0007      DO 10 I=2,NUM
0008      UB(I)=UA(I)+(UB(I-1)*UX-VB(I-1)*VX)
0009      10 VB(I)=VA(I)+(VB(I-1)*UX+UB(I-1)*VX)
0010      UPX=UB(NUM)
0011      VPX=VB(NUM)
0012      RETURN
0013      END

```


TABLE F.III (Continued)

```

0001      SUBROUTINE CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,V
C      *****
C      *
C      * GIVEN THREE APPROXIMATIONS X(N-2), X(N-1), AND X(N), SUBROUTINE CALC
C      * APPROXIMATES THE POLYNOMIAL BY A QUADRATIC AND SOLVES FOR THE ZERO OF
C      * THE QUADRATIC CLOSEST TO X(N). THIS ZERO IS THE NEW APPROXIMATION
C      * X(N+1) TO THE ZERO OF THE POLYNOMIAL.
C      *
C      *****
0002      1PX3,UX4,VX4,UQ4,VQ4,UH3,VH3)
0003      DOUBLE PRECISION ARG1,ARG2
      DOUBLE PRECISION UPX3,VPX3,UPX2,VPX2,UX1,VX1,UX2,VX2,UX3,VX3,UPX1,
      1VPX1,UH3,VH3,UH2,VH2,UQ3,VQ3,UD,VD,UB,V8,UC,VC,UDISC,VOISC,UCCC,VC
      2CC,UDEN1,VDEN1,UDEN2,VDEN2,UQ4,VQ4,UX4,VX4,EPSRT,EPSQ,EPS,UDDD,VDD
      3D,AAA,B8B,RAD,UAAA,VAAA,UB8B,VB8B
0004      DOUBLE PRECISION THETA,ANGLE,UTEST,VTEST
0005      DOUBLE PRECISION EPS1,EPSM
0006      COMMON EPSRT,EPS1,EPS,EPSQ,EPSM,IO2,MAX
0007      UH3=UX3-UX2
0008      VH3=VX3-VX2
0009      UH2=UX2-UX1
0010      VH2=VX2-VX1
0011      B8B=UH2*UH2+VH2*VH2
0012      UQ3=(UH3*UH2+VH3*VH2)/B8B
0013      VQ3=(VH3*UH2-UH3*VH2)/B8B
0014      UDDD=1.0+UQ3
0015      VDDD=VQ3
0016      UD=(UPX3-(UDDD*UPX2-VDDD*VPX2))+(UQ3*UPX1-VQ3*VPX1)
0017      VD=(VPX3-(VDDD*UPX2+UDDD*VPX2))+(VQ3*UPX1+UQ3*VPX1)
0018      UAAA=2.0*UQ3
0019      VAAA=2.0*VQ3
0020      UAAA=UAAA+1.0
0021      UB8B=UDDD*UDDD-VDDD*VDDD
0022      VB8B=VDDD*UDDD+UDDD*VDDD
0023      UCCC=UQ3*UQ3-VQ3*VQ3
0024      VCCC=VQ3*UQ3+UQ3*VQ3
0025      UB=(UAAA*UPX3-VAAA*VPX3)-(UB8B*UPX2-VB8B*VPX2)+(UCCC*UPX1-VCCC*V
      1PX1)
0026      VB=(VAAA*UPX3+UAAA*VPX3)-(VB8B*UPX2+UB8B*VPX2)+(VCCC*UPX1+UCCC*V
      1PX1)
0027      UC=UDDD*UPX3-VDDD*VPX3
0028      VC=VDDD*UPX3+UDDD*VPX3
0029      UDISC=(UB*UB-VB*VB)-(4.0*(UD*UC-VD*VC))
0030      VOISC=(2.0*(VB*UB))-(4.0*(VD*UC+UD*VC))
0031      AAA=DSQRT(UDISC*UDISC+VOISC*VOISC)
0032      IF(AAA.EQ.0.0) GO TO 5
0033      GO TO 7
0034      5 THETA=0.0
0035      GO TO 9
0036      7 THETA=DATAN2(VOISC,UDISC)
0037      9 RAD=DSQRT(AAA)
0038      ANGLE=THETA/2.0
0039      UTEST=RAD*DCOS(ANGLE)
0040      VTEST=RAD*DSIN(ANGLE)
0041      UDEN1=UB+UTEST
0042      VDEN1=VB+VTEST
0043      UDEN2=UB-UTEST
0044      VDEN2=VB-VTEST

```

TABLE F.III (Continued)

```

0045      ARG1=UDEN1*UDEN1+VDEN1*VDEN1
0046      ARG2=UDEN2*UDEN2+VDEN2*VDEN2
0047      AAA=DSQRT(ARG1)
0048      BBB=DSQRT(ARG2)
0049      IF(AAA.LT.BBB) GO TO 10
0050      IF(AAA.EQ.0.0) GO TO 60
0051      UAAA=-2.0*UC
0052      VAAA=-2.0*VC
0053      UQ4=(UAAA*UDEN1+VAAA*VDEN1)/ARG1
0054      VQ4=(VAAA*UDEN1-UAAA*VDEN1)/ARG1
0055      GO TO 50
0056  10 IF(BBB.EQ.0.0) GO TO 60
0057      UAAA=-2.0*UC
0058      VAAA=-2.0*VC
0059      UQ4=(UAAA*UDEN2+VAAA*VDEN2)/ARG2
0060      VQ4=(VAAA*UDEN2-UAAA*VDEN2)/ARG2
0061      GO TO 50
0062  50 UX4=UX3+(UH3*UQ4-VH3*VQ4)
0063      VX4=VX3+(VH3*UQ4+UH3*VQ4)
0064      RETURN
0065  60 UQ4=1.0
0066      VQ4=0.0
0067      GO TO 50
0068      END

```

TABLE F.III (Continued)

```

0001      SUBROUTINE COMSQT(UX,VX,UY,VY)
C      *****
C      *
C      * THIS SUBROUTINE COMPUTES THE SQUARE ROOT OF A COMPLEX NUMBER.
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UY,VY,DUMMY,R,AAA,BBB
0003      R=DSQRT(UX*UX+VX*VX)
0004      AAA=DSQRT(DABS((R+UX)/2.0))
0005      BBB=DSQRT(DABS((R-UX)/2.0))
0006      IF(VX) 10,20,30
0007      10 UY=AAA
0008      VY=-1.0*BBB
0009      GO TO 100
0010      20 IF(UX) 40,50,60
0011      30 UY=AAA
0012      VY=BBB
0013      GO TO 100
0014      40 DUMMY=DABS(UX)
0015      UY=0.0
0016      VY=DSQRT(DUMMY)
0017      GO TO 100
0018      50 UY=0.0
0019      VY=0.0
0020      GO TO 100
0021      60 DUMMY=DABS(UX)
0022      UY=DSQRT(DUMMY)
0023      VY=0.0
0024      100 RETURN
0025      END

```

APPENDIX G

REPEATED G.C.D. - NEWTON'S METHOD

1. Use of the Program

A double precision FORTRAN IV program using the repeated G.C.D. method with Newton's method as a supporting method is presented here. Flow charts for this program are given in Figure G.2 while Table G.III gives a FORTRAN IV listing of this program. Single precision variables are listed in Table G.II. The single precision variables are used in the flow charts and the corresponding double precision variables can be obtained from Table G.II.

This program is designed to solve polynomials having degree less than or equal to 25. In order to solve polynomials of degree N where $N > 25$, the data statement and array dimensions given in Table G.I must be changed.

In this program both the leading coefficient and the constant coefficient are assumed to be non-zero.

TABLE G.I

PROGRAM CHANGES NECESSARY TO SOLVE POLYNOMIALS OF DEGREE
GREATER THAN 25 BY THE REPEATED G.C.D. - NEWTON'S METHOD

Main Program

Data Entry/1H1,1H2,...,1H9,2H10,2H11,...,2HXX/where XX = N+1

UP (N+1), VP (N+1)
UAPP (N), VAPP (N)
UD0 (N+1), VD0 (N+1)
UDD0 (N+1), VDD0 (N+1)
UD1 (N+1), VD1 (N+1)
UD2 (N+1), VD2 (N+1)
UDD1 (N+1), VDD1 (N+1)
UG (N+1), VG (N+1)
UD3 (2N+1), VD3 (2N+1)
UD4 (2N+1), VD4 (2N+1)
UZROS (N), VZROS (N)
UAP (N), VAP (N)
UROOT (N), VROOT (N)
NULT (N)
ENTRY (N+1)

Subroutine PROD

UH (2N+1), VH (2N+1)
UF (N+1), VF (N+1)
UG (N+1), VG (N+1)

Subroutine ZROS

UAPP (N), VAPP (N)
UROOT (N), VROOT (N)
UQ (N+1), VQ (N+1)
UQQ (N+1), VQQ (N+1)
UAP (N), VAP (N)
UQD (N+1), VQD (N+1)
ENTRY (N+1)
UROOTS (N), VROOTS (N)

Subroutines GENAPP, GCD, NEWTON, DIVIDE,
HORNER, and DERIV

See corresponding subroutine in Table E.I.

Subroutine QUAD

UROOT (N), VROOT (N)
UA (N+1), VA (N+1)

2. Input Data for Repeated G.C.D. - Newton's Method

The input data for repeated G.C.D. - Newton's method is prepared as described for G.C.D. - Newton's method in Appendix E, § 2 except that the item EPS4 on the control card (Figure E.2) is omitted. An example control card for the repeated G.C.D. - Newton's method is given in Figure G.1.

3. Variables Used in Repeated G.C.D. - Newton's Method

The definitions of variables used in repeated G.C.D. - Newton's method are given in Table G.II. For definitions of variables not listed in this table, see the main program or corresponding subprogram of Table E.VI. The notation and symbols used are defined in Appendix E, § 3.

4. Description of Program Output

The number of the polynomial, control data, degree and coefficients of the polynomial are printed as described in Appendix E, § 4.

All roots of multiplicity one are extracted first. Following the first row of asterixes, the message "THE FOLLOWING POLYNOMIAL, $G(X)$, CONTAINS ALL THE ROOTS OF $P(X)$ WHICH HAVE MULTIPLICITY 1." This is followed by the coefficients of $G(X)$ with the leading coefficient listed first. If there are no roots of multiplicity one, then the message "NO ROOTS OF MULTIPLICITY ONE" is printed.

The roots of $G(X)$ are printed under the heading "ROOTS OF $G(X)$." These are the roots obtained before the attempt to improve accuracy. The initial approximations producing convergence to the corresponding root are printed under the heading "INITIAL APPROXIMATION." The

message "RESULTS OF SUBROUTINE QUAD" means that the corresponding root was obtained from subroutine QUAD.

The roots found as a result of attempting to improve accuracy are printed under the heading "ROOTS OF P(X)." Their multiplicity is given under the heading "MULTIPLICITIES." The initial approximation is printed above where "NO INITIAL APPROXIMATION" means the same as "RESULTS OF SUBROUTINE QUAD."

A line of asterixes is then printed. This procedure is then repeated for the roots of multiplicity 2,3,4, etc. until all roots have been found.

5. Informative Messages and Error Messages

The informative messages and error messages for repeated G.C.D. - Newton's method are given below. For those not listed, see Appendix E, § 5.

"NOT ALL ROOTS OF THE ABOVE POLYNOMIAL, G, WERE FOUND." This message indicates that some of the roots of the polynomial $G(X)$ were not extracted.

"QUAD FOUND XXX TO BE A MULTIPLE ROOT." XXX represents the value of the root found as a multiple root by Subroutine QUAD.

TABLE G.II

REPEATED GCD - NEWTON'S METHOD

<u>Single Precision</u>		<u>Double Precision</u>		<u>Disposition</u>	<u>Description</u>
<u>Variable</u>	<u>Type</u>	<u>Variable</u>	<u>Type</u>	<u>of Argument</u>	
Main Program					
KD	I	KD	I		Number of distinct roots found
K	I	K	I		Number of roots found
J1	I	J1	I		Multiplicity of given root
D0	C	UD0, VD0	D		Array of coefficients of original polynomial
ND0	I	ND0	I		Degree of original polynomial
DD0	C	UDD0, VDD0	D		Array of coefficients of derivative of D0(X) i.e. D0'(X)
NDD0	I	NDD0	I		Degree of DD0(X)
D1	C	UD1, VD1	D		Array of coefficients of g.c.d. of D0(X) and DD0(X)
ND1	I	ND1	I		Degree of D1(X)
DD1	C	UDD1, VDD1	D		Array of coefficients of derivative of D1(X) i.e. D1'(X)
NDD1	I	NDD1	I		Degree of DD1(X)
D2	C	UD2, VD2	D		Array of coefficients of g.c.d. of D1(X) and DD1(X)
ND2	I	ND2	I		Degree of D2(X)
D3	C	UD3, VD3	D		Array of coefficients of the product of D0(X) and D2(X)
ND3	I	ND3	I		Degree of D3(X)
D4	C	UD4, VD4	D		Array of coefficients of the square of D1(X)
ND4	I	ND4	I		Degree of D4(X)
G	C	UG, VG	D		Array of coefficients of the quotient D3(X)/D4(X)
NG	I	NG	I		Degree of G(X)
ZROS	C	UZROS, VZROS	D		Array of roots of G(X)
Subroutine ZROS					
APROX	C	UAPROX, VAPROX	D	R	Starting approximation (initial or altered)

TABLE G.II (Continued)

<u>Single Precision</u> <u>Variable</u>	<u>Type</u>	<u>Double Precision</u> <u>Variable</u>	<u>Type</u>	<u>Disposition</u> <u>of Argument</u>	<u>Description</u>
					Subroutine PROD
M	I	M	I	E	Degree of polynomial to be multiplied
F	C	UF, VF	D	E	Array of coefficients of polynomial to be multiplied
N	I	N	I	E	Degree of polynomial to be multiplied
G	C	UG, VG	D	E	Array of coefficients of polynomial to be multiplied
MN	I	MN	I	R	Degree of product polynomial H(X)
H	C	UH, VH	D	R	Array of coefficients of product polynomial
LIMIT	I	LIMIT	I		Number of coefficients of polynomial F(X)
K	I	K	I		Counter

MAIN PROGRAM




Figure G.2. Flow Charts for Repeated G.C.D.-Newton's Method







Figure G.2. (Continued)

PROD



QUAD






Figure G.2. (Continued)

DIVIDE



HORNER



DERIV





Figure G.2. (Continued)

GENAPP



ALTER





Figure G.2. (Continued)

QUAD



NEWTON




Figure G.2. (Continued)




Figure G.2. (Continued)

COMSQT




Figure G.2. (Continued)

TABLE G.III

PROGRAM FOR REPEATED G.C.D.-NEWTON'S METHOD

```

C *****
C *
C * DOUBLE PRECISION PROGRAM FOR THE REPEATED G.C.D. - NEWTON'S METHOD *
C *
C *
C * THIS METHOD REPEATEDLY FINDS THE GREATEST COMMON DIVISOR OF TWO *
C * POLYNOMIALS IN ORDER TO EXTRACT THE ZEROS IN GROUPS ACCORDING TO *
C * MULTIPLICITY USING NEWTON'S METHOD. ALL ZEROS OF MULTIPLICITY 1 *
C * ARE EXTRACTED FOLLOWED BY THOSE OF MULTIPLICITY 2, ETC. *
C *
C *****
0001 DOUBLE PRECISION EPS1, EPS2, EPS3, UP, VP, UAPP, VAPP, UDO, VDO, UDDO, VDDO,
      1UD1, VD1, UD2, VD2, UDD1, VDD1, UG, VG, UD3, VD3, UD4, VD4, UZROS, VZROS, UAP, VA
      2P, UROOT, VROOT, DENOM
0002 DOUBLE PRECISION XSTART
0003 DOUBLE PRECISION XEND
0004 DIMENSION ANAME(2), UP(26), VP(26), UAPP(25), VAPP(25), UDO(26), VDO(26),
      1, UDDO(26), VDDO(26), UD1(26), VD1(26), UD2(26), VD2(26), UDD1(26), VDD1(2
      26), UG(26), VG(26), UD3(51), VD3(51), UD4(51), VD4(51), UZROS(25), VZROS(2
      35), UAP(25), VAP(25), UROOT(25), VROOT(25), MULTI(25), ENTRY(26)
0005 COMMON EPS1, EPS2, EPS3, IO2, MAX
0006 DATA ASTER/4H****/
0007 DATA PNAME, GNAME/2HP(, 2HG(/, D1NAME/3HD1(/
0008 DATA ENTRY/1H1, 1H2, 1H3, 1H4, 1H5, 1H6, 1H7, 1H8, 1H9, 2H10, 2H11, 2H12, 2H13
      1, 2H14, 2H15, 2H16, 2H17, 2H18, 2H19, 2H20, 2H21, 2H22, 2H23, 2H24, 2H25, 2H26/
0009 DATA ANAME(1), ANAME(2)/4HNEWT, 4HONS /
0010 IO1=5
0011 IO2=6
0012 1 READ(IO1, 1000) NOPOLY, NP, NAPP, MAX, EPS1, EPS2, EPS3, XSTART, XEND, KCHEC
      1K
0013 IF(KCHECK.EQ.1) STOP
0014 WRITE(IO2, 1020) ANAME(1), ANAME(2), NOPOLY
0015 WRITE(IO2, 2000) NAPP
0016 WRITE(IO2, 2010) MAX
0017 WRITE(IO2, 2070) EPS1
0018 WRITE(IO2, 2020) EPS2
0019 WRITE(IO2, 2080) EPS3
0020 WRITE(IO2, 2040) XSTART
0021 WRITE(IO2, 2050) XEND
0022 WRITE(IO2, 2060)
0023 KKK=NP+1
0024 NNN=KKK+1
0025 DO 5 I=1, KKK
0026 JJJ=NNN-I
0027 5 READ(IO1, 1010) UP(JJJ), VP(JJJ)
0028 IF(NAPP.NE.0) GO TO 22
0029 NAPP=NP
0030 CALL GENAPP(UAPP, VAPP, NAPP, XSTART)
0031 GO TO 23
0032 22 READ(IO1, 1015) (UAPP(I), VAPP(I), I=1, NAPP)
0033 23 WRITE(IO2, 1030) NP
0034 KKK=NP+1
0035 NNN=KKK+1
0036 DO 8 I=1, KKK
0037 JJJ=NNN-I
0038 8 WRITE(IO2, 1040) PNAME, ENTRY(JJJ), UP(JJJ), VP(JJJ)
0039 K=0
0040 KD=0

```

TABLE G.III (Continued)

```

0041      J1=1
0042      KKK=NP+1
0043      DO 10 I=1, KKK
0044      UDO(I)=UP(I)
0045      10 VDO(I)=VP(I)
0046      NDO=NP
0047      CALL DERIV(NDO, UDO, VDO, NDDO, UDDO, VDDO)
0048      CALL GCD(NDO, UDO, VDO, NDDO, UDDO, VDDO, ND1, UD1, VD1)
0049      20 WRITE(IO2, 3000) (ASTER, I=1, 33)
0050      IF(ND1.LE.1) GO TO 30
0051      GO TO 40
0052      30 UD2(I)=1.0
0053      VD2(I)=0.0
0054      ND2=0
0055      GO TO 50
0056      40 CALL DERIV(ND1, UD1, VD1, NDD1, UDD1, VDD1)
0057      CALL GCD(ND1, UD1, VD1, NDD1, UDD1, VDD1, ND2, UD2, VD2)
0058      50 IF(ND0+ND2.LE.2*ND1) GO TO 60
0059      GO TO 70
0060      60 WRITE(IO2, 1025) J1
0061      GO TO 170
0062      70 IF(ND1.EQ.0) GO TO 80
0063      GO TO 90
0064      80 KKK=NDO+1
0065      DO 85 I=1, KKK
0066      UG(I)=UDO(I)
0067      85 VG(I)=VDO(I)
0068      NG=NDO
0069      GO TO 110
0070      90 IF(ND2.EQ.0) GO TO 115
0071      CALL PROD(NDO, UDO, VDO, ND2, UD2, VD2, ND3, UD3, VD3)
0072      100 CALL PROD(ND1, UD1, VD1, ND1, UD1, VD1, ND4, UD4, VD4)
0073      CALL DIVIDE(ND3, UD3, VD3, ND4, UD4, VD4, NG, UG, VG)
0074      110 WRITE(IO2, 1035) J1
0075      KKK=NG+1
0076      NNN=KKK+1
0077      DO 112 I=1, KKK
0078      JJJ=NNN-I
0079      112 WRITE(IO2, 1040) GNAME, ENTRY(JJJ), UG(JJJ), VG(JJJ)
0080      CALL ZEROS(NG, UG, VG, NAPP, UAPP, VAPP, J, UZROS, VZROS, JAP, UAP, VAP, ENTRY
1, XSTART, XEND)
0081      IF(J.EQ.0) GO TO 150
0082      WRITE(IO2, 1180)
0083      IF(JAP.EQ.0) GO TO 120
0084      GO TO 130
0085      115 KKK=NDO+1
0086      DO 116 I=1, KKK
0087      UD3(I)=UDO(I)
0088      116 VD3(I)=VDO(I)
0089      ND3=NDO
0090      GO TO 100
0091      120 KKK=JAP+1
0092      WRITE(IO2, 1085) (I, UZROS(I), VZROS(I), J1, I=KKK, J)
0093      GO TO 140
0094      130 WRITE(IO2, 1190) (I, UZROS(I), VZROS(I), J1, UAP(I), VAP(I), I=1, JAP)
0095      IF(JAP.LT.J) GO TO 120
0096      140 IF(J.EQ.NG) GO TO 155
0097      150 WRITE(IO2, 1095)

```

TABLE G.III (Continued)

```

0098      IF(J.EQ.0) GO TO 170
0099      155 DO 160 I=1,J
0100          UROOT(KD+I)=UZROS(I)
0101          VROOT(KD+I)=VZROS(I)
0102      160 MULT(KD+I)=J1
0103          K=(J*J1)+K
0104          KD=KD+J
0105          IF(K.GE.NP) GO TO 1
0106      170 J1=J1+1
0107          IF(ND1.LE.1) GO TO 200
0108          DO 180 I=1,ND1
0109              UDO(I)=UD1(I)
0110              VDO(I)=VD1(I)
0111              UDDO(I)=UDD1(I)
0112      180 VDDO(I)=VDD1(I)
0113              UDO(ND1+1)=UD1(ND1+1)
0114              VDO(ND1+1)=VD1(ND1+1)
0115              NDO=ND1
0116              NDDO=NDD1
0117              KKK=ND2+1
0118              DO 190 I=1,KKK
0119                  UD1(I)=UD2(I)
0120      190 VD1(I)=VD2(I)
0121                  ND1=ND2
0122                  GO TO 29
0123      200 IF(ND1.EQ.0) GO TO 1
0124          KD=KD+1
0125          DENOM=UD1(2)*UD1(2)+VD1(2)*VD1(2)
0126          UROOT(KD)=(-UD1(1)*UD1(2)-VD1(1)*VD1(2))/DENOM
0127          VROOT(KD)=(-VD1(1)*UD1(2)+UD1(1)*VD1(2))/DENOM
0128          MULT(KD)=J1
0129          WRITE(102,3000) (ASTER,I=1,33)
0130          WRITE(102,1035) J1
0131          KKK=ND1+1
0132          NNN=KKK+1
0133          DO 210 I=1,KKK
0134              JJJ=NNN-I
0135      210 WRITE(102,1100) D1NAME,ENTRY(JJJ),UD1(JJJ),VD1(JJJ)
0136          WRITE(102,1180)
0137          WRITE(102,1085) KD,UROOT(KD),VROOT(KD),J1
0138          GO TO 1
0139      1020 FORMAT(1H1,10X,48HREPEATED USE OF THE GREATEST COMMON DIVISOR AND
1,A4,A4,58H METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIA
2LS/11X,18HPOLYNOMIAL NUMBER ,I2///)
0140      1025 FORMAT(///1X,25HNO ROOTS OF MULTIPLICITY ,I2///)
0141      1035 FORMAT(///1X,87HTHE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE R
1OOTS OF P(X) WHICH HAVE MULTIPLICITY ,I2///)
0142      1085 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,18X,25H
1NO INITIAL APPROXIMATIONS)
0143      1095 FORMAT(///1X,51HNOT ALL ROOTS OF THE ABOVE POLYNOMIAL,G, WERE FOUN
1D//)
0144      1000 FORMAT(3(I2,1X),9X,I3,1X,3(D6.0,1X),20X,2(D7.0,1X),11)
0145      1010 FORMAT(2D30.0)
0146      1015 FORMAT(2D30.0)
0147      1030 FORMAT(1X,22HTHE DEGREE OF P(X) IS ,I2,22H THE COEFFICIENTS ARE//
1)
0148      1040 FORMAT(2X,A2,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0149      1100 FORMAT(2X,A3,A2,4H) = ,D23.16,3H + ,D23.16,2H I)

```

TABLE G.III (Continued)

```

0150      1180 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES,17X,21HINITIAL
0151      1190 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,7X,I2,7X,D23.
          116,3H + ,D23.16,2H I)
0152      2000 FORMAT(1X,41HNUMBER OF INITIAL APPROXIMATIONS GIVEN. ,I2)
0153      2010 FORMAT(1X,29HMAXIMUM NUMBER OF ITERATIONS.,11X,I3)
0154      2020 FORMAT(1X,21HTEST FOR CONVERGENCE.,13X,D9.2)
0155      2040 FORMAT(1X,23HRADIUS TO START SEARCH.,11X,D9.2)
0156      2050 FORMAT(1X,21HRADIUS TO END SEARCH.,13X,D9.2)
0157      2060 FORMAT(//1X)
0158      2070 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE GCD. ,D9.2)
0159      2080 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE QUAD. ,D9.2)
0160      3000 FORMAT(/////1X,A3,32A4)
0161      END

```


TABLE G.III (Continued)

```

0001      SUBROUTINE PROD(M,UF,VF,N,UG,VG,MN,UH,VH)
C      *****
C      *
C      * GIVEN POLYNOMIALS R(X) AND S(X), THIS SUBROUTINE COMPUTES THE
C      * COEFFICIENTS OF THE PRODUCT POLYNOMIAL T(X) = R(X).S(X).
C      *
C      *****
0002      DOUBLE PRECISION UH,VH,UF,VF,UG,VG
0003      DIMENSION UH(51),VH(51),UF(26),VF(26),UG(26),VG(26)
0004      MN=M+N
0005      KKK=MN+1
0006      DO 100 I=1,KKK
0007          K=I
0008          UH(I)=0.0
0009          VH(I)=0.0
0010          IF(I.LE.M+1) GO TO 10
0011          LIMIT=M+1
0012          GO TO 20
0013      10 LIMIT=I
0014      20 DO 50 J=1,LIMIT
0015          IF(K.GT.N+1) GO TO 50
0016          IF(J+K.EQ.I+1) GO TO 40
0017          GO TO 50
0018      40 UH(I)=UH(I)+(UF(J)*UG(K)-VF(J)*VG(K))
0019          VH(I)=VH(I)+(VF(J)*UG(K)+UF(J)*VG(K))
0020      50 K=K-1
0021      100 CONTINUE
0022      RETURN
0023      END

0001      SUBROUTINE GENAPP(APPR,APPI,NAPP,XSTART)
C      *****
C      *
C      * SUBROUTINE GENAPP GENERATES N INITIAL APPROXIMATIONS, WHERE N IS THE
C      * DEGREE OF THE ORIGINAL POLYNOMIAL.
C      *
C      *****
0002      DOUBLE PRECISION APPR,APPI,XSTART,BETA, EPS1,EPS2,EPS3
0003      DIMENSION APPR(25),APPI(25)
0004      COMMON EPS1,EPS2,EPS3,IO2,MAX
0005      IF(XSTART.EQ.0.0) XSTART=0.5
0006      BETA=0.2617994
0007      DO 10 I=1,NAPP
0008          APPR(I)=XSTART*DCOS(BETA)
0009          APPI(I)=XSTART*DSIN(BETA)
0010          BETA=BETA+0.5235988
0011      10 XSTART=XSTART+0.5
0012      RETURN
0013      END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE ALTER(XOLDR,XOLDI,NALTER,ITIME)
C      *****
C      *
C      * SUBROUTINE ALTER ALTERS THE INITIAL APPROXIMATIONS WHICH PRODUCE NO
C      * CONVERGENCE TO A ZERO. THIS IS DONE A MAXIMUM OF 5 TIMES FOR EACH ROOT.
C      *
C      *****
0002      DOUBLE PRECISION XOLDR,XOLDI,ABXOLD,BETA,EPS1,EPS2,EPS3
0003      COMMON EPS1,EPS2,EPS3,IO2,MAX
0004      IF(ITIME.NE.0) GO TO 5
0005      ITIME =1
0006      WRITE(IO2,1010) MAX
0007      5 IF(NALTER.EQ.0) GO TO 10
0008      WRITE(IO2,1000) XOLDR,XOLDI
0009      GO TO 20
0010      10 ABXOLD=DSQRT((XOLDR*XOLDR)+(XOLDI*XOLDI))
0011      BETA=DATAN2(XOLDI,XOLDR)
0012      WRITE(IO2,1020) XOLDR,XOLDI
0013      20 NALTER=NALTER+1
0014      IF(NALTER.GT.5) RETURN
0015      GO TO (30,40,30,40,30),NALTER
0016      30 XOLDR=-XOLDR
0017      XOLDI=-XOLDI
0018      GO TO 50
0019      40 BETA=BETA+1.0471976
0020      XOLDR=ABXOLD*DCOS(BETA)
0021      XOLDI=ABXOLD*DSIN(BETA)
0022      50 RETURN
0023      1000 FORMAT(1X,D23.16,3H + ,D23.16,2H I,10X,21HALTERED APPROXIMATION)
0024      1010 FORMAT(///1X,54HNO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AF
           ITER ,I3,12H ITERATIONS.//)
0025      1020 FORMAT(1X,D23.16,3H + ,D23.16,2H I,10X,21HINITIAL APPROXIMATION)
0026      END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE ZEROS(NQ,UQ,VQ,NAPP,UAPP,VAPP,J,UROOT,VROOT,JAP,UAP,VAP
          1,ENTRY,XSTART,XEND)
C      *****
C      *
C      * NEWTONS METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C      * POLYNOMIAL OF MAXIMUM DEGREE 25 BY COMPUTING A SEQUENCE OF APPROX-
C      * IMATIONS CONVERGING TO A ZERO OF THE POLYNOMIAL USING THE ITERATION
C      * FORMULA
C      *
C      *           $X(N+1) = X(N) - P(X(N)) / P'(X(N))$ 
C      *
C      *****
0002      DOUBLE PRECISION UAPP,VAPP,UROOT,VROOT,UZRO,VZRO,UQ,VQ,UDUMMY,VDUM
          1MY,UQQ,VQQ,UAP,VAP,UQD,VQD,UROOTS,VROOTS,EPS1,EPS2,EPS3,UAPROX,VAP
          2ROX
0003      DOUBLE PRECISION XEND,XSTART
0004      DIMENSION UAPP(25),VAPP(25),UROOT(25),VROOT(25),UQ(26),VQ(26),UQQ(
          126),VQQ(26),UAP(25),VAP(25),UQD(26),VQD(26),ENTRY(26),UROOTS(25),V
          2ROOTS(25)
0005      COMMON EPS1,EPS2,EPS3,I02,MAX
0006      DATA QQNAME,QNAME/3HQQ/,2HQ(/
0007      LOGICAL CONV
0008      J=0
0009      ITIME=0
0010      IF(NQ.GE.3) GO TO 85
0011      GO TO 110
0012      85 KKK=NQ+1
0013      DO 90 I=1,KKK
0014      UQQ(I)=UQ(I)
0015      90 VQQ(I)=VQ(I)
0016      NQ=NQ
0017      GO TO 120
0018      110 CALL QUAD(NQ,UQ,VQ,J,UROOT,VROOT)
0019      JAP=0
0020      GO TO 310
0021      120 DO 200 I=1,NAPP
0022      IALTER=0
0023      UAPROX=UAPP(I)
0024      VAPROX=VAPP(I)
0025      130 CALL NEWTON(UAPROX,VAPROX,NQQ,UQQ,VQQ,UZRO,VZRO,CONV)
0026      IF(CONV) GO TO 160
0027      CALL ALTER(UAPP(I),VAPP(I),IALTER,ITIME)
0028      IF(IALTER.GT.5) GO TO 200
0029      UAPROX=UAPP(I)
0030      VAPROX=VAPP(I)
0031      GO TO 130
0032      160 J=J+1
0033      UROOT(J)=UZRO
0034      VROOT(J)=VZRO
0035      UAP(J)=UAPROX
0036      VAP(J)=VAPROX
0037      CALL HORNER(UZRO,VZRO,NQQ,UQQ,VQQ,UQD,VQD,UDUMMY,VDUMMY)
0038      DO 180 I1=1,NQQ
0039      UQQ(I1)=UQD(I1+1)
0040      180 VQQ(I1)=VQD(I1+1)
0041      NQ=NQ-1
0042      IF(NQQ.GE.3) GO TO 200
0043      JAP=J
0044      GO TO 220

```

TABLE G.III (Continued)

```

0045      200 CONTINUE
0046          IF(J.GE.NQ) GO TO 205
0047          IF(XEND.EQ.0.0) GO TO 205
0048          IF(XSTART.GT.XEND) GO TO 205
0049          NAPP=NQ
0050          CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0051          GO TO 120
0052      205 IF(NQQ.LE.2) GO TO 210
0053          WRITE(102,1200)
0054          KKK=NQQ+1
0055          NNN=KKK+1
0056          DO 157 L=1,KKK
0057              JJJ=NNN-L
0058      157 WRITE(102,1100) QNAME,ENTRY(JJJ),UQQ(JJJ),VQQ(JJJ)
0059      210 IF(J.EQ.0) GO TO 310
0060          JAP=J
0061          GO TO 230
0062      220 CALL QUAD(NQQ,UQQ,VQQ,J,UROOT,VROOT)
0063      230 WRITE(102,1132)
0064          WRITE(102,1133) (I,UROOT(I),VROOT(I),UAP(I),VAP(I),I=1,JAP)
0065          IF(JAP.LT.J) GO TO 235
0066          GO TO 240
0067      235 KKK=JAP+1
0068          WRITE(102,1134) (I,UROOT(I),VROOT(I),I=KKK,J)
0069      240 J1=0
0070          DO 300 I=1,J
0071              CALL NEWTON(UROOT(I),VROOT(I),NQ,UQ,VQ,UZRO,VZRO,CONV)
0072              IF(CONV) GO TO 280
0073              WRITE(102,1140) I,UROOT(I),VROOT(I),MAX,NQ
0074              KKK=NQ+1
0075              NNN=KKK+1
0076              DO 242 L=1,KKK
0077                  JJJ=NNN-L
0078      242 WRITE(102,1040) QNAME,ENTRY(JJJ),UQ(JJJ),VQ(JJJ)
0079                  IF(I.LT.JAP) GO TO 241
0080                  IF(I.EQ.JAP) GO TO 250
0081                  GO TO 300
0082      241 KKK=JAP-1
0083                  DO 245 I1=I,KKK
0084                      UAP(I1)=UAP(I1+1)
0085                      VAP(I1)=VAP(I1+1)
0086      250 JAP=JAP-1
0087                  GO TO 300
0088      280 J1=J1+1
0089              UROOTS(J1)=UZRO
0090              VROOTS(J1)=VZRO
0091      300 CONTINUE
0092          J=J1
0093          IF(J.EQ.0) GO TO 305
0094          DO 303 I=1,J
0095              UROOT(I)=UROOTS(I)
0096      303 VROOT(I)=VROOTS(I)
0097          GO TO 310
0098      305 WRITE(102,1150) NQ
0099          KKK=NQ+1
0100          NNN=KKK+1
0101          DO 306 L=1,KKK
0102              JJJ=NNN-L

```

TABLE G.III (Continued)

```

0103      306 WRITE(102,1040) QNAME,ENTRY(JJJ),UQ(JJJ),VQ(JJJ)
0104      310 RETURN
0105      1200 FORMAT(///1X,70HCOEFFICIENTS OF THE DEFLATED POLYNOMIAL FOR WHICH
           1NO ZEROS WERE FOUND.//)
0106      1132 FORMAT(///1X,13HROOTS OF G(X),84X,21HINITIAL APPROXIMATION//)
0107      1133 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,17X,D23.16,3H
           1 + ,D23.16,2H I)
0108      1134 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,22X,26HRESULT
           1S OF SUBROUTINE QUAD)
0109      1140 FORMAT(///,1X,40HNO ROOTS FOR INITIAL APPROXIMATION ROOT(,I2,4H) =
           1 ,D23.16,3H + ,D23.16,2H I/6H AND ,I3,40H ITERATIONS ON THE POLYN
           2OMIAL OF DEGREE ,I2,18H WITH COEFFICIENTS//)
0110      1150 FORMAT(///,1X,45HNO ROOTS FOR THE POLYNOMIAL Q(X) OF DEGREE = ,I2,
           138H WITH GENERATED INITIAL APPROXIMATIONS//)
0111      1040 FORMAT(2X,A2,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0112      1100 FORMAT(2X,A3,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0113      END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE GCD(N,UR,VR,M,US,VS,M1,USS,VSS)
C      *****
C      *
C      * GIVEN POLYNOMIALS P(X) AND DP(X) WHERE DEG. DP(X) IS LESS THAN DEG.
C      * P(X), SUBROUTINE GCD COMPUTES THE GREATEST COMMON DIVISOR OF P(X) AND
C      * DP(X).
C      *
C      *****
0002      DOUBLE PRECISION USSSSS,VSSSSS
0003      DOUBLE PRECISION UR,VR,US,VS,USS,VSS,URR,VRR,UD,VD,UT,VT,EPSLON,EP
0004      IS2,EPS3,BBB
0005      DIMENSION UR(26),VR(26),US(26),VS(26),USS(26),VSS(26),URR(26),VRR(
0006      126),UT(26),VT(26)
0007      COMMON EPSLON,EPS2,EPS3,I02,MAX
0008      N1=N
0009      M1=M
0010      KKK=N+1
0011      DO 20 I=1,KKK
0012      URR(I)=UR(I)
0013      VRR(I)=VR(I)
0014      KKK=M+1
0015      DO 25 I=1,KKK
0016      USS(I)=US(I)
0017      VSS(I)=VS(I)
0018      30 BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0019      UD=(URR(N1+1)*USS(M1+1)+VRR(N1+1)*VSS(M1+1))/BBB
0020      VD=(USS(M1+1)*VRR(N1+1)-URR(N1+1)*VSS(M1+1))/BBB
0021      KKK=N1+1-M1
0022      DO 40 I=KKK,N1
0023      UT(I)=URR(I)-(UD*USS(I-N1+M1)-VD*VSS(I-N1+M1))
0024      VT(I)=VRR(I)-(UD*VSS(I-N1+M1)+VD*USS(I-N1+M1))
0025      IF(M1.EQ.N1) GO TO 70
0026      KKK=N1-M1
0027      DO 60 I=1,KKK
0028      UT(I)=URR(I)
0029      VT(I)=VRR(I)
0030      60 VT(I)=VRR(I)
0031      70 DO 90 I=1,N1
0032      BBB=DSQRT(UT(N1+1-I)*UT(N1+1-I)+VT(N1+1-I)*VT(N1+1-I))
0033      IF(BBB.GT.EPSLON) GO TO 100
0034      CONTINUE
0035      DO 95 I=1,M1
0036      BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0037      USSSSS=(USS(I)*USS(M1+1)+VSS(I)*VSS(M1+1))/BBB
0038      VSSSSS=(VSS(I)*USS(M1+1)-USS(I)*VSS(M1+1))/BBB
0039      USS(I)=USSSSS
0040      VSS(I)=VSSSSS
0041      USS(M1+1)=1.0
0042      VSS(M1+1)=0.0
0043      GO TO 200
0044      100 K=N1-I
0045      IF(K.EQ.0) GO TO 170
0046      IF(K.LT.M1) GO TO 140
0047      KKK=K+1
0048      DO 130 J=1,KKK
0049      URR(J)=UT(J)
0050      VRR(J)=VT(J)
0051      N1=K
0052      GO TO 30

```

TABLE G.III (Continued)

```

0050      140 KKK=K+1
0051          DO 150 J=1, KKK
0052              URR(J)=USS(J)
0053              VRR(J)=VSS(J)
0054              USS(J)=UT(J)
0055      150 VSS(J)=VT(J)
0056          KKK=K+2
0057          NNN=M1+1
0058          DO 160 J=KKK, NNN
0059              URR(J)=USS(J)
0060      160 VRR(J)=VSS(J)
0061          N1=M1
0062          M1=K
0063          GO TO 30
0064      170 USS(1)=1.0
0065          VSS(1)=0.0
0066          M1=0
0067      200 RETURN
0068          END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE NEWTON(UX,VX,N,UP,VP,UXO,VXO,CONV)
C      *****
C      *
C      * THIS SUBROUTINE CALCULATES A NEW APPROXIMATION FROM THE OLD APPROX-
C      * IMATION BY USING THE ITERATION FORMULA
C      *  $X(N+1) = X(N) - P(X(N))/P'(X(N))$ .
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UP,VP,UXO,VXO,UB,VB,UDPXO,VDPXO,UPXO,VPXO,U
0003      IDIFF,VDIFF,EPS1,EPSLON,EPS3,AAA,BBB
0004      DOUBLE PRECISION DDD
0005      DOUBLE PRECISION ABPXO
0006      DIMENSION UP(26),VP(26),UB(26),VB(26)
0007      COMMON EPS1,EPSLON,EPS3,IO2,MAX
0008      LOGICAL CONV
0009      UXO=UX
0010      VXO=VX
0011      DO 10 I=1,MAX
0012      CALL HORNER(UXO,VXO,N,UP,VP,UB,VB,UDPXO,VDPXO)
0013      UPXO=UB(1)
0014      VPXO=VB(1)
0015      DDD=DSQRT(UDPXO*UDPXO+VDPXO*VDPXO)
0016      IF(DDD.NE.0.0) GO TO 5
0017      ABPXO=DSQRT(UPXO*UPXO+VPXO*VPXO)
0018      IF(ABPXO.EQ.0.0) GO TO 20
0019      GO TO 15
0020      5 BBB=UDPXO*UDPXO+VDPXO*VDPXO
0021      UDIFF=(UPXO*UDPXO+VPXO*VDPXO)/BBB
0022      VDIFF=(VPXO*UDPXO-UPXO*VDPXO)/BBB
0023      UXO=UXO-UDIFF
0024      VXO=VXO-VDIFF
0025      AAA=DSQRT(UDIFF*UDIFF+VDIFF*VDIFF)
0026      BBB=DSQRT(UXO*UXO+VXO*VXO)
0027      IF(BBB.EQ.0.0) GO TO 10
0028      IF(AAA/BBB.LT.EPSLON) GO TO 20
0029      10 CONTINUE
0030      15 CONV=.FALSE.
0031      RETURN
0032      20 CONV=.TRUE.
0033      RETURN
0034      END

```


TABLE G.III (Continued)

```

0001      SUBROUTINE DIVIDE(N,UP,VP,M,UD,VD,K,UQ,VQ)
C      *****
C      *
C      * GIVEN TWO POLYNOMIALS F(X) AND G(X), SUBROUTINE DIVIDE COMPUTES THE
C      * QUOTIENT POLYNOMIAL H(X) = F(X)/G(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UD,VD,UQ,VQ,UTERM,VTERM,UDUMMY
0003      DIMENSION UP(26),VP(26),UD(26),VD(26),UQ(26),VQ(26)
0004      K=N-M
0005      UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0006      UQ(K+1)=(UP(N+1)*UD(M+1)+VP(N+1)*VD(M+1))/UDUMMY
0007      VQ(K+1)=(VP(N+1)*UD(M+1)-UP(N+1)*VD(M+1))/UDUMMY
0008      IF(K.EQ.0) GO TO 100
0009      J=-1
0010      DO 50 I=1,K
0011      J=J+1
0012      UTERM=UP(N-J)
0013      VTERM=VP(N-J)
0014      KK=K+1
0015      NNN=M-J
0016      DO 40 M1=NNN,M
0017      IF(KK.GT.1) GO TO 10
0018      GO TO 45
0019      10 IF(M1.GE.1) GO TO 20
0020      GO TO 40
0021      20 UTERM=UTERM-(UQ(KK)*UD(M1)-VQ(KK)*VD(M1))
0022      VTERM=VTERM-(UQ(KK)*VD(M1)+VQ(KK)*UD(M1))
0023      40 KK=KK-1
0024      45 UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0025      UQ(K+1-I)=(UTERM*UD(M+1)+VTERM*VD(M+1))/UDUMMY
0026      50 VQ(K+1-I)=(VTERM*UD(M+1)-UTERM*VD(M+1))/UDUMMY
0027      100 RETURN
0028      END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE HORNER(UX,VX,N,UP,VP,UB,VB,UC,VC)
C      *****
C      *
C      * HORNER'S METHOD COMPUTES THE VALUE OF THE POLYNOMIAL P(X) AT A
C      * POINT D AND ITS DERIVATIVE AT D. SYNTHETIC DIVISION IS USED TO
C      * DEFLATE THE POLYNOMIAL BY DIVIDING OUT THE FACTOR (X - D).
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UP,VP,UB,VB,UC,VC
0003      DOUBLE PRECISION UDUMMY,VDUMMY
0004      DIMENSION UP(26),VP(26),UB(26),VB(26)
0005      UB(N+1)=UP(N+1)
0006      VB(N+1)=VP(N+1)
0007      UB(N)=(UX*UB(N+1)-VX*VB(N+1))+UP(N)
0008      VB(N)=(UX*VB(N+1)+VX*UB(N+1))+VP(N)
0009      UC=UB(N+1)
0010      VC=VB(N+1)
0011      KKK=N-1
0012      DO 10 I=1,KKK
0013      UB(KKK+1-I)=(UX*UB(KKK+2-I)-VX*VB(KKK+2-I))+UP(KKK+1-I)
0014      VB(KKK+1-I)=(UX*VB(KKK+2-I)+VX*UB(KKK+2-I))+VP(KKK+1-I)
0015      UDUMMY=UX*UC-VX*VC
0016      VDUMMY=UX*VC+VX*UC
0017      UC=UDUMMY+UB(KKK+2-I)
0018      10 VC=VDUMMY+VB(KKK+2-I)
0019      RETURN
0020      END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE QUAD(N,UA,VA,J,UROOT,VROOT)
C      *****
C      *
C      * SUBROUTINE QUAD SOLVES DIRECTLY FOR THE ZEROS AND THEIR MULTIPLICITIES *
C      * OF EITHER A QUADRATIC POLYNOMIAL OR A LINEAR FACTOR. SOLUTION OF THE *
C      * QUADRATIC IS DONE USING THE QUADRATIC FORMULA. *
C      *
C      *****
0002      DOUBLE PRECISION EPS1, EPS2, EPSLON, UROOT, VROOT, UA, VA, UDISC, VDISC, UD
0003      1, VD, DDD, UTEMP, VTEMP, BBB
0004      DIMENSION UROOT(25), VROOT(25), UA(26), VA(26)
0005      COMMON EPS1, EPS2, EPSLON, IO2, MAX
0006      IF(N.GT.1) GO TO 10
0007      J=J+1
0008      BBB=UA(2)*UA(2)+VA(2)*VA(2)
0009      UROOT(J)=- (UA(1)*UA(2)+VA(1)*VA(2))/BBB
0010      VROOT(J)=- (VA(1)*UA(2)-UA(1)*VA(2))/BBB
0011      GO TO 100
0012      10 UDISC=(UA(2)*UA(2)-VA(2)*VA(2))-(4.0*(UA(3)*UA(1)-VA(3)*VA(1)))
0013      VDISC=(2.0*UA(2)*VA(2))-(4.0*(UA(3)*VA(1)+VA(3)*UA(1)))
0014      UD=2.0*UA(3)
0015      VD=2.0*VA(3)
0016      DDD=DSQRT(UDISC*UDISC+VDISC*VDISC)
0017      IF(DDD.LT.EPSLON) GO TO 20
0018      CALL COMSQT(UDISC,VDISC,UTEMP,VTEMP)
0019      BBB=UD*UD+VD*VD
0020      UROOT(J+1)=((-UA(2)+UTEMP)*UD+(-VA(2)+VTEMP)*VD)/BBB
0021      VROOT(J+1)=((-VA(2)+VTEMP)*UD-(-UA(2)+UTEMP)*VD)/BBB
0022      UROOT(J+2)=((-UA(2)-UTEMP)*UD+(-VA(2)-VTEMP)*VD)/BBB
0023      VROOT(J+2)=((-VA(2)-VTEMP)*UD-(-UA(2)-UTEMP)*VD)/BBB
0024      J=J+2
0025      GO TO 100
0026      20 J=J+1
0027      BBB=UD*UD+VD*VD
0028      UROOT(J)=(-UA(2)*UD-VA(2)*VD)/BBB
0029      VROOT(J)=(-VA(2)*UD+UA(2)*VD)/BBB
0030      WRITE(IO2,1000) UROOT(J),VROOT(J)
0031      1000 FORMAT(//1X,11HQUAD FOUND ,D23.16,3H + ,D23.16,2H I,22H TO BE A M
0032      1ULTIPLE ROOT//)
0031      100 RETURN
0032      END

```

TABLE G.III (Continued)

```

0001      SUBROUTINE DERIV(N,UP,VP,M,UA,VA)
C      *****
C      *
C      * GIVEN A POLYNOMIAL P(X), SUBROUTINE DERIV COMPUTES THE COEFFICIENTS OF
C      * ITS DERIVATIVE P'(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UA,VA,AAA
0003      DIMENSION UP(26),VP(26),UA(26),VA(26)
0004      KKK=N+1
0005      DO 10 I=2,KKK
0006      AAA=I-1
0007      UA(I-1)=AAA*UP(I)
0008      10 VA(I-1)=AAA*VP(I)
0009      M=N-1
0010      RETURN
0011      END

```

```

0001      SUBROUTINE COMSQT(UX,VX,UY,VY)
C      *****
C      *
C      * THIS SUBROUTINE COMPUTES THE SQUARE ROOT OF A COMPLEX NUMBER.
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UY,VY,DUMMY,R,AAA,BBB
0003      R=DSQRT(UX*UX+VX*VX)
0004      AAA=DSQRT(DABS((R+UX)/2.0))
0005      BBB=DSQRT(DABS((R-UX)/2.0))
0006      IF(VX) 10,20,30
0007      10 UY=AAA
0008      VY=-1.0*BBB
0009      GO TO 100
0010      20 IF(UX) 40,50,60
0011      30 UY=AAA
0012      VY=BBB
0013      GO TO 100
0014      40 DUMMY=DABS(UX)
0015      UY=0.0
0016      VY=DSQRT(DUMMY)
0017      GO TO 100
0018      50 UY=0.0
0019      VY=0.0
0020      GO TO 100
0021      60 DUMMY=DABS(UX)
0022      UY=DSQRT(DUMMY)
0023      VY=0.0
0024      100 RETURN
0025      END

```

APPENDIX H

REPEATED G.C.D. - MULLER'S METHOD

1. Use of the Program

A double precision FORTRAN IV program using the repeated G.C.D. method with Muller's method as a supporting method is presented here. Flow charts for this program are given in Figure H.1 while Table H.III gives a FORTRAN IV listing of this program.

This program is designed to solve polynomials having degree less than or equal to 25. In order to solve polynomials of degree N where $N > 25$, the data statement and array dimensions given in Table H.I must be changed.

In this program both the leading coefficient and the constant coefficient are assumed to be non-zero.

TABLE H.I

PROGRAM CHANGES NECESSARY TO SOLVE POLYNOMIALS OF DEGREE
GREATER THAN 25 BY THE REPEATED G.C.D. - MULLER'S METHOD

Main Program

Data Entry/1H1,1H2,...,1H9,2H10,2H11,...,2HXX/where XX = N+1

UAPP(N,3), VAPP(N,3)
URAPP(N,3), URAPP(N,3)
UP(N+1), VP(N+1)
MULT(N)
UDD0(N+1), VDD0(N+1)
UD1(N+1), VD1(N+1)
UDD1(N+1), VDD1(N+1)
UD2(N+1), VD2(N+1)
UG(N+1), VG(N+1)
UD3(2N+1), VD3(2N+1)
UD4(2N+1), VD4(2N+1)
UAP(N+1), VAP(N+1)
UZROS(N), VZROS(N)
UROOT(N), VROOT(N)
UDO(N+1), VDO(N+1)
ENTRY(N+1)

Subroutines PROD, QUAD

See corresponding subroutine in Table G.I.

Subroutines DERIV, GCD, and DIVIDE

See corresponding subroutine in Table E.I.

Subroutines MULLER, GENAPP, BETTER and HORNER

See corresponding subroutine in Table F.I.

2. Input Data for Repeated G.C.D. - Muller's Method

The input data to the repeated G.C.D. - Muller's method is the same as for the repeated G.C.D. - Newton's method as described in Appendix G, § 2.

3. Variables Used in Repeated G.C.D. - Muller's Method

The variables used in this program are referenced in Table H.II. The notation and symbols used in the referenced tables are described in Appendix E, § 3.

TABLE H.II
VARIABLES USED IN REPEATED G.C.D. - MULLER'S METHOD

Main Program and Subroutine PROD

See Table G.II.

Subroutines QUAD, DERIV, GCD, DIVIDE, and COMSQT

See corresponding subroutine in Table E.VI.

Subroutines CALC, MULLER, GENAPP, ALTER, BETTER,
TEST, and HORNER

See corresponding subroutine in Table F.II.

4. Description of Program Output

The output for this program is the same as that for repeated G.C.D. - Newton's method as described in Appendix G, § 4. Only one initial approximation, X_0 , (not three) is printed. The other two required by Muller's method are $.9X_0$ and $1.1X_0$. The message "SOLVED BY DIRECT METHOD" means that the corresponding root was obtained by Subroutine QUAD.

5. Informative Messages and Error Messages

Descriptions of the informative messages and error messages printed by this program can be found either in Appendix E, § 5, Appendix F, § 5, or Appendix G, § 5.

MAIN PROGRAM




Figure H.1. Flow Charts for Repeated G.C.D.-Muller's Method




Figure H.1. (Continued)

MULLER





Figure H.1. (Continued)

PROD



QUAD





Figure H.1. (Continued)

DIVIDE



DERIV




Figure H.1. (Continued)





Figure H.1. (Continued)

CALC



TEST





Figure H.1. (Continued)

GENAPP



ALTER




Figure H.1. (Continued)

BETTER

HORNER




Figure H.1. (Continued)

COMSQT




Figure H.1. (Continued)

TABLE H.III

PROGRAM FOR REPEATED G.C.D.-MULLER'S METHOD

```

C *****
C *
C * DOUBLE PRECISION PROGRAM FOR THE REPEATED G.C.D. - MULLER'S METHOD
C *
C *
C * THIS METHOD REPEATEDLY FINDS THE GREATEST COMMON DIVISOR OF TWO
C * POLYNOMIALS IN ORDER TO EXTRACT THE ZEROS IN GROUPS ACCORDING TO
C * MULTIPLICITY USING NEWTON'S METHOD. ALL ZEROS OF MULTIPLICITY 1
C * ARE EXTRACTED FOLLOWED BY THOSE OF MULTIPLICITY 2, ETC.
C *
C *****
0001 DOUBLE PRECISION EPS1, EPS2, EPS3, UP, VP, UAPP, VAPP, UDO, VDO, UDDO, VDDO,
      1UD1, VD1, UD2, VD2, UDD1, VDD1, UG, VG, UD3, VD3, UD4, VD4, UZROS, VZROS, UAP, VA
      2P, UROOT, VROOT, DENOM
0002 DOUBLE PRECISION XSTART
0003 DOUBLE PRECISION XEND
0004 DOUBLE PRECISION URAPP, VRAPP
0005 DOUBLE PRECISION EPS4
0006 DIMENSION UAPP(25,3), VAPP(25,3), URAPP(25,3), VRAPP(25,3)
0007 DIMENSION UP(26), VP(26), MULT(25), UDDO(26), VDDO(26), UD1(26), VD1(26),
      1, UDD1(26), VDD1(26), UD2(26), VD2(26), UG(26), VG(26), UD3(51), VD3(51), U
      2D4(51), VD4(51), UAP(26), VAP(26), UZROS(25), VZROS(25), UROOT(25), VROOT
      3(25), ANAME(2), UDO(26), VDO(26), ENTRY(26)
0008 COMMON EPS1, EPS2, EPS3, EPS4, IO2, MAX
0009 DATA PNAME, GNAME/2HP1, 2HG1/, B1NAME/3HD1(/
0010 DATA ASTER/4H****/
0011 DATA ENTRY/1H1, 1H2, 1H3, 1H4, 1H5, 1H6, 1H7, 1H8, 1H9, 2H10, 2H11, 2H12, 2H13
      1, 2H14, 2H15, 2H16, 2H17, 2H18, 2H19, 2H20, 2H21, 2H22, 2H23, 2H24, 2H25, 2H26/
0012 DATA ANAME(1), ANAME(2)/4HMULL, 4HERS /
0013 IO1=5
0014 IO2=6
0015 1 READ(IO1, 1000) NOPOLY, NP, NAPP, MAX, EPS1, EPS2, EPS3, XSTART, XEND, KCHEC
      1K
0016 IF(KCHECK.EQ.1) STOP
0017 WRITE(IO2, 1020) ANAME(1), ANAME(2), NOPOLY
0018 WRITE(IO2, 2000) NAPP
0019 WRITE(IO2, 2010) MAX
0020 WRITE(IO2, 2070) EPS1
0021 WRITE(IO2, 2020) EPS2
0022 WRITE(IO2, 2080) EPS3
0023 WRITE(IO2, 2040) XSTART
0024 WRITE(IO2, 2050) XEND
0025 WRITE(IO2, 2060)
0026 KKK=NP+1
0027 NNN=KKK+1
0028 DO 5 I=1, KKK
0029   JJJ=NNN-I
0030 5 READ(IO1, 1010) UP(JJJ), VP(JJJ)
0031 IF(NAPP.NE.0) GO TO 22
0032 NAPP=NP
0033 CALL GENAPP(UAPP, VAPP, NAPP, XSTART)
0034 GO TO 23
0035 22 READ(IO1, 1015) (UAPP(I,2), VAPP(I,2), I=1, NAPP)
0036 23 WRITE(IO2, 1030) NP
0037 KKK=NP+1
0038 NNN=KKK+1
0039 DO 8 I=1, KKK
0040   JJJ=NNN-I

```

TABLE H.III (Continued)

```

0041      8 WRITE(102,1040) PNAME,ENTRY(JJJ),UP(JJJ),VP(JJJ)
0042      K=0
0043      KD=0
0044      J1=1
0045      KKK=NP+1
0046      DO 10 I=1,KKK
0047      UDO(I)=UP(I)
0048      10 VDO(I)=VP(I)
0049      NDO=NP
0050      CALL DERIV(NDO,UDO,VDO,NDDO,UDDO,VDDO)
0051      CALL GCD(NDO,UDO,VDO,NDDO,UDDO,VDDO,ND1,UD1,VD1)
0052      20 WRITE(102,3000) (ASTER,I=1,33)
0053      IF(ND1.LE.1) GO TO 30
0054      GO TO 40
0055      30 UD2(1)=1.0
0056      VD2(1)=0.0
0057      ND2=0
0058      GO TO 50
0059      40 CALL DERIV(ND1,UD1,VD1,NDD1,UDD1,VDD1)
0060      CALL GCD(ND1,UD1,VD1,NDD1,UDD1,VDD1,ND2,UD2,VD2)
0061      50 IF(NDO+ND2.LE.2*ND1) GO TO 60
0062      GO TO 70
0063      60 WRITE(102,1025) J1
0064      GO TO 170
0065      70 IF(ND1.EQ.0) GO TO 80
0066      GO TO 90
0067      80 KKK=NDO+1
0068      DO 85 I=1,KKK
0069      UG(I)=UDO(I)
0070      85 VG(I)=VDO(I)
0071      NG=NDO
0072      GO TO 110
0073      90 IF(ND2.EQ.0) GO TO 115
0074      CALL PROD(NDO,UDO,VDO,ND2,UD2,VD2,ND3,UD3,VD3)
0075      100 CALL PROD(ND1,UD1,VD1,ND1,UD1,VD1,ND4,UD4,VD4)
0076      CALL DIVIDE(ND3,UD3,VD3,ND4,UD4,VD4,NG,UG,VG)
0077      110 WRITE(102,1035) J1
0078      KKK=NG+1
0079      NNN=KKK+1
0080      DO 112 I=1,KKK
0081      JJJ=NNN-I
0082      112 WRITE(102,1040) GNAME,ENTRY(JJJ),UG(JJJ),VG(JJJ)
0083      KKK=NG+1
0084      DO 113 I=1,KKK
0085      UAP(I)=UG(KKK+1-I)
0086      113 VAP(I)=VG(KKK+1-I)
0087      CALL MULLER(NG,UAP,VAP,NAPP,UAPP,VAPP,J,UZROS,VZROS,JAP,XSTART,XEN
10,NOPOLY,URAPP,VRAPP)
0088      IF(J.EQ.0) GO TO 150
0089      WRITE(102,1180)
0090      IF(JAP.EQ.0) GO TO 120
0091      GO TO 130
0092      115 KKK=NDO+1
0093      DO 116 I=1,KKK
0094      UD3(I)=UDO(I)
0095      116 VD3(I)=VDO(I)
0096      ND3=NDO
0097      GO TO 100

```

TABLE H.III (Continued)

```

0098      120 KKK=JAP+1
0099      WRITE(IO2,1085) (I,UZROS(I),VZROS(I),J1,I=KKK,J)
0100      GO TO 140
0101      130 DO 135 I=1,JAP
0102      135 WRITE(IO2,1190) I,UZROS(I),VZROS(I),J1,URAPP(I,2),VRAPP(I,2)
0103      IF(JAP.LT.J) GO TO 120
0104      140 IF(J.EQ.NG) GO TO 155
0105      150 WRITE(IO2,1095)
0106      IF(J.EQ.0) GO TO 170
0107      155 DO 160 I=1,J
0108      UROOT(KD+I)=UZROS(I)
0109      VROOT(KD+I)=VZROS(I)
0110      160 MULT(KD+I)=J1
0111      K=(J*J1)+K
0112      KD=KD+J
0113      IF(K.GE.NP) GO TO 1
0114      170 J1=J1+1
0115      IF(ND1.LE.1) GO TO 200
0116      DO 180 I=1,ND1
0117      UDO(I)=UD1(I)
0118      VDO(I)=VD1(I)
0119      UDDO(I)=UDD1(I)
0120      180 VDDO(I)=VDD1(I)
0121      UDO(ND1+1)=UD1(ND1+1)
0122      VDO(ND1+1)=VD1(ND1+1)
0123      NDO=ND1
0124      NDDO=NDD1
0125      KKK=ND2+1
0126      DO 190 I=1,KKK
0127      UD1(I)=UD2(I)
0128      190 VD1(I)=VD2(I)
0129      ND1=ND2
0130      GO TO 20
0131      200 IF(ND1.EQ.0) GO TO 1
0132      KD=KD+1
0133      DENOM=UD1(2)*UD1(2)+VD1(2)*VD1(2)
0134      UROOT(KD)=(-UD1(1)*UD1(2)-VD1(1)*VD1(2))/DENOM
0135      VROOT(KD)=(-VD1(1)*UD1(2)+UD1(1)*VD1(2))/DENOM
0136      MULT(KD)=J1
0137      WRITE(IO2,3000) (ASTER,I=1,33)
0138      WRITE(IO2,1035) J1
0139      KKK=ND1+1
0140      NNN=KKK+1
0141      DO 210 I=1,KKK
0142      JJJ=NNN-I
0143      210 WRITE(IO2,1100) D1NAME,ENTRY(JJJ),UD1(JJJ),VD1(JJJ)
0144      WRITE(IO2,1180)
0145      WRITE(IO2,1085) KD,UROOT(KD),VROOT(KD),J1
0146      GO TO 1
0147      1020 FORMAT(1H1,10X,48HREPEATED USE OF THE GREATEST COMMON DIVISOR AND
1A4,A4,58H METHOD TO EXTRACT ROOTS AND MULTIPLICITIES OF POLYNOMIAL
2LS/11X,18HPOLYNOMIAL NUMBER ,I2///)
0148      1025 FORMAT(///1X,25HNO ROOTS OF MULTIPLICITY ,I2//)
0149      1035 FORMAT(///1X,87HTHE FOLLOWING POLYNOMIAL, G(X), CONTAINS ALL THE R
10OTS OF P(X) WHICH HAVE MULTIPLICITY ,I2//)
0150      1085 FORMAT(2X,5HROOT(I,I2,4H) = ,D23.16,3H + ,D23.16,2H I,8X,I2,9X,25HN
10 INITIAL APPROXIMATIONS)
0151      1095 FORMAT(///1X,51HNOT ALL ROOTS OF THE ABOVE POLYNOMIAL,G, WERE FOUN

```

TABLE H.III (Continued)

```

10//)
0152 1000 FORMAT(3(I2,1X),9X,I3,1X,3(D6.0,1X),20X,2(D7.0,1X),I1)
0153 1010 FORMAT(2D30.0)
0154 1015 FORMAT(2D30.0)
0155 1030 FORMAT(1X,22H THE DEGREE OF P(X) IS ,I2,22H THE COEFFICIENTS ARE//
1)
0156 1040 FORMAT(2X,A2,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0157 1100 FORMAT(2X,A3,A2,4H) = ,D23.16,3H + ,D23.16,2H I)
0158 1180 FORMAT(///1X,13HROOTS OF P(X),52X,14HMULTIPLICITIES,17X,21HINITIAL
1 APPROXIMATION//)
0159 1190 FORMAT(2X,5HROOT(,I2,4H) = ,D23.16,3H + ,D23.16,2H I,8X,I2,8X,D23.
116,3H + ,D23.16,2H I)
0160 2000 FORMAT(1X,41HNUMBER OF INITIAL APPROXIMATIONS GIVEN. ,I2)
0161 2010 FORMAT(1X,29HMAXIMUM NUMBER OF ITERATIONS.,11X,I3)
0162 2020 FORMAT(1X,21HTEST FOR CONVERGENCE.,13X,D9.2)
0163 2040 FORMAT(1X,23HRADIUS TO START SEARCH.,11X,D9.2)
0164 2050 FORMAT(1X,21HRADIUS TO END SEARCH.,13X,D9.2)
0165 2060 FORMAT(//1X)
0166 2070 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE GCD. ,D9.2)
0167 2080 FORMAT(1X,34HTEST FOR ZERO IN SUBROUTINE QUAD. ,D9.2)
0168 3000 FORMAT(/////1X,A3,32A4)
0169 END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE PROD(M,UF,VF,N,UG,VG,MN,UH,VH)
C      *****
C      *
C      * GIVEN POLYNOMIALS R(X) AND S(X), THIS SUBROUTINE COMPUTES THE
C      * COEFFICIENTS OF THE PRODUCT POLYNOMIAL T(X) = R(X).S(X).
C      *
C      *****
0002      DOUBLE PRECISION UH,VH,UF,VF,UG,VG
0003      DIMENSION UH(51),VH(51),UF(26),VF(26),UG(26),VG(26)
0004      MN=M+N
0005      KKK=MN+1
0006      DO 100 I=1,KKK
0007          K=I
0008          UH(I)=0.0
0009          VH(I)=0.0
0010          IF(I.LE.M+1) GO TO 10
0011          LIMIT=M+1
0012          GO TO 20
0013      10  LIMIT=I
0014      20  DO 50 J=1,LIMIT
0015          IF(K.GT.N+1) GO TO 50
0016          IF(J+K.EQ.I+1) GO TO 40
0017          GO TO 50
0018      40  UH(I)=UH(I)+(UF(J)*UG(K)-VF(J)*VG(K))
0019          VH(I)=VH(I)+(VF(J)*UG(K)+UF(J)*VG(K))
0020      50  K=K-1
0021      100 CONTINUE
0022      RETURN
0023      END

```


TABLE H.III. (Continued)

```

0001      SUBROUTINE QUAD(N,UA,VA,J,UROOT,VROOT)
C      *
C      * SUBROUTINE QUAD SOLVES DIRECTLY FOR THE ZEROS AND THEIR MULTIPLICITIES *
C      * OF EITHER A QUADRATIC POLYNOMIAL OR A LINEAR FACTOR. SOLUTION OF THE *
C      * QUADRATIC IS DONE USING THE QUADRATIC FORMULA. *
C      *
C      *
0002      DOUBLE PRECISION EPS1,EPS2,EPSLON,UROOT,VROOT,UA,VA,UDISC,VDISC,UD
0003      1,VD,DDD,UTEMP,VTEMP,BBB
0004      DOUBLE PRECISION EPS4
0005      DIMENSION UROOT(25),VROOT(25),UA(26),VA(26)
0006      COMMON EPS1,EPS2,EPSLON,EPS4,IO2,MAX
0007      IF(N.GT.1) GO TO 10
0008      J=J+1
0009      BBB=UA(2)*UA(2)+VA(2)*VA(2)
0010      UROOT(J)=-{UA(1)*UA(2)+VA(1)*VA(2)}/BBB
0011      VROOT(J)=-{VA(1)*UA(2)-UA(1)*VA(2)}/BBB
0012      GO TO 100
0013      10 UDISC={UA(2)*UA(2)-VA(2)*VA(2)}-{4.0*{UA(3)*UA(1)-VA(3)*VA(1)}}
0014      VDISC={2.0*UA(2)*VA(2)}-{4.0*{UA(3)*VA(1)+VA(3)*UA(1)}}
0015      UD=2.0*UA(3)
0016      VD=2.0*VA(3)
0017      DDD=DSQRT(UDISC*UDISC+VDISC*VDISC)
0018      IF(DDD.LT.EPSLON) GO TO 20
0019      CALL COMSQT(UDISC,VDISC,UTEMP,VTEMP)
0020      BBB=UD*UD+VD*VD
0021      UROOT(J+1)={(-UA(2)+UTEMP)*UD+(-VA(2)+VTEMP)*VD}/BBB
0022      VROOT(J+1)={(-VA(2)+VTEMP)*UD-(-UA(2)+UTEMP)*VD}/BBB
0023      UROOT(J+2)={(-UA(2)-UTEMP)*UD+(-VA(2)-VTEMP)*VD}/BBB
0024      VROOT(J+2)={(-VA(2)-VTEMP)*UD-(-UA(2)-UTEMP)*VD}/BBB
0025      J=J+2
0026      GO TO 100
0027      20 J=J+1
0028      BBB=UD*UD+VD*VD
0029      UROOT(J)={-UA(2)*UD-VA(2)*VD}/BBB
0030      VROOT(J)={-VA(2)*UD+UA(2)*VD}/BBB
0031      WRITE(IO2,1000) UROOT(J),VROOT(J)
1000  FORMAT(//1X,11HQUAD FOUND ,D23.16,3H + ,D23.16,2H I,22H TO BE A M
1001  ULTIPLE ROOT//)
0032      100 RETURN
0033      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE DERIV(N,UP,VP,M,UA,VA)
C      *****
C      *
C      * GIVEN A POLYNOMIAL P(X), SUBROUTINE DERIV COMPUTES THE COEFFICIENTS OF
C      * ITS DERIVATIVE P'(X).
C      *
C      *****
0002      DOUBLE PRECISION UP,VP,UA,VA,AAA
0003      DIMENSION UP(26),VP(26),UA(26),VA(26)
0004      KKK=N+1
0005      DO 10 I=2,KKK
0006      AAA=I-1
0007      UA(I-1)=AAA*UP(I)
0008      10 VA(I-1)=AAA*VP(I)
0009      M=N-1
0010      RETURN
0011      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE GCD(N,UR,VR,M,US,VS,M1,USS,VSS)
C      *****
C      *
C      * GIVEN POLYNOMIALS P(X) AND DP(X) WHERE DEG. DP(X) IS LESS THAN DEG.
C      * P(X), SUBROUTINE GCD COMPUTES THE GREATEST COMMON DIVISOR OF P(X) AND
C      * DP(X).
C      *
C      *****
0002      DOUBLE PRECISION USSSSS,VSSSSS
0003      DOUBLE PRECISION UR,VR,US,VS,USS,VSS,URR,VRR,UD,VD,UT,VT,EPSLON,EP
0004      1S2,EPS3,EPS4,BBB
0004      DIMENSION UR(26),VR(26),US(26),VS(26),USS(26),VSS(26),URR(26),VRR(
0005      126),UT(26),VT(26)
0005      COMMON EPSLON,EPS2,EPS3,EPS4,IO2,MAX
0006      N1=N
0007      M1=M
0008      KKK=N+1
0009      DO 20 I=1,KKK
0010      URR(I)=UR(I)
0011      20 VRR(I)=VR(I)
0012      KKK=M+1
0013      DO 25 I=1,KKK
0014      USS(I)=US(I)
0015      25 VSS(I)=VS(I)
0016      30 BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0017      UD=(URR(N1+1)*USS(M1+1)+VRR(N1+1)*VSS(M1+1))/BBB
0018      VD=(USS(M1+1)*VRR(N1+1)-URR(N1+1)*VSS(M1+1))/BBB
0019      KKK=N1+1-M1
0020      DO 40 I=KKK,N1
0021      UT(I)=URR(I)-(UD*USS(I-N1+M1)-VD*VSS(I-N1+M1))
0022      40 VT(I)=VRR(I)-(UD*VSS(I-N1+M1)+VD*USS(I-N1+M1))
0023      IF(M1.EQ.N1) GO TO 70
0024      KKK=N1-M1
0025      DO 60 I=1,KKK
0026      UT(I)=URR(I)
0027      60 VT(I)=VRR(I)
0028      70 DO 90 I=1,N1
0029      BBB=DSQRT(UT(N1+1-I)*UT(N1+1-I)+VT(N1+1-I)*VT(N1+1-I))
0030      IF(BBB.GT.EPSLON) GO TO 100
0031      90 CONTINUE
0032      DO 95 I=1,M1
0033      BBB=USS(M1+1)*USS(M1+1)+VSS(M1+1)*VSS(M1+1)
0034      USSSSS=(USS(I)*USS(M1+1)+VSS(I)*VSS(M1+1))/BBB
0035      VSSSSS=(VSS(I)*USS(M1+1)-USS(I)*VSS(M1+1))/BBB
0036      USS(I)=USSSSS
0037      95 VSS(I)=VSSSSS
0038      USS(M1+1)=1.0
0039      VSS(M1+1)=0.0
0040      GO TO 200
0041      100 K=N1-I
0042      IF(K.EQ.0) GO TO 170
0043      IF(K.LT.M1) GO TO 140
0044      KKK=K+1
0045      DO 130 J=1,KKK
0046      URR(J)=UT(J)
0047      130 VRR(J)=VT(J)
0048      N1=K
0049      GO TO 30

```

TABLE H.III (Continued)

```

0050      140 KKK=K+1
0051          DO 150 J=1, KKK
0052              URR(J)=USS(J)
0053              VRR(J)=VSS(J)
0054              USS(J)=UT(J)
0055      150 VSS(J)=VT(J)
0056          KKK=K+2
0057          NNN=M1+1
0058          DO 160 J=KKK, NNN
0059              URR(J)=USS(J)
0060      160 VRR(J)=VSS(J)
0061          M1=M1
0062          M1=K
0063          GO TO 30
0064      170 USS(1)=1.0
0065          VSS(1)=0.0
0066          M1=0
0067      200 RETURN
0068          END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE DIVIDE(N,UP,VP,M,UD,VD,K,UQ,VQ)
C      *
C      *
C      * GIVEN TWO POLYNOMIALS F(X) AND G(X), SUBROUTINE DIVIDE COMPUTES THE
C      * QUOTIENT POLYNOMIAL H(X) = F(X)/G(X).
C      *
C      *
0002      DOUBLE PRECISION UP,VP,UD,VD,UQ,VQ,UTERM,VTerm,UDUMMY
0003      DIMENSION UP(26),VP(26),UD(26),VD(26),UQ(26),VQ(26)
0004      K=N-M
0005      UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0006      UQ(K+1)=(UP(N+1)*UD(M+1)+VP(N+1)*VD(M+1))/UDUMMY
0007      VQ(K+1)=(VP(N+1)*UD(M+1)-UP(N+1)*VD(M+1))/UDUMMY
0008      IF(K.EQ.0) GO TO 100
0009      J=-1
0010      DO 50 I=1,K
0011      J=J+1
0012      UTERM=UP(N-J)
0013      VTERM=VP(N-J)
0014      KK=K+1
0015      NNN=M-J
0016      DO 40 M1=NNN,M
0017      IF(KK.GT.1) GO TO 10
0018      GO TO 45
0019      10 IF(M1.GE.1) GO TO 20
0020      GO TO 40
0021      20 UTERM=UTERM-(UQ(KK)*UD(M1)-VQ(KK)*VD(M1))
0022      VTERM=VTERM-(UQ(KK)*VD(M1)+VQ(KK)*UD(M1))
0023      40 KK=KK-1
0024      45 UDUMMY=UD(M+1)*UD(M+1)+VD(M+1)*VD(M+1)
0025      UQ(K+1-I)=(UTERM*UD(M+1)+VTERM*VD(M+1))/UDUMMY
0026      50 VQ(K+1-I)=(VTERM*UD(M+1)-UTERM*VD(M+1))/UDUMMY
0027      100 RETURN
0028      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE CONSQT(UX,VX,UY,VY)
C      *****
C      *
C      * THIS SUBROUTINE COMPUTES THE SQUARE ROOT OF A COMPLEX NUMBER.
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UY,VY,DUMMY,R,AAA,BBB
0003      R=DSQRT(UX*UX+VX*VX)
0004      AAA=DSQRT(DABS((R+UX)/2.0))
0005      BBB=DSQRT(DABS((R-UX)/2.0))
0006      IF(VX) 10,20,30
0007      10 UY=AAA
0008      VY=-1.0*BBB
0009      GO TO 100
0010      20 IF(UX) 40,50,60
0011      30 UY=AAA
0012      VY=BBB
0013      GO TO 100
0014      40 DUMMY=DABS(UX)
0015      UY=0.0
0016      VY=DSQRT(DUMMY)
0017      GO TO 100
0018      50 UY=0.0
0019      VY=0.0
0020      GO TO 100
0021      60 DUMMY=DABS(UX)
0022      UY=DSQRT(DUMMY)
0023      VY=0.0
0024      100 RETURN
0025      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,V
          1PX3,UX4,VX4,UQ4,VQ4,UH3,VH3)
C      *****
C      *
C      * GIVEN THREE APPROXIMATIONS X(N-2), X(N-1), AND X(N), SUBROUTINE CALC
C      * APPROXIMATES THE POLYNOMIAL BY A QUADRATIC AND SOLVES FOR THE ZERO OF
C      * THE QUADRATIC CLOSEST TO X(N). THIS ZERO IS THE NEW APPROXIMATION
C      * X(N+1) TO THE ZERO OF THE POLYNOMIAL.
C      *
C      *****
0002      DOUBLE PRECISION ARG1,ARG2
0003      DOUBLE PRECISION UPX3,VPX3,UPX2,VPX2,UX1,VX1,UX2,VX2,UX3,VX3,UPX1,
          1VPX1,UH3,VH3,UH2,VH2,UQ3,VQ3,UD,VD,UB,VB,UC,VC,UDISC,VDISC,UCCC,VC
          2CC,UDEN1,VDEN1,UDEN2,VDEN2,UQ4,VQ4,UX4,VX4,EPSRT,EPS0,EPS,UDDD,VDD
          3D,AAA,BBB,RAD,UAAA,VAAA,UBBB,VBBB
0004      DOUBLE PRECISION THETA,ANGLE,UTEST,VTEST
0005      DOUBLE PRECISION EPS1
0006      COMMON EPS1,EPS,EPS0,EPSRT,I02,MAX
0007      UH3=UX3-UX2
0008      VH3=VX3-VX2
0009      UH2=UX2-UX1
0010      VH2=VX2-VX1
0011      BBB=UH2*UH2+VH2*VH2
0012      UQ3=(UH3*UH2+VH3*VH2)/BBB
0013      VQ3=(VH3*UH2-UH3*VH2)/BBB
0014      UDDD=1.0+UQ3
0015      VDDD=VQ3
0016      UD=(UPX3-(UDDD*UPX2-VDDD*VPX2))+{(UQ3*UPX1-VQ3*VPX1)}
0017      VD=(VPX3-(VDDD*UPX2+UDDD*VPX2))+{(VQ3*UPX1+UQ3*VPX1)}
0018      UAAA=2.0*UQ3
0019      VAAA=2.0*VQ3
0020      UAAA=UAAA+1.0
0021      UBBB=UDDD*UDDD-VDDD*VDDD
0022      VBBB=VDDD*UDDD+UDDD*VDDD
0023      UCCC=UQ3*UQ3-VQ3*VQ3
0024      VCCC=VQ3*UQ3+UQ3*VQ3
0025      UB=(UAAA*UPX3-VAAA*VPX3)-(UBBB*UPX2-VBBB*VPX2))+{UCCC*UPX1-VCCC*V
          1PX1}
0026      VB=(VAAA*UPX3+UAAA*VPX3)-(VBBB*UPX2+UBBB*VPX2))+{VCCC*UPX1+UCCC*V
          1PX1}
0027      UC=UDDD*UPX3-VDDD*VPX3
0028      VC=VDDD*UPX3+UDDD*VPX3
0029      UDISC=(UB*UB-VB*VB)-(4.0*(UD*UC-VD*VC))
0030      VDISC=(2.0*(VB*UB))-(4.0*(VD*UC+UD*VC))
0031      AAA=DSQRT(UDISC*UDISC+VDISC*VDISC)
0032      IF(AAA.EQ.0.0) GO TO 5
0033      GO TO 7
0034      5 THETA=0.0
0035      GO TO 9
0036      7 THETA=ATAN2(VDISC,UDISC)
0037      9 RAD=DSQRT(AAA)
0038      ANGLE=THETA/2.0
0039      UTEST=RAD*DCOS(ANGLE)
0040      VTEST=RAD*DSIN(ANGLE)
0041      UDEN1=UB+UTEST
0042      VDEN1=VB+VTEST
0043      UDEN2=UB-UTEST
0044      VDEN2=VB-VTEST

```

TABLE H.III (Continued)

```

0045      ARG1=UDEN1*UDEN1+VDEN1*VDEN1
0046      ARG2=UDEN2*UDEN2+VDEN2*VDEN2
0047      AAA=DSQRT(ARG1)
0048      BBB=DSQRT(ARG2)
0049      IF(AAA.LT.BBB) GO TO 10
0050      IF(AAA.EQ.0.0) GO TO 60
0051      UAAA=-2.0*UC
0052      VAAA=-2.0*VC
0053      UQ4=(UAAA*UDEN1+VAAA*VDEN1)/ARG1
0054      VQ4=(VAAA*UDEN1-UAAA*VDEN1)/ARG1
0055      GO TO 50
0056 10 IF(BBB.EQ.0.0) GO TO 60
0057      UAAA=-2.0*UC
0058      VAAA=-2.0*VC
0059      UQ4=(UAAA*UDEN2+VAAA*VDEN2)/ARG2
0060      VQ4=(VAAA*UDEN2-UAAA*VDEN2)/ARG2
0061      GO TO 50
0062 50 UX4=UX3+(UH3*UQ4-VH3*VQ4)
0063      VX4=VX3+(VH3*UQ4+UH3*VQ4)
0064      RETURN
0065 60 UQ4=1.0
0066      VQ4=0.0
0067      GO TO 50
0068      END

```


TABLE H.III (Continued)

```

0001      SUBROUTINE MULLER(NP,UA,VA,NAPP,UAPP,VAPP,NROOT,UROOT,VROOT,IROOT,
          1XSTART,XEND,NOPOLY,URAPP,VRAPP)
C      *****
C      *
C      * MULLER'S METHOD EXTRACTS THE ZEROS AND THEIR MULTIPLICITIES OF A
C      * POLYNOMIAL OF MAXIMUM DEGREE 25. THROUGH THREE GIVEN POINTS THE
C      * POLYNOMIAL IS APPROXIMATED BY A QUADRATIC. THE ZERO OF THE QUADRATIC
C      * CLOSEST TO THE OLD APPROXIMATION IS TAKEN AS THE NEW APPROXIMATION.
C      * IN THIS MANNER A SEQUENCE IS OBTAINED CONVERGING TO A ZERO.
C      *
C      *****
0002      DOUBLE PRECISION UPX3,VPX3,UPX2,VPX2,UROOT,VROOT,UX1,VX1,UAPP,VAPP
          1,UX2,VX2,UWORK,VWORK,UX3,VX3,UB,VB,UX4,VX4,UA,VA,UPX1,VPX1,URAPP,V
          2RAPP,UPX4,VPX4,EPSRT,EPSD,EPS,CCC,EPSM,UH3,VH3,UQ4,VQ4,ABPX4,ABPX3
          3,QQQ,XSTART,XEND
0003      DIMENSION UROOT(25),VROOT(25),MULT(25),UAPP(25,3),VAPP(25,3),UWORK
          1(26),VWORK(26),UB(26),VB(26),UA(26),VA(26),URAPP(25,3),VRAPP(25,3)
0004      LOGICAL CONV
0005      COMMON EPSM,EPS,EPSD,EPSRT,IQ2,MAX
0006      DATA PNAME,DNAME/2HP(,2HD(/
0007      EPSM=0.0D00
0008      EPSRT=0.999
0009      NROOT=0
0010      IROOT=0
0011      IPATH=1
0012      NOMULT=0
0013      NALTER=0
0014      ITIME=0
0015      IAPP=1
0016      ITER=1
0017      IF(NAPP.NE.0) GO TO 18
0018      NAPP=NP
0019      CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0020      GO TO 27
0021      18 DO 25 I=1,NAPP
0022          UAPP(I,1)=0.9*UAPP(I,2)
0023          VAPP(I,1)=0.9*VAPP(I,2)
0024          UAPP(I,3)=1.1*UAPP(I,2)
0025          VAPP(I,3)=1.1*VAPP(I,2)
0026      27 KKK=NP+1
0027          DO 30 I=1,KKK
0028              UWORK(I)=UA(I)
0029              VWORK(I)=VA(I)
0030      30 NWORK=NP
0031      40 UX1=UAPP(IAPP,1)
0032          VX1=VAPP(IAPP,1)
0033          UX2=UAPP(IAPP,2)
0034          VX2=VAPP(IAPP,2)
0035          UX3=UAPP(IAPP,3)
0036          VX3=VAPP(IAPP,3)
0037      CALL HORNER(NWORK,UWORK,VWORK,UX1,VX1,UB,VB,UPX1,VPX1)
0038      CALL HORNER(NWORK,UWORK,VWORK,UX2,VX2,UB,VB,UPX2,VPX2)
0039      CALL HORNER(NWORK,UWORK,VWORK,UX3,VX3,UB,VB,UPX3,VPX3)
0040      50 CALL CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UX
          14,VX4,UQ4,VQ4,UH3,VH3)
0041      60 CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0042          ABPX4=DSQRT(UPX4*UPX4+VPX4*VPX4)
0043          ABPX3=DSQRT(UPX3*UPX3+VPX3*VPX3)

```

TABLE H.III (Continued)

```

0044      IF(ABPX3.EQ.0.0) GO TO 70
0045      QQQ=ABPX4/ABPX3
0046      IF(QQQ.LE.10.) GO TO 70
0047      UQ4=0.5*UQ4
0048      VQ4=0.5*VQ4
0049      UX4=UX3+(UH3*UQ4-VH3*VQ4)
0050      VX4=VX3+(VH3*UQ4+UH3*VQ4)
0051      GO TO 60
0052  70 CALL TEST(UX3,VX3,UX4,VX4,CONV)
0053      IF(CONV) GO TO 120
0054      IF(ITER.LT.MAX) GO TO 110
0055      CALL ALTER(UAPP(IAPP,1),VAPP(IAPP,1),UAPP(IAPP,2),VAPP(IAPP,2),UAP
1P(IAPP,3),VAPP(IAPP,3),NALTER,ITIME)
      IF(NALTER.GT.5) GO TO 75
      ITER=1
      GO TO 40
0056
0057  75 IF(IAPP.LT.NAPP) GO TO 100
0058      IF(XEND.EQ.0.0) GO TO 77
0059      IF(XSTART.GT.XEND) GO TO 77
0060      NAPP=NP
0061      CALL GENAPP(UAPP,VAPP,NAPP,XSTART)
0062      IAPP=0
0063      GO TO 100
0064
0065  77 WRITE(IO2,1090)
0066      KKK=NWORK+1
0067      WRITE(IO2,1035) (DNAME,J,UWORK(J),VWORK(J),J=1,KKK)
0068
0069  80 IF(NROOT.EQ.0) GO TO 90
0070      IF(IPATH.EQ.1) GO TO 82
0071  81 IPATH=2
0072      CALL BETTER(UA,VA,NP,UROOT,VROOT,NROOT,URAPP,VRAPP,IROOT,MULT)
0073      RETURN
0074
0075  82 IF(NROOT.EQ.0) GO TO 90
0076      IF(IROOT.EQ.0) GO TO 85
0077      WRITE(IO2,1080)
0078      DO 55 I=1,IROOT
0079  55 WRITE(IO2,1085) I,UROOT(I),VROOT(I),URAPP(I,2),VRAPP(I,2)
0080      IF(IROOT.LT.NROOT) GO TO 85
0081      GO TO 87
0082  85 KKK=IROOT+1
0083      WRITE(IO2,1086) (I,UROOT(I),VROOT(I),I=KKK,NROOT)
0084  87 IF(IPATH.EQ.1) GO TO 81
0085      RETURN
0086
0087  90 WRITE(IO2,1070) NOPOLY
0088      RETURN
0089
0090  100 IAPP=IAPP+1
0091      ITER=1
0092      NALTER=0
0093      GO TO 40
0094
0095  120 NROOT=NROOT+1
0096      IROOT=NROOT
0097      MULT(NROOT)=1
0098      NOMULT=NOMULT+1
0099      UROOT(NROOT)=UX4
0100      VROOT(NROOT)=VX4
      URAPP(NROOT,1)=UAPP(IAPP,1)
      VRAPP(NROOT,1)=VAPP(IAPP,1)
      URAPP(NROOT,2)=UAPP(IAPP,2)
      VRAPP(NROOT,2)=VAPP(IAPP,2)

```

TABLE H.III (Continued)

```

0101      URAPP(NROOT,3)=UAPP(IAPP,3)
0102      VRAPP(NROOT,3)=VAPP(IAPP,3)
0103      125 IF(NOMULT.LT.NP) GO TO 130
0104      GO TO 80
0105      130 CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0106      NWORK=NWORK-1
0107      KKK=NWORK+1
0108      DO 140 I=1,KKK
0109      UWORK(I)=UB(I)
0110      140 VWORK(I)=VB(I)
0111      CALL HORNER(NWORK,UWORK,VWORK,UX4,VX4,UB,VB,UPX4,VPX4)
0112      CCC=DSQRT(UPX4*UPX4+VPX4*VPX4)
0113      IF(CCC.LT.EPSM) GO TO 150
0114      IF(NWORK.GT.2) GO TO 75
0115      IROOT=NROOT
0116      KKK=NWORK+1
0117      DO 145 I=1,KKK
0118      UB(I)=UWORK(KKK+1-I)
0119      145 VB(I)=VWORK(KKK+1-I)
0120      CALL QUAD(NWORK,UB,VB,NROOT,UROOT,VROOT)
0121      GO TO 80
0122      150 MULT(NROOT)=MULT(NROOT)+1
0123      NOMULT=NOMULT+1
0124      GO TO 125
0125      110 UX1=UX2
0126      VX1=VX2
0127      UX2=UX3
0128      VX2=VX3
0129      UX3=UX4
0130      VX3=VX4
0131      UPX1=UPX2
0132      VPX1=VPX2
0133      UPX2=UPX3
0134      VPX2=VPX3
0135      UPX3=UPX4
0136      VPX3=VPX4
0137      ITER=ITER+1
0138      GO TO 50
0139      1090 FORMAT(///,1X,65HCOEFFICIENTS OF DEFLATED POLYNOMIAL FOR WHICH NO
1ZEROS WERE FOUND//)
0140      1080 FORMAT(///1X,13HROOTS OF G(X),83X,21HINITIAL APPROXIMATION//)
0141      1070 FORMAT(///,43H NO ZEROS WERE FOUND FOR POLYNOMIAL NUMBER ,I2)
0142      1086 FORMAT(2X,5HROOT(I,12,4H) = ,D23.16,3H + ,D23.16,2H I,19X,23HSOLVED
1 BY DIRECT METHOD)
0143      1035 FORMAT(3X,A2,I2,4H) = ,D23.16,3H + ,D23.16,2H I)
0144      1085 FORMAT(2X,5HROOT(I,12,4H) = ,D23.16,3H + ,D23.16,2H I,18X,D23.16,3H
1 + ,D23.16,2H I)
0145      1000 FORMAT(3(I2,1X),9X,13,8X,3(D6.0,1X),13X,2(D7.0,1X),11)
0146      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE GENAPP(APPR,APPI,NAPP,XSTART)
C      *****
C      *
C      * SUBROUTINE GENAPP GENERATES N INITIAL APPROXIMATIONS, WHERE N IS THE *
C      * DEGREE OF THE ORIGINAL POLYNOMIAL. *
C      *
C      *****
0002      DOUBLE PRECISION APPR,APPI,XSTART,EPS1,EPS2,EPS3,BETA
0003      DOUBLE PRECISION EPSM
0004      DIMENSION APPR(25,3),APPI(25,3)
0005      COMMON EPSM,EPS1,EPS2,EPS3,IO2,MAX
0006      IF(XSTART.EQ.0.0) XSTART=0.5
0007      BETA=0.2617994
0008      DO 10 I=1,NAPP
0009      APPR(I,2)=XSTART*DCOS(BETA)
0010      APPI(I,2)=XSTART*DSIN(BETA)
0011      BETA=BETA+0.5235988
0012      10 XSTART=XSTART+0.5
0013      DO 20 I=1,NAPP
0014      APPR(I,1)=0.9*APPR(I,2)
0015      APPI(I,1)=0.9*APPI(I,2)
0016      APPR(I,3)=1.1*APPR(I,2)
0017      20 APPI(I,3)=1.1*APPI(I,2)
0018      RETURN
0019      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE ALTER(X1R,X1I,X2R,X2I,X3R,X3I,NALTER,ITIME)
C      *
C      * SUBROUTINE ALTER ALTERS THE INITIAL APPROXIMATIONS WHICH PRODUCE NO
C      * CONVERGENCE TO A ZERO. THIS IS DONE A MAXIMUM OF 5 TIMES FOR EACH ROOT.
C      *
C      *
0002      DOUBLE PRECISION X1R,X1I,X2R,X2I,X3R,X3I,EPS1,EPS2,EPS3,R,BETA
0003      DOUBLE PRECISION EPSM
0004      COMMON EPSM, EPS1, EPS2, EPS3, IO2, MAX
0005      IF(ITIME.NE.0) GO TO 5
0006      ITIME=1
0007      WRITE(IO2,1010) MAX
0008      5 IF(NALTER.EQ.0) GO TO 10
0009      WRITE(IO2,1000) X1R,X1I,X2R,X2I,X3R,X3I
0010      GO TO 20
0011      10 R=DSQRT(X2R*X2R+X2I*X2I)
0012      BETA=DATAN2(X2I,X2R)
0013      WRITE(IO2,1020) X1R,X1I,X2R,X2I,X3R,X3I
0014      20 NALTER=NALTER+1
0015      IF(NALTER.GT.5) RETURN
0016      GO TO (30,40,30,40,30),NALTER
0017      30 X2R=-X2R
0018      X2I=-X2I
0019      GO TO 50
0020      40 BETA=BETA+1.0471976
0021      X2R=R*DCOS(BETA)
0022      X2I=R*DSIN(BETA)
0023      50 X1R=0.9*X2R
0024      X1I=0.9*X2I
0025      X3R=1.1*X2R
0026      X3I=1.1*X2I
0027      RETURN
0028      1000 FORMAT(1X,5HX1 = ,D23.16,3H + ,D23.16,2H I,10X,22HALTERED APPROXIM
1ATIONS/1X,5HX2 = ,D23.16,3H + ,D23.16,2H I/1X,5HX3 = ,D23.16,3H +
2,D23.16,2H I/)
0029      1020 FORMAT(1H0,5HX1 = ,D23.16,3H + ,D23.16,2H I,10X,22INITIAL APPROXIM
1ATIONS/1X,5HX2 = ,D23.16,3H + ,D23.16,2H I/1X,5HX3 = ,D23.16,3H +
2 ,D23.16,2H I/)
0030      1010 FORMAT(///1X,54HNO CONVERGENCE FOR THE FOLLOWING APPROXIMATIONS AF
1TER ,I3,12H ITERATIONS.//)
0031      END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE BETTER(UA,VA,NP,UROOT,VROOT,NROOT,URAPP,VRAPP,IROOT,MUL
          1T)
C          *****
C          *
C          * SUBROUTINE BETTER ATTEMPTS TO IMPROVE THE ACCURACY OF THE ZEROS FOUND *
C          * BY USING THEM AS INITIAL APPROXIMATIONS WITH MULLER'S METHOD APPLIED TO *
C          * THE FULL, UNDEFLATED POLYNOMIAL. *
C          *
0002      *****
          DOUBLE PRECISION UROOT,VROOT,UA,VA,UBAPP,VBAPP,UX1,VX1,UX2,VX2,UX3
          1,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UB,VB,UROOTS,VROOTS,EPSRT,UX4,V
          2X4,URAPP,VRAPP,EPSO,EPS,UQ4,VQ4,UH3,VH3
0003      DOUBLE PRECISION EPSM
0004      LOGICAL CONV
0005      DIMENSION UROOT(25),VROOT(25),UA(26),VA(26),UBAPP(25,3),VBAPP(25,3
          1),UB(26),VB(26),UROOTS(25),VROOTS(25),URAPP(25,3),VRAPP(25,3),MULT
          3(25)
0006      COMMON EPSM,EPS,EPSO,EPSRT,I02,MAX
0007      IF(NROOT.LE.1) RETURN
0008      L=0
0009      DO 10 I=1,NROOT
0010          UBAPP(I,1)=UROOT(I)*EPSRT
0011          VBAPP(I,1)=VROOT(I)*EPSRT
0012          UBAPP(I,2)=UROOT(I)
0013          VBAPP(I,2)=VROOT(I)
0014          UBAPP(I,3)=UROOT(I)*(2.0-EPSRT)
0015      10 VBAPP(I,3)=VROOT(I)*(2.0-EPSRT)
0016      DO 100 J=1,NROOT
0017          UX1=UBAPP(J,1)
0018          VX1=VBAPP(J,1)
0019          UX2=UBAPP(J,2)
0020          VX2=VBAPP(J,2)
0021          UX3=UBAPP(J,3)
0022          VX3=VBAPP(J,3)
0023      ITER=1
0024      CALL HORNER(NP,UA,VA,UX1,VX1,UB,VB,UPX1,VPX1)
0025      CALL HORNER(NP,UA,VA,UX2,VX2,UB,VB,UPX2,VPX2)
0026      20 CALL HORNER(NP,UA,VA,UX3,VX3,UB,VB,UPX3,VPX3)
0027      CALL CALC(UX1,VX1,UX2,VX2,UX3,VX3,UPX1,VPX1,UPX2,VPX2,UPX3,VPX3,UX
          14,VX4,UQ4,VQ4,UH3,VH3)
0028      30 CALL TEST(UX3,VX3,UX4,VX4,CONV)
0029      IF(CONV) GO TO 50
0030      IF(ITER.LT.MAX) GO TO 40
0031      WRITE(I02,1000) J,UROOT(J),VROOT(J),MAX
0032      WRITE(I02,1010) UX4,VX4
0033      IF(J.LT.IROOT) GO TO 33
0034      IF(J.EQ.IROOT) GO TO 35
0035      GO TO 100
0036      33 KKK=IROOT-1
0037      DO 34 K=J,KKK
0038          URAPP(K,1)=URAPP(K+1,1)
0039          VRAPP(K,1)=VRAPP(K+1,1)
0040          URAPP(K,2)=URAPP(K+1,2)
0041          VRAPP(K,2)=VRAPP(K+1,2)
0042          URAPP(K,3)=URAPP(K+1,3)
0043      34 VRAPP(K,3)=VRAPP(K+1,3)
0044      35 IROOT=IROOT-1
0045      GO TO 100

```

TABLE H.III. (Continued)

```

0046      40 UX1=UX2
0047          VX1=VX2
0048          UX2=UX3
0049          VX2=VX3
0050          UX3=UX4
0051          VX3=VX4
0052          UPX1=UPX2
0053          VPX1=VPX2
0054          UPX2=UPX3
0055          VPX2=VPX3
0056          ITER=ITER+1
0057          GO TO 20
0058      50 L=L+1
0059          UROOTS(L)=UX4
0060          VROOTS(L)=VX4
0061      100 CONTINUE
0062          IF(L.EQ.0) GO TO 120
0063          DO 110 I=1,L
0064              UROOT(I)=UROOTS(I)
0065          110 VROOT(I)=VROOTS(I)
0066          NROOT=L
0067          RETURN
0068      120 NROOT=0
0069          RETURN
0070      1000 FORMAT(///42H IN THE ATTEMPT TO IMPROVE ACCURACY, ROOT(,I2,4H) = ,
          1D23.16,3H + ,D23.16,2H I/24H DID NOT CONVERGE AFTER ,I3,11H ITERAT
          2IONS)
0071      1010 FORMAT(30H THE PRESENT APPROXIMATION IS ,D23.16,3H + ,D23.16,2H I/
          1/)
0072          END

```

TABLE H.III (Continued)

```

0001      SUBROUTINE TEST(UX3,VX3,UX4,VX4,CONV)
C      *****
C      *
C      * SUBROUTINE TEST CHECKS FOR CONVERGENCE OF THE SEQUENCE OF APPROX-
C      * IMATIONS BY TESTING THE EXPRESSION
C      * ABSOLUTE VALUE OF (X(N+1)-X(N))/ABSOLUTE VALUE OF X(N+1).
C      * WHEN IT IS AS SMALL AS DESIRED, CONVERGENCE IS OBTAINED.
C      *
C      *****
0002      DOUBLE PRECISION UX3,VX3,UX4,VX4,EPSRT,EPSO,EPS,AAA,UDUMMY,VDUMMY,
10      DENOM
0003      DOUBLE PRECISION EPSM
0004      LOGICAL CONV
0005      COMMON EPSM,EPS,EPSO,EPSRT,IOZ,MAX
0006      UDUMMY=UX4-UX3
0007      VDUMMY=VX4-VX3
0008      AAA=DSQRT(UDUMMY*UDUMMY+VDUMMY*VDUMMY)
0009      DENOM=DSQRT(UX4*UX4+VX4*VX4)
0010      IF(DENOM.LT.EPSO) GO TO 20
0011      IF(AAA/DENOM.LT.EPS) GO TO 10
0012      5 CONV=.FALSE.
0013      GO TO 100
0014      10 CONV=.TRUE.
0015      GO TO 100
0016      20 IF(AAA.LT.EPSO) GO TO 10
0017      GO TO 5
0018      100 RETURN
0019      END

0001      SUBROUTINE HORNER(NA,UA,VA,UX,VX,UB,VB,UPX,VPX)
C      *****
C      *
C      * HORNER'S METHOD COMPUTES THE VALUE OF THE POLYNOMIAL P(X) AT A POINT D.
C      * SYNTHETIC DIVISION IS USED TO DEFLATE THE POLYNOMIAL BY DIVIDING OUT THE
C      * FACTOR (X-D).
C      *
C      *****
0002      DOUBLE PRECISION UX,VX,UPX,VPX,UB,VB,UA,VA
0003      DIMENSION UA(26),VA(26),UB(26),VB(26)
0004      UB(1)=UA(1)
0005      VB(1)=VA(1)
0006      NUM=NA+1
0007      DO 10 I=2,NUM
0008      UB(I)=UA(I)+(UB(I-1)*UX-VB(I-1)*VX)
0009      10 VB(I)=VA(I)+(VB(I-1)*UX+UB(I-1)*VX)
0010      UPX=UB(NUM)
0011      VPX=VB(NUM)
0012      RETURN
0013      END

```