

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

INTERAGENCY REPORT USGS-183

A SYSTEM OF REGIONAL AGRICULTURAL LAND USE MAPPING
TESTED AGAINST SMALL SCALE APOLLO 9 COLOR INFRARED PHOTOGRAPHY
OF THE IMPERIAL VALLEY (CALIFORNIA)

by
Claude W. Johnson
Leonard W. Bowden
Robert W. Pease
Department of Geography
University of California
Riverside, California

November 1969

Prepared by the U.S. Geological Survey (USGS) for the National Aeronautics and Space Administration (NASA) under NASA Contract No. W-12570, Task No. 160-75-01-32-10. Work performed by the University of California for the USGS Geographic Applications Program under USGS Contract No. 14-08-0001-10674. **This report will not be resubmitted to NTIS by NASA.**

Plate 1.

Color infrared (CIR) photograph of the Imperial Valley with field pattern map overprint. The photo was taken from the Apollo 9 earth orbiting satellite on March 12, 1969 from an altitude of $150\pm$ miles (241.5 km). (Taken with a Hasselblad camera with 80 mm focal length lens using a Wratten 15 filter). The print is a 12 times enlargement from the original. The problem of rectifying an orthographic photo to a polyconic map projection can be seen by the out-of-register area in the upper right corner. The Apollo 9 CIR photo was used to test the agricultural land use interpretation and analysis system described in this paper.




TABLE OF CONTENTS

Plate I	<i>i</i>
Table of Contents	<i>ii-v</i>
Abstract.	<i>vi</i>
I Introduction	1
A. Objective	2
B. Study Areas	2
II Data Source and Technical Evaluation	4
A. Image Resolution	4
B. Image Quality	6
III Analysis of Crop Color Variations Recorded on CIR Imagery.	9
A. Changes in Color Response Related to Season	10
1. Effects of Differences in Crop Maturity	10
2. Effects of Soil Exposure	11
3. Influences of Farming Practices	11
4. Effects of Crop Shading on Amount of Background Infrared Illuminance	12
B. Variations of Crop Color on CIR Imagery with Changes in Resolution	13
IV Design of Agricultural Land Use Interpretation and Mapping Systems	15
A. Crop Signature Tables	16
1. Color Signatures	17
2. Environmental Factors	18
a. Physical Factors	18
b. Cultural-Economic Factors	20
3. Farming Patterns	20

TABLE OF CONTENTS (Contd)

B.	Worldwide Agricultural Land Use Code	22
1.	Requirements	23
2.	A User Oriented Land Use Code	23
3.	Hierarchical Five Digit Level Code	25
C.	Design and Preparation of Local Base Maps	27
1.	Selection and Use of Control Map	27
2.	Adaptation of Coordinate Systems	28
a.	Geographical Coordinate System Modification	28
b.	Local Grid System	30
V	Correlation of Photographic Interpretation with Actual Ground Data	32
A.	Assumptions and Methods of Correlation	32
1.	<u>A priori</u> Knowledge of Test Site Area	32
2.	Ground Surveys for Correlation Studies	33
3.	Representative Selection of Survey Data	34
B.	Correlation of Apollo 9 Crop Identification with Ground Survey Data	35
1.	Specific Crop Identification (Fifth Digit Level).	37
2.	Primary Agricultural Class Categorization (Third Digit Level)	38
3.	Agricultural Land Use Not Identifiable or Not Detected	39
C.	Correlation of Apollo 9 Crop Acreage Summary with Reported Crop Acreage	39
1.	Specific Crop Type Acreage	40
2.	Primary Agricultural Class Acreage	44
3.	Total Cultivated and Available Cropland Acreage	45
D.	Cosistency of Error Between Correlations	45

TABLE OF CONTENTS (Contd)

VI Conclusion	47
Bibliography.	49
Appendix A Description of Imperial Valley.	50
1. Geographical	50
2. Climate	51
Appendix B Interpretation, Mapping, and Analysis Procedures	55
1. Base Map Preparation	56
2. Development of Crop Calendar for the Date of the Imagery	56
3. Image Projection or Base Map Overlay or Overprint	56
4. Establishing Crop Color Signatures from Ground Survey Data	59
5. Determine and Record Individual Field Unit Location	59
6. Determine and Record Individual Field Unit Color on CIR Image	60
7. Perform Computer Crop Identification	60
8. Prepare Final Agricultural Land Use Map and Statistical Data	62
Appendix C	
1. Crop Calendar for the Imperial Valley	63
2. Worldwide Agricultural Land Use Code	64-75
3. Agricultural Land Use Map of Imperial Valley	76
4. Ground Survey Routes for Apollo 9 Correlation Studies	77
Appendix D Thematic Land Use Maps of the Imperial Valley From March 12, 1969 Apollo 9 Photography	78-94
Table I Source of Imagery (Imperial Valley)	5
Table II Correlation of Apollo 9 Land Use Identification with Ground Survey Data	36

TABLE OF CONTENTS (Contd)

Table III Analysis of Errors on Field Crop Identification. 37

Table IV Summary of Correlation Between Reported and Detected . .41-43

Table V Comparison of Errors of Selected Field Crops 46

Table VI-1 Climatic Data for the Imperial Valley. 52

Table VI-2 Climatic Data for the Imperial Valley. 53

Table VII List of Major Crops Under Cultivation on March 12
(Imperial Valley). 57

Figure 1 Sensitometry of SO-180 Color Infrared Film Used in the
Apollo 9 Photography 8

Plate 1 Apollo 9 Imperial Valley CIR Photo with Map Overprint. . . i

Plate 2 Example of 7 Crop Colors Recorded on CIR Imagery 95

Plate 3 Example of Alfalfa Field in 3 Stages of Cutting with 3
Colors 95

Plate 4 Example of Six Different Alfalfa Fields in Different
Stages of Maturity 95

Plate 5 Example of Two Different Crops Showing the Same Color. . . 95

Plate 6 Example of Sugar Beet Field at High Resolution 96

Plate 7 Example of Same Sugar Beet Field at Medium Resolution. . . 96

Plate 8 Example of Sugar Beet Field After Harvest 96

Plate 9 Ground CIR Photo of Sugar Beets Showing Background
Difference 96

ABSTRACT

Interpretation results of the small scale CIR photography of the Imperial Valley (California) taken on March 12, 1969 by the Apollo 9 earth orbiting satellite have shown that world wide agricultural land use mapping can be accomplished from satellite CIR imagery if sufficient a priori information is available for the region being mapped. Correlation of results with actual data is encouraging although the accuracy of identification of specific crops from the single image is poor. The poor results can be partly attributed to only one image taken during mid-season when the three major crops were reflecting approximately the same and their CIR image appears to indicate the same crop type. However, some incapacity can be attributed to lack of understanding of the subtle variations of visual and infrared color reflectance of vegetation and surrounding environment. Analysis of integrated color variations of the vegetation and background environment recorded on CIR imagery is discussed.

Problems associated with the color variations may be overcome by development of a semi-automatic processing system which considers individual field units or cells. Design criteria for semi-automatic processing system are outlined.

Section I

Introduction

Studies using color infrared (CIR) photography of the Imperial Valley (California) at several different scales and ground resolutions have produced three equally important conclusions on the development of agricultural land use photo interpretation systems. The results are summarized as follows:

1. Definitive agricultural land use mapping of a very large region can be accomplished from interpretation of small scale CIR satellite photography. Analysis utilizing the unit record system described below has shown a capability to define and classify a unit as small as 10 acres (4⁺ hectares). Results show that the accuracy of interpretation based upon crop CIR color signature from small scale imagery is dependent upon availability of a priori information of the region, detection of other identification factors on the image, and the careful selection of dates of sequential CIR imagery.

2. The ability to identify agricultural crops on small scale imagery can be attributed, in part, to predictable color variations occurring among crop types. These variations, however, are also seasonally related which introduces another set of variables that makes color signatures inconsistent. This creates identification problems when color comparisons are being made between known and unknown examples of the same crop on the same image. Considerable analysis, is still required before automatic equipment can be programmed to detect the subtle variations occurring within the same crop type.

Additional variations in crop color are created by technical problems of film quality control, variabilities at time of exposure, storage history of the film before processing, and the processing itself. These latter variations cause differences to appear between films of different coating batches rather than on images contained within the same film roll.

3. A method for overcoming problems being encountered due to variations of the crop image color is to design semi-automatic processing systems to utilize data recorded on the image, and to employ previously acquired data. The system designed in this study requires that each field unit or cell on the image be examined individually to determine color record, location, and any other data present. Referral is then made by the computer to previously stored data of that field and region to assist in final identification. It should be noted that efficient processing of sequential photography will require a system of individual unit or cell comparison if agriculture is to be analyzed. Maintaining field unit records enhances the ability to perform special regional agricultural analysis and distribution studies as well as to simplify preparation of regional agricultural land use maps or summaries. (Table IV, Appendix D)

A. Objective

The study emphasizes the design and development of a computer oriented agricultural land use mapping system. However, the role that the crop color and tone variations play, especially on small scale CIR imagery, requires considerable analysis be made of the many variations that occur when recording agricultural crops on CIR film. Of necessity the analysis of crop color variations will precede the description of the design and development of an agricultural land use mapping system. Testing of the system with the small scale Apollo 9 CIR imagery of the Imperial Valley concludes the report.

B. Study Area

Selection of the study area was predetermined by: (1) available imagery; (2) the high probability of obtaining future satellite imagery; (3) a need for

an area with considerable diversification of agricultural crop types; and (4) accessibility of the area to the investigators to enable frequent ground and/or field surveys to verify and/or correct results being obtained.

Two areas met the established requirements: (1) The Coachella Valley centered around Indio, California, and (2) the Imperial Valley between the Salton Sea and the California-Mexican border. The first is dominated by tree crops (citrus and dates) and grapes. A review of the annual crop reports of the Imperial Valley Irrigation District shows that there are 48 different types of crops normally grown in the Imperial Valley. Of these 48 crop types there are 34 which are reported as having more than 100 acres (40⁺ hectares) under cultivation annually and of these latter 34 crop types there are 18 crops which may be considered as the major crops since they are reported as having more than 1000 acres (404.7 hectares) under cultivation annually. In addition, the Imperial Valley contains more than 100 cattle feed lots and supports most of Southern California's production of beef cattle. Of the two, the Imperial Valley was selected as the area of study because the closer Coachella Valley, with its more permanent crop types, had already been mapped, one of the experimental goals of the project, but complete mapping of the Imperial Valley was not as yet accomplished. A description of the geography and the climate of the Imperial Valley may be found in Appendix A.

Section II

Date Source and Technical Evaluation

Table I outlines the source of imagery used to conduct this study. High and medium scale photography was obtained by the NASA Mission 73 flights conducted over the Southern California Test Site 130. The small scale satellite imagery was obtained by the Apollo 9 earth orbiting satellite.

A. Image Resolution

Preparation of the base map was made primarily from the June 11, 1968 imagery with a nominal ground resolution of 1.3 feet (40 cm) at a scale of 1:16,000. While this scale on the 9 inch (23 cm) format proved to be a workable image for hand mapping, the 1:4,000 scale image taken by the NASA mission 73 flight on May 21, 1968 was too large to map efficiently at the 1:62,500 scale. The latter imagery was also too large to provide useful comparison between crop types because quite often only a portion of a single field crop appeared on the 9 inch format. Each frame at the latter scale covered little more than a quarter section (160 acres, 64 hectares) and experience has shown visual interpretation can best be accomplished with at least ten different fields with several crop types appearing in the same exposure. The fine resolution (often as detailed as 4 inches or 10 cm) does permit close inspection of the cultural features and farm management practices, and provides confirmation of details that are not as obvious on smaller scale imagery, but it provided scanty readability of crop color tones. The inadequate readability was caused by separation of the signature inputs (vegetation and visible soil) that combine to make the normal crop color seen in medium and small scale imagery.

TABLE I SOURCE OF IMAGERY - (IMPERIAL VALLEY)

Date	% Cover	Originating Agency*	Platform	Altitude (feet)	Sensor	Focal Length (mm)	Filter Combination	Shutter f-stop	Shutter Speed	Nominal Resolution (meter)(ft.)	Scale
13 May 68	11	WAS	Apache	10,000	T-11	254	WR15+80B	f-5	1/200	0.25	0.82 1:10,000
13 May 68	1	UCR(WAS)	Apache	10,000	35mm	50	WR15+80B	f-6	1/200	1.52	5.00 1:63,000
21 May 68	11	NASA/USGS	CV240A	2,000	RC-8	153	WR15	f-6.8	1/350	0.10	0.33 1:4,000
11 June 68	60	WAS	Apache	16,000	T-11	254	WR15+80B	f-8	1/250	0.40	1.30 1:16,000
8 Mar 69	6	UCR	Cherokee	10,000	35mm	50	WR15+80B	f-8	1/250	1.52	5.00 1:63,000
8 Mar 69	30	UCR	Cml 727	40,000	35mm	50	WR15+80B(80C)	f-8	1/250	7.60	25.00 1:250,000
12 Mar 69	100	NASA	Apollo 9	792,000	Hasselblad	80	WR15	f-8	1/250	80.00	230.00 1:3million
10 May 69	0.5	UCR	Cherokee	10,000	35mm	50	WR15+80B	f-8	1/250	1.52	5.00 1:63,000

* WAS = Western Aerial Survey, Inc. of Riverside

UCR = University of California, Riverside faculty and investigators using handheld cameras in private aircraft or commercial airliners.

B. Image Quality

Technical evaluation of the CIR film used in the 1968 missions has been discussed by Bowden in "Mission 73 Summary and Data Catalog", Technical Letter NASA 132. It is noted again that the recommended W80B filter was used in combination with the W15 filter on certain flights to equalize the yellow and magenta dye layers in the film in order to prevent predominance of cyan in the image. (Pease, Bowden, 1969). The May 13 imagery was overexposed. From this and other experiments it has been found that a shutter setting of f-8 at a speed of 1/250th of a second is a suitable setting for 8443 film using 15 + 80B filters under normal lighting conditions (high sun angle) in desert environment of Southern California. Compensation should be made for abnormal conditions such as low sun angle or large expanses of white sand. CIR photography made with 35 mm handheld cameras at 10 and 20 thousand feet have also yielded valuable information. These are near obliques with resolution sufficient to perform most types of agricultural land use studies. The small expense of both film and platform for 35 mm flights is a significant consideration when planning repeated aerial surveys for studies needing time-lapsed imagery.

The Apollo 9 imagery, taken with only a w 15 filter, was found to be atypical CIR film on postflight inspection. The magenta dye-forming layer had a lower than usual gamma which with the EA-4 processing performed by the Manned Spacecraft Center, was lower in sensitivity than the cyan layer at densities below 0.8. Fortunately, for crop targets, this equalized layer sensitivities with only the Wratten 15 filter. This negated the cyan

color balance that frequently ensues from airluminance which from high altitudes destroys the red of many vegetation targets. Figure 1 graphically indicates the difference between the sensitometry of the Apollo 9 (S0 - 180) film and normal type 8443 CIR film, processed in both the recommended Kodak E-3 chemicals and the EA-4 used. (Pease 1969). This is but one example of the problems produced by the several variables which exist with the use of CIR imagery.

If interpretation systems are to depend on color as a crop signature, more stringent quality control must be instituted in all steps of the system to reduce the variables. The current inclusion by MSC of a step wedge or "frisket" prior to and after each series of exposures is helpful in noting variations in film, exposure, and processing. This is important since several days may elapse between exposure and processing. Ideally a standard ground target should also be exposed at the start of each series, but this becomes impractical for satellite photographs. In the absence of such a ground target, an available reference may be a black-top surface such as a highway or parking lot which is essentially neutral or a soil surface of known reflectance may also be used as a neutral target.¹

¹Further discussion may be found in Egan, 1969.




Figure 1. Sensitometry of the S0-180 color infrared film used in the Apollo 9 mission with EA-4 processing compared to normal type 8443 CIR film with recommended E-3 processing. It is to be noted that the magenta layer of the S0-180 has a lower gamma than normal (less slope to curve) and thus becomes less sensitive than the infrared sensitive cyan layer at a density of 0.8 (point A). With the two layer sensitivities close to equal at densities below 1.0, there was little shift to a cyan color balance due to air luminance for low density vegetation targets. Most crops, therefore, recorded with a good red record. Dark vegetation targets, such as conifer forest on adjacent mountains, with densities close to 2.0 recorded as blue. Sensitivity of the yellow layer could also have been brought to point A with a CC30M compensation filter.

The S0-180 curves based upon sensitometric testing by NASA, Manned Spacecraft Center.

Section III

Analysis of Crop Color Variations Recorded on CIR Imagery

Analysis of Crop Color Variations Recorded on CIR Imagery

Many crop color responses recorded on CIR film follow predictable changes (i.e., cereal grain crops turn yellow when they mature.) which are most useful in identifying crops, especially when utilizing time-lapse imagery. Unfortunately, predictable crop color changes do not always follow a uniform crop-wide pattern. Because of time variations in farm management practices the same crop has been detected to record a different color response within the same field (Plate 3) and between different fields (Plate 4) while different crops have been noted with the same color response (Plate 5). Changes occurring within the vegetation and the surrounding environment cause the variations in color response, but the recording of the response on CIR imagery is further complicated by the variation in the integration of color responses caused by changes in the ground resolution and/or scale of the image. A factorial analysis of the crop response variations may assist in improving the ability to identify agricultural crops.

A. Changes in Color Response Related to Season

The predictable variations in crop color response related to the growth cycle have been found to occur in:

- (1) differences in the crop maturity,
- (2) the ratio of soil exposure to vegetative color,
- (3) the influence of farming practices (irrigation, cultivation, etc.) on the color response of the soil and vegetation, and
- (4) the effect of crop shading on the amount of background infrared illumination.

1. Effects of Differences in Crop Maturity on the Color Response

One of the most important factors in agricultural crop identification is the change of the color response with change in time. The transition of cereal grain crops from the true vegetative color of blue-green to the ripe golden yellow of barley or the bronze color of mature wheat yields false color differences on CIR photography useful in detecting cereal grains at most resolutions. Detection of the temporal change of crop colors requires sequential exposures of the same target throughout the various phenologic stages of a crop's growth. The CIR image color for ripening grain, for example, changes from the deep red of the vegetative stage to the lighter pink of the ripening grain and finally to a yellow, light tan color or yellowish green, depending on auxiliary filtration and processing (Plate 2). Dieter Steiner (1969) found that the ideal time period between successive images for detection of the change in crop color response was three weeks with at least three successive images. Verbal discussions with David Simonett suggested that the timing sequence is tied to the growth cycle in three divisions. In mid-season many crops are at a phenological stage in which their infrared color response appears quite similar and they are difficult to differentiate. During the earlier vegetative stage differences in crops may be detected by differences in planting times. The third division provides the greatest potential in crop differentiation with some crops having ripened to different colors, or showing the effects of different types of harvesting (Plates 7 and 8).

2. Effects of Soil Exposure on the Color Response

The ratio of soil exposure to agricultural vegetative cover changes with the maturity of the crop (Plate 9). A crop that completely covers a field will yield a pure false color on the CIR image; if the crop is in a vegetative stage this will be red. The other extreme may be found in newly seeded fields in which no crop color appears and only the soil reflectance, modified by moisture, creates the color response recorded on the CIR image. The resultant image recorded on the CIR film will be an integrated color combining respective proportions of the color response of the exposed soil and the crop cover. However, the amount of color integration is dependent upon the resolution. At maximum resolution (ground photo, Plate 9) the crop color is separated from the soil color and two distinct non-integrated colors are recorded. Low resolution as exemplified by the Apollo 9 image (Plate 1) yields the maximum integration of combined colors.

3. Influences of Farming Practices on Color Response

Farm management practices are also important contributors to color variations. A frequent practice within a given agricultural area is to stagger planting of annual crops which thus presents several stages of growth of the same crop on the same image. Perennial crops (i.e., alfalfa) may have several cuttings throughout the growing season (Plate 3) and in one instance an image was found with alfalfa fields in six different tones of red from light pink to deep red (Plate 4). The application of fertilizer has a marked effect on the image of the crop by changing the vigor of growth which can be detected on CIR images by deepening the shade of red.

Either irrigation or cultivation will change the amount of soil moisture exposed to the sensor and the integrated color will vary in both hue and tone with the change in the visible moisture conditions. Spectral reflectance characteristics show that dry soil has a much higher reflectivity than wet soil at all wavelengths. The greatest difference occurs in the near-infrared (900-1000 nm) (Condit, 1969). Dry soil, lighter in color, will tend to decrease the overall density of the integrated crop color while wet soil with lower reflectivity will increase the density of the color yielding a darker tone with a shift toward a cyan balance. Examples of the moisture difference phenomena was noted throughout the Apollo 9 photo. The seeded fields were easily detected because of the dark bluish (cyan) color of the very wet soil. Certain cultivated growing crops such as onions showed a combination of modulation of the dark bluish soil background and the vivid red of the vegetation with the result being a magenta color. Differences in bare fields were noted wherein fallow fields were recorded on the CIR film as a neutral color (with EA-4 processing) and the plowed fields yielded a very distinctive turquoise blue.

4. Effect of Crop Shading on Amount of Background Infrared Illuminance

Current investigation related to the IR reflectance characteristics of vegetation (Pease) suggests that vegetation is significantly transparent to near infrared radiation. Thus, vegetation with a shadow substrata will appear darker than the same vegetation with a light or illuminated background. The phenomenon is most marked when the background is illuminated directly by the sun and simply shows through the crop tissue. This effect is dependent on the stage of growth and condition of the vegetation and will

decrease with multiple layering of crop leaves. Soil within an individual field that has been modified by irrigation water will present an even more complicated reflectance pattern. The exposed soil will dry sooner than the soil which is shaded by the vegetation. The variation in the wetness of the soil in the same field may cause a difference in the uniformity of color and certainly can cause a difference in the colors of two fields of the same crop at the same stage of maturity. An adjustment in the scale or resolution will not necessarily reduce the effect of the light contribution of the soil, but the phenomenon may further explain the differences frequently noted in the color records of two identical crops.

B. Variations of Crop Color on CIR Imagery with Changes in Resolution

The change in the color record is quite apparent as the scale and resolution is increased from that of the Apollo 9 image (1:3,000,000 scale; 80 meters resolution) to the large scale and high resolution of the low altitude NASA Mission 73 flight (1:4,000 scale; 10 cm resolution). The high resolution has a most deleterious effect on the readability of crop colors which, as noted in section III-A-2 above is apparently caused by the plant color being visually separated from the associated ground color. The separation effect in high resolution CIR imagery suggests that the color record of an agricultural crop on medium and low resolution CIR imagery must, therefore, be an integrated color reading which is obtained from the combined reflectance of both target vegetation and visible soil background. The plant images on high resolution photography are usually represented by small dots (Plates 5 and 6) which make it difficult to

obtain an average densitometric reading of the entire field. Average values from high resolution imagery may require several readings to be taken for different image elements of each individual field, a process that is not compatible with automatic processing.

Adulteration of the pure vegetation color occurs at least as soon as image resolution is reduced to the CIR film resolution (40 lines/mm) which no longer permits separation of the reflectances of soil and vegetation (Plates 6 and 7). Color blending will continue to increase as the resolution is decreased. At some upper limit the agricultural crops in adjacent fields will also present a cross-modulation between fields and the blended margins will include colors from both. The Apollo 9 image (Plate 1) displays examples of this latter effect. Adjacent fields of the same crop type simply continue the color of each other, but adjacent fields with different crops create a compounded adulteration of the color records of both cells.

Experience during this study has shown that the highest resolution for good color integration on CIR imagery is 40 cm which was noted on Mission 73 at a scale of 1:16,000 (Plate 2).

Section IV

Design of Agricultural Land Use Interpretation and Analysis Systems²

The variations in crop color responses originating in the micro-environment of the individual field suggests that any system designed to process agricultural CIR imagery will need to examine each individual field unit or cell. Likewise, the uniqueness of the macro-environment of a region, as noted in the Imperial Valley, suggests that each agricultural region will also have to be examined individually. The environmental differences between regions, both physical and cultural, are so complex that neither a manual nor an automatic universal worldwide systems designed at this time to consider each variable would be practical. The alternative presented here is to use the results of the Imperial Valley study as design criteria for a basic interpretation and analysis system which then, perhaps, can be modified to fit each agricultural area. The basic modification for applying the system to other areas would be the preparation of an appropriate table of crop signatures, develop surrogates of local conditions, and preparation of a local area base map showing individual field patterns. A further expansion of the system would be the design of a world-wide agricultural land use code which could be used in its entirety or only in those parts applicable to the area under study.

The design criteria that follows is intended for a system that would either manually or automatically examine each field or cell, record the

²The term system is being used rather than methods because it is more indicative of automatic or computer processing techniques and it indicates a repeatable systematic process for each step performed during the interpretation and analysis sequence.

field location index, measure and record the field color response, and then refer to crop signature tables and the field historical data to determine the probable agricultural land use type. Upon complete examination of the entire region various summary reports could be prepared. This type of system requires the preparation of certain tables before any examination of the imagery can begin. The table of crop signatures is one of the first steps before any attempt can be made to perform agricultural land use interpretation.

A. Crop Signature Tables

It has been stated by Bomberger (1960) that agricultural land use interpretation can be made on the basis of tone, texture, pattern, shape, size, and topographic site. Experimentation with satellite color photography has shown that many of these signature factors are lost with the decrease in resolution; however, the color has remained. While, as Strandberg (1968) suggests, color will enhance the resolution capability of the imagery, it is still not sufficient on Apollo 9 imagery to provide much more than gross shape and size information. CIR imagery enhances the separation of cultivated vegetation, but it creates the additional problem of crop color variation discussed in section III. The interpretation of the vegetative color with image elements other than the crops needs additional signature data to help identify individual crop types. These data can be provided by incorporating additional environmental signature facts, both physical and cultural, in the signature table. The complete table should include a list of farming patterns peculiar to each crop type. The design criteria for each of these three signature elements is as follows:

1. Color Signatures

Lack of quality control and sufficient color variation data requires that, at present, color signature tables must be prepared from each set of images. Preparation of a uniform table of color signatures for agricultural crops is impossible at present because of: (1) the previously mentioned lack of adequate quality control in the CIR film manufacture, image exposure, and development processes; and (2) the lack of adequate specific information on the variation in color throughout the growth cycle of each crop. To prepare a table from existing imagery, the ground survey work must be accomplished at the time of the flight to obtain positive identification of representative crops and a sampling for each crop type in various stages of growth.

Despite factors producing color variations, there is still a high percentage of crop types which present a uniform color on a single image - annual crops such as cereal grains, sugar beets, cotton, melons and tomatoes (Plate 2). The uniformity of color within crop type on a single image offers some hope of success in the use of automatic electronic image analyzing devices which are being developed to interpret an entire image without the necessity for examining individual fields. Lack of uniformity in some crops (i.e., alfalfa), however, suggests that it will continue to be necessary to examine certain fields individually using auxiliary signature data before final identification can be made.

An essential requirement in maintaining uniformity in color signature readings is a measuring device which will maintain the same color standard throughout all the imagery. A readout device may be an electronic

scanner which feeds the information to other automatic processing equipment or it may be a manual readout color densitometer which separates the three dye-layer modulations into numerical density readings. For application to individual field analysis the latter instrument may be a desirable adjunct in that sets of precise numerical values may be obtained for each field. Recording numerical values for each dye-layer density will enable crop color signatures to be expressed as a ratio between two of the three density values. The color signature tables should be expressed as upper and lower ratio limits to insure inclusion of the same type crop even when there are slight color variations. If these limits are too broad for the number of crops to be evaluated in a single series of images, then the color signature table should be prepared with subdivisions of crop types and note made regarding overlap of characteristics. The subdivision should consider possible stages of growth and upper and lower limits for each of the stages established. Crops which cannot be identified by color signature tables should be referred to other signature factors for further processing.

2. Environmental Factors

Clues other than color signatures may occur on the CIR imagery or may be available as auxiliary data to reduce the number of identification possibilities. Auxiliary data are those data which occur as physical or cultural factors of the geographic target area.

a. Physical Factors

Among the most useful physical factors is the climate. The agronomist will often divide the cultivated plants into warm weather and cool weather crops. For a given geographical location, the climatic factor

alone enables us to reduce the 1070 cultivated plants listed by Sturtevant (1958) to a more manageable number. Of the usual climatic data for the Imperial Valley, temperature is most useful. Modern technology in irrigation practices has diminished the importance of precipitation there as is exemplified by more than 284 miles of main canals and 1497 miles of laterals. In the Imperial Valley, the temperature extremes have the greatest influence on the crop calendar. (The climatic data for the Imperial Valley is shown in Appendix A-2.) The average 316 frost free days is a poor indication of the growing season in the Imperial Valley. During the winter months, when frost is apt to occur, most of the crops under cultivation, such as lettuce and carrots, have moderate frost tolerances. Hence, the growing season in the Imperial Valley is much closer to 365 days than the frost free calendar would indicate. The climatic factors can usually be summarized and defined in an expanded crop signature table that includes a crop calendar. (Refer to Appendix C-1 for the Imperial Valley Crop Calendar.)

Another physical factor directly connected with climate, but much simpler to define, is the geographical location or limits where various species of vegetation can be cultivated under natural climatic conditions. Date palms for example, have an extreme northern hemisphere limit of approximately 34° latitude. Likewise, bananas have a climatic zone limit that can be described by geographical coordinates. Most of the economic crops can similarly be delimited.

Other indirect physical factors that determine the type of crops which may be planted in the area are local factors such as soil conditions

and importation of water with a high salinity content. The latter problem exists in the Imperial Valley, but is countered by a vast underground tile drainage network that permits ground leaching to remove excess salts. Whenever special local conditions can be stated in definitive form, they may be used to reduce the number of possible crops to be identified in an area.

b. Cultural-Economic Factors

The type of crops produced in a given area or country can also be prejudged to some extent by cultural practices. The diet of the populations of Asia, for example, involves the use of large amounts of rice. In Tongo, one suspects that an abundance of yams are produced. Likewise, in the United States, an investigator would anticipate considerable cropland subsidizing the raising of beef cattle. Diet habits connected with the cultural history of an area supports the idea that tables of crop signatures should be modified for each of the cultural regions of the world to make the job simpler.

Location of the croplands with respect to their markets, cost competition between crops, land costs and export crops are economic factors that should also be considered as crop identification surrogates.

The physical and cultural environmental factors can usually be found summarized in an annual crop calendar for the local area. The selection of crops which appear on the calendar are greatly affected by physical factors, but within the frame of possibilities offered by the physical environment are local cultural factors which dictate crops planted.

3. Farming Patterns

The major divisions of agricultural productions (Field Crops, Vegetable Crops, Fruit and Nut Crops, Livestock and Animal Production, and

Horticultural Specialties) each have distinctive patterns resulting from farming practices. Field crops are generally noted for their extensive farming practices, and the majority of crops which are planted by broadcasting or drilling the seed are identified as field crops. Vegetable crops normally consist of intensive farming practices with mechanized crops planted mostly in rows. The rows are for either irrigation or the necessary crop care of periodic cultivation, fertilizing, spraying, and manual or mechanized harvesting. However, certain field crops are planted in rows also for the same type of care, so the fact a field contains rows does not necessarily identify the type agriculture being practiced. However, the rows of specific crops are often distinctive. For example, spacing of watermelon rows is generally 9 feet (2.74 meters) in the Imperial Valley while Field crops of cotton and sugar beets and many of the vegetable crops are only 42 inches (107 cm.) apart. Within the vegetable crops, the number of rows within a hummock or hill is found to be different. Carrots, in contrast to watermelons, may be planted in closely spaced multiple rows within double rows one foot (30 cm.) apart on either side of the hummock.

The size of a field or the uninterrupted cover area within a field is very indicative of the type of agricultural crop as well as the harvesting method employed. Most large field crops can be harvested mechanically and the only limit on the field size is land ownership and practicality. Grain fields may have an unbroken cover for miles in a dry farming area. In irrigated areas the length of a grain field is limited by ownership and the fall of the land to contain or spread irrigation waters or run sprinkler

systems. Vegetable crops are quite different in size requirements. Manual harvesting has limited the length of an unbroken row of vegetables to the distance a human can carry the product. Hand picked canteloupe are usually limited to rows of a distance that a picker can carry a full bag before unloading. Mechanized harvesting may impose size limits and also spacing limits. Conveyor belts on a semi-automatic watermelon harvester will reach across only 9 rows. Therefore, the tenth row in a watermelon field is bare to facilitate the movement of a harvester. Because vegetables must be irrigated, the size of the field is limited to the fall of the land for flood irrigation or the practical length of pipe for sprinkler irrigation.

Farming practices are thus a very distinctive aid in interpreting agricultural land use. While they all may not be present in a particular image because of the resolution, those which are present can be used effectively to categorize the land use or crop types.

B. Worldwide Agricultural Land Use Code

The development of the coding system for any land use mapping is the keystone to the entire interpretation and analysis process. The success of the system will stand or fall on the adequacy of the classification divisions to accept and distinguish as many land use types as possible and its ability to present the data to the user in a form that will enable solutions to as many problems or questions about the area as possible. These questions may be those of the regional geographer who desires to determine the economic culture of the area, the market analyst who wants to know how many refrigerated rail cars will be needed, or even the hydrologic engineer who must plan for the delivery of water for the season.

1. Requirements

The four prime requirements for an Agricultural Land Use coding classification system are as follows:

- (a) be able to accept and distinguish as many agricultural land use types as possible throughout the world without overlapping, duplicate or ambiguous categories;
- (b) be acceptable in classification divisions and titles to a wide spectrum of users;
- (c) be compatible with or adaptable to computer or machine processing techniques; and
- (d) permit the interpreter to categorize all land use types with the highest possible accuracy consistent with the resolution of the imagery being examined.

2. A User Oriented Land Use Code

Requirement (d) above, suggests a hierarchical classification system wherein the categories become more generalized as the resolution becomes less distinct. Acceptability of titles to the major users, laymen, suggests commonly accepted terms of the farmer be used rather than the biological titles of the scientist. Computer or machine processing suggests the use of pure decimal numerals rather than a mixture of letters and numerals. A system that has been developed, is gaining wide acceptance, and meets three of the above four requirements has the unassuming title of Standard Land Use Code. The code was prepared in 1965 by the Urban Renewal Agency in conjunction with the Bureau of Public Roads. The Standard Code, as originally prepared, falls short of providing for complete worldwide agricultural classes and lacks

sufficient number of divisions for the majority of users. Understandably, the code was prepared for use within the United States - primarily for urban and regional planners. However, the preface to the publication states that users should expand or adapt the code to individual needs. The San Diego County planning commission has made such an adaptation to fit the categories found in their county (San Diego County 1968)).

Probably no single code can be produced for all users. On the other hand, experience in the Imperial Valley suggests a single code can be produced to provide for the planner viewing the scene from the ground as well as the interpreter utilizing aircraft or even satellite imagery taken from 1.0 miles. The Code will require one more digit than is used in the standard code and possibly an additional suffix for the specialist who desires to know if the crop is irrigated or dry, or a cash or a feed crop, or state of crop maturity.

The standard land use code does utilize agronomic and horticultural names and is a classification commonly used by farmers and laymen. The specific crop type of barley (fifth digit level), for example, fits into the hierarchical classification in the secondary agricultural land use of cereal and grain crops (fourth level), in the primary agricultural land use category of Field Crop (third level), in the rural land use of Agriculture (second level), in the general land use of Resources and Production (first level). An alternative agricultural land use classification system might be to categorize crops by the distinguishing features or patterns of the objects being detected. Although such a system is very convenient for the interpreter, he is only doing part of the task. Such signature classification

might place the row crops of sugar beets, cotton, most vegetables, and perhaps some fruit and nut crops all in one category. A user of such a list would have a difficult time making any judgements about a region if he were told it contained 100,000 acres of row crops. The interpreter must use the pattern signatures of a crop, such as the fact it is planted in rows, to assist in establishing the utility of the crop classification or the activity undertaken. If the land is supporting 30,000 acres of forage crops, 20,000 acres of grain crops, 50,000 acres of fiber crops, and contains 100 cattle feed lots, he can make a considerable judgment about the agricultural base and economy of the area.

Because many users need to know how many acres of a specific type of crop are under cultivation in order to plan for harvesting, storage, and transportation, or to make farm loans with reasonable safety, or for many other planning purposes, the code needs to specify the individual crop. The standard code has, therefore, been expanded to five digits from the original four. Duplication of numeral codes to indicate cash crops or grain crops of the same type, or irrigated vs. dry farming have been eliminated. These items can be identified by a letter suffix by those users who need further detail.

3. Hierarchical Five-Digit Level Code

The Urban Renewal Agency (URA) standard land use code employs a four digit level code. The first digit on the left is the most generalized category and the fourth digit (on the right) is the most detailed activity. In the five digit adaptation the fifth digit lists the specific crop type. Appendix C-2 contains the complete Agricultural five digit level land use

code developed for this study. The digit level categories are summarized as follows:

<u>Digit Level</u>	<u>Category</u>	<u>Code and Description Example</u>
First	General Land use	8 Resource Production and Extraction
Second	Rural Land use	81 Agricultural
Third	Primary Agricultural Land use	811 Field and Seed Crops
Fourth	Secondary Agricultural Land use	8111 Cereal and Grain Crops
Fifth	Specific Crop Type	81111 Barley

Establishment of the five digit-level code as an hierarchical system permits the interpreter to attempt specific crop identification without jeopardizing the accuracy of the classification system at some higher level; i.e., the error of classifying wheat (81118) as barley (81111) does not affect the accuracy of classifying cereal grains (8111). Reciprocally, the hierarchical system facilitates a more accurate specific type classification by providing for an initial division into broad categories of land use types. Field crops (811) are generally comprised of large acreage or extensive farms, while vegetable crops (812) are the smaller or more intensive farms.

The resolution of the imagery being examined will influence the digit-level accuracy. CIR imagery of very low resolution will permit distinction of agricultural land (81) noted by the regular red toned patterns of the cultivated lands. Increase of resolution to that of Apollo 9 (80 meters)

reveals field patterns and sizes that will generally indicate intensive vs extensive farming practices and in conjunction with auxiliary environmental data may permit third digit-level (811) distinction of crops. Resolution requirements for fourth and fifth-digit levels were not determined during this study. However, it was found that specific crop types (fifth digit-level) were identified with nearly 100% accuracy from imagery with 40 cm resolution,³ but it is quite possible that a lower resolution will still permit equal accuracy at this fifth digit-level.

C. Design and Preparation of Local Base Maps

Current efforts in remote sensing land use mapping indicates that the procedure (manual or automatic) will be facilitated by the use of a base map which contains the field patterns of the agricultural crop lands under investigation.

1. Selection and Use of Control Map

Convenience in availability and the detail of preparation has made topographic maps the most commonly used base map in rural land use mapping. However, the very fact of the great amount of detail often makes these maps difficult to use when the final land use map is prepared. Also the normal topographic maps do not contain all of the field divisions that are normally encountered. An alternative method is to prepare a new base map of the area from the imagery being utilized and use the local topographic map as a control for accurate planimetric representation.

A convenient scale for agricultural land use mapping in the U.S. has been found to be 1:62,500. Fortunately most of the United States

³Mission 73, Western Aerial Surveys, June 1968.

agricultural areas have been mapped by USGS at this scale. Ideally the base map for aerial or satellite imagery should be orthophoto maps, but satisfactory maps can be prepared using modern automatic cartographic equipment or the somewhat less expensive simultaneous film-map viewers. Lacking the above type equipment, a method was devised for this study will to reduce the filmed image to map scale by means of a photo enlarger/reducer. Strip sections from the control map coincident with the flight line of the imagery were traced on acetate overlays. The filmed image was then projected onto the tracing for completion of the field patterns with correction and rectification for film and platform distortion being made as necessary. The final base map was then prepared by transferring these strip overlays onto a full size overlay of the control map.

2. Adaptation of Coordinate Systems

Detailed agricultural land use, unlike some other types of land use, is often a dynamic process, changing daily. To observe the quantity of changes occurring over an extended period of time will require the use of automatic data processing. To facilitate such processing it will be necessary to index the fields to provide for retrieval of all previously data for any given field. Indexing must provide for both worldwide location and local identification.

a. Geographic Coordinate System Modification

The early scientists who created the geographic coordinate system did not have modern day computers in mind when the system was established. Consequently, any computations of distances or the recording of locations requires a considerable number of calculations and digits. Much

of the calculations and numbering can be simplified by the use of decimal fractions of each degree of latitude or longitude. To assist in computer processing, both the local area coordinates and individual field coordinates, when used, have been established in this study by decimal fractions of each degree. It would also be desirable to perform area computations utilizing the geographic coordinates. The odd shaped fields that are encountered throughout the world make such computations difficult and many problems need to be solved before complete machine computations can be accomplished.

The field shape problem is not so great in the Imperial Valley since most of the fields are rectangular. By using the decimal geographic coordinates at each corner of a field the distance between points and the acreage can be calculated by a computer with only one additional factor included in the machine instructions. The factor would be the length of one degree at any given parallel on the earth grid.

The decision to locate the geographic coordinates of a field at the corner rather than the center of the field may prove a handicap when less rectangular shaped fields are encountered. For the present, geographic coordinates of the local region, as well as the fields, have been indicated for the Greenwich/Equator (G/E) corner of the region or field. The G/E corner is defined as that corner nearest the 0° or Greenwich longitude and 0° or equatorial latitude. In the northern hemisphere - west of Greenwich - the coordinate is in the southeast corner of the field.

b. Local Grid System

Often it may be desirable to indicate geographical coordinates for just the local region and use a local index system for individual field location. If land tenure does not vary to any large degree and once the acreage for a given field has been determined, the index will remain relatively constant. A basic land unit of 160 acres (65 hectares) exists throughout the Imperial Valley. Historically, the land unit dates to the original homesteading of the Valley in 1901 when the Desert Lands Act permitted the settling of 160 acres. Unfortunately, the survey in existence at the time was not accurate and the correct term for a basic land unit in the southern half of the valley is 160 acres more or less. Later purchases of the railroad and school lands and the land in the northern part of the valley were made after a corrected survey and the acreage had been more accurately determined. The variations between the southern half of the valley can be seen on maps in the appendices. The 160 acres representing a quarter of the one square mile (2.6 sq km) sections of the Township and Range cadastral survey provide a uniform one-half mile (0.8 km) to the side square pattern throughout the valley. Consequently, a cartesian coordinate index system consisting of one-half mile divisions provide a naturally derived grid. It was necessary to use quarter-mile divisions in the west central area around Brawley which was displaced one-quarter mile in a North-South direction because of survey adjustments and inaccuracies. Where the basic land unit of 160 acres has been divided into smaller fields a letter suffix has been added to the index number. The suffixes start with the letter 'A' in the upper right hand corner and proceed counter-clockwise around the 160 acre unit.

Similar local grid systems in all parts of the world should be facilitated by use of basic land unit theory. In addition to historical background in land tenure, political factors are also a consideration. In the western United States, the 1902 Land Reclamation Act limiting the distribution of water from federal reclamation projects to owners of 160 acres has a pronounced effect on the land unit. Government control and relocation of agricultural units in areas such as the Mexicali Valley of Baja California have an effect on the land unit. Other factors no doubt influence the establishment of a basic land unit. Many of these conditions become apparent once the field areas are mapped and a study is made to determine the local pattern.

A step by step outline of the procedures to be followed in the interpretation, mapping and analysis of CIR imagery is contained in Appendix B.

Section V

Correlation of Photographic Interpretation with Actual Ground Data

Receipt of the Apollo 9 CIR photography provided a timely input for testing the interpretive and analysis system outlined in the preceding section and appendix B. The sequence of imagery inputs from aircraft overflights of the Southern California test site enabled the NASA mission 73 imagery to be used for the development of interpretation procedural methods, preparation of the base map, and familiarization training. This preparation permitted the Apollo 9 imagery to be used to provide testing of the processing system as well as testing of small scale CIR imagery.

A. Assumptions and Methods of Correlation

The availability of Crop Calendars for many agricultural areas suggests that considerable knowledge is available about agricultural areas of the world. Crop Calendars can be used to reduce the crops which must be considered in a given area at a given time. The accuracy of determining agricultural land use can be considerably improved if the area is thought of as a region of known culture rather than an unknown hostile military target. Development of surrogates of local peculiarities will greatly increase the accuracy of results.

1. A priori Knowledge of the Test Site Area

In developing the processing system which led to the interpretation of the Apollo 9 photography many ground surveys of the Imperial Valley were conducted and several discussions were held with local agricultural advisors. The result was an accumulation of information pertaining

to the farming practices in the Imperial Valley which included the knowledge that certain crops are localized in production, (i.e., asparagus crops are centered around the local cooperative association in the central area of the southwest corner of the valley and carrots are localized around Holtville, "The Carrot Capital of America", etc.). Additional essential preinformation for the Apollo 9 photo interpretation was made available from Mission 73 analysis with nearly 75 percent of the previous crops being identified. Thus, the plausibility of crop sequences was ascertained for many of the fields during the interpretation of the Apollo 9 photo. In other areas, localized a priori knowledge may not be available to the interpreter, but much of information can be developed from repeated flights. Hence, the test of the processing system on the Apollo 9 photo provided simulated conditions that could well exist after several missions had been flown over the area.

2. Ground Surveys for Correlation Studies

To provide accurate data for the development of crop signatures for the Apollo 9 photo, a ground survey of Imperial Valley was conducted coincident with the first day of Apollo 9 overflights. On this initial survey, 586 fields were examined and identified representing 7.5 percent of the 7801 fields subsequently identified.

After approximately one-half of the satellite photos had been interpreted, a second ground survey was conducted on May 10 with the data already obtained reconstructed to the conditions existing on March 12, the date of the Apollo 9 photo. The second survey examined 447 fields (5.7 percent of total). A third survey was made on May 21 for purposes not

connected with the Apollo 9 photo analysis, but data was obtained that could be reconstructed back to March 12 thus providing 463 more fields (6 percent of the total) for identification correlation analysis. The three surveys provided a total of 1507 fields (19 percent of total) visually identified by ground inspection. The field sample data has been correctly indicated on the final Agricultural Land Use Map of the Imperial Valley. (Appendix C-3).⁴

3. Representative Selection of Survey Data

Like most statistical correlations, the question arose whether the 13 percent sample represented by the latter two surveys was representative of the field population? The selection of a truly random sample in the Imperial Valley was biased by the usual human choice of crops planted in a given location as well as physical limitations as to where a particular type crop may be grown. No attempt was made to select random samples by the usual statistical methods. The first training survey was not used in the correlation analysis, therefore the only effect it had on the correlation was to eliminate some of the possible samples from duplicative efforts. The two ground surveys that were used for correlation analysis were not completely random in selection, nor were they completely planned. Both were taken in the more southern portion of the valley in order to obtain a representative cross section of crops. The large acreage fields (160 acres) north of the east section were particularly avoided because they are limited to the field crops of alfalfa, barley, sugar beets, and cotton. Survey time was better spent in areas of smaller acreage fields

⁴ Readers who desire this ground information may find the route of each survey track listed in Appendix C-4.

and more diverse crop types. An indication the latter two surveys were a representative sample was that correlation analysis performed on them separately resulted in almost identical findings. A later survey, not included in this study, was given a test correlation with the interpreted data and the results were again identical. While the above did not yield statistically derived confidence limits or significance, the evidence provides a basis for believing the conclusions drawn from the correlation are valid.

B. Correlation of Apollo 9 Crop Identification with Ground Survey Data

The arid conditions of the Imperial Valley make it a simple matter to distinguish the agricultural land from the non-agricultural land. To produce vegetation and to continue its growth requires the importation and application of water. Consequently, a glance at the Apollo 9 photograph of the Imperial Valley illustrates the growing croplands in their red false color image. Even cropland that has no visible vegetation can be detected by the surrogates of surface moisture through irrigation of newly seeded fields, or the exposure of soil moisture through plowing. The ease with which agricultural land can be detected within the Imperial Valley on CIR imagery makes the detection accuracy at the second-digit level of the Agricultural land use code nearly 100 percent. The correlation analysis of this study is then concerned with the accuracy of the third digit level (primary agricultural class) and the fifth digit level (specific crop type).⁵

⁵An Imperial Valley and Coachella Valley agricultural land use map produced by Richard Francaviglia from Gemini V normal color imagery was limited to three categories, i.e., permanent crops, fallow land, and field crops. For further information see Bowden, 1967 and Nunelly, 1968.

TABLE 11

CORRELATION OF LAND-USE IDENTIFICATION WITH GROUND SURVEY CHECK: APOLLO 9 CIR IMPERIAL VALLEY IMAGE (MAR 11, 1969)

AGRICULTURAL CLASSIFICATION	IDENTIFICATION ACCURACY WITHIN CLASSIFICATION (Digit Level)								
	PRIMARY CLASSIFICATION (Third Digit Level)			SECONDARY CLASSIFICATION (Fourth Digit Level)			SPECIFIC CROP (Fifth Digit Level)		
	SAMPLE SIZE	ACCUMULATIVE CORRECTLY IDENTIFIED	%	SAMPLE SIZE	ACCUMULATIVE CORRECTLY IDENTIFIED	%	SAMPLE SIZE	NUMBER CORRECTLY IDENTIFIED	%
<u>AGRICULTURE</u>	81								
<u>FIELD CROPS</u>	811	537	88.6	150	62	41.3	140	53	37.9
<u>GRAIN CROPS</u>	8111	(79)			(56)		10	0	0.0
Barley	81111	(9)		319	200	62.7			
Wheat	81118	(10)			(7)		11	0	0.0
<u>FORAGE CROPS</u>	8113	(264)		132	(193)		308	193	62.7
Short Grass (Rye)	81131	(130)		5	(27)		132	27	20.5
Alfalfa	81133	(4)			0		5	0	0.0
<u>SUGAR CROPS</u>	8114				(0)		606	289	47.6
Sugar Beets	81142						5	0	0.0
<u>OIL CROPS</u>	8117						606	273	45.0
Flax	81172						5	0	0.0
<u>FIELD CROP TOTALS</u>		23	57.5	40	23	57.5	40	23	57.5
<u>VEGETABLE CROPS</u>	812			18	13	72.2	18	13	72.2
<u>SALAD CROPS</u>	8123			10	7	70.0	10	7	70.0
Lettuce	81231			12	3	25.0	10	7	70.0
<u>ROOT CROPS</u>	8127			40	23	57.5	40	23	57.5
Carrots	81272						12	3	25.0
<u>BULB CROPS</u>	8128						12	3	25.0
Onions	81284						40	23	57.5
<u>VEGETABLE CROP TOTALS</u>		247	92.9	266	247	92.9	266	247	92.9
<u>NON-PRODUCING & TRANSITION LAND</u>	818			32	26	81.3	32	26	81.3
FALLOW	8181			130	120	92.3	130	120	92.3
<u>PLOWED</u>	8182			1	1	100.0	1	1	100.0
<u>LEACHED</u>	8183			39	37	94.9	39	37	94.9
<u>ABANDONED</u>	8184			64	63	98.4	64	63	98.4
<u>PREPARED & SEEDED</u>	8186			266	247	92.9	266	247	92.9
<u>NON-PROD & TRANS LAND TOTALS</u>		807	88.5	912	807	88.5	912	807	88.5
<u>GRAND TOTALS</u>				912	559	61.2	912	543	59.5

1. Correlation of Specific Crop Identification (Fifth Digit Level)

Table II summarizes the results of the correlation between the Apollo 9 interpreted data with the ground survey data from 912 fields (13 percent of the field population total). The overall accuracy of 59.5 percent for the Specific Crop identification is not only a poor result, but it also is a poor indicator of the actual results. The excellent results obtained in the Non-producing and Transition crop lands is masked by the poor results obtained from the three major crops (barley, alfalfa, and sugar beets) in production on the date of the Apollo 9 flight. The vegetable crops surveyed are too few to be of statistical significance. A primary question is, 'why is the identification accuracy of field crops so low (45 percent) and, in particular, why is the sugar beet accuracy only 20.5 percent?'

An analysis of errors on the misidentification of the three main field crops show the following results:

Table III: Analysis of Errors on Field Crop Identification

<u>Crop Type</u>	<u>Identified as (%)</u> :				
	<u>Errors</u>	<u>Barley</u>	<u>Alfalfa</u>	<u>Sugar Beets</u>	<u>Plowed Land</u>
Barley	87		58.3	22.6	
Alfalfa	115	40.4		24.5	18.1
Sugar Beets	105	29.5	59.9		

Several reasons may exist for such a large percentage of error with the most likely being that the maturity of the three crops on March 12 is such that they generally present the same small scale CIR color image. The dark red did not necessarily equate to one specific type crop with light reds showing another type crop. No doubt, moisture conditions in the individual fields created color differences among a specific crop type - especially alfalfa. The 18.1 percent of the alfalfa fields, misidentified as plowed land, implies moisture was a major factor in causing identification errors. Another factor was the unusually large number of weeds in the sugar beet fields which simulated alfalfa. The abnormal and untimely winter rains of 1968-69 germinated a large weed crop which in some cases entirely covered the sugar beets. The large error of field crop identification indicates that additional information is required at this time of year to provide specific crop information.

2. Correlation of Primary Agricultural Categorization

The majority of errors within the field crop class were within the class itself so that the overall differentiation of Primary agricultural classes was a reasonable 88.5 percent. There were at least three contributing factors: (1) The separation of the Non-producing and Transition Crop Land was obvious by the non-red (blue or neutral color) of the field area. (2) The vegetable crops were smaller size acreages (predominantly 40 acres, but never in excess of 80 acres) and familiarization with the area provided clues as to the location of the vegetable crops. (3) The two main vegetable crops (lettuce and onions) each had an easily identifiable color that enabled even the small intensive type acreages to be classified.

3. Agricultural Land Use Not Identifiable or Not Detectable

Several agricultural land use categories within the Imperial Valley were not recognizable on the Apollo 9 imagery. In general, small areas (less than 20 acres (8 hectares)) could not be established as distinctive from other croplands. However, three types of land use concerned (feed lots, tree crops, and asparagus) are permanent type crops or land use and once established the identification will remain constant for several years. These three land uses have been located on the land-use map (Appendix C-3) by ground survey and have not been included in the correlation statistics presented here. Inspection of images of several feed lots on the Apollo 9 photo reveals a color that matches plowed or seeded fields. Tree crop areas in the Imperial Valley are too small and too few to establish a color signature for the Apollo 9 image.⁶ Asparagus fields are found in a variety of stages in March with too few fields of similar maturity to establish a consistent color signature.

C. Correlation of Apollo 9 Acreage Summary with Reported Crop Acreage

Three times a year the Imperial Irrigation District prepares a Report of Crops Growing. Fortunately, one of the periodic reporting dates is March 15, which permitted an exact acreage correlation to be made. The Irrigation District Report is prepared from visual reports by 30 zanjeros or "ditch riders" who control water deliveries to each field. There is a possibility of errors in the reporting system, but over the years it has become accepted by the valley farmers as "the reliable report".

⁶ However, the nearby Coachella Valley has an abundance of date palms, citrus groves, and vineyards which are detectable on CIR from aircraft altitudes and generally identifiable on the Apollo 9 imagery.

1. Correlation of Specific Crop Type Acreage (Fifth Digit Level)

Specific crop type acreages are compared in Table IV. The summary for grain crops show that it was impossible to differentiate the wheat or oats from barley. In fact, the only wheat or oats identified on the Apollo 9 photograph were those fields visually checked by ground survey. The forage crop summary reveals a definite prejudice in the interpretation evidenced by 18 percent more alfalfa acreage being identified than actually existed at the time. Alfalfa represents 34 percent of the total crop acreage in the valley, hence an interpreter would be inclined to lean toward alfalfa identification when in doubt. Another factor in the excess of alfalfa acreage is the suspected effect on the color of sugar beets by weed cover in the fields. The table reveals alfalfa overestimated by nearly the same acreage that sugar beets are underestimated.

Vegetable crop accuracy attained on asparagus and other crops is not shown since most were identified from ground survey data, or from previous image interpretation. The only three vegetable crops with sufficient acreage to make a valid comparison are lettuce, carrots and onions. The high percentage of accuracy in identifying lettuce is a confirmation of the distinctive red record that lettuce presents in March in the Imperial Valley. On the other hand, carrots are near the end of the harvest season and considerable error can occur between the date of ground observation and the date of the Irrigation Districts report. Lack of sufficient initial ground survey data to establish a firm color record for carrots on the Apollo 9 image may also be a cause of the differences in the carrot acreage. Onion crops have a distinctive color on the Apollo 9 image, but many of the onion fields are

TABLE IV
SUMMARY OF CORRELATION BETWEEN REPORTED AND DETECTED
AGRICULTURAL CROP ACREAGE IN THE IMPERIAL VALLEY (March 15, 1969)

Summary: Agricultural Land Class	Acreage		Difference		Percent Accuracy *
	Reported	Detected	Over *	Short *	
811 Field Crops	315,858	292,440		23,418	93
812 Vegetable Crops	21,162	20,574		588	97
813 Fruit & Nut Crops	2,309	668			
8186 Prepared & Seeded	51,160	50,980		180	97
816 Pasture Land	682	120			
Unidentified Ag. Land		6,980			
<u>Total Land With Growing Crops</u>	<u>391,171</u>	<u>371,752</u>		<u>19,419</u>	<u>94.8</u>
818 Non-Productive Ag. Land		93,180			
<u>Total Available Ag. Land</u>	<u>474,437</u>	<u>464,932</u>		<u>9,505</u>	<u>98.0</u>
<u>Individual Crops:</u>					
811 Field Crops					
111 Barley	72,829	68,350		4,479	94
115 Oats	2,423	140			
118 Wheat	9,932	1,100			
Total Grain Crops	85,184	69,590			
131 Short Grass (Rye, Bermuda)	14,714	4,490		10,224	31
133 Alfalfa	134,692	163,420	28,728		82
Total Forage Crops	148,714	167,910			
141 Sugar Cane	2				
142 Sugar Beets	79,679	54,350		25,329	68
Total Sugar Crops	79,681	54,350			
172 Flax	2,279	590		1,689	26
<u>TOTAL FIELD CROPS</u>	<u>315,828</u>	<u>292,440</u>			
812 Vegetable Crops					
211 Asparagus	2,832	3,100			
221 Beans, Green (Fava)	85	85			
223 Peas, Green	74				
224 Okra	51				
231 Lettuce	7,601	7,840	239		97
232 Celery	4	4			
234 Parsley	1				
236 Chicory	3	3			
237 Mustard	32				
Total Salad Crops	7,641	7,847			

* Only valid comparisons of individual crops have been made. No comparison has been attempted for those crops in which the majority of the detected acreage reported is the result of ground survey information (i.e., Wheat and Oats).

<u>Individual Crops (cont.):</u>	<u>Acreage</u>		<u>Difference</u>		<u>Percent Accuracy *</u>
	<u>Reported</u>	<u>Detected</u>	<u>Over *</u>	<u>Short *</u>	
241 Broccoli	289	72			
243 Cabbage	224	240			
244 Cauliflower	40				
Total Cole Crops	553	312			
272 Carrots	4,127	5,580	1,453		74
274 Potatoes (Chinese)	3				
Total Root Crops	4,130	5,580			
282 Garlic	391				
284 Onions	5,405	3,650		1,755	68
Total Bulb Crops	5,796	3,650			
<u>TOTAL VEGETABLE CROPS</u>	<u>21,162</u>	<u>20,574</u>		588	97
<u>813 Fruit & Nut Crops</u>					
314 Grapes	2				
322 Apricots	19				
330 Citrus, Undifferentiated	324	600			
331 Grapefruit	507				
333 Lemon	276				
335 Orange	611				
337 Tangerine	400				
Total Citrus Crops	2,118	600			
342 Dates	93	60			
363 Pecans	77	8			
<u>TOTAL FRUIT & NUT CROPS</u>	<u>2,309</u>	<u>668</u>			
<u>860 Prepared and Seeded Land</u>					
113 Corn	80				
114 Sorghum, Grain	9,510				
132 Tall Grass (Sudan)	1,145				
151 Cotton	20,888				
Total Seeded Field Crops	31,623				
250 Melons, Undifferentiated	1,053				
251 Cantaloupes	11,861				
252 Cucumbers	50				
253 Crenshaw melons	22				
255 Squash	336				
256 Watermelons	3,709				
Total Cucurbit Crops	17,031				
<u>TOTAL PREPARED & SEEDED LAND</u>	<u>51,160</u>	<u>50,980</u>	180		99.6

* Only valid comparisons of individual crops have been made. No comparison has been attempted for those crops in which the majority of the detected acreage reported is the result of ground survey information (i.e., Wheat and Oats).

	<u>Acreage</u>	
<u>Other Agricultural Land</u>	<u>Reported</u>	<u>Detected</u>
610 Pasture Land	682	120
Unidentified Ag. Land		6,980
<u>818 Non-Productive Land</u>		
810 Fallow Land		35,110
820 Plowed Land		35,560
830 Land Being Reclaimed: Leached		1,540
840 Abandoned Ag. Land		20,620
850 Harvested Land		<u>350</u>
<u>TOTAL NON-PRODUCTIVE AG. LAND</u>		93,180

small and difficult to detect which is the reason for most of the difference in onion acreage comparison.

Ground inspection permits the field surveyor to see certain type crops that have been planted (i.e., melon plants under caps), but such fields present only a single bluish color tone on the Apollo 9 image. The result of soil reflection modified by irrigation. Therefore, these crops were identified by Irrigation District report have been grouped under the Apollo 9 classification of prepared or seeded. The 99.6 percent accuracy for this class of land use verifies how easy it is on CIR imagery to identify fields that have been seeded and are under irrigation.

The Irrigation District reports non-productive land only once per year so that no comparison can be made for the month of March. However, the detection of this acreage by its light, almost neutral, color enables an overall comparison to be made of land available for farming within the Imperial Valley.

2. Correlation of Primary Agricultural Class Acreage (Third Digit Level)

The summary of acreage by agricultural class confirms some of the findings presented in the specific crop identification correlation discussed above. Although specific crop type cannot always be identified, the class (i.e., field crop, vegetable crop, seeded land) maintains consistency in detection as indicated by over 90 percent accuracy in establishing acreage for the three major classes of agricultural land use, field, vegetable, transition and non-productive.

3. Correlation of Total Cultivated and Available Cropland Acreage

A slight difference of 5.2 percent between CIR detected acreage under cultivation and reported acreage is a tolerable error of the interpretation system. The method used in the detected acreage summary is to list the total gross acreage of a crop as estimated from the base map and automatically deduct 10 percent from the gross acreage for roads, canals, drains, service areas, and farm areas. Ten percent has been established within the Imperial Valley as the average acreage that is lost from the production of each field for these various purposes. Also errors may occur because the fields in the southern half of the valley are not always exact divisors of 160 acres (the basic land unit) due to errors in the survey system existing at the time the land was homesteaded.

The total estimate of 464,932 acres (188,158 hectares) of agricultural land within the Imperial Valley is within 2 percent of that reported available during the year 1968 (474,437 acres (189,775 hectares)). The latter estimate is well within accuracy tolerance of the processing system.

D. Consistency of Error Between Correlations

The major sources of error in the two correlations above were found in differentiating between the same crop types (alfalfa, sugar beets, and barley) and in the same proportions in both correlations. A comparison of errors for these three crops in the two correlations is shown below:

Table V. Comparison of Errors of Selected Field Crops

	<u>Identification Correlation</u>		<u>Acreage Correlation</u>	
	<u>Percent Over</u>	<u>Percent Short</u>	<u>Acreage Over</u>	<u>Acreage Short</u>
111 Barley		18.3		4,479
133 Alfalfa	53.7		28,728	
142 Sugar beets		35.4		25,329

The discrepancies in the identification correlation of Table V represents the difference remaining after adjustment for cross-identification, (i.e., 58.3 percent of the barley fields surveyed were identified as alfalfa while only 40 percent of the alfalfa fields were cross identified as barley, leaving a total shortage of 18.3 percent in barley field identification.)

The inability to differentiate accurately between these major crops reaffirms the problem of color variations for specific crops. Improvement in identification accuracy from satellite CIR imagery will occur with better standardization of CIR color records. However, improvement may be obtained from better resolution or by time-lapse photography.

Section VI

Conclusion

The production of a complete agricultural land use map from small scale satellite imagery of a region such as the Imperial Valley is most encouraging (Appendix C-3). Two factors permitted the attainment of this goal: (1) CIR imagery that provided distinctive color records of the separate crop types, and (2) availability of a priori information. Without either the task would have been impossible.

Lack of accuracy in differentiating individual crop types was due to: (1) lack of sufficient research into the subtle variations of individual crop colors recorded on CIR imagery at Apollo 9 resolution, and (2) lack of sequential or time-lapse imagery designed to take advantage of seasonal crop color variations. Availability of time oriented imagery would improve not only the ability to distinguish between crop color signatures, but would also assist identification by providing a permanent history of the individual field units.

The laborious task of completing the test mapping in this study has shown that timely agricultural land use mapping from volumes of imagery that may soon be obtained from satellites will require automatic or semi-automatic processing utilizing an appropriate land classification system. Repeated surveys of a region will be facilitated by the maintenance of a historical file of individual field units or cells, especially when utilizing sequential or time-lapsed imagery. Certain criteria for a semi-automatic system have been outlined here with the intent that a computer can be employed in the system to provide data storage and retrieval as well as performing the identification task from input data read directly from the imagery.

Because of the dependence of automatic image readout systems, including electronic image analyzers, on the image color variations for performing crop identifications, continued research must be conducted into the crop color variations that are being recorded on CIR and B/W multispectral imagery. Hopefully, results of further investigation will permit electronic analysis equipment to differentiate in spite of inconsistent subtle image color variations for a particular crop. Subsequently, an entire image of a region, like the Imperial Valley, can thus be reliably interpreted by fully automatic methods in a very few minutes.

BIBLIOGRAPHY

- Bomberger, E. H., et al., 1960. "Photo Interpretation in Agriculture," Manual of Photographic Interpretation, (American Society of Photogrammetry, Washington, D.C.).
- Bowden, L. W., N. J. W. Thrower and C. E. Tiedemann, 1967. "Status Report on Remote Sensing of Southern California and Related Environments," Status Report I USDI Contract 12-08-0001-10674, October, (University of California, Riverside and Los Angeles, California).
- Condit, H. R., "Spectral Reflectance of Soil and Sand," 1969, New Horizons in Color Aerial Photography. Seminar Proceedings of the American Society of Photogrammetry and the Society of Photographic Scientist and Engineers, June, (American Society of Photogrammetry, Washington, D.C.), pp. 3-17.
- Egan, W. G., 1969. "Practical Calibration and Control Techniques for Type 8443 and Ektachrome Films," Grumman Research Department Memorandum RM-443J, (Grumman Aircraft Engineering Corporation, Bethpage, New York).
- Jaques, H. E., 1958. "How to Know the Economic Plants," (Wm. C. Brown Co. Publishers, Dubuque, Iowa) p.2.
- Nunelly, N. R. and R. F. Witmer, 1968. "A Strategy for Developing Classifications of Land Use as Interpreted from Remote Sensing Imagery," ASP/ACSM Technical Papers on the 28th Meeting, (American Society of Photogrammetry, Washington, D.C.).
- Pease, R. W. and L. W. Bowden, 1969. "Making Color Infrared Film a More Effective High Altitude Remote Sensor," Remote Sensing of the Environment, Vol. 1, No. 1, 1 March 1969 (American Elsevier Publishing Co., Inc., New York, N.Y.).
- Steiner, D., 1969. "Using the Time Dimension for Automated Crop Surveys from Space," American Society of Photogrammetry Technical Papers From the 35th Annual Meeting, March 9-14, 1969, (ASP, Washington, D.C.) pp. 286-299.
- Strandberg, C. H., 1969. "The Language of Color," Manual of Color Aerial Photography, (American Society of Photogrammetry, Washington, D.C.).
- San Diego County Planning Department, 1969. "Standard Land Use Code, San Diego Region, Topical-Descriptive Index," Comprehensive Planning Program, (Planning Department, County of San Diego, California).
- Urban Renewal Administration, 1965. Standard Land Use Coding Manual, First Edition, (Supt. of Documents, USGO, Washington, D.C., 20402).

Appendix A

Description of Imperial Valley

1. Geographical

Geographically the Imperial Valley is bounded between latitudes 32.7° North and 33.3° North and longitudes 115.3° West and 115.8° West. An area of 34 statute miles (54 km) east and west and 48 statute miles (77 km) north and south, it encompasses 655,680 acres (262,272 hectares) between the highline irrigation canals which bound the east and west sides of the valley. Within the above acreage there is a total of 474,437 (189,775 hectares) farmable acres with an additional 14,716 acres (5,866 hectares) devoted to farms in homes, feed lots, cotton gins, experimental areas, and agricultural industrial areas. Another 71,818 acres (28,727 hectares) are in drains, canals, rivers, railroads, and roads. Urban recreation areas (both parks and lakes), and rural schools account for 12,498 acres (5,000 hectares). The remaining acreage is mostly in undeveloped land.

The East Highline Canal trends along the sea level contour. The All-American Canal, along the U.S.-Mexican border, drops from 35 feet (10 meters) above sea level at the junction with the East Highline Canal to 10 feet (3 meters) below sea level at the junction with the Westside Main Canal. The Westside Main Canal drops from 10 feet (3 meters) below sea level at the international border to 175 feet (52 meters) below sea level in the northwest corner of the valley. The lowest elevation in the valley is the southern shoreline of the Salton Sea which fluctuates between -231 feet (-69 meters) and -234 feet (-70 meters). The high water level occurs in April and the low water mark in October. Contour lines reveal a gentle slope from the southeast corner to the northwest corner.

The slope has an average fall of 7 feet per mile (2.1 meters per km). The northeast corner near Niland, however, is steeply sloping with an average fall of 80 feet per mile (24.4 meters per km).

2. Climate

The climate is sub-tropical arid.⁷ Slightly less than 3 inches (76 mm) of annual average rainfall occurs between August and April with 0.75 inch (19 mm) occurring in August and September as summer thunderstorms or "cloudbursts." The effect of this meager rain is slight as reflected by the barren desert surrounding the valley. Consequently, moisture content of soil and fields is controlled entirely through irrigation which provides a completely artificial moisture environment for the study area. Table VI outlines the Valley's climate data. Normal mean annual temperature of 72°F (22.2°C) only hints at the extreme summer temperatures. The mean daily maximum temperatures during the months of June through September ranges from 103°F (39.4°C) to 108°F (42.2°C) with daily maximums of 115°F (46.1°C) being normal and random maximums of 122°F (50.4°C) not uncommon. High evaporation rates, over 12 inches (30 cm) for one summer month, help account for the stable level of the Salton Sea even though the sea is a catchment for all excess irrigation water.

From the time the first irrigation waters flowed into the valley on June 21, 1901 the area has presented an ideal site for study of isolated agricultural land use. Lush fields present a striking contrast to barren desert as many satellite photos have shown. Except for three winter

⁷The climate is classified as sub-tropical rather than tropical even though the mean of the coldest month is over 50°F because frost occurs in more than 40 nights during the winter.

TABLE VI-1 CLIMATIC DATA FOR THE IMPERIAL VALLEY

Section I. Annual Normals

	<u>Brawley</u>	<u>Callexico</u>	<u>El Centro</u>	<u>Imperial</u>
Average Frost Free Days	317		316	
Mean Date of First Frost	Dec. 11		Dec. 28*	Dec. 28*
Mean Date of Last Frost	Jan. 28		Jan. 31*	Jan. 23*
Actual Number of Frost Nights	48	37	42	51
Mean Number of Days 90°F or over	182		183	177
Mean Number of Days 90°F or under	7		15	5
Highest Temperature	122(50.4°C)	117(47.6°C)	122(50.4°C)	117(47.6°C)
Mean Max. Temp.	88.3(31.5°C)	86.4(30.5°C)	89.6(32.3°C)	88.2(31.5°C)
Mean Min. Temp.	55.6(13.2°C)	55.7(13.2°C)	55.9(13.4°C)	56.8(13.9°C)
Record Mean	71.9(22.3°C)	71.0(21.8°C)	72.8(22.9°C)	72.6(22.7°C)
Lowest	19(-7.3°C)	21(-6.2°C)	16(-9.0°C)	26(-3.4°C)
Degree Days (Nov-Mar)	1216		1137	1086
Degree Days (Apr-Oct)	16		8	3
Mean Annual Precipitation	2.57(65.3mm)	2.93(74.4mm)	2.81(71.4mm)	2.84(72.1mm)
Average Number of Days .01 Precip. or more (0.25mm)	11	13	15	
Average Number of Days .10 Precip. or more (2.54mm)	2	2	6	5
Average Number of Days .50 Precip. or more	0	0	0	0

*1966 ONLY

TABLE VI-2 CLIMATIC DATA FOR THE IMPERIAL VALLEY
(Temperatures in °F, Precipitation in inches)

Section 2. Monthly Normals

	Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec
<u>Brawley</u>												
Highest Temp	87	91	104	108	118	118	122	120	121	111	98	122
Mean Max Temp	68.4	73.2	79.0	86.7	94.7	103.2	107.8	106.6	101.8	90.4	78.6	69.5
Mean Min Temp	37.5	41.7	47.8	53.7	60.3	66.9	75.4	75.7	68.7	56.5	44.6	38.9
Mean Temp	53.0	56.2	63.4	70.3	77.4	85.2	91.6	91.1	85.2	73.5	61.6	54.1
Lowest Temp	19	25	29	35	41	49	55	59	50	34	26	20
Mean Precip	0.35	0.36	0.19	0.11	0.04	0.02	0.05	0.23	0.25	0.24	0.14	0.59
<u>Calxico</u>												
Highest Temp	85	93	100	104	116	117	117	117	113	106	99	89
Mean Max Temp	67.5	73.3	78.6	85.5	92.2	101.3	103.9	103.4	98.6	88.0	76.7	67.3
Mean Min Temp	38.9	43.4	47.7	53.3	58.7	67.4	75.9	74.5	67.7	56.0	45.7	39.0
Mean Temp	53.2	58.4	63.2	69.4	75.3	84.3	89.8	89.0	83.2	72.0	61.1	53.2
Lowest Temp	21	28	32	39	41	54	59	55	48	36	28	23
Mean Precip	0.47	0.41	0.24	0.08	0.04	0.02	0.08	0.53	0.20	0.18	0.24	0.44
<u>El Centro</u>												
Highest Temp	86	90	102	109	116	121	122	122	120	112	96	92
Mean Max Temp	69.1	73.3	80.5	88.3	96.0	104.0	109.2	107.7	103.6	92.2	80.0	71.4
Mean Min Temp	37.8	41.7	47.2	53.7	60.7	66.7	75.2	75.6	69.0	57.5	45.2	40.2
Mean Temp	53.4	57.5	63.8	71.0	78.3	85.3	92.2	91.7	86.3	75.1	62.6	55.8
Lowest Temp	16	23	29	33	44	47	52	58	52	39	26	24
Mean Precip	0.30	0.46	0.23	0.07	T	0.02	0.10	0.30	0.38	0.25	0.08	0.62
<u>Imperial</u>												
Highest Temp	86	92	93	103	111	117	117	115	113	108	92	90
Mean Max Temp	69.7	73.4	78.6	85.9	93.3	102.5	106.6	104.5	102.7	91.6	78.1	71.1
Mean Min Temp	41.1	44.1	48.6	55.5	61.3	69.0	77.2	76.6	71.5	60.5	47.4	41.6
Mean Temp	54.1	58.0	63.6	70.9	77.8	85.1	91.8	90.8	86.0	74.8	62.3	55.9
Lowest Temp	26	29	33	41	47	53	63	60	54	44	32	26
Mean Precip	0.37	0.43	0.18	0.14	0.01	0.00	0.07	0.40	0.37	0.27	0.11	0.49

months the valley is almost 100% cloud free and these three winter months
months the valley is almost 100% cloud free and these three winter months
have more than 80% of the possible sunshine. Target timetables for imaging
have more than 80% of the possible sunshine. Target timetables for imaging
overflights will seldom have any problems with cloud cover.
overflights will seldom have any problems with cloud cover.

Appendix B

Interpretation, Mapping, and Analysis Procedures

Traditionally, land use mapping has been a long, tedious, manual process. Attempts are being made to reduce the task to a completely automatic process through the use of image or analysis equipment. The previously mentioned individual crop color variations often prevent automatic equipment from achieving satisfactory results. One may speculate that this problem will require some agricultural fields to always be examined individually. In fact, individual field examination is the basis for the system described. It is a system which can evolve from a manual method and be developed into a semiautomatic and perhaps automatic system. The system is not completely dependent on color signatures but takes advantage of environmental factors as well as other information available about agricultural practices in the study area. Basically, the process requires data to be read from the image by a human equipment operator, fed to the computer, and the computer (on the basis of information previously provided) will determine the type of land use. The procedure is:

- (1) Prepare base map,
- (2) Establish Crop Calendar for the date of the imagery,
- (3) Overlay base map on imagery or project imagery onto the base map,
- (4) Establish Crop Signature color values from ground survey test fields,
- (5) Determine and record field location,
- (6) Determine and record field color values by use of color i.e., densitometer or similar device,
- (7) Perform computer crop identification,
- (8) Prepare final agricultural land use map i.e., manually or by computer graphics.

Additional comments and discussion of the eight basic steps:

1. Base Map Preparation

Preparation of the local base map has been previously discussed.

2. Development of Crop Calendar for the Date of the Image

One of the more important steps in delimiting the number of crops which must be examined is to determine those crops under cultivation on the date of the imagery. The calendar is best established from the annual crop calendar for the local agricultural area being examined. (Table VII, is a List of Major Crops Under Cultivation on March 12 prepared for the date of the Apollo 9 photo of the Imperial Valley.) In the Imperial Valley there are approximately 48 different crops being cultivated throughout the year, but seldom are there more than half of these under cultivation at any one time. Crop calendars must be used with caution since they are only a "close approximation" of the average conditions and a shift in climate or market conditions can alter cultivation or harvesting plans. (i.e., sugar beets are shown on the Imperial Valley crop calendar as being completely harvested by mid-July. Yet in 1969 there were still sugar beets in the ground on the last day of July. This was caused by a three week trucker strike and mechanical failure in the Union Sugar Plant. In 1968, the planting of cotton was delayed 6 to 8 weeks because of predicted insect invasion.)

3. Image Projection or Base Map Overlay

Many errors have occurred and much time has been lost in performing land use mapping from imagery by attempting to view the image transparency

TABLE VII

List of Major Crops
Under Cultivation on March 12 in Imperial Valley
(From the Crop Calendar of the Imperial Valley)

A. Crops Showing Vegetative Cover

Field Crops

- 111 Barley (Mid-Season Stage)
- 118 Wheat (Mid-Season Stage)
- 131 Short Grasses (Rye, Bermuda)
- 133 Alfalfa
- 142 Sugar Beets (Near Full Maturity)
- 172 Flax (Blooming)

Vegetable Crops

- 211 Asparagus (Being Harvested)
- 231 Lettuce (Harvested, Being Plowed Under)
- 232 Celery
- 272 Carrots (End of Harvest Season)
- 282 Garlic
- 284 Onions (Under Harvest)

Fruit and Nut Crops

- 330 Citrus Fruit, Undifferentiated
- 342 Dates
- 363 Pecans (Blooming)

B. Crops Planted or Seeded and Irrigated, but Not Showing Vegetative Cover

Field Crops

- 114 Grain Sorghum (Planting Season One Week Old)
- 131 Tall Grasses (Sudan) (Planting Season Two Weeks Old)
- 151 Cotton (Fields Prepared, but Not Yet Seeded)

Vegetable Crops (Fields Seeded and Irrigated)

- 250 Melons, Undifferentiated
- 251 Cantaloupes (Planted with Caps On)
- 256 Watermelons (Planted with Caps On)
- 263 Tomatoes (Planted)

on a separate light table and transfer the interpreted data to some location on a base map. Efficient manual or automatic processing dictates that the image and the map be made coincident. Several methods exist for accomplishing this technique either by projecting the image onto the base map or overlaying the base map on the image depending which is larger and which system is more convenient. Several map-image projectors are on the market that can accomplish the coincidence and utilizing x-y recording plotters which can be programmed to read particular map coordinates from the image. The latter devices can be used to develop an automatic readout system. Plate 1 illustrates the method used in this study to project the image on the base map for interpretation of the Apollo 9 imagery of the Imperial Valley. Bringing the image into register with the base map was accomplished and slight rectification of the field patterns in each of the four quadrants of the map made the map and image coincident. An image edge enhancement resulted that permitted the eye to distinguish individual field colors which in the image alone often blended with the color of adjacent fields making precise determination difficult to achieve.

When the image and projections are planimetrically different, some difficulty will be experienced in registering a projected image onto a base map. Aerial photographic images are essentially orthographic map projections in which the distortion increases radially from the center of the image. The USGS topographic control map used and many maps used for similar studies are polyconic or like projections in which the distortion begins at a central meridian and increase outward in an east-west or north-south direction but not omni-directional. The disparities would be

small over small areas of the earth, but increase as the area increases. The enlarged 70 mm. Apollo photograph, covering an area approximately 100 miles (161 km) on each side, required very little rectification to bring it into register with the base map.

4. Establishing Crop Color Signature from Ground Survey Data

Sufficient research and experimental data has not been obtained to establish uniform crop color signature values at present. The values will have to be established from the imagery being analyzed. Enough information has been obtained to establish gross relative color differences between crop types, but it still is necessary to establish a single reference level of color tone from the particular imagery being analyzed. However, the task is not insurmountable. Initial color densitometric readings taken of a new set of imagery should be of ground survey test sites. The test fields will serve as the "training" areas for the interpreter (manual or automatic). Data from these readings can then be used to establish the color values placed in the Crop Signature tables of the computer program for crop identification.

5. Determine and Record Individual Field Location

The dual coordinate system established on the local base map as previously described permits two methods of field location to be entered into the computer data. When image data reduction is being performed manually, the field location can be recorded by the less complex local grid system. When data readout is being performed by use of x-y coordinate equipment, it may be desirable to record the geographic coordinates of the field which provides the computer with sufficient data to perform acreage calculations.

6. Determine and Record Individual Field Color Values

To maintain color consistency, color measurements should be performed with a color photo densitometer or similar device. An arrangement should be made whereby, if three color readings are necessary for each field, they should be made at the same time that the field coordinates are obtained. The three readings must all be of an identical spot using the different color separation filters (green, red, and blue) of the densitometer. Regular photo densitometers have spot sizes from 1 to 4 mm. If the field size is more than 5 times larger (15-25 mm.) than the densitometer spot size, it may be prudent to take at least three separate spot readings in a field and average color values.

7. Perform Computer Crop Identification

The above procedures have been designed so that recorded data for each field may be provided as a computer input for subsequent identification. To perform computer identification, pre-programming of the computer with identification tables is necessary. A minimum number and size of table can be achieved by utilizing the local crop calendar. The crop calendar consolidates most of the cultural and physical environmental factors into a concise list of the type of crops grown in the area and the dates these crops can possibly be grown. In the past, interpreters have qualitatively used the crop calendar in their work, but in establishing a computer program a more conscious effort must be employed to insure consideration of all factors and reduce expensive processing time. Utilizing a table search method for computer identification, the procedure may logically begin with the most important factor in CIR

imagery - the crop color. The color tone is matched as loosely as possible to those provided by the color table and a tentative identification established. As the environmental determinants have been considered in the crop calendar, the only other table necessary in the computer processing is a table of farming practice factors.

Resolution and available detail on the image will determine how many farming pattern factors can be used. Normally field size and shape will be available to help differentiate between crops. Other factors, such as number of rows per given distance or width of flood borders, will provide further delimiting information. If sequential imagery has been obtained and recorded for the area, referral to previous crop type may help to determine plausibility of the tentatively identified crop. For example, in the Imperial Valley, the cotton crop is not picked until after mid-October and since sugar beets must be planted before this date, it is not plausible that sugar beets would follow a cotton crop. Other similar crop rotations exist and are a helpful delimiter in crop identification.


There will usually be crops which the computer identification will indicate as two or three possibilities. The program may be designed to show weighted possibilities with the best choice indicated. Hopefully, few identification problems will force the interpreter to give individual attention before a final decision can be made. Based upon previous experience with other types of computer identification systems, unidentified crops should be about 3%.

8. Prepare Final Agricultural Land Use Map and Statistical Data

Two normally desired outputs, a statistical map and summary, can both be accomplished by a computer. A statistical summary printout can be prepared in almost any form desired. Section V lists a few such summaries. Computer mapping is not quite as common, but with the advent of computer graphics, more and more map displays will be produced. Current geographical computer mapping is being accomplished mostly by line printing using different symbols to produce a form of choropleth map which is not entirely satisfactory for land use mapping. With the aid of computer auxiliary equipment such as the Calcomp Plotter, a regular outline map can be prepared with the land use code indicated in each field.

A compromise between a completely computer prepared map and a completely hand drawn map, is to have the computer plotting paper produced with the outline base map as a background. Using the locally devised grid system, the computer plotter can then be instructed to insert the agricultural land use code in the appropriate fields. The advantage of the latter system is the ease with which a series of topical maps can be prepared for each crop type or crop class for study of distribution patterns. Preparation of such topical maps would also provide valuable support to the image enhancement and analysis equipment investigations providing the necessary target patterns for matching distinct color patterns of the enhancement equipment. (See Appendix D)

APPENDIX C-1 CROP CALENDAR FOR THE IMPERIAL VALLEY



APPENDIX C-2 AGRICULTURAL LAND USE CODE
 (A HORTICULTURAL/AGRONOMIC CLASSIFICATION)
 5 DIGIT CODE LISTING

8 RESOURCE PRODUCTION AND EXTRACTION

80000 RESOURCE PRODUCTION AND EXTRACTION, UNDIFFERENTIATED

81 AGRICULTURE

81000 AGRICULTURE, UNDIFFERENTIATED

811 FIELD AND SEED CROPS (EXTENSIVE FARMING)

8110 UNDIFFERENTIATED FIELD AND SEED CROPS

81100 FIELD AND SEED CROPS, UNDIFFERENTIATED

8111 CEREAL AND GRAIN CROPS

81110 CEREAL AND GRAIN CROPS, UNDIFFERENTIATED

81111 BARLEY

81112 RUCKWHEAT

81113 CORN(MAIZE)

81114 SORGHAM, GRAIN

81115 OATS

81116 RICE

81117 RYE

81118 WHEAT

81119 CEREAL AND GRAIN CROPS, OTHER DIFFERENTIATED

8112 LEGUMES FOR SEED CROPS

81120 LEGUMES FOR SEED)DRY*, UNDIFFERENTIATED

81121 BEANS, FIELD (DRIED BEFORE HARVEST)

81122 PEAS, FIELD (DRIED BEFORE HARVEST-INCLUDES-CHICKPEAS +COWPEAS

81123 LENTILS

81124 BEANS, LIMA (DRIED BEFORE HARVEST)

81125 PEANUTS

81126 SOYBEAN (FOOD)

81127

81128

81129 LEGUMES FOR SEED)DRY*, OTHER DIFFERENTIATED

8113 FORAGE CROPS (NON GRAINS)

81130 FORAGE CROPS, UNDIFFERENTIATED

81131 GRASSES, SHORT (I.E. BERMUDA, BLUEGRASS, TIMOTHY, ETC.)

81132 GRASSES, TALL (I.E. SUDAN/SORGHAM, CORN, ETC.)

81133 LEGUMES (I.E. ALFALEA, CLOVER, VETCH, ETC.)

81134 ROOTS (I.E. JERUSALEM ARTICHOKEs, MANGELS(PREFTS), ETC.)

81135

81136

81137

81138

81139 FORAGE CROPS, OTHER DIFFERENTIATED

8114 SUGAR CROPS

81140 SUGAR, CROPS, UNDIFFERENTIATED
81141 SUGAR CANE
81142 SUGAR BEETS
81143
81144
81145
81146
81147
81148
81149 SUGAR CROPS, OTHER DIFFERENTIATED FIELD CROPS

8115 FIBER CROPS

81150 FIBER CROPS, UNDIFFERENTIATED
81151 COTTON
81152 FIBER FLAX
81153 HEMP
81154 JUTE
81155 MILKWEED
81156 RAFFIA
81157 RAMIE
81158 RATTAN
81159 FIBER CROPS, OTHER DIFFERENTIATED

8116 BEVERAGE, DRUG, FLAVORING, OR SPICE CROPS

81160 BEVERAGE, SPICE, FLAVORING OR DRUG CROPS, UNDIFFERENTIATED
81161 BEVERAGE CROPS, UNDIFFERENTIATED
81162 COCO
81163 COFFEE
81164 TEA
81165 BEVERAGE CROPS, OTHER DIFFERENTIATED
81166 SPICE CROPS
81167 FLAVORING CROPS
81168 DRUG CROPS
81169 BEVERAGE, DRUG, FLAVORING, OR SPICE CROP, OTHER DIFFERENTIATED

8117 OIL CROPS

81170 OIL CROPS, UNDIFFERENTIATED
81171 CASTOR BEAN
81172 FLAX, SEED
81173 PERRILIA
81174 SAFFLOWER
81175 SESAME
81176 SOYBEAN (OIL)
81177
81178
81179 OIL CROPS, OTHER DIFFERENTIATED

8118 RUBBER CROPS

81180 RUBBER CROPS, UNDIFFERENTIATED FIELD
81181 GUAYULE
81182 KOK-SAGYZ
81183
81184

81185

81186

81187

81188

81189 RUBBER CROPS, OTHER DIFFERENTIATED

8119 OTHER DIFFERENTIATED FIELD AND SEED CROPS

81190 FIELD AND SEED CROPS, OTHER DIFFERENTIATED

812 VEGETABLE CROPS (INTENSIVE FARMING)

8120 UNDIFFERENTIATED VEGETABLE CROPS

81200 VEGETABLE CROPS, UNDIFFERENTIATED

8121 PERENNIAL VEGETABLE CROPS

81210 PERENNIAL VEGETABLE CROPS, UNDIFFERENTIATED

81210 VEGETABLE CROPS, PERENNIAL, UNDIFFERENTIATED

81211 ASPARAGUS

81212 ARTICHOKE, GLOBE

81213 HORSE RADISH

81214 RHUBARB

81215

81216

81217

81218

81219 PERENNIAL VEGETABLE CROPS, OTHER DIFFERENTIATED

81219 VEGETABLE CROPS, PERENNIAL, OTHER DIFFERENTIATED

8122 GREEN LEGUME (POD) CROPS

81220 LEGUME CROPS, GREEN, UNDIFFERENTIATED

81221 BEANS, GREEN (SNAP, POLE, KENTUCKY WONDERS, STRING, ETC)

81222 BEANS, GREEN LIMA

81223 PEAS, GREEN

81224 OKRA

81225 BLACK EYED PEAS (CANNED GREEN)

81226

81227

81228

81229 LEGUME CROPS, GREEN, OTHER DIFFERENTIATED

8123 SALAD AND GREENS CROPS

81230 SALAD AND GREENS CROPS, UNDIFFERENTIATED

81231 LETTUCE (SALAD)

81232 CELERY (SALAD)

81233 CRESS (SALAD)

81234 PARSLEY (SALAD)

81235 CHINESE CABBAGE (SALAD)

81236 CHARD AND KALE (GREENS)

81237 MUSTARD GREENS

81238 SPINACH GREENS

81239 SALAD AND GREEN CROPS, OTHER UNDIFFERENTIATED

8124 COLE CROPS

81240 COLE CROPS, UNDIFFERENTIATED

81241 BROCCOLI

81242 BRUSSELS SPROUTS

81243 CABBAGE

81244 CAULIFLOWER

81245 COLLARDS

81246 KOHLRABI

81247

81248

81249 COLE CROPS, OTHER DIFFERENTIATED

8125 CURCUBITS (VINE) CROPS

- 81250 CURCUBITS (VINE) CROPS, UNDIFFERENTIATED
- 81251 CANTALOUPE
- 81252 CUCUMBERS
- 81253 MELONS (OTHER THAN CANTALOUPE)
- 81254 PUMPKINS
- 81255 SQUASHES
- 81256 WATERMELONS
- 81257
- 81258
- 81259 OTHER DIFFERENTIATED CURCUBITS (VINE) CROPS

8126 SOLANACEOUS CROPS

- 81260 SOLANACEOUS CROPS, UNDIFFERENTIATED
- 81261 EGGPLANT
- 81262 PEPPERS
- 81263 TOMATOES
- 81264
- 81265
- 81266
- 81267
- 81268
- 81269 SOLANACEOUS CROPS, OTHER DIFFERENTIATED

8127 ROOT AND TUBER CROPS

- 81270 ROOT AND TUBER CROPS, UNDIFFERENTIATED
- 81271 BEETS (OTHER THAN SUGAR BEETS)
- 81272 CARROTS
- 81273 PARSNIPS
- 81274 POTATOES (TUBER)
- 81275 RADISHES
- 81276 RUTABAGAS
- 81277 SWEET POTATOES
- 81278 TURNIPS
- 81279 ROOT AND TUBER CROPS, OTHER DIFFERENTIATED

8128 BULB CROPS

- 81280 BULB CROPS, UNDIFFERENTIATED
- 81281 CHIVES
- 81282 GARLIC
- 81283 LEAKS
- 81284 ONIONS
- 81285
- 81286
- 81287
- 81288
- 81289 BULB CROPS, OTHER DIFFERENTIATED

8129 OTHER DIFFERENTIATED VEGETABLE CROPS

- 81290 VEGETABLE CROPS, OTHER DIFFERENTIATED

813 FRUIT AND NUT CROPS

8130 UNDIFFERENTIATED FRUITS OR NUTS

81300 FRUIT AND NUT CROPS, UNDIFFERENTIATED

8131 SMALL FRUITS

81310 FRUITS, SMALL, UNDIFFERENTIATED

81310 SMALL FRUITS, UNDIFFERENTIATED

81311 BRAMBLES (BLACKBERRY, DEWBERRY, BOYSONBERRY, RASPBERRY)

81312 PLUMBERRY (HUCKLEBERRY) AND CRANBERRY

81313 CURRANT AND GOOSEBERRY

81314 GRAPE

81315 STRAWBERRIES

81316

81317

81318

81319 FRUIT, SMALL, OTHER DIFFERENTIATED

81319 SMALL FRUIT, OTHER DIFFERENTIATED

8132 DECIDUOUS TREE FRUITS

81320 DECIDUOUS TREE FRUIT, UNDIFFERENTIATED

81320 FRUIT, TREE, DECIDUOUS, UNDIFFERENTIATED

81321 APPLE (INCLUDES- CRABAPPLE AND QUINCE)

81322 APRICOT

81323 CHERRY

81324 FIG

81325 NECTARINE

81326 PEACH

81327 PEAR

81328 PLUM (INCLUDES PRUNE*)

81329 DECIDUOUS TREE FRUIT, OTHER DIFFERENTIATED

81329 FRUIT, TREE, DECIDUOUS, OTHER DIFFERENTIATED

8133 CITRUS TREE FRUITS

81330 CITRUS TREE FRUIT, UNDIFFERENTIATED

81331 GRAPEFRUIT

81332 KUMQUAT

81333 LEMON

81334 LIME

81335 ORANGE

81336 TANGALO

81337 TANGERINE

81339

81339 CITRUS TREE FRUIT, OTHER DIFFERENTIATED

8134 MISCELLANEOUS EVERGREEN TREE FRUIT

81340 EVERGREEN TREE FRUIT, UNDIFFERENTIATED

81340 FRUIT, TREE, EVERGREEN, UNDIFFERENTIATED

81341 AVACADO

81342 DATE

81343 MANGO

81344 OLIVE

81345 PAPAYA

81346

81347
81348
81349 EVERGREEN TREE FRUIT, OTHER DIFFERENTIATED
81349 FRUIT, TREE, EVERGREEN, OTHER DIFFERENTIATED

8135 HERBACEOUS PERENNIAL FRUITS

81350 FRUIT, PERENNIAL, HERBACEOUS, UNDIFFERENTIATED
81350 HERBACEOUS PERENNIAL FRUIT, UNDIFFERENTIATED
81351 BANANA
81352 GUAVA
81353 PINAPPLE
81354
81356
81357
81358
81359 FRUIT, PERENNIAL, HERBACEOUS, OTHER DIFFERENTIATED
81359 HERBACEOUS PERENNIAL FRUIT, OTHER DIFFERENTIATED

8136 DECIDUOUS NUTS

81360 DECIDUOUS NUTS, UNDIFFERENTIATED
81360 NUTS, DECIDUOUS, UNDIFFERENTIATED
81361 ALMOND
81362 FILBERT (HAZELNUT)
81363 PECAN
81364 PISTACHIO
81365 WALNUT
81366 NUTMEG
81367
81368
81369 DECIDUOUS NUTS, OTHER DIFFERENTIATED
81369 NUTS, DECIDUOUS, OTHER DIFFERENTIATED

8137 EVERGREEN NUTS

81370 EVERGREEN NUTS, UNDIFFERENTIATED
81370 NUTS, EVERGREEN, UNDIFFERENTIATED
81371 BRAZIL
81372 CASHEW
81373 COCONUT
81374 LITCHI (LYCHEE)
81375 MACADAMIA
81376
81377
81378
81379 EVERGREEN NUTS, OTHER DIFFERENTIATED
81379 NUTS, EVERGREEN, OTHER DIFFERENTIATED

8138

8139 OTHER DIFFERENTIATED FRUIT AND NUT CROPS

81390 FRUIT AND NUT CROPS, OTHER DIFFERENTIATED

814 LIVESTOCK

8140 UNDIFFERENTIATED LIVESTOCK

81400 LIVESTOCK, UNDIFFERENTIATED

8141 BEEF CATTLE (OTHER THAN DAIRY-INCLUDES FEED LOTS)

81410 BEEF CATTLE (OTHER THAN DAIRY- INCLUDES FEED LOTS)

8142 HORSES

81420 HORSES

8143 SWINE

81430 SWINE

8144 DAIRIES AND DAIRY FEEDING

81440 DAIRIES AND DAIRY FEEDING

8145 SHEEP

81450 SHEEP

8146 GOAT

81460 GOAT

8147

8148

8149 OTHER DIFFERENTIATED LIVESTOCK

81490 LIVESTOCK, OTHER DIFFERENTIATED

815 ANIMAL SPECIALTIES

8150 UNDIFFERENTIATED SMALL ANIMAL PRODUCTION

81500 ANIMAL PRODUCTIONS, SMALL, UNDIFFERENTIATED

81500 SMALL ANIMAL PRODUCTION, UNDIFFERENTIATED

8151 CHICKEN (MEAT)

81510 CHICKEN (MEAT)

8152 CHICKEN (EGG)

81520 CHICKEN (EGG PRODUCTION)

8153 TURKEY

81530 TURKEY

8154 OTHER DIFFERENTIATED POULTRY

81540 POULTRY, OTHER DIFFERENTIATED

8155 RABBITS

81550 RABBITS

8156 APARIES

81560 APARIES

8157
8158

8159 OTHER DIFFERENTIATED ANIMAL SPECIALTIES
81590 ANIMAL SPECIALTIES, OTHER DIFFERENTIATED

816 PASTURE AND RANGELAND

8160 UNDIFFERENTIATED PASTURE AND RANGELAND
81600 PASTURE AND RANGELAND, UNDIFFERENTIATED

8161 PASTURE
81610 PASTURE

8162 RANGELAND
81620 RANGELAND

817 HORTICULTURAL SPECIALTIES

8170 UNDIFFERENTIATED HORTICULTURAL SPECIALTIES
81700 HORTICULTURAL SPECIALTIES, UNDIFFERENTIATED

8171 CUT FLOWERS STOCK, COVERED
81710 FLOWERS, COVERED CUT STOCK

8172 CUT FLOWERS STOCK, OPEN FIELD
81720 FLOWERS, OPEN FIELD CUT STOCK

8173 OTHER DIFFERENTIATED NURSARY STOCK
81730 NURSARY STOCK, OTHER DIFFERENTIATED

8174
8175
8176
8177
8178

8179 OTHER DIFFERENTIATED HORTICULTURAL SPECIALTIES
81790 HORTICULTURAL SPECIALTIES, OTHER DIFFERENTIATED

818 NON-PRODUCING AND TRANSITION CROP LAND

8180 UNDIFFERENTIATED NON-PRODUCING CROP LAND

81800 NON-PRODUCING CROP LAND, UNDIFFERENTIATED

8181 FALLOW CROP LAND

81810 FALLOW CROP LAND (NON-GROWING, NON-PASTURE, OR BARE)

8182 PLOWED CROP LAND

81820 PLOWED CROP LAND

8183 LEACHED CROP LAND

81830 LEACHED CROP LAND

8184 ABANDONED CROP LAND (TREE, VINE, HERBACEOUS, GRASS, FIELD, ETC.)

81840 ABANDONED CROP LAND (TREE, VINE, HERBACEOUS, GRASS, FIELD, ET

8185 HARVESTED FIELD (STUBBLE, INCLUDES CROP LAND OPEN TO CATTLE GRAZIN

81850 HARVESTED FIELD (STUBBLE, INCLUDES CROP LAND OPEN TO CATTLE)

8186 PREPARED CROP LAND (READY FOR SEEDING, OR SEEDED, AND/OR IRRIGATED

81860 PREPARED LAND (READY FOR SEEDING, OR SEEDED, AND/OR IRRIGATED

8187

8188

8189 OTHER DIFFERENTIATED NON-PRODUCING CROP LAND

81890 NON-PRODUCING CROP LAND, OTHER DIFFERENTIATED

819 OTHER AGRICULTURAL ACTIVITY

8190 UNDIFFERENTIATED OTHER AGRICULTURAL ACTIVITY

81900 AGRICULTURAL ACTIVITIES, OTHER, UNDIFFERENTIATED

81900 OTHER AGRICULTURAL ACTIVITY, UNDIFFERENTIATED

8191 TREE FARMS

81910 TREE FARMS

8199 OTHER DIFFERENTIATED OTHER AGRICULTURAL ACTIVITIES

81990 OTHER AGRICULTURAL ACTIVITIES, OTHER DIFFERENTIATED

82 AGRICULTURAL RELATED ACTIVITIES

8200 UNDIFFERENTIATED AGRICULTURAL RELATED ACTIVITIES

82000 AGRICULTURAL RELATED ACTIVITIES, UNDIFFERENTIATED

821 AGRICULTURAL PROCESSING

8210 UNDIFFERENTIATED AGRICULTURAL PROCESSING

82100 AGRICULTURAL PROCESSING, UNDIFFERENTIATED

8211 COTTON GINNING AND COMPRESSING

82110 COTTON GINNING AND COMPRESSING

8212 GRIST MILLING SERVICES

82120 GRIST MILLING SERVICES

8213 CORN SHELLING, HAY BALING, THRESHING SERVICES

82130 CORN SHELLING, HAY BALING, THRESHING SERVICES

8214 CONTRACT SORTING, GRADING, AND PACKAGING SERVICES

82140 CONTRACT SORTING, GRADING, AND PACKAGING SERVICES

8215 EGG PROCESSING

82150 EGG PROCESSING

8216

8217

8218

8219 OTHER DIFFERENTIATED AGRICULTURAL PROCESSING SERVICES

82190 AGRICULTURAL PROCESSING SERVICES, OTHER DIFFERENTIATED

822

823

824

825

826

827

828

829 OTHER AGRICULTURAL RELATED SERVICES

8290 OTHER AGRICULTURAL RELATED SERVICES, UNDIFFERENTIATED

82900 AGRICULTURAL RELATED SERVICES, OTHER, UNDIFFERENTIATED

82900 OTHER AGRICULTURAL RELATED SERVICES, UNDIFFERENTIATED

8291

8292 SPRAYING, DUSTING, PRUNING, PLANTING, TREE SURGERY

82920 SPRAYING, DUSTING, PRUNING, PLANTING, TREE SURGERY

8293 HORTICULTURAL SERVICES (INCLUDES - EXPERIMENTAL STATIONS*

82930 HORTICULTURAL SERVICES (INCLUDES-EXPERIMENTAL STATIONS)

8294

8295

8296

8297

8298

8299 OTHER DIFFERENTIATED OTHER AGRICULTURAL RELATED SERVICES

82990 AGRICULTURAL RELATED SERVICES, OTHER DIFFERENTIATED

82990 OTHER AGRICULTURAL RELATED SERVICES, OTHER DIFFERENTIATED

91 UNDEVELOPED AND UNUSED LAND AREA

91 UNDEVELOPED AND UNUSED LAND AREA

910 UNDEVELOPED AND UNUSED LAND AREA


9100 UNDIFFERENTIATED UNDEVELOPED AND UNUSED LAND AREA

91000 UNDEVELOPED AND UNUSED LAND AREA, UNDIFFERENTIATED

Appendix C-3 Agricultural Land Use Map of Imperial Valley

(This appendix will be found in the inside pocket of the back cover).

GROUND SURVEY ROUTES



AGRICULTURAL
LAND USE

APOLLO 9 MISSION 12 MARCH 69


IMPERIAL VALLEY


BY PLANS & CONTROL, DEPARTMENT OF AGRICULTURE, UNIVERSITY OF CALIFORNIA, STANFORD
THIS MAPLET WAS MADE BY JOHN WOODRUFF, AGRICULTURAL ENGINEER, STATION 1000, UNIVERSITY OF CALIFORNIA





Appendix D Thematic Land Use Maps of the Imperial Valley from March 12, 1969
Apollo 9 Photography


1. Barley, Alfalfa, and Sugar Beets	79
2. Barley	80
3. Alfalfa	81
4. Sugar Beets	82
5. Grasses (Exclusive of Barley and Alfalfa)	83
6. All Vegetables	84
7. Asparagus	85
8. Carrots	86
9. Lettuce	87
10. Onions	88
11. Plowed, Prepared or Seeded	89
12. Plowed	90
13. Prepared or Seeded	91
14. Fallow	92
15. Abandoned	93
16. Feed Lots	94












AGRICULTURAL LAND-USE CODE

TYPE OR	FIELD CODE	MAP TYPE
001	00	WATER
002	01	WATER
003	02	WATER
004	03	WATER
005	04	WATER
006	05	WATER
007	06	WATER
008	07	WATER
009	08	WATER
010	09	WATER
011	10	WATER
012	11	WATER
013	12	WATER
014	13	WATER
015	14	WATER
016	15	WATER
017	16	WATER
018	17	WATER
019	18	WATER
020	19	WATER
021	20	WATER
022	21	WATER
023	22	WATER
024	23	WATER
025	24	WATER
026	25	WATER
027	26	WATER
028	27	WATER
029	28	WATER
030	29	WATER
031	30	WATER
032	31	WATER
033	32	WATER
034	33	WATER
035	34	WATER
036	35	WATER
037	36	WATER
038	37	WATER
039	38	WATER
040	39	WATER
041	40	WATER
042	41	WATER
043	42	WATER
044	43	WATER
045	44	WATER
046	45	WATER
047	46	WATER
048	47	WATER
049	48	WATER
050	49	WATER
051	50	WATER
052	51	WATER
053	52	WATER
054	53	WATER
055	54	WATER
056	55	WATER
057	56	WATER
058	57	WATER
059	58	WATER
060	59	WATER
061	60	WATER
062	61	WATER
063	62	WATER
064	63	WATER
065	64	WATER
066	65	WATER
067	66	WATER
068	67	WATER
069	68	WATER
070	69	WATER
071	70	WATER
072	71	WATER
073	72	WATER
074	73	WATER
075	74	WATER
076	75	WATER
077	76	WATER
078	77	WATER
079	78	WATER
080	79	WATER
081	80	WATER
082	81	WATER
083	82	WATER
084	83	WATER
085	84	WATER
086	85	WATER
087	86	WATER
088	87	WATER
089	88	WATER
090	89	WATER
091	90	WATER
092	91	WATER
093	92	WATER
094	93	WATER
095	94	WATER
096	95	WATER
097	96	WATER
098	97	WATER
099	98	WATER
100	99	WATER
101	00	WATER
102	01	WATER
103	02	WATER
104	03	WATER
105	04	WATER
106	05	WATER
107	06	WATER
108	07	WATER
109	08	WATER
110	09	WATER
111	10	WATER
112	11	WATER
113	12	WATER
114	13	WATER
115	14	WATER
116	15	WATER
117	16	WATER
118	17	WATER
119	18	WATER
120	19	WATER
121	20	WATER
122	21	WATER
123	22	WATER
124	23	WATER
125	24	WATER
126	25	WATER
127	26	WATER
128	27	WATER
129	28	WATER
130	29	WATER
131	30	WATER
132	31	WATER
133	32	WATER
134	33	WATER
135	34	WATER
136	35	WATER
137	36	WATER
138	37	WATER
139	38	WATER
140	39	WATER
141	40	WATER
142	41	WATER
143	42	WATER
144	43	WATER
145	44	WATER
146	45	WATER
147	46	WATER
148	47	WATER
149	48	WATER
150	49	WATER
151	50	WATER
152	51	WATER
153	52	WATER
154	53	WATER
155	54	WATER
156	55	WATER
157	56	WATER
158	57	WATER
159	58	WATER
160	59	WATER
161	60	WATER
162	61	WATER
163	62	WATER
164	63	WATER
165	64	WATER
166	65	WATER
167	66	WATER
168	67	WATER
169	68	WATER
170	69	WATER
171	70	WATER
172	71	WATER
173	72	WATER
174	73	WATER
175	74	WATER
176	75	WATER
177	76	WATER
178	77	WATER
179	78	WATER
180	79	WATER
181	80	WATER
182	81	WATER
183	82	WATER
184	83	WATER
185	84	WATER
186	85	WATER
187	86	WATER
188	87	WATER
189	88	WATER
190	89	WATER
191	90	WATER
192	91	WATER
193	92	WATER
194	93	WATER
195	94	WATER
196	95	WATER
197	96	WATER
198	97	WATER
199	98	WATER
200	99	WATER
201	00	WATER
202	01	WATER
203	02	WATER
204	03	WATER
205	04	WATER
206	05	WATER
207	06	WATER
208	07	WATER
209	08	WATER
210	09	WATER
211	10	WATER
212	11	WATER
213	12	WATER
214	13	WATER
215	14	WATER
216	15	WATER
217	16	WATER
218	17	WATER
219	18	WATER
220	19	WATER
221	20	WATER
222	21	WATER
223	22	WATER
224	23	WATER
225	24	WATER
226	25	WATER
227	26	WATER
228	27	WATER
229	28	WATER
230	29	WATER
231	30	WATER
232	31	WATER
233	32	WATER
234	33	WATER
235	34	WATER
236	35	WATER
237	36	WATER
238	37	WATER
239	38	WATER
240	39	WATER
241	40	WATER
242	41	WATER
243	42	WATER
244	43	WATER
245	44	WATER
246	45	WATER
247	46	WATER
248	47	WATER
249	48	WATER
250	49	WATER
251	50	WATER
252	51	WATER
253	52	WATER
254	53	WATER
255	54	WATER
256	55	WATER
257	56	WATER
258	57	WATER
259	58	WATER
260	59	WATER
261	60	WATER
262	61	WATER
263	62	WATER
264	63	WATER
265	64	WATER
266	65	WATER
267	66	WATER
268	67	WATER
269	68	WATER
270	69	WATER
271	70	WATER
272	71	WATER
273	72	WATER
274	73	WATER
275	74	WATER
276	75	WATER
277	76	WATER
278	77	WATER
279	78	WATER
280	79	WATER
281	80	WATER
282	81	WATER
283	82	WATER
284	83	WATER
285	84	WATER
286	85	WATER
287	86	WATER
288	87	WATER
289	88	WATER
290	89	WATER
291	90	WATER
292	91	WATER
293	92	WATER
294	93	WATER
295	94	WATER
296	95	WATER
297	96	WATER
298	97	WATER
299	98	WATER
300	99	WATER

SPECIFIC CROPS: GRASSES (EXCL. BARLEY & ALF.)


AGRICULTURAL LAND USE
 APOLLO 9 MISSION 12 MARCH 69
IMPERIAL VALLEY



SPECIFIC CROPS: ASPARAGUS

AGRICULTURAL LAND-USE CODES

000	FIELD CROPS	000	WATER
010	WHEAT	010	RESERVOIR
020	BARLEY	020	LAKE
030	RYE	030	SWAMP
040	ORCHARD	040	WETLAND
050	VEGETABLE CROPS	050	WATERWAY
060	CORN	060	RAILROAD
070	SUGAR BEET	070	ROAD
080	SOYBEAN	080	TRAIL
090	WATERWAY	090	UNDEVELOPED
100	VEGETABLE CROPS	100	WATERWAY
110	CORN	110	RAILROAD
120	SUGAR BEET	120	ROAD
130	SOYBEAN	130	TRAIL
140	WATERWAY	140	UNDEVELOPED
150	VEGETABLE CROPS	150	WATERWAY
160	CORN	160	RAILROAD
170	SUGAR BEET	170	ROAD
180	SOYBEAN	180	TRAIL
190	WATERWAY	190	UNDEVELOPED
200	VEGETABLE CROPS	200	WATERWAY
210	CORN	210	RAILROAD
220	SUGAR BEET	220	ROAD
230	SOYBEAN	230	TRAIL
240	WATERWAY	240	UNDEVELOPED
250	VEGETABLE CROPS	250	WATERWAY
260	CORN	260	RAILROAD
270	SUGAR BEET	270	ROAD
280	SOYBEAN	280	TRAIL
290	WATERWAY	290	UNDEVELOPED
300	VEGETABLE CROPS	300	WATERWAY
310	CORN	310	RAILROAD
320	SUGAR BEET	320	ROAD
330	SOYBEAN	330	TRAIL
340	WATERWAY	340	UNDEVELOPED
350	VEGETABLE CROPS	350	WATERWAY
360	CORN	360	RAILROAD
370	SUGAR BEET	370	ROAD
380	SOYBEAN	380	TRAIL
390	WATERWAY	390	UNDEVELOPED
400	VEGETABLE CROPS	400	WATERWAY
410	CORN	410	RAILROAD
420	SUGAR BEET	420	ROAD
430	SOYBEAN	430	TRAIL
440	WATERWAY	440	UNDEVELOPED
450	VEGETABLE CROPS	450	WATERWAY
460	CORN	460	RAILROAD
470	SUGAR BEET	470	ROAD
480	SOYBEAN	480	TRAIL
490	WATERWAY	490	UNDEVELOPED
500	VEGETABLE CROPS	500	WATERWAY
510	CORN	510	RAILROAD
520	SUGAR BEET	520	ROAD
530	SOYBEAN	530	TRAIL
540	WATERWAY	540	UNDEVELOPED
550	VEGETABLE CROPS	550	WATERWAY
560	CORN	560	RAILROAD
570	SUGAR BEET	570	ROAD
580	SOYBEAN	580	TRAIL
590	WATERWAY	590	UNDEVELOPED
600	VEGETABLE CROPS	600	WATERWAY
610	CORN	610	RAILROAD
620	SUGAR BEET	620	ROAD
630	SOYBEAN	630	TRAIL
640	WATERWAY	640	UNDEVELOPED
650	VEGETABLE CROPS	650	WATERWAY
660	CORN	660	RAILROAD
670	SUGAR BEET	670	ROAD
680	SOYBEAN	680	TRAIL
690	WATERWAY	690	UNDEVELOPED
700	VEGETABLE CROPS	700	WATERWAY
710	CORN	710	RAILROAD
720	SUGAR BEET	720	ROAD
730	SOYBEAN	730	TRAIL
740	WATERWAY	740	UNDEVELOPED
750	VEGETABLE CROPS	750	WATERWAY
760	CORN	760	RAILROAD
770	SUGAR BEET	770	ROAD
780	SOYBEAN	780	TRAIL
790	WATERWAY	790	UNDEVELOPED
800	VEGETABLE CROPS	800	WATERWAY
810	CORN	810	RAILROAD
820	SUGAR BEET	820	ROAD
830	SOYBEAN	830	TRAIL
840	WATERWAY	840	UNDEVELOPED
850	VEGETABLE CROPS	850	WATERWAY
860	CORN	860	RAILROAD
870	SUGAR BEET	870	ROAD
880	SOYBEAN	880	TRAIL
890	WATERWAY	890	UNDEVELOPED
900	VEGETABLE CROPS	900	WATERWAY
910	CORN	910	RAILROAD
920	SUGAR BEET	920	ROAD
930	SOYBEAN	930	TRAIL
940	WATERWAY	940	UNDEVELOPED
950	VEGETABLE CROPS	950	WATERWAY
960	CORN	960	RAILROAD
970	SUGAR BEET	970	ROAD
980	SOYBEAN	980	TRAIL
990	WATERWAY	990	UNDEVELOPED





**AGRICULTURAL
LAND USE**


APOLLO 9 MISSION 12 MARCH 69


IMPERIAL VALLEY

U.S. GEOLOGICAL SURVEY, DEPARTMENT OF COMMERCE, UNIVERSITY OF CALIFORNIA, RIVERSIDE
SCALE: 1:50,000 (AS SHOWN ON MAP)









AGRICULTURAL LAND-USE CODE

CODE	FIELD CROPS	VEGETABLE CROPS	FRUIT AND NUT CROPS	OTHER
001	WHEAT	001	APPLES	001
002	BARLEY	002	ORANGES	002
003	RYE	003	LIMES	003
004	TRITICUM	004	LEMONS	004
005	MAIZE	005	PEACHES	005
006	SORGHUM	006	APRICOTS	006
007	WHEAT	007	PLUMS	007
008	BARLEY	008	CHERRIES	008
009	RYE	009	ALMONDS	009
010	TRITICUM	010	PEANUTS	010
011	MAIZE	011	WALNUTS	011
012	SORGHUM	012	CASHEWS	012
013	WHEAT	013	COCONUTS	013
014	BARLEY	014	PINEAPPLES	014
015	RYE	015	MANGOS	015
016	TRITICUM	016	GUAVAS	016
017	MAIZE	017	PAPAYAS	017
018	SORGHUM	018	AVOCADOS	018
019	WHEAT	019	PEACHES	019
020	BARLEY	020	APRICOTS	020
021	RYE	021	PLUMS	021
022	TRITICUM	022	CHERRIES	022
023	MAIZE	023	ALMONDS	023
024	SORGHUM	024	PEANUTS	024
025	WHEAT	025	WALNUTS	025
026	BARLEY	026	CASHEWS	026
027	RYE	027	COCONUTS	027
028	TRITICUM	028	PINEAPPLES	028
029	MAIZE	029	MANGOS	029
030	SORGHUM	030	GUAVAS	030
031	WHEAT	031	PAPAYAS	031
032	BARLEY	032	AVOCADOS	032
033	RYE	033	PEACHES	033
034	TRITICUM	034	APRICOTS	034
035	MAIZE	035	PLUMS	035
036	SORGHUM	036	CHERRIES	036
037	WHEAT	037	ALMONDS	037
038	BARLEY	038	PEANUTS	038
039	RYE	039	WALNUTS	039
040	TRITICUM	040	CASHEWS	040
041	MAIZE	041	COCONUTS	041
042	SORGHUM	042	PINEAPPLES	042
043	WHEAT	043	MANGOS	043
044	BARLEY	044	GUAVAS	044
045	RYE	045	PAPAYAS	045
046	TRITICUM	046	AVOCADOS	046
047	MAIZE	047	PEACHES	047
048	SORGHUM	048	APRICOTS	048
049	WHEAT	049	PLUMS	049
050	BARLEY	050	CHERRIES	050
051	RYE	051	ALMONDS	051
052	TRITICUM	052	PEANUTS	052
053	MAIZE	053	WALNUTS	053
054	SORGHUM	054	CASHEWS	054
055	WHEAT	055	COCONUTS	055
056	BARLEY	056	PINEAPPLES	056
057	RYE	057	MANGOS	057
058	TRITICUM	058	GUAVAS	058
059	MAIZE	059	PAPAYAS	059
060	SORGHUM	060	AVOCADOS	060
061	WHEAT	061	PEACHES	061
062	BARLEY	062	APRICOTS	062
063	RYE	063	PLUMS	063
064	TRITICUM	064	CHERRIES	064
065	MAIZE	065	ALMONDS	065
066	SORGHUM	066	PEANUTS	066
067	WHEAT	067	WALNUTS	067
068	BARLEY	068	CASHEWS	068
069	RYE	069	COCONUTS	069
070	TRITICUM	070	PINEAPPLES	070
071	MAIZE	071	MANGOS	071
072	SORGHUM	072	GUAVAS	072
073	WHEAT	073	PAPAYAS	073
074	BARLEY	074	AVOCADOS	074
075	RYE	075	PEACHES	075
076	TRITICUM	076	APRICOTS	076
077	MAIZE	077	PLUMS	077
078	SORGHUM	078	CHERRIES	078
079	WHEAT	079	ALMONDS	079
080	BARLEY	080	PEANUTS	080
081	RYE	081	WALNUTS	081
082	TRITICUM	082	CASHEWS	082
083	MAIZE	083	COCONUTS	083
084	SORGHUM	084	PINEAPPLES	084
085	WHEAT	085	MANGOS	085
086	BARLEY	086	GUAVAS	086
087	RYE	087	PAPAYAS	087
088	TRITICUM	088	AVOCADOS	088
089	MAIZE	089	PEACHES	089
090	SORGHUM	090	APRICOTS	090
091	WHEAT	091	PLUMS	091
092	BARLEY	092	CHERRIES	092
093	RYE	093	ALMONDS	093
094	TRITICUM	094	PEANUTS	094
095	MAIZE	095	WALNUTS	095
096	SORGHUM	096	CASHEWS	096
097	WHEAT	097	COCONUTS	097
098	BARLEY	098	PINEAPPLES	098
099	RYE	099	MANGOS	099
100	TRITICUM	100	GUAVAS	100

FEED LOTS

AGRICULTURAL LAND USE

APOLLO 9 MISSION 12 MARCH 69

IMPERIAL VALLEY

BY CLAUDE W. JENNINGS, DEPARTMENT OF GEOGRAPHY, UNIVERSITY OF CALIFORNIA, RIVERSIDE

THIS MAP IS A REPRODUCTION OF THE ORIGINAL DRAWING MADE BY THE AUTHOR IN 1968

SCALE: 1:50,000

PROJECTION: UTM

UNIT: METERS

DATE: 1969

REVISION: 1

AGRICULTURAL LAND USE

IMPERIAL VALLEY

APOLLO 9 MISSION 12 MARCH 69

BY CLAUDE W. JENNINGS, DEPARTMENT OF GEOGRAPHY, UNIVERSITY OF CALIFORNIA, RIVERSIDE

THIS MAP IS A REPRODUCTION OF THE ORIGINAL DRAWING MADE BY THE AUTHOR IN 1968

SCALE: 1:50,000

PROJECTION: UTM

UNIT: METERS

DATE: 1969

REVISION: 1

AGRICULTURAL LAND USE

IMPERIAL VALLEY

APOLLO 9 MISSION 12 MARCH 69

BY CLAUDE W. JENNINGS, DEPARTMENT OF GEOGRAPHY, UNIVERSITY OF CALIFORNIA, RIVERSIDE

THIS MAP IS A REPRODUCTION OF THE ORIGINAL DRAWING MADE BY THE AUTHOR IN 1968


SCALE: 1:50,000

PROJECTION: UTM

UNIT: METERS

DATE: 1969

REVISION: 1




AGRICULTURAL LAND-USE CODE SHEET

CODE	FIELD SYMBOL	DESCRIPTION
001	[Symbol]	FIELD CROPS
002	[Symbol]	WHEAT
003	[Symbol]	BARLEY
004	[Symbol]	RYE
005	[Symbol]	ORCHARD
006	[Symbol]	WATERMELON
007	[Symbol]	CUCUMBER
008	[Symbol]	PUMPKIN
009	[Symbol]	PEPPER
010	[Symbol]	EGGPLANT
011	[Symbol]	CARROT
012	[Symbol]	CELERY
013	[Symbol]	SPINACH
014	[Symbol]	LETTUCE
015	[Symbol]	ONION
016	[Symbol]	POTATO
017	[Symbol]	BEAN
018	[Symbol]	PEANUT
019	[Symbol]	ALMOND
020	[Symbol]	ORANGE
021	[Symbol]	LEMON
022	[Symbol]	LIME
023	[Symbol]	GRAPE
024	[Symbol]	DATE
025	[Symbol]	AVOCADO
026	[Symbol]	APPLE
027	[Symbol]	PEAR
028	[Symbol]	CHERRY
029	[Symbol]	PLUM
030	[Symbol]	PRUNE
031	[Symbol]	ALMOND
032	[Symbol]	WALNUT
033	[Symbol]	CHESTNUT
034	[Symbol]	PALM
035	[Symbol]	COCONUT
036	[Symbol]	DATE PALM
037	[Symbol]	AVOCADO PALM
038	[Symbol]	ORANGE PALM
039	[Symbol]	LEMON PALM
040	[Symbol]	LIME PALM
041	[Symbol]	GRAPE PALM
042	[Symbol]	DATE PALM
043	[Symbol]	AVOCADO PALM
044	[Symbol]	ORANGE PALM
045	[Symbol]	LEMON PALM
046	[Symbol]	LIME PALM
047	[Symbol]	GRAPE PALM
048	[Symbol]	DATE PALM
049	[Symbol]	AVOCADO PALM
050	[Symbol]	ORANGE PALM
051	[Symbol]	LEMON PALM
052	[Symbol]	LIME PALM
053	[Symbol]	GRAPE PALM
054	[Symbol]	DATE PALM
055	[Symbol]	AVOCADO PALM
056	[Symbol]	ORANGE PALM
057	[Symbol]	LEMON PALM
058	[Symbol]	LIME PALM
059	[Symbol]	GRAPE PALM
060	[Symbol]	DATE PALM
061	[Symbol]	AVOCADO PALM
062	[Symbol]	ORANGE PALM
063	[Symbol]	LEMON PALM
064	[Symbol]	LIME PALM
065	[Symbol]	GRAPE PALM
066	[Symbol]	DATE PALM
067	[Symbol]	AVOCADO PALM
068	[Symbol]	ORANGE PALM
069	[Symbol]	LEMON PALM
070	[Symbol]	LIME PALM
071	[Symbol]	GRAPE PALM
072	[Symbol]	DATE PALM
073	[Symbol]	AVOCADO PALM
074	[Symbol]	ORANGE PALM
075	[Symbol]	LEMON PALM
076	[Symbol]	LIME PALM
077	[Symbol]	GRAPE PALM
078	[Symbol]	DATE PALM
079	[Symbol]	AVOCADO PALM
080	[Symbol]	ORANGE PALM
081	[Symbol]	LEMON PALM
082	[Symbol]	LIME PALM
083	[Symbol]	GRAPE PALM
084	[Symbol]	DATE PALM
085	[Symbol]	AVOCADO PALM
086	[Symbol]	ORANGE PALM
087	[Symbol]	LEMON PALM
088	[Symbol]	LIME PALM
089	[Symbol]	GRAPE PALM
090	[Symbol]	DATE PALM
091	[Symbol]	AVOCADO PALM
092	[Symbol]	ORANGE PALM
093	[Symbol]	LEMON PALM
094	[Symbol]	LIME PALM
095	[Symbol]	GRAPE PALM
096	[Symbol]	DATE PALM
097	[Symbol]	AVOCADO PALM
098	[Symbol]	ORANGE PALM
099	[Symbol]	LEMON PALM
100	[Symbol]	LIME PALM
101	[Symbol]	GRAPE PALM
102	[Symbol]	DATE PALM
103	[Symbol]	AVOCADO PALM
104	[Symbol]	ORANGE PALM
105	[Symbol]	LEMON PALM
106	[Symbol]	LIME PALM
107	[Symbol]	GRAPE PALM
108	[Symbol]	DATE PALM
109	[Symbol]	AVOCADO PALM
110	[Symbol]	ORANGE PALM
111	[Symbol]	LEMON PALM
112	[Symbol]	LIME PALM
113	[Symbol]	GRAPE PALM
114	[Symbol]	DATE PALM
115	[Symbol]	AVOCADO PALM
116	[Symbol]	ORANGE PALM
117	[Symbol]	LEMON PALM
118	[Symbol]	LIME PALM
119	[Symbol]	GRAPE PALM
120	[Symbol]	DATE PALM
121	[Symbol]	AVOCADO PALM
122	[Symbol]	ORANGE PALM
123	[Symbol]	LEMON PALM
124	[Symbol]	LIME PALM
125	[Symbol]	GRAPE PALM
126	[Symbol]	DATE PALM
127	[Symbol]	AVOCADO PALM
128	[Symbol]	ORANGE PALM
129	[Symbol]	LEMON PALM
130	[Symbol]	LIME PALM
131	[Symbol]	GRAPE PALM
132	[Symbol]	DATE PALM
133	[Symbol]	AVOCADO PALM
134	[Symbol]	ORANGE PALM
135	[Symbol]	LEMON PALM
136	[Symbol]	LIME PALM
137	[Symbol]	GRAPE PALM
138	[Symbol]	DATE PALM
139	[Symbol]	AVOCADO PALM
140	[Symbol]	ORANGE PALM
141	[Symbol]	LEMON PALM
142	[Symbol]	LIME PALM
143	[Symbol]	GRAPE PALM
144	[Symbol]	DATE PALM
145	[Symbol]	AVOCADO PALM
146	[Symbol]	ORANGE PALM
147	[Symbol]	LEMON PALM
148	[Symbol]	LIME PALM
149	[Symbol]	GRAPE PALM
150	[Symbol]	DATE PALM
151	[Symbol]	AVOCADO PALM
152	[Symbol]	ORANGE PALM
153	[Symbol]	LEMON PALM
154	[Symbol]	LIME PALM
155	[Symbol]	GRAPE PALM
156	[Symbol]	DATE PALM
157	[Symbol]	AVOCADO PALM
158	[Symbol]	ORANGE PALM
159	[Symbol]	LEMON PALM
160	[Symbol]	LIME PALM
161	[Symbol]	GRAPE PALM
162	[Symbol]	DATE PALM
163	[Symbol]	AVOCADO PALM
164	[Symbol]	ORANGE PALM
165	[Symbol]	LEMON PALM
166	[Symbol]	LIME PALM
167	[Symbol]	GRAPE PALM
168	[Symbol]	DATE PALM
169	[Symbol]	AVOCADO PALM
170	[Symbol]	ORANGE PALM
171	[Symbol]	LEMON PALM
172	[Symbol]	LIME PALM
173	[Symbol]	GRAPE PALM
174	[Symbol]	DATE PALM
175	[Symbol]	AVOCADO PALM
176	[Symbol]	ORANGE PALM
177	[Symbol]	LEMON PALM
178	[Symbol]	LIME PALM
179	[Symbol]	GRAPE PALM
180	[Symbol]	DATE PALM
181	[Symbol]	AVOCADO PALM
182	[Symbol]	ORANGE PALM
183	[Symbol]	LEMON PALM
184	[Symbol]	LIME PALM
185	[Symbol]	GRAPE PALM
186	[Symbol]	DATE PALM
187	[Symbol]	AVOCADO PALM
188	[Symbol]	ORANGE PALM
189	[Symbol]	LEMON PALM
190	[Symbol]	LIME PALM
191	[Symbol]	GRAPE PALM
192	[Symbol]	DATE PALM
193	[Symbol]	AVOCADO PALM
194	[Symbol]	ORANGE PALM
195	[Symbol]	LEMON PALM
196	[Symbol]	LIME PALM
197	[Symbol]	GRAPE PALM
198	[Symbol]	DATE PALM
199	[Symbol]	AVOCADO PALM
200	[Symbol]	ORANGE PALM

PLOWED
PREPARED OR
SEEDED

AGRICULTURAL
LAND USE
APOLLO 9 MISSION 12 MARCH 69
IMPERIAL VALLEY
BY LANCE W. GARDNER, DEPARTMENT OF SOILS, UNIVERSITY OF CALIFORNIA, BERKELEY
1969 EDITION. SCALE: 1:50,000. PHOTOGRAPHIC SOURCE: NASA, TM 69-104, PART 1, FIG. 104-1




AGRICULTURAL LAND USE CODE (80)

CODE	FIELD CODE	DESCRIPTION
001	00	PLANTED
002	01	PLANTED
003	02	PLANTED
004	03	PLANTED
005	04	PLANTED
006	05	PLANTED
007	06	PLANTED
008	07	PLANTED
009	08	PLANTED
010	09	PLANTED
011	10	PLANTED
012	11	PLANTED
013	12	PLANTED
014	13	PLANTED
015	14	PLANTED
016	15	PLANTED
017	16	PLANTED
018	17	PLANTED
019	18	PLANTED
020	19	PLANTED
021	20	PLANTED
022	21	PLANTED
023	22	PLANTED
024	23	PLANTED
025	24	PLANTED
026	25	PLANTED
027	26	PLANTED
028	27	PLANTED
029	28	PLANTED
030	29	PLANTED
031	30	PLANTED
032	31	PLANTED
033	32	PLANTED
034	33	PLANTED
035	34	PLANTED
036	35	PLANTED
037	36	PLANTED
038	37	PLANTED
039	38	PLANTED
040	39	PLANTED
041	40	PLANTED
042	41	PLANTED
043	42	PLANTED
044	43	PLANTED
045	44	PLANTED
046	45	PLANTED
047	46	PLANTED
048	47	PLANTED
049	48	PLANTED
050	49	PLANTED
051	50	PLANTED
052	51	PLANTED
053	52	PLANTED
054	53	PLANTED
055	54	PLANTED
056	55	PLANTED
057	56	PLANTED
058	57	PLANTED
059	58	PLANTED
060	59	PLANTED
061	60	PLANTED
062	61	PLANTED
063	62	PLANTED
064	63	PLANTED
065	64	PLANTED
066	65	PLANTED
067	66	PLANTED
068	67	PLANTED
069	68	PLANTED
070	69	PLANTED
071	70	PLANTED
072	71	PLANTED
073	72	PLANTED
074	73	PLANTED
075	74	PLANTED
076	75	PLANTED
077	76	PLANTED
078	77	PLANTED
079	78	PLANTED
080	79	PLANTED
081	80	PLANTED
082	81	PLANTED
083	82	PLANTED
084	83	PLANTED
085	84	PLANTED
086	85	PLANTED
087	86	PLANTED
088	87	PLANTED
089	88	PLANTED
090	89	PLANTED
091	90	PLANTED
092	91	PLANTED
093	92	PLANTED
094	93	PLANTED
095	94	PLANTED
096	95	PLANTED
097	96	PLANTED
098	97	PLANTED
099	98	PLANTED
100	99	PLANTED
101	00	PLANTED
102	01	PLANTED
103	02	PLANTED
104	03	PLANTED
105	04	PLANTED
106	05	PLANTED
107	06	PLANTED
108	07	PLANTED
109	08	PLANTED
110	09	PLANTED
111	10	PLANTED
112	11	PLANTED
113	12	PLANTED
114	13	PLANTED
115	14	PLANTED
116	15	PLANTED
117	16	PLANTED
118	17	PLANTED
119	18	PLANTED
120	19	PLANTED
121	20	PLANTED
122	21	PLANTED
123	22	PLANTED
124	23	PLANTED
125	24	PLANTED
126	25	PLANTED
127	26	PLANTED
128	27	PLANTED
129	28	PLANTED
130	29	PLANTED
131	30	PLANTED
132	31	PLANTED
133	32	PLANTED
134	33	PLANTED
135	34	PLANTED
136	35	PLANTED
137	36	PLANTED
138	37	PLANTED
139	38	PLANTED
140	39	PLANTED
141	40	PLANTED
142	41	PLANTED
143	42	PLANTED
144	43	PLANTED
145	44	PLANTED
146	45	PLANTED
147	46	PLANTED
148	47	PLANTED
149	48	PLANTED
150	49	PLANTED
151	50	PLANTED
152	51	PLANTED
153	52	PLANTED
154	53	PLANTED
155	54	PLANTED
156	55	PLANTED
157	56	PLANTED
158	57	PLANTED
159	58	PLANTED
160	59	PLANTED
161	60	PLANTED
162	61	PLANTED
163	62	PLANTED
164	63	PLANTED
165	64	PLANTED
166	65	PLANTED
167	66	PLANTED
168	67	PLANTED
169	68	PLANTED
170	69	PLANTED
171	70	PLANTED
172	71	PLANTED
173	72	PLANTED
174	73	PLANTED
175	74	PLANTED
176	75	PLANTED
177	76	PLANTED
178	77	PLANTED
179	78	PLANTED
180	79	PLANTED
181	80	PLANTED
182	81	PLANTED
183	82	PLANTED
184	83	PLANTED
185	84	PLANTED
186	85	PLANTED
187	86	PLANTED
188	87	PLANTED
189	88	PLANTED
190	89	PLANTED
191	90	PLANTED
192	91	PLANTED
193	92	PLANTED
194	93	PLANTED
195	94	PLANTED
196	95	PLANTED
197	96	PLANTED
198	97	PLANTED
199	98	PLANTED
200	99	PLANTED

PLOWED

AGRICULTURAL
LAND USE
APOLLO 9 MISSION 12 MARCH 69
IMPERIAL VALLEY

U.S. DEPARTMENT OF AGRICULTURE, AGRICULTURAL RESEARCH SERVICE
LAND USE MAPS OF THE IMPERIAL VALLEY - DISTRIBUTION RECORD - 1969




AGRICULTURAL LAND-USE CODE 100

FIELD CODE	FIELD TYPE
001	AGRICULTURAL LAND
002	AGRICULTURAL LAND
003	AGRICULTURAL LAND
004	AGRICULTURAL LAND
005	AGRICULTURAL LAND
006	AGRICULTURAL LAND
007	AGRICULTURAL LAND
008	AGRICULTURAL LAND
009	AGRICULTURAL LAND
010	AGRICULTURAL LAND
011	AGRICULTURAL LAND
012	AGRICULTURAL LAND
013	AGRICULTURAL LAND
014	AGRICULTURAL LAND
015	AGRICULTURAL LAND
016	AGRICULTURAL LAND
017	AGRICULTURAL LAND
018	AGRICULTURAL LAND
019	AGRICULTURAL LAND
020	AGRICULTURAL LAND
021	AGRICULTURAL LAND
022	AGRICULTURAL LAND
023	AGRICULTURAL LAND
024	AGRICULTURAL LAND
025	AGRICULTURAL LAND
026	AGRICULTURAL LAND
027	AGRICULTURAL LAND
028	AGRICULTURAL LAND
029	AGRICULTURAL LAND
030	AGRICULTURAL LAND
031	AGRICULTURAL LAND
032	AGRICULTURAL LAND
033	AGRICULTURAL LAND
034	AGRICULTURAL LAND
035	AGRICULTURAL LAND
036	AGRICULTURAL LAND
037	AGRICULTURAL LAND
038	AGRICULTURAL LAND
039	AGRICULTURAL LAND
040	AGRICULTURAL LAND
041	AGRICULTURAL LAND
042	AGRICULTURAL LAND
043	AGRICULTURAL LAND
044	AGRICULTURAL LAND
045	AGRICULTURAL LAND
046	AGRICULTURAL LAND
047	AGRICULTURAL LAND
048	AGRICULTURAL LAND
049	AGRICULTURAL LAND
050	AGRICULTURAL LAND
051	AGRICULTURAL LAND
052	AGRICULTURAL LAND
053	AGRICULTURAL LAND
054	AGRICULTURAL LAND
055	AGRICULTURAL LAND
056	AGRICULTURAL LAND
057	AGRICULTURAL LAND
058	AGRICULTURAL LAND
059	AGRICULTURAL LAND
060	AGRICULTURAL LAND
061	AGRICULTURAL LAND
062	AGRICULTURAL LAND
063	AGRICULTURAL LAND
064	AGRICULTURAL LAND
065	AGRICULTURAL LAND
066	AGRICULTURAL LAND
067	AGRICULTURAL LAND
068	AGRICULTURAL LAND
069	AGRICULTURAL LAND
070	AGRICULTURAL LAND
071	AGRICULTURAL LAND
072	AGRICULTURAL LAND
073	AGRICULTURAL LAND
074	AGRICULTURAL LAND
075	AGRICULTURAL LAND
076	AGRICULTURAL LAND
077	AGRICULTURAL LAND
078	AGRICULTURAL LAND
079	AGRICULTURAL LAND
080	AGRICULTURAL LAND
081	AGRICULTURAL LAND
082	AGRICULTURAL LAND
083	AGRICULTURAL LAND
084	AGRICULTURAL LAND
085	AGRICULTURAL LAND
086	AGRICULTURAL LAND
087	AGRICULTURAL LAND
088	AGRICULTURAL LAND
089	AGRICULTURAL LAND
090	AGRICULTURAL LAND
091	AGRICULTURAL LAND
092	AGRICULTURAL LAND
093	AGRICULTURAL LAND
094	AGRICULTURAL LAND
095	AGRICULTURAL LAND
096	AGRICULTURAL LAND
097	AGRICULTURAL LAND
098	AGRICULTURAL LAND
099	AGRICULTURAL LAND
100	AGRICULTURAL LAND

PREPARED OR SEEDED

AGRICULTURAL LAND USE
 APOLLO 9 MISSION 12 MARCH 69
IMPERIAL VALLEY

BY CLAUDE W. JORDAN, DEPARTMENT OF SOILS, UNIVERSITY OF CALIFORNIA, RIVERSIDE
 1969 SOURCE: DATA BY AIRCRAFT PHOTOGRAMMETRY - PHOTO INTERPRETATION ASSISTED BY DATA FROM AERIAL PHOTOGRAMMETRY



AGRICULTURAL LAND-USE CODE KEY

CODE NO.	FIELD CROPS	VEGETABLE CROPS	LIVESTOCK	OTHER
001	BARLEY	ARTICHOKES	CATTLE	WATER
002	WHEAT	ASPARAGUS	HOGS	WATERWAYS
003	ORZONIA	BROCCOLI	SHEEP	WATERWAYS
004	MAIZE	CABBAGE	GOATS	WATERWAYS
005	RYE	CARROTS	PORCUPINES	WATERWAYS
006	BARLEY	CUCUMBERS	CHICKENS	WATERWAYS
007	WHEAT	EGGPLANT	DUCKS	WATERWAYS
008	ORZONIA	KALE	GESE	WATERWAYS
009	MAIZE	LETTUCE	GUINEA FOWL	WATERWAYS
010	RYE	PUMPKIN	TRAIL	WATERWAYS
011	BARLEY	SPINACH	WATERWAYS	WATERWAYS
012	WHEAT	SWEET CORN	WATERWAYS	WATERWAYS
013	ORZONIA	TOMATOES	WATERWAYS	WATERWAYS
014	MAIZE	TURNIPS	WATERWAYS	WATERWAYS
015	RYE	WATERWAYS	WATERWAYS	WATERWAYS
016	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
017	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
018	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
019	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
020	RYE	WATERWAYS	WATERWAYS	WATERWAYS
021	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
022	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
023	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
024	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
025	RYE	WATERWAYS	WATERWAYS	WATERWAYS
026	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
027	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
028	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
029	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
030	RYE	WATERWAYS	WATERWAYS	WATERWAYS
031	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
032	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
033	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
034	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
035	RYE	WATERWAYS	WATERWAYS	WATERWAYS
036	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
037	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
038	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
039	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
040	RYE	WATERWAYS	WATERWAYS	WATERWAYS
041	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
042	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
043	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
044	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
045	RYE	WATERWAYS	WATERWAYS	WATERWAYS
046	BARLEY	WATERWAYS	WATERWAYS	WATERWAYS
047	WHEAT	WATERWAYS	WATERWAYS	WATERWAYS
048	ORZONIA	WATERWAYS	WATERWAYS	WATERWAYS
049	MAIZE	WATERWAYS	WATERWAYS	WATERWAYS
050	RYE	WATERWAYS	WATERWAYS	WATERWAYS

AGRICULTURAL LAND USE
 APOLLO 9 MISSION 12 MARCH 69
IMPERIAL VALLEY
 BY CLAUDE H. JOHNSON, DEPARTMENT OF SOILS, UNIVERSITY OF CALIFORNIA, RIVERSIDE
 THIS MAP WAS MADE AS PART OF THE AGRICULTURAL LAND USE SURVEY OF THE IMPERIAL VALLEY, CALIFORNIA

Plate 2 (Upper left)

Examples of varieties of crop colors on CIR imagery. (Original scale of 1:16,000 and resolution of 40 cm taken with filter combination of W 15 plus 8)B). The scale, resolution, and number of fields available on the same frame for comparison presents an excellent image for agricultural land use interpretation.

Identification Key to Plate 2.

- 111 Barley (or wheat)
- 133 Alfalfa (all stages of maturity)
- 151 Cotton
- 251 Cantaloupe
- 263 Tomatoes
- 410 Feed Lot (Beef cattle)
- 810 Fallow Land
- 820 Plowed Land
- 850 Harvest Land (Stubble)

	151		151		263
111	133				
111	850		820		151
111			850		
	133		133		151
					810
	410				151
		2	5	1	133 151

Plate 3 (Upper Right)

The 160 acre alfalfa crop in the center field of the picture, shown in three stages of cutting, is an example of CIR image color variation within the same field. The lower 3/5ths of the field has been raked into windrows for drying. The middle 3/10ths of the field has been cut and is waiting to be raked. The upper 1/10th is fully matured standing alfalfa.

Plate 4 (Lower left)

Examples of the same type crop (alfalfa) showing variations in color on CIR imagery due to different stages of growth.





Identification Key to Plate 4.

- 1 Cut and pastured Alfalfa field
- 2 Medium regrowth field of Alfalfa
- 3 Advanced regrowth field of Alfalfa
- 4 & 5 Mature Alfalfa field under irrigation
- 6 Fully matured Alfalfa field

	2	
	6	3
5	1	4

Plate 5 (Lower right)

Example of low altitude, high resolution CIR imagery showing two different crop types reflecting the same false color. Center crop is mature sugar beets. Crop in upper right corner is young sorgham.








Plate 6 (Upper left)

Example of the effect of high resolution on CIR imagery. The false red color indicating the infrared reflectance of the vegetation (sugar beets) is separated from the brownish-yellow color of the dead leaves and soil background. Photo was taken at 2000 foot altitude with a 6 inch focal length RC-8 camera on May 21, 1968. (Ground resolution approximately 10 cm.)

Plate 7 (Upper left)

Same sugar beet field as in plate 6 showing how the false red vegetation color is integrated with the brownish-yellow background color on CIR imagery at medium resolution. Photo taken on May 24, 1968.

Plate 8 (Lower left)

The effect of time-lapse imagery is shown in this image of the same sugar beet field as in plates 6 and 7 taken at medium resolution 21 days later (June 11, 1968). The fallow brown soil color indicates the field has now been harvested. The faded or cloudy appearance of the print is due to smoke drifting under the flight path from the burning of barley stubble on the day of the mission.

Plate 9 (Lower right)

Ground CIR photo of a mature sugar beet field taken in May 1968 showing the reduced amount of infrared reflectance from the leaves of the plant. Many leaves which appear green in normal color show no IR reflectance in this picture and blend into the background color of the soil. The combined color of the dead leaves and the background soil will integrate with the IR reflectance to present an integrated color on medium resolution imagery which appears as a light pink compared to the vivid red that is seen on the ground photo.

STANDARD TITLE PAGE FOR TECHNICAL REPORTS	1. Report No. USGS-DO-71-009	2. Govt. Accession No.	3. Recipient's Catalog No.
4. Title and Subtitle A System of Regional Agricultural Land Use Mapping Tested Against Small-Scale Apollo 9 Color Infrared Photography of the Imperial Valley (California)		5. Report Date November 1969	
7. Author(s) Claude W. Johnson, Leonard W. Bowden, and Robert W. Pease, Univ. of California, Riverside		6. Performing Organization Code	
9. Performing Organization Name and Address U.S. Geological Survey Geographical Applications Program Washington, D.C. 20242		8. Performing Organization Report No. TR-USGS-183	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546		10. Project Task Work Unit No. 160-75-01-32-10	
15. Supplementary Notes Work performed by the University of California for the USGS Geographic Applications Program under USGS contract no. 14-08-0001-10674		11. Contract Grant No. NASA W-12570 USGS 14-08-0001-10674	
16. Abstracts Interpretation results of the small scale CIR photography of the Imperial Valley (California) taken on March 12, 1969 by the Apollo 9 earth orbiting satellite have shown that world wide agricultural land use mapping can be accomplished from satellite CIR imagery if sufficient <u>a priori</u> information is available for the region being mapped. Correlation of results with actual data is encouraging although the accuracy of identification of specific crops from the single image is poor. The poor results can be partly attributed to only one image taken during mid-season when the three major crops were reflecting approximately the same and their CIR image appears to indicate the same crop type. However, some incapacity can be attributed to lack of understanding of the subtle variations of visual and infrared color reflectance of vegetation and surrounding environment. Analysis of integrated color variations of the vegetation and background environment recorded on CIR imagery is discussed. Problems associated with the color variations may be overcome by development of a semi-automatic processing system which considers individual field units or cells.		13. Type of Report & Period Covered Final report	
17. Key Words and Document Analysis. (a). Descriptors Land use, Mapping, Spaceborne photography, Infrared photography		14. Sponsoring Agency Code	
17b. Identifiers/Open-Ended Terms Apollo 9		<h1>CASE FILE COPY</h1>	
17c. COSATI Field/Group 8B, 8F, 14E			
18. Distribution Statement Unclassified - Unlimited		19. Security Class (This Report) UNCLASSIFIED	21. No. of Pages 107
		20. Security Class. (This Page) UNCLASSIFIED	22. Price \$3.00