






PLASMA PHYSICS GROUP



UNIVERSITY OF CALIFORNIA
LOS ANGELES



NASA CR-
122447

AD741 681

Get DRA

(NASA CR-122447) ELECTROSTATIC INSTABILITY
OF RING CURRENT PROTONS BEYOND THE
PLASMAPAUSE DURING INJECTION EVENTS
Coroniti, et al (California Univ.) F.V.
1972 17 p

REC
NASA
IMP

CSC 03B G3/13
May

Unclass
15856

N72-27347

*Catalyzer : see
page 7 for support*

ELECTROSTATIC INSTABILITY OF RING CURRENT
PROTONS BEYOND THE PLASMAPAUSE DURING
INJECTION EVENTS

F. V. Coroniti¹, R. W. Fredricks²,
Roscoe White¹

May, 1972

PPG-119

¹Physics Department, University of California, Los Angeles,
California 90024

²Space Sciences Department, TRW Systems Group, Redondo
Beach, California 90278

ELECTROSTATIC INSTABILITY OF RING CURRENT PROTONS BEYOND THE PLASMAPAUSE DURING INJECTION EVENTS

ABSTRACT

The stability of ring current protons with an injection spectrum modeled by an $m = 2$ mirror distribution function has been examined for typical ring current parameters. It is found that the "high frequency loss cone mode" with $\omega_{ci} < \omega < \omega_{pi}$ can be excited at wave numbers $k\lambda_{Di} \sim 0.1 - 0.5$, at frequencies $\omega \sim (0.2 - 0.6) \omega_{pi}$ and with growth rates up to $\gamma/\omega_{pi} \sim 0.03$. These waves interact with the main body of the proton distribution, and propagate nearly perpendicular [$k_{||}/k \sim (m_e/m_i)^{1/2}$] to the local magnetic field. Cold particle partial densities tend to reduce the growth rate in such a way that the waves are quenched at or near to the plasmopause boundary. Wave e-folding lengths are comparable to $0.1 R_e$, compared to the value of $\sim 4 R_e$ found for ion cyclotron waves at the same plasma conditions. The electrostatic waves investigated should lead to significant $v_{\perp}$ diffusion, and thus act to fill in the empty loss cone of the initial injection spectrum on time scales of 10^3 to $\sim 10^2$ seconds. Their detailed properties as a precipitator and diffusor mechanism require analysis beyond the scope of the present report.

ELECTROSTATIC INSTABILITY OF RING CURRENT PROTONS BEYOND THE PLASMAPAUSE DURING INJECTION EVENTS

During substorms and magnetic storms ring current protons of 5 to 50 keV energies (Frank, 1967) are injected deep into the magnetosphere, presumably by rapid internal convection from the plasma sheet. Russell and Thorne (1970) found that the innermost penetration of the dusk-midnight asymmetric ring current was about $0.5 R_e$ inside the plasmopause. Whether convection causes protons to flow to the plasmopause or the plasmopause location is determined by the ring current interaction with cold ionospheric plasma (Vasyliunas, 1972) is unclear. Cornwall, Coroniti, and Thorne (1970; referred herein as CCT) suggested that if the ring current protons penetrated the plasmopause, intense resonant electromagnetic ion cyclotron turbulence would lead to their rapid precipitation loss. CCT also argued that since in the low density ($N \sim 10$ particles/cm³), high- β ($\beta = 8\pi P/B^2$) region just outside the plasmasphere, ion cyclotron instability for ring current energies requires a large pitch angle anisotropy; the ring current in this region should be stable to electromagnetic cyclotron waves. Cornwall and Shulz (1972) calculated the ion cyclotron growth rate γ_{cyc} for a hot ring current plasma with $\beta = 1$, and anisotropy $A = \frac{T_{\perp} - T_{\parallel}}{T_{\parallel}}$ ($T_{\perp}, T_{\parallel}$ are the perpendicular and parallel temperatures for a bi-Maxwellian distribution), and no cold or plasmaspheric plasma. They found the maximum growth rate $\gamma_{cyc}/\Omega_{i,e} = 5 \times 10^{-3}$ ($\Omega_{i,e} = eB/m_{i,e}c$, the ion cyclotron-frequency). Estimating the wave group speed as the Alfvén speed, $c_A = B/\sqrt{4\pi N m_i}$, and for $L = 4$ ring current parameters, the e-folding or growth length along the field line is $\sim c_A/\gamma_{cyc} \sim 4 R_e$. Hence, the ion cyclotron wave is effectively stable just outside the plasmopause due to convective wave losses, which compensate wave growth.

An electrostatic instability of ring current protons was first suggested by Swift (1968) as a mechanism to produce VLF chorus. He investigated the case of wave growth rates less than the ion gyrofrequency, in which case harmonics of the ion gyrofrequency are excited. This is also a case in which the ion gyrofrequency (or its harmonic being excited) exceeds the lower hybrid resonance frequency. Thus, Swift's calculation is for a plasma regime quite distinct from that which we investigate in the present note.

Recently Frank (private communication) on Injun V detected the direct injection of ring current protons during the expansion phase of a magnetospheric substorm. The pitch angle distribution was strongly peaked at 90° with virtually no proton flux in the loss cone. Hence the ring current injection spectrum contains abundant free energy to drive plasma instabilities. In this note we investigate the stability of a high frequency electrostatic ion loss cone mode for ring current parameters appropriate to the high-beta region outside the plasmopause. We find that waves with frequencies around the ion plasma frequency $\omega_{pi} = \left(\frac{4\pi N e^2}{m_i} \right)^{1/2}$ can have growth rates at least two to three orders of magnitude greater than those of the ion cyclotron wave, for the same ring current distribution.

For the ring current protons we choose a loss-cone distribution (Dory et al., 1965) given by

$$f_{rc}(v_{\perp}, v_{\parallel}) = \frac{N(1-r)}{\pi^{3/2} a^3 m_i} \left(\frac{v_{\perp}}{a} \right)^{2m} \exp \left[- \frac{v_{\perp}^2 + v_{\parallel}^2}{a^2} \right]$$

where N is the total plasma density and r is the ratio of the cold proton density to the total density, N . $f_{rc}(v_{\perp}, v_{\parallel})$ has a positive slope $\partial f_{rc} / \partial v_{\perp} > 0$

between $v_{\perp} = 0$ and $v_{\perp} = \sqrt{m_i a^2}$. For this distribution the parallel temperature is $T_{\parallel} = \frac{m_i a^2}{2}$ and the perpendicular temperature is $T_{\perp} = (m+1) \frac{m_i a^2}{2}$; hence the anisotropy is $A = \frac{T_{\perp} - T_{\parallel}}{T_{\parallel}} = m$. We include an isotropic cold ion component in order to model the possible mixture of cold plasmaspheric and ring current protons which might exist just outside the storm-time plasmapause. For the electrons we take a single isotropic Maxwellian given by

$$f_e(v) = \frac{N}{\pi^{3/2} b^3} \exp[-v^2/b^2],$$

so that b is the electron thermal speed. We examine the stability of this plasma to high frequency (ω) electrostatic waves near the ion plasma frequency ω_{pi} and wave numbers k near the ion Debye length $\lambda_{Di} \equiv a/\omega_{pi}$, $k\lambda_{Di} \lesssim 1$. In computing the plasma dielectric response, we can approximate the proton trajectories as being essentially straight-line orbits provided that the wave growth rate $\gamma = \text{Im}\omega$ satisfies $\gamma > \Omega_i$ (Post and Rosenbluth, 1966). We note that even if the plasma $\beta \gtrsim 1$, for these frequencies and wave numbers the electrostatic approximation is valid since $kc/\omega_{pe} \gg 1$ ($\omega_{pe} = [4\pi Ne^2/m_e]^{1/2}$ is the electron plasma frequency).

Choosing $A = m = 2$ in the ring current distribution function, the dispersion relation can be written as

$$\begin{aligned} \epsilon(\omega, k) = & 1 + \frac{\omega_{pe}^2}{\Omega_e^2} \sin^2 \theta - \frac{\omega_{pe}^2}{k^2 b^2} Z' \left(\frac{\omega}{k_{\parallel} b} \right) - \frac{\omega_{pi}^2 r}{\omega^2} \\ & - \frac{\omega_{pi}^2 (1-r)}{k^2 a^2} \left[Z' \left(\frac{\omega}{ka} \right) + \frac{\sin^2 \theta}{2} Z'' \left(\frac{\omega}{ka} \right) + \frac{\sin^4 \theta}{32} Z^{(4)} \left(\frac{\omega}{ka} \right) \right]. \end{aligned}$$

$k_{\parallel}$ ($k_{\perp}$) is the parallel (perpendicular) component of the wave propagation vector

$\underline{k}$, $\sin \theta = k_{\perp}/k$. $Z(s)$ is the plasma dispersion function of complex argument (Fried and Conte, 1961), and prime denotes differentiation with respect to s . The dispersion relation was solved for the real (ω), and imaginary (γ) part of the complex wave frequency with the parameters $\omega_{pe}^2/\Omega_e^2 = 3$, $\cos \theta = \left(\frac{m_e}{m_i}\right)^{1/2}$, variable r , and $\frac{m_i a^2}{m_e b^2} = 25$, which might correspond to 25 keV ring current protons and 1 keV electrons. Thus the waves are propagating and have electric field polarizations almost perpendicular to the magnetic field. Figure 1 is a plot of ω/ω_{pi} vs $k\lambda_{Di}$ and γ/ω_{pi} vs ω/ω_{pi} for $r = 0, 0.5$, and 0.9 . The maximum growth rate occurs for $r = 0$ at $\omega/\omega_{pi} \sim 0.4$ and $\frac{\gamma_{max}}{\omega_{pi}} = 0.03$. The frequency range for unstable growth is $0.2 \leq \omega/\omega_{pi} \leq 0.5$. For $r = 0.05$, the maximum growth rate is reduced by about a factor of 2, and for $r = 0.9$ the waves are virtually stable. Hence the electrostatic ion loss cone waves should be predominantly unstable in the low density region outside the plasmopause and should be stable in the cold, high density plasmasphere.

For $r = 0$ and other parameters fixed, changing the propagation angle to $\cos \theta = \frac{1}{2} \left(\frac{m_e}{m_i}\right)^{1/2}$ yielded $\gamma_{max}/\omega_{pi} = 0.03$ at $\omega/\omega_{pi} = 0.3$; for $\cos \theta = \frac{1}{4} \left(\frac{m_e}{m_i}\right)^{1/2}$, $\gamma_{max}/\omega_{pi} = 0.02$ at $\omega/\omega_{pi} = 0.2$. Hence γ_{max} is relatively insensitive to propagation angle. If the ratio $m_i a^2/m_e b^2$ is lowered from 25 to 16 with other parameters fixed, $\gamma_{max}/\omega_{pi} = 0.026$ at $\omega/\omega_{pi} = 0.5$, indicating that the instability should occur for reasonable range of ring current energies. Finally reducing ω_{pe}^2/Ω_e^2 to unity, with other parameters fixed, yielded $\gamma_{max}/\omega_{pi} = 0.037$ at $\omega/\omega_{pi} = 0.53$. Hence the region of maximum wave growth might occur off the equatorial plane where lower ω_{pe}^2/Ω_e^2 would be expected if N/B is roughly constant along the field line.

Whether or not instability is achieved in a finite dimensional system depends on the path integrated growth across the system, or in the magnetosphere parallel to the lines of force. The total number of wave amplitude exponentiations can be estimated as $\ell\gamma/v_{g\parallel}$, where ℓ is the effective length along the magnetic field over which instability occurs and $v_{g\parallel}$ is the parallel wave group speed. Since $k_{\parallel}/k_{\perp} \ll 1$, we can estimate $v_{g\parallel}$ to lowest order in $k_{\parallel}/k_{\perp}$ from the dispersion relation as

$$v_{g\parallel} = - \frac{\partial \epsilon / \partial k_{\parallel}}{\partial \epsilon / \partial \omega} \approx \frac{\frac{\omega^2}{k_{\perp}^2 b^2} Z' \left(\frac{\omega}{k_{\parallel} b} \right) \frac{\omega}{k_{\perp}^2 b}}{\frac{\omega^2}{k_{\perp}^2 b^2} Z' \left(\frac{\omega}{k_{\parallel} b} \right) \frac{1}{k_{\parallel} b}} = \frac{\omega}{k_{\parallel}} = \frac{\omega}{\omega_{pi} \cdot k \lambda_{Di}} \cdot \frac{k_{\perp}}{k_{\parallel}} a.$$

For the maximally unstable mode with $\omega/\omega_{pi} = 0.4$, $k\lambda_{Di} = 0.5$, and $k/k_{\parallel} = \left(\frac{m_e}{m_i}\right)^{1/2}$, we find $v_{g\parallel} \approx 30 a$. Imposing the condition that the wave amplitude exponentiate at least 10 times to be considered unstable and taking $a \approx 2 \times 10^8$ cm/sec and $\gamma \sim (0.01-0.03) \omega_{pi}$ gives $\ell \approx 10 v_{g\parallel} / \gamma \approx 2$ to $2/3 R_e$ for the length of the growth region. Hence the electrostatic loss cone waves are unstable even though the wave energy is rapidly convected along the lines of force.

Since the unstable free energy derives from the positive slope in the ring current proton perpendicular velocity distribution, the nonlinear turbulent relaxation will be predominantly a diffusion toward lower $v_{\perp}$ for $0 \leq v_{\perp} \leq (m_i a^2)^{1/2}$, i.e., toward the atmospheric loss cone (Galeev, 1966). We have not solved the nonlinear diffusion problem to determine the turbulent electric field energy density and the time evolution of the distribution function so that only a crude estimate of the wave amplitude can be made. Two dimensional

numerical simulations of electrostatic ion waves indicate that, as a rough rule-of-thumb, wave saturation occurs when the wave-particle trapping frequency $(ekE'/m_i)^{1/2}$, based on the rms electric field E' , becomes comparable with the linear growth rate (Forslund and Shonk, 1970; Davidson et al., 1970). Using this criteria as an estimate $(ekE'/m_i)^{1/2} \approx \gamma$, for $\gamma/\omega_{pi} \approx 0.01 - 0.03$, we find $E' \sim 10 - 10^2$ mV/meter.

To estimate the diffusion time for a proton to reach the loss cone, we construct a phenomenological diffusion coefficient for perpendicular velocities $D = \frac{(\Delta v_\perp)^2}{2\Delta t}$, where $\Delta v_\perp \sim \frac{eE'}{m_i} \Delta t$ is the scattering step in $v_\perp$ and Δt is the wave-particle correlation time. Assuming $\Delta t \sim \gamma^{-1}$, we find $D \sim \frac{\omega_{pi}^2 a^2}{2k^2 \lambda_{Di}^2} (\gamma/\omega_{pi})^3$. For $\gamma/\omega_{pi} \approx 0.01 - 0.03$, the time for a proton to diffuse the characteristic velocity a is 3×10^3 to 10^2 seconds. If the 10^2 second (higher E') time scale occurs, then a rapid relaxation of the ring current anisotropy and significant proton precipitation should result. For the longer time scale (lower E'), however, the injected protons could drift out of the unstable region (e.g., their drift orbits might cross the plasmopause) before significant perpendicular diffusion occurred. In this case, the electrostatic turbulence would only partially reduce the anisotropy, and little precipitation would result. Thus whether or not the electrostatic ion loss cone turbulence outside the plasmopause substantially reduces the ring current proton fluxes via precipitation is very much an open question.

In conclusion, the injection of highly anisotropic ring current protons will destabilize the electrostatic ion loss cone mode in the zero or small cold ion plasma density regions outside the plasmopause. Since the electromagnetic ion cyclotron wave is virtually stable in this region, electrostatic waves

will dominate any turbulent relaxation of the proton anisotropy and any proton precipitation. Several important questions, however, remain unanswered. Further instability investigations are needed to determine the threshold ring current density, energy, and anisotropy for instability. This information is necessary to determine at what L-value the injected protons first become unstable. The nonlinear turbulence theory must be investigated to establish the wave saturation level, the time scale for diffusive relaxation of the anisotropy, and the extent to which electrostatic waves contribute to precipitation losses.

ACKNOWLEDGMENTS

It is a pleasure to acknowledge several beneficial discussions with Professors L. A. Frank, C. F. Kennel, R. M. Thorne, J. M. Cornwall, and Drs. D. J. Williams and F. L. Scarf.

This work was partially supported at UCLA by the Office of Naval Research, Grant ^AN00014-69-0200-4023; the National Science Foundation Grant GP-22817; the Atomic Energy Commission, Contract AT(04-3)-34, Project No. 157; and the National Aeronautics and Space Administration, Contracts NGR-05-007-190 and NGR-05-007-116. Support at TRW was under Contract No. NASW-2357.

REFERENCES

- Cornwall, J. M., F. V. Coroniti, and R. M. Thorne, Turbulent loss of ring current protons, J. Geophys. Res., 75, 4699 (1970).
- Cornwall, J. M., and Michael Shulz, Electromagnetic ion-cyclotron instabilities in a multicomponent plasma, J. Geophys. Res., 76, 7791 (1971).
- Davidson, R. C., N. A. Krall, K. Papadopoulos, R. Shanny, Electron heating by electron-ion beam instabilities, Phys. Rev. Letters, 24, 579 (1970).
- Dory, R. A., G. E. Guest, and E. G. Harris, Unstable electrostatic plasma waves propagating perpendicular to a magnetic field, Phys. Rev. Letters, 14, 131 (1965).
- Forslund, D. W., and C. R. Shonk, Two-dimensional two-species numerical simulation of electrostatic counter-streaming ion-beam instabilities, Phys. Rev. Letters, 25, 1699 (1970).
- Frank, L. A., On the extraterrestrial ring current during geomagnetic storms, J. Geophys. Res., 72, 3753 (1967).
- Fried, B. D., and S. D. Conte, The Plasma Dispersion Function, Academic Press, New York (1961).
- Galeev, A. A., Quasi-linear theory of the loss-cone mode, J. Plasma Phys., 1, 105 (1967).
- Post, R. F., and M. N. Rosenbluth, Electrostatic instabilities in mirror-confined plasmas, Phys. Fluids, 9, 730 (1966).



Russell, C. T., and R. M. Thorne, On the structure of the inner magnetosphere, Cosmic Electrodyn., 1, 67 (1970).

Swift, Daniel W., A new interpretation of VLF chorus, J. Geophys. Res., 73, 7447 (1968).

Vasyliunas, V. M., The inter-relationship of magnetospheric processes, in Earth's Particles and Fields, D. Reidel Publishing Co., Dordrecht-Holland, B. M. McCormac, editor (in press), 1972.

FIGURE CAPTION

Figure 1. Growth rate versus frequency and frequency versus wave number for the high frequency electrostatic mode driven by an $m = 2$ proton distribution in the case $T_i/T_e = 25$, $\omega_{pe}^2/\Omega_e^2 = 3$, $k_{\parallel}/k = (m_e/m_i)^{1/2}$ for cold-to-total number density ratios $r = 0, 0.5$, and 0.9 . Details in text.



UCLA PLASMA PHYSICS GROUP REPORTS

* Published by Experimental Group

† Published by Theoretical Group

- R-1 "Propagation of Ion Acoustic Waves Along Cylindrical Plasma Columns", A.Y. Wong (July 1965)*
- R-2 "Stability Limits for Longitudinal Waves in Ion Beam-Plasma Interaction", B.D. Fried and A.Y. Wong (August 1965)*
- R-3 "The Kinetic Equation for an Unstable Plasma in Parallel Electric and Magnetic Fields", B.D. Fried and S.L. Ossakow (November 1965)†
- R-4 "Low-Frequency Spatial Response of a Collisional Electron Plasma", B.D. Fried, A.N. Kaufman and D.L. Sachs (August 1965)†
- R-5 "Effects of Collisions on Electrostatic Ion Cyclotron Waves", A.Y. Wong, D. Judd and F. Hai (December 1965)*
- R-6 "Interaction Between Ion Beams and Plasmas", R. Rowberg, A.Y. Wong and J.M. Sellen (April 1966)*
- R-7 "Observation of Cyclotron Echoes from a Highly Ionized Plasma", D.E. Kaplan and R.M. Hill (May 1966)*
- R-8 "Excitation and Damping of Drift Waves", A.Y. Wong and R. Rowberg (July 1966)*
- R-9 "The Guiding Center Approximation in Lowest Order", Alfredo Baños, Jr. (September 1966)†
- R-10 "Plasma Streaming into a Magnetic Field", S.L. Ossakow (November 1966)†
- R-11 "Cooperative Effects in Plasma Echo Phenomena", A.Y. Wong (March 1967)*
- R-12 "A Quantum Mechanical Study of the Electron Gas Via the Test Particle Method", M.E. Rensink (March 1967)
- R-13 "Linear and Nonlinear Theory of Grid Excitation of Low Frequency Waves in a Plasma", G.L. Johnston (April 1967)
- R-14 "The Expansion and Diffusion of an Isolated Plasma Column", J. Hyman (May 1967)
- R-15 "Two-pole Approximation for the Plasma Dispersion Function", B.D. Fried, C.L. Hedrick and J. McCune (August 1967)†
- R-16 "Experimental Investigation of Electron Runaway Phenomena", J.S. DeGroot (August 1967)
- R-17 "Parametric Coupling Between Drift Waves", F. Hai, R. Rowberg and A.Y. Wong (October 1967)*
- R-18 "Cyclotron Echoes from Doppler Effects", A.Y. Wong (March 1968)
- R-19 "Ion Wave Echoes", D.R. Baker, N.R. Ahern and A.Y. Wong (November 1967)*
- R-20 "Cyclotron Echoes in Plasmas", D. Judd, Thesis (March 1968)
- R-21 "Test Particle Theory for Quantum Plasmas", M.E. Rensink (October 1967)†
- R-22 "Artificial Van Allen Belt", Charles F. Kennel (November 1967)
- R-23 "Landau Damping of Ion Acoustic Waves in a Cesium Plasma with Variable Electron-Ion Temperature Ratio", K.B. Rajangam (October 1967)
- R-24 "The Inhomogeneous Two-Stream Instability", G. Knorr (September 1967)
- R-25 "Magnetic Turbulence in Shocks", C.F. Kennel and H.E. Petschek (December 1967)†
- R-26 "Small Amplitude Waves in High Beta Plasmas", V. Formisano and C. Kennel (February 1968)†

- R-27 "Low Beta Plasma Penetration Across a Magnetic Field", B.D. Fried and S. Ossakow (March 1968)†
- R-28 "Annual Status Report", February 1, 1967-January 31, 1968, Principal Investigators A. Baños, Jr., B.D. Fried, C.F. Kennel
- R-29 "The Theorist's Magnetosphere", C. Kennel (April 1968)
- R-30 "Electromagnetic Pitch Angle Instabilities in Space", C.F. Kennel and F.L. Scarf (April 1968)†
- R-31 "Electromagnetic Echoes in Collisionless Plasmas", A.Y. Wong (April 1968)*
- R-32 "Parametric Excitation of Drift Waves in a Resistive Plasma", G. Weyl and M. Goldman (June 1968)†
- R-33 "Parametric Excitation from Thermal Fluctuations at Plasma Drift Wave Frequencies", A.Y. Wong, M.V. Goldman, F. Hai, R. Rowberg (May 1968)*
- R-34 "Current Decay in a Streaming Plasma Due to Weak Turbulence", S.L. Ossakow and B.D. Fried (June 1968)†
- R-35 "Temperature Gradient Instabilities in Axisymmetric Systems", C.S. Liu (August 1968)†
- R-36 "Electron Cyclotron Echo Phenomena in a Hot Collisionless Plasma", D. Judd (August 1968)
- R-37 "Transverse Plasma Wave Echoes", B.D. Fried and Craig Olson (October 1968)†
- R-38 "Low Frequency Interchange Instabilities of the Ring Current Belt", C.S. Liu (January 1969)†
- R-39 "Drift Waves in the Linear Regime", R.E. Rowberg and A.Y. Wong (February 1969)*
- R-40 "Parametric Mode-Mode Coupling Between Drift Waves in Plasmas", F. Hai and A.Y. Wong (January 1969)*
- R-41 "Nonlinear Oscillatory Phenomena with Drift Waves in Plasmas", F. Hai and A.Y. Wong (September 1970)
- R-42 "Ion-Burst Excited by a Grid in a Plasma", H. Ikezi and R.J. Taylor (February 1969)
- R-43 "Measurements of Diffusion in Velocity Space from Ion-Ion Collisions", A. Wong and D. Baker (March 1969)*
- R-44 "Nonlinear Excitation in the Ionosphere", A.Y. Wong (March 1969)
- R-45 "Observation of First-Order Ion Energy Distribution in Ion Acoustic Waves", H. Ikezi and R. Taylor (March 1969)*
- R-46 "A New Representative for the Conductivity Tensor of a Collisionless Plasma in a Magnetic Field", B.D. Fried and C. Hedrick (March 1969)†
- R-47 "Direct Measurements of Linear Growth Rates and Nonlinear Saturation Coefficients", A.Y. Wong and F. Hai (April 1969)*
- R-48 "Electron Precipitation Pulsations", F. Coroniti and C.F. Kennel (April 1969)†
- R-49 "Auroral Micropulsation Instability", F. Coroniti and C.F. Kennel (May 1969)†
- R-50 "Effect of Fokker-Planck Collisions on Plasma Wave Echoes", G. Johnston (June 1969)†
- R-51 "Linear and Nonlinear Theory of Grid Excitation of Low Frequency Waves in a Plasma", G. Johnston (July 1969)
- R-52 "Theory of Stability of Large Amplitude Periodic (BGK) Waves in Collisionless Plasmas", M.V. Goldman (June 1969)†
- R-53 "Observation of Strong Ion Wave-Wave Interaction", R. Taylor and H. Ikezi (August 1969)
- R-55 "Optical Mixing in a Magnetoactive Plasma", G. Weyl (August 1969)†
- R-56 "Trapped Particles and Echoes", A.Y. Wong and R. Taylor (October 1969)*

- R-57 "Formation and Interaction of Ion-Acoustic Solitons", H. Ikezi, R.J. Taylor and D.R. Baker (July 1970)*
- R-58 "Observation of Collisionless Electrostatic Shocks", R. Taylor, D. Baker and H. Ikezi (December 1969)*
- R-59 "Turbulent Loss of Ring Current Protons", J.M. Cornwall, F.V. Coroniti and R.M. Thorne (January 1970)†
- R-60 "Efficient Modulation Coupling Between Electron and Ion Resonances in Magnetoactive Plasmas", A. Wong, D.R. Baker, N. Booth (December 1969)*
- R-61 "Interaction of Quasi-Transverse and Quasi-Longitudinal Waves in an Inhomogeneous Vlasov Plasma", C.L. Hedrick (January 1970)
- R-62 "Observation of Strong Ion-Acoustic Wave-Wave Interaction", R.J. Taylor and H. Ikezi (January 1970)
- R-63 "Perturbed Ion Distributions in Ion Waves and Echoes", H. Ikezi and R. Taylor (January 1970)*
- R-64 "Propagation of Ion Cyclotron Harmonic Wave", E.R. Ault and H. Ikezi (November 1970)
- R-65 "The Analytic and Asymptotic Properties of the Plasma Dispersion Function", A. Baños, Jr. and G. Johnston (February 1970)
- R-66 "Effect of Ion-Ion Collision and Ion Wave Turbulence on the Ion Wave Echo", Dan Baker (June 1970)
- R-67 "Dispersion Discontinuities of Strong Collisionless Shocks", F.V. Coroniti (March 1970)†
- R-68 "An Ion Cyclotron Instability", E.S. Weibel (April 1970)†
- R-69 "Turbulence Structure of Finite-Beta Perpendicular Fast Shocks", F.V. Coroniti (April 1970)†
- R-70 "Steepening of Ion Acoustic Waves and Formation of Collisionless Electrostatic Shocks", R. Taylor (April 1970)
- R-71 "A Method of Studying Trapped Particles Behavior in Magnetic Geometries", C.S. Liu and A.Y. Wong (April 1970)*
- R-72 "A Note on the Differential Equation $g'' + x^2 g = 0$ ", E.S. Weibel (April 1970)
- R-73 "Plasma Response to a Step Electric Field Greater than the Critical Runaway Field, With and Without an Externally Applied Magnetic Field", J.E. Robin (June 1970)
- R-74 "The UC Mathematical On-Line Systems as a Tool for Teaching Physics", B.D. Fried and R. White (August 1970)†
- R-75 "High Frequency Hall Current Instability", K. Lee, C.F. Kennel, J.M. Kindel (August 1970)†
- R-76 "Laminar Wave Train Structure of Collisionless Magnetic Slow Shocks", F.V. Coroniti (September 1970)†
- R-77 "Field Aligned Current Instabilities in the Topside Ionosphere", J.M. Kindel and C.F. Kennel (August 1970)†
- R-78 "Spatial Cyclotron Damping", Craig Olson (September 1970)
- R-79 "Electromagnetic Plasma Wave Propagation Along a Magnetic Field", C.L. Olson (September 1970)†
- R-80 "Electron Plasma Waves and Free-Streaming Electron Bursts", H. Ikezi, P.J. Barrett, R.B. White and A.Y. Wong (November 1970)*
- R-81 "Relativistic Electron Precipitation During Magnetic Storm Main Phase", R.M. Thorne and C.F. Kennel (November 1970)†
- R-82 "A Unified Theory of SAR Arc Formation at the Plasmopause", J.M. Cornwall, F.V. Coroniti and R.M. Thorne (November 1970)†
- R-83 "Nonlinear Collisionless Interaction between Electron and Ion Modes in Inhomogeneous Magnetoactive Plasmas", N. Booth (December 1970)*

- R-84 "Observations of Parametrically Excited Ion Acoustic Waves", R. Stenzel (March 1971)
- R-85 "Remote Double Resonance Coupling of Radar Energy to Ionospheric Irregularities", C.F. Kennel (January 1971)+
- R-86 "Ion Acoustic Waves in a Multi-Ion Plasma", B.D. Fried, R. White, T. Samec (January 1971)+
- R-87 "Current-Driven Electrostatic and Electromagnetic Ion Cyclotron Instabilities", D.W. Forslund, C.F. Kennel, J. Kindel (February 1971)
- R-88 "Locating the Magnetospheric Ring Current", C.F. Kennel and Richard Thorne (March 1971)
- R-89 "Ion Acoustic Instabilities Due to Ions Streaming Across Magnetic Field", P.J. Barrett, R.J. Taylor (March 1971)
- R-90 "Evolution of Turbulent Electronic Shocks", A.Y. Wong and R. Means (July 1971)*
- R-91 "Density Step Production of Large Amplitude Collisionless Electrostatic Shocks and Solitons", David B. Cohen (June 1971)
- R-92 "Turbulent Resistivity, Diffusion and Heating", B.D. Fried, C.F. Kennel, K. MacKenzie, F.V. Coroniti, J.M. Kindel, R. Stenzel, R.J. Taylor, R.B. White, A.Y. Wong, W. Bernstein, J.M. Sellen, Jr., D. Forslund and R.Z. Sagdeev (June 1971)
- PPG-93 "Nonlinear Evolution and Saturation of an Unstable Electrostatic Wave", B.D. Fried, C.S. Liu, R.W. Means and R.Z. Sagdeev (August 1971)
- PPG-94 "Cross-Field Current-Driven Ion Acoustic Instability", P.J. Barrett, B.D. Fried, C.F. Kennel, J.M. Sellen and R.J. Taylor (December 1971)
- R-95 "3-D Velocity Space Diffusion in Beam-Plasma Interaction without Magnetic Field", P.J. Barrett, D. Gresillon and A.Y. Wong (September 1971)
- PPG-96 "Daytime Auroral Oval Plasma Density and Conductivity Enhancements due to Magnetosheath Electron Precipitation", C.F. Kennel and M.H. Rees (September 1971)
- PPG-97 "Collisionless Wave-Particle Interactions Perpendicular to the Magnetic Field", A.Y. Wong, D.L. Jassby (September 1971)
- PPG-98 "Magnetospheric Substorms", F.V. Coroniti and C.F. Kennel (September 1971)
- PPG-99 "Magnetopause Motions, DP-2, and the Growth Phase of Magnetospheric Substorms", F.V. Coroniti and C.F. Kennel (September 1971)
- PPG-100 "Structure of Ion Acoustic Solitons and Shock Waves in a Two-Component Plasma", R.B. White, B.D. Fried and F.V. Coroniti (September 1971)
- PPG-101 "Solar Wind Interaction with Lunar Magnetic Field", G. Siscoe (Meteorology Dept.) and Bruce Goldstein (JPL) (November 1971)
- PPG-102 "Changes in Magnetospheric Configuration During Substorm Growth Phase", F.V. Coroniti and C.F. Kennel (November 1971)
- PPG-103 "Trip Report - 1971 Kiev Conference on Plasma Theory and Visits to Lebedev and Kurchatov Institutes", B.D. Fried (October 1971)
- PPG-104 "Pitch Angle Diffusion of Radiation Belt Electrons within the Plasmasphere", Lawrence R. Lyons, Richard M. Thorne, Charles F. Kennel (January 1972)
- PPG-105 "Remote Feedback Stabilization of a High-Beta Plasma", Francis F. Chen, Daniel Jassby and M. Marhic (December 1971)