



THE PENNSYLVANIA  
STATE UNIVERSITY

# IONOSPHERIC RESEARCH

Scientific Report 397

## THE REACTION OF $O(^1D)$ WITH $H_2O$ AND THE REACTION OF $OH$ WITH $C_3H_6$

by


R. Simonaitis and Julian Heicklen

September 5, 1972

*The Research in this document was supported by the  
National Science Foundation under Grant GA-12385 and  
in part by the National Aeronautics and Space Admin-  
istration under Grant NGL 39-009-003.*

IONOSPHERE RESEARCH LABORATORY

(NASA-CR-130976) THE REACTION OF  $O(^1D)$  WITH  $H_2O$  AND THE REACTION OF  $OH$  WITH  $C_3H_6$  (Pennsylvania State Univ.) 26 p HC  
\$3.50 CSCI 04A N73-18398  
G3/13 64236  
Unclas



University Park, Pennsylvania

PSU - IRL - SCI 397

Scientific Report 397

"The Reaction of  $O(^1D)$  with  $H_2O$  and The Reaction of  $OH$  with  $C_3H_6$ "

by

R. Simonaitis and Julian Heicklen

September 5, 1972

"The Research in this document was supported by the National Science Foundation under Grant GA-12385 and in part by the National Aeronautics and Space Administration under Grant NGL 39-009-003"

Submitted by: Julian Heicklen  
Julian Heicklen, Professor of Chemistry  
Project Supervisor

Approved by: John S. Nisbet  
John S. Nisbet, Director  
Ionosphere Research Laboratory

Ionosphere Research Laboratory

The Pennsylvania State University

University Park, Pennsylvania 16802

## ACKNOWLEDGEMENTS

We thank Dr. Marcel Nicolet for many useful discussions. This work was supported by the Atmospheric Sciences Section of the National Science Foundation through Grant No. GA 12385 and the National Aeronautics and Space Administration under Grant No. NGL-39-009-003 for which we are grateful.

# TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT . . . . .	i
LIST OF TABLES . . . . .	iii
LIST OF FIGURES . . . . .	iv
ABSTRACT . . . . .	v
INTRODUCTION . . . . .	1
EXPERIMENTAL . . . . .	4
RESULTS . . . . .	6
DISCUSSION . . . . .	12
Determination of $k_1/k_4$ . . . . .	13
Determination of $k_2/k_3$ . . . . .	15
REFERENCES . . . . .	18

# LIST OF TABLES


Table	Page
I. Competitive Reaction of $O(^1D)$ with $N_2O$ and $H_2O$ at $25^\circ C$ .	7
II. $N_2O$ Photolysis at 2139A in the Presence of $H_2O$ -CO- $C_3H_6$ Mixtures . . . . .	8
III. Photolysis of $N_2O$ at 2139A in the Presence of $H_2O$ -TMP Mixtures at $100^\circ C$ . . . . .	11

# LIST OF FIGURES

Figure		Page
1	Plot of $R\{N_2\}/R\{CO_2\}(1 + \alpha)$ vs. $[N_2O]/[H_2O]$ in the photolysis of $N_2O$ in the presence of $H_2O$ and $CO$ . . . . .	14
2	Plot of $\Phi' \{CO_2\}^{-1}$ vs. $[C_3H_6]/[CO]$ in the photolysis of $N_2O$ in the presence of $H_2O$ , $CO$ , and $C_3H_6$ . . . . .	17

# Abstract

N<sub>2</sub>O was photolyzed at 2139Å to produce O(<sup>1</sup>D) atoms in the presence of H<sub>2</sub>O and CO. The O(<sup>1</sup>D) atoms react with H<sub>2</sub>O to produce HO radicals, as measured by CO<sub>2</sub> production from the reaction of OH with CO. The relative importance of the various possible O(<sup>1</sup>D)-H<sub>2</sub>O reactions is



The relative rate constant for O(<sup>1</sup>D) removal by H<sub>2</sub>O compared to that by N<sub>2</sub>O is 2.1 in good agreement with that found earlier in our laboratory.

In the presence of C<sub>3</sub>H<sub>6</sub>, the OH can be removed by reaction with either CO or C<sub>3</sub>H<sub>6</sub>.



From the CO<sub>2</sub> yield,  $k_3/k_2 = 75.0$  at 100°C and 55.0 at 200°C. When these values are combined with the value of  $k_2 = 7.0 \times 10^{-13} \exp(-1100/RT)$  cm<sup>3</sup>/sec,  $k_3 = 1.36 \times 10^{-11} \exp(-100/RT)$  cm<sup>3</sup>/sec. At 25°C,  $k_3$  extrapolates to  $1.1 \times 10^{-11}$  cm<sup>3</sup>/sec.



## INTRODUCTION

The reaction between  $O(^1D)$  and  $H_2O$  is the principle source of OH radicals in the stratosphere and it is one of the key reactions in assessing the effects on the  $O_3$  layer of introducing  $H_2O$  into the stratosphere by supersonic aircraft. From laboratory measurements it is known that the reaction of  $O(^1D)$  with  $H_2O$  gives OH radicals.

McGrath and Norrish<sup>1,2</sup> demonstrated the presence of OH radicals in the flash photolysis of  $O_3$ - $H_2O$  mixtures and proposed the reaction



In further experiments, Basco and Norrish<sup>3</sup> demonstrated that up to 2 quanta of vibrational energy could be present in the OH radical. More recently Biedenkapp, Hartzhorn, and Bair<sup>4</sup> in a study of the same system also observed vibrationally excited OH. However, the efficiency of reaction (1a) is not known, although it is generally assumed to be high. Other possible reactions are



A recent report<sup>5</sup> has indicated that reaction 1b is unimportant.

The absolute total rate constant for the reaction of  $O(^1D)$  with  $H_2O$  has been measured by Bair and coworkers<sup>4</sup> by flash photolysis of ozone- $H_2O$  mixtures to be  $3.1 \times 10^{-11}$  cm<sup>3</sup>/sec. However, this number differs by an order of magnitude, when compared with the rate constant for  $O(^1D) + N_2O$ , with the relative value of 1.5 obtained by Scott and Cvetanović<sup>6</sup> and 2.2 obtained by Lissi and Heicklen.<sup>7</sup>

In order to ascertain the relative importance of reactions 1a-1d, and to resolve the discrepancy in the rate measurements, we have carried




out a detailed study of this reaction.

The  $O(^1D)$  atoms were generated by the photolysis of  $N_2O$  at 2139Å radiation in the presence of  $H_2O$ ,  $CO$ , and in some cases  $C_3H_6$ . The  $O(^1D)$  atoms react with either  $N_2O$  or with  $H_2O$  and the resulting  $OH$  radicals from the latter reaction are scavenged by  $CO$  in accordance with reaction 2



The resulting  $CO_2$  production is therefore a measure of  $OH$  production. The presence of small amounts of propylene scavenges any  $O(^3P)$  atoms that may be formed in reaction 1b, thus preventing  $CO_2$  formation via



Actually, the rate constant for reaction 3



is much faster than for reaction 2. Consequently it is difficult to adjust the  $[CO]/[C_3H_6]$  ratio such that  $CO$  scavenges only  $OH$  and propylene scavenges only  $O(^3P)$ . The procedure adopted therefore was to vary the  $[C_3H_6]/[CO]$  ratio and extrapolate to zero. This procedure turns out to be useful in more ways than one, since the competition between reactions 2 and 3 may be studied. In order to obtain an accurate measure of reaction 1b, scavenging experiments with 2-trifluoromethylpropylene (TMP) were carried out. The reaction between TMP and  $O(^3P)$  atoms is known to give only two addition products: 2-trifluoromethylpropionaldehyde (A), and 2-trifluoromethylpropylene oxide (E) with quantum yields of 0.40 and 0.60, respectively, independent of conditions.<sup>8,9</sup> Consequently measurement of these products should provide an accurate measure of reaction 1b providing that the reaction of  $OH$  with TMP does not give the same products. The importance of reaction 1d was determined by measuring the hydrogen

yield. A direct measure of reaction 1c by the present experiments was not possible.

The reaction between OH and olefin is of interest in photochemical smog. This reaction has been proposed as an important chain propagating step in the photochemical oxidation of NO to NO<sub>2</sub>.<sup>10</sup> The reaction between OH and olefins was studied by Wilson and Westenberg<sup>11</sup> in a flow-discharge system using ESR detection, by Greiner<sup>12</sup> using flash photolysis and ultraviolet absorption spectroscopy for OH detection, and by Morris, et al.<sup>13</sup> in a discharge flow system using mass spectrometric detection. Avramenko and Kolesnikova<sup>14</sup> have also studied OH reactions with olefins, but their water discharge method has been shown to be unsuitable for kinetic studies.<sup>15</sup> In the above methods fast secondary reactions complicate the studies. These complications can be easily avoided in steady-state systems such as that employed in the present study. Unfortunately only relative rate constants may be obtained by this procedure. However, the rate constant for reaction 2 is well known,<sup>16</sup> thus absolute rate constants may be computed.

## EXPERIMENTAL

A conventional high-vacuum line utilizing Teflon stopcocks with Viton "O" rings was used. The reaction vessel was a cylindrical quartz cell 10 cm long and 5 cm in diameter. The cell was enclosed in a wire-wound aluminum block furnace, the temperature of which was controlled to  $\pm 1^\circ\text{C}$  by a Cole-Parmer Proportio Null Regulator Series 1300.

The  $\text{N}_2\text{O}$  and CO used were Matheson C. P. grade. The  $\text{N}_2\text{O}$  was purified by passage over ascarite and degassed at  $-196^\circ\text{C}$ . Gas chromatographic analysis indicated no detectable impurities. In particular  $\text{CO}_2$  and  $\text{N}_2$  were absent. The CO was purified by passage over glass beads and several traps at  $-196^\circ\text{C}$ , degassing at  $-196^\circ\text{C}$  and distillation from liquid Argon. The CO thus purified was free of  $\text{CO}_2$  but contained 540 ppm of  $\text{N}_2$ . The  $\text{N}_2$  yield in any photolysis experiment was appropriately corrected for this background  $\text{N}_2$ . The propylene (Matheson) was purified by degassing at  $-196^\circ\text{C}$ . Small amounts of tap water were degassed at  $-96^\circ\text{C}$  prior to use in order to remove dissolved air and  $\text{CO}_2$ . The 2-trifluoromethylpropene (TMP) was obtained from Peninsular ChemResearch Inc., and it was purified by distillation from a trap maintained at  $-130^\circ\text{C}$  to a trap maintained at  $-160^\circ\text{C}$ .

All gas pressures including  $\text{H}_2\text{O}$  below its room temperature vapor pressure were measured with either a McCloud gauge or a Hg manometer used in conjunction with a cathetometer. The high  $\text{H}_2\text{O}$  pressures were not measured directly, but calculated assuming the perfect gas law. Our procedure was to introduce  $\text{H}_2\text{O}$  below its room temperature vapor pressure into the line, measure its pressure on the Hg manometer, and condense it into the reaction vessel. The stopcock to the insulated reaction vessel was closed and the temperature raised to  $100^\circ\text{C}$  or  $200^\circ\text{C}$ . From

the known volume ratios the  $H_2O$  pressure was calculated.

Irradiation was from a Phillips Zn resonance lamp Typ 93106E. The effective radiation was at 2139Å. After irradiation the noncondensables were collected with a Toepler pump and analyzed for  $N_2$ , and in a few cases for  $H_2$ , by gas chromatography using a 10-ft. long by 1/4 in. diameter column packed with 5A molecular sieves. The carrier gas for  $N_2$  analysis was He, whereas for  $H_2$  the carrier gas was Argon. A second fraction volatile at  $-96^\circ C$  was removed and analyzed for  $CO_2$  using a 24-ft. long by 1/4-in. column packed with Porapak Q operating at  $25^\circ C$ . In the runs with TMP this fraction was analyzed for 2-trifluoromethylpropionaldehyde and 2-trifluoromethylpropylene oxide on a 1/4-in. diameter by 10-ft. long column at  $30^\circ C$  packed with 20% Kel-F oil No. 3 on chromosorb P.

At  $100^\circ C$  a very small amount of  $CO_2$  ( $\sim 0.03 \mu/min.$ ) was produced in the dark. At  $200^\circ C$  the dark reaction was somewhat larger ( $\sim 0.15 \mu/min.$ ). The correction for the dark reaction was small, usually less than a few percent.

In most runs, particularly at the lower values of  $[C_3H_6]$ , appreciable amounts of  $C_3H_6$  were consumed. The amount remaining was determined at the end of the run by gas chromatographic analysis and the average value of  $[C_3H_6]$  was used in the computations. The amount of propylene consumed was never more than 30% of the initial amount and was usually less than 10%, so that the use of an average value introduces no appreciable error.

## RESULTS

The results for the set of experiments carried out to determine the competition between  $\text{N}_2\text{O}$  and  $\text{H}_2\text{O}$  for the  $\text{O}(^1\text{D})$  atom are presented in Table I.  $\text{N}_2\text{O}$  was irradiated in the presence of  $\text{H}_2\text{O}$  and  $\text{CO}$ . The products observed were  $\text{N}_2$ ,  $\text{CO}_2$ ,  $\text{O}_2$ , and  $\text{H}_2$ , but only the  $\text{N}_2$  and  $\text{CO}_2$  were measured. The ratio  $R\{\text{CO}_2\}/R\{\text{N}_2\}$  increases as the  $[\text{N}_2\text{O}]/[\text{H}_2\text{O}]$  ratio decreases, reaching a value of 1.71 at our lowest  $[\text{N}_2\text{O}]/[\text{H}_2\text{O}]$  ratio of 0.152. The  $R\{\text{CO}_2\}/R\{\text{N}_2\}$  ratio is otherwise independent of  $\text{N}_2\text{O}$  or  $\text{H}_2\text{O}$  pressure, temperature (one run at  $100^\circ\text{C}$ ), absorbed light intensity (changed by a factor of  $\sim 7$ ), and the time of irradiation.

The experiments in the presence of  $\text{C}_3\text{H}_6$  were carried out in excess  $\text{H}_2\text{O}$  in order to minimize the deactivation of  $\text{O}(^1\text{D})$  by  $\text{N}_2\text{O}$  and  $\text{CO}$ . Also, in order to achieve high pressures of  $\text{H}_2\text{O}$ , the minimum temperature employed was  $100^\circ\text{C}$ . Some experiments were also performed at  $200^\circ\text{C}$ . The only products measured were  $\text{CO}_2$  and  $\text{N}_2$ . The  $R\{\text{CO}_2\}/R\{\text{N}_2\}$  ratio is independent of the time of irradiation, the absorbed light intensity (changed by a factor of  $\sim 8$ ), the  $\text{N}_2\text{O}$  pressure or the  $[\text{N}_2\text{O}]/[\text{H}_2\text{O}]$  ratio.  $R\{\text{CO}_2\}/R\{\text{N}_2\}$  depends only on the  $[\text{C}_3\text{H}_6]/[\text{CO}]$  ratio and the temperature. It increases with a decreasing  $[\text{C}_3\text{H}_6]/[\text{CO}]$  ratio and reaches a value of 1.24 at our lowest  $[\text{C}_3\text{H}_6]/[\text{CO}]$  ratio of 0.0046.  $R\{\text{CO}_2\}/R\{\text{N}_2\}$  increases slightly as the temperature is raised from  $100^\circ\text{C}$  to  $200^\circ\text{C}$ . It should be evident that at large  $[\text{H}_2\text{O}]/[\text{N}_2\text{O}]$  ratios ( $> 10$ )  $R\{\text{CO}_2\}/R\{\text{N}_2\}$  is the quantum yield of  $\text{CO}_2$ ,  $\Phi\{\text{CO}_2\}$ , since in the primary photolytic act  $\Phi\{\text{N}_2\} = 1.00^{17-21}$  and all the  $\text{O}(^1\text{D})$  atoms ( $> 95\%$ ) react with  $\text{H}_2\text{O}$ .

Some  $\text{CO}_2$  was also produced from the reaction of  $\text{OH}$  with  $\text{C}_3\text{H}_6$ . At  $100^\circ\text{C}$  the quantum yield of  $\text{CO}_2$ ,  $\Phi\{\text{CO}_2\}$ , due to this reaction is 0.025 and at  $200^\circ\text{C}$  it is 0.048.

TABLE I: COMPETITIVE REACTION OF  $O(^1D)$  WITH  $N_2O$  AND  $H_2O$  AT  $25^\circ C$

$\frac{[N_2O]}{[H_2O]}$	$[N_2O],$ Torr	$[H_2O],$ Torr	$[CO],$ Torr	Irradiation Time, min.	$10^9 R$ $\{CO_2\},$ M/Sec	$10^9 R$ $\{N_2\}$ M/Sec	$\frac{R\{CO_2\}}{R\{N_2\}}$
17.6	53.0	3.00	8.50	32.0	1.85	9.6	0.192
16.6	49.5	3.00	3.20	12.0	4.75	25.8	0.185
14.9	44.7	3.00	2.30	10.0	5.70	27.8	0.204
14.7	57.5	3.90	2.60	25.0	1.77	10.3	0.173
11.5	57.4	5.00	2.60	12.0	7.62	30.2	0.253
9.30	56.0	6.00	4.00	90.0	1.25	3.52	0.353
7.30	55.5	7.60	2.40	23.0	2.64	9.4	0.281
6.45	51.5	8.00	7.5	90.0	2.59	6.33	0.410
5.22	45.5	8.50	4.00	95.0	1.45	3.18	0.460
3.72 <sup>a</sup>	59.5	16.0	10.5	51.0	2.11	4.11	0.510
3.33	55.0	16.5	8.5	10.0	4.93	8.25	0.600
3.30	53.0	16.0	10.0	30.0	4.61	9.5	0.488
0.193	3.25	16.8	10.5	180.0	0.98	0.604	1.64
0.152	2.85	18.7	10.5	172.0	1.03	0.606	1.71

a) This run done at  $100^\circ C$ .

TABLE II: N<sub>2</sub>O PHOTOLYSIS AT 2139A IN THE PRESENCE OF H<sub>2</sub>O-CO-C<sub>3</sub>H<sub>6</sub> MIXTURES

$\frac{[C_3H_6]^a}{[CO]}$	$\frac{[N_2O],}{Torr}$	$\frac{[H_2O],}{Torr}$	$\frac{[CO],}{Torr}$	$\frac{[C_3H_6]^a,}{Torr}$	$\frac{Irradiation}{Time, min.}$	$\frac{10^9 R\{CO_2\},}{M/sec}$	$\frac{10^9 R\{N_2\},}{M/sec}$	$\frac{R\{CO_2\}}{R\{N_2\}}$
Temp = 100°C								
0.157	30.0	370	14.0	2.20	60.0	0.57	3.30	0.172
0.110	15.5	490	21.0	2.30	80.0	1.02	3.30	0.310
0.102	3.40	350	24.0	2.45	180.0	0.110	0.505	0.217
0.061	39.0	400	34.5	2.10	77.0	1.44	3.98	0.362
0.0605	10.8	490	43.0	2.60	90.0	0.88	2.39	0.368
0.0365	21.5	400	66.0	2.40	90.0	0.115	0.261	0.442
0.0520	5.10	357	53.5	2.80	210.0	0.93	1.87	0.500
0.025	9.90	490	100	2.50	90.0	1.45	2.20	0.661
0.0161	26.7	400	46.5	0.75	63.0	3.86	5.02	0.770
0.129	28.2	350	58.0	0.75	65.0	2.23	2.14	1.04
0.0122	27.2	350	59.0	0.72	60.0	2.67	2.62	1.02
0.0133	8.80	560	51.0	0.68	90.0	2.26	2.47	0.92
0.0108	16.2	725	60.0	0.65	89.0	1.32	1.50	0.88
0.0094	12.9	340	53.0	0.50	85.0	0.84	0.92	0.92

TABLE II: N<sub>2</sub>O PHOTOLYSIS AT 2139A IN THE PRESENCE OF H<sub>2</sub>O-CO-C<sub>3</sub>H<sub>6</sub> MIXTURES (concluded)

$\frac{[C_3H_6]^a}{[CO]}$	$\frac{[N_2O],}{Torr}$	$\frac{[H_2O],}{Torr}$	$\frac{[CO],}{Torr}$	$\frac{[C_3H_6],}{Torr}$	$\frac{Irradiation}{Time, min.}$	$\frac{10^9 R\{CO_2\},}{M/sec}$	$\frac{10^9 R\{N_2\},}{M/sec}$	$\frac{R\{CO_2\}}{R\{N_2\}}$
Temp = 100°C								
0.0062	30.0	400	106	0.65	45.0	4.92	5.62	0.88
0.0054	27.3	350	108	0.58	60.0	2.26	2.49	0.91
0.0046	8.90	520	117	0.54	90.0	2.68	2.16	1.24
Temp = 200°C								
0.151	32.1	440	18.5	2.80	80.0	0.53	1.85	0.288
0.058	28.2	486	46.5	2.70	91.0	1.14	2.08	0.555
0.0348	30.5	462	69	2.40	87.0	1.17	1.76	0.665
0.0212	33.5	440	108	2.30	90.0	1.51	1.84	0.820

a) Average pressure of C<sub>3</sub>H<sub>6</sub> during run.



The results for the direct determination of the efficiency of reaction 1b are presented in Table III. The products measured in the presence of TMP are 2-trifluoromethylpropionaldehyde (A) and 2-trifluoromethylpropylene oxide (E). These same products are produced when  $O(^3P)$  atoms add to TMP; however the ratio  $\phi\{E\}/\phi\{A\} = 1.50^{8,9}$  in that system is very different from that observed in the present system.

Irradiation at 2139Å of 75 torr of  $N_2O$  in the presence of 660 torr of  $H_2O$  at  $100^\circ C$  for 90 minutes gave traces of  $H_2$ . However the yield was so small that the quantum yield of 0.004 must be taken as an upper limit for reaction 1d, since it is difficult to exclude other minor sources of  $H_2$ .

TABLE III: PHOTOLYSIS OF N<sub>2</sub>O AT 2139 Å IN THE PRESENCE OF H<sub>2</sub>O-TMP MIXTURES


AT 100°C

<u>[N<sub>2</sub>O],</u> <u>Torr</u>	<u>[H<sub>2</sub>O],</u> <u>Torr</u>	<u>[TMP],</u> <u>Torr</u>	<u>Irradiation</u> <u>Time, min.</u>	$10^9$ <u><math>\dot{R}\{N_2\}</math></u> <u>M/sec</u>	<u><math>\Phi\{E\}^a</math></u>	<u><math>\Phi\{A\}^a</math></u>	<u><math>\frac{\Phi\{E\}}{\Phi\{A\}}</math></u>
38.0	370	2.00	50.0	2.98	0.020	0.120	0.17
38.0	340	1.25	90.0	3.57	0.025	0.138	0.18

a)  $\Phi\{X\} = \dot{R}\{X\}/\dot{R}\{N_2\}$

# DISCUSSION

Photolysis of  $\text{N}_2\text{O}$  at 2139A in the presence of  $\text{H}_2\text{O}$ ,  $\text{CO}$  and  $\text{C}_3\text{H}_6$  can be analyzed by the following scheme:



The formation of  $\text{O}({}^1\text{D})$  with unit efficiency in the photolysis of  $\text{N}_2\text{O}$  at 2139A is well known.<sup>17-21</sup> The reaction of  $\text{O}({}^1\text{D})$  with  $\text{N}_2\text{O}$  has been extensively studied.<sup>22,23</sup> The exclusive reactions are 4a and 4b with  $k_{4a}/k_{4b} = 0.67^{23}$  when the  $\text{O}({}^1\text{D})$  atoms are produced from  $\text{N}_2\text{O}$  photolysis. Reaction 5 has been studied in this laboratory.<sup>24</sup> The ratio of rate coefficients  $k_1/k_5 = 2.6 \exp(-1200/RT)$  where  $k_1 \equiv k_{1a} + k_{1b} + k_{1c} + k_{1d}$ , and  $R$  is in calories/mole-°K. The reactions of  $\text{OH}$  with  $\text{C}_3\text{H}_6$  and  $\text{CO}$  have already been mentioned in the Introduction. The reaction of  $\text{O}({}^3\text{P})$  with  $\text{C}_3\text{H}_6$  can produce  $\text{CO}_2$  a small fraction of the time.<sup>25</sup> However since  $\text{O}({}^3\text{P})$  production turns out to be negligible, the amount of  $\text{CO}_2$ , if any, from Reaction 6 can be ignored.

# Determination of $k_1/k_4$ :

$k_1/k_4$  may be determined from the variation of  $R\{CO_2\}/R\{N_2\}$  with  $[N_2O]/[H_2O]$  from experiments in the absence of  $C_3H_6$ . The relevant reactions are reactions 1 and 4 followed by reaction 2. The following expression is obtained from this scheme:

$$\left( \frac{1}{1 + \alpha} \right) \frac{R\{N_2\}}{R\{CO_2\}} = \frac{1}{2} \frac{k_1}{k_{1a}} \left( 1 + \frac{k_4 [N_2O]}{k_1 [H_2O]} \right) \quad I$$

$$\text{where } \alpha \equiv \frac{k_{4a} [N_2O]}{k_4 [N_2O] + k_1 [H_2O]}$$

Actually in some runs reaction 5 may quench as much as 10% of the  $O(^1D)$ , thus reducing the yield of  $CO_2$ . However, the reduction may be more than compensated by reformation of  $CO_2$  via the  $O(^3P) + CO + M$  reaction. An accurate correction for this effect could be made except for the fact that at low CO pressures some of the  $O(^3P)$  atoms will be irreversibly lost at the walls.<sup>26</sup> These effects probably nearly compensate so that the error introduced by neglecting the correction for  $O(^1D)$  deactivation by CO cannot be more than 5%.

$\alpha$  may be computed by an iterative procedure from the known value  $k_{4a}/k_{4b}$  and from a rough initial value for  $k_1/k_4$  obtained at low  $[N_2O]/[H_2O]$ . A plot of  $[1/(1 + \alpha)] R\{N_2\}/R\{CO_2\}$  vs.  $[N_2O]/[H_2O]$  should be linear with an intercept of  $k_1/2k_{1a}$  and a ratio of slope to intercept of  $k_4/k_1$ . The plot is presented in Figure 1. It is reasonably linear with an intercept of 0.50 and a slope of 0.236 independent of temperature in the range of 25°C - 100°C.

The value for  $k_1/k_4$  found from the ratio of intercept to slope is 2.13. This value is in reasonable agreement with the value of 1.50 obtained by Scott and Cvetanović,<sup>6</sup> and in excellent agreement with the results of 2.2 found by Lissi and Heicklen.<sup>7</sup>




Figure 1

The intercept gives  $k_{1a}/k_1 = 1.0$ , so that reactions 1b, 1c, and 1d must be unimportant (<10% total). The quantum yields for A and E presented in Table III give an upper limit for reaction 1b. From the ratio of  $\Phi\{E\}/\{A\} = 0.18$  it is clear that most of the aldehyde does not arise from  $O(^3P)$  atom addition, but from the reaction with OH radicals, since this ratio is 1.50 for  $O(^3P)$  atom addition.<sup>8,9</sup> Aldehydes are major products in the reaction of OH radicals with  $C_2H_4$ ,  $C_3H_6$  and presumably with other olefins.<sup>13</sup> The small yield of the epoxide however could come from  $O(^3P)$  or from OH addition. From the epoxide quantum yield an upper limit for  $\Phi\{O(^3P)\}$  is 0.040, in agreement with the findings of Paraskevopoulos and Cvetanović.<sup>5</sup>

The quantum yield of  $H_2$  gives an upper limit of 0.004 for reaction 1d. A direct determination of  $H_2O_2$  yield was not possible, but the fact that the intercept in Figure 1 is 1/2 and the fact that this is also the case in the presence of  $C_3H_6$  (see below) indicate that the quantum yield for  $H_2O_2$  production cannot be much greater than 0.05 and probably is zero.

#### Determination of $k_2/k_3$ :

Based on the reaction scheme of reactions 1-7, the following expression can be obtained

$$\Phi' \{CO_2\}^{-1} = \frac{k_1}{2k_{1a}} \left( 1 + \frac{k_3 [C_3H_6]}{k_2 [CO]} \right) \quad II$$

where  $\Phi' \{CO_2\} \equiv \beta\Phi\{CO_2\} - \gamma$ .

$\Phi' \{CO_2\}$  is essentially the quantum yield of  $CO_2$ ,  $\Phi\{CO_2\}$ , corrected for some quenching of  $O(^1D)$  by  $N_2O$  and CO ( $\beta$ ) and for the production of some  $CO_2$  by reaction 3a ( $\gamma$ ). Both corrections are small. The correction term  $\beta$  is given by:

$$\beta \equiv 1 + \frac{k_4}{k_1} \frac{[N_2O]}{[H_2O]} + \frac{k_5}{k_1} \frac{[CO]}{[H_2O]}$$

and  $\gamma$  is given by:

$$\gamma = \frac{2k_{1a}}{k_1} \frac{k_{3a}/k_3}{1 + k_2[CO]/k_3[C_3H_6]}$$

$\beta$  may be readily evaluated using  $k_4/k_1 = 0.425$  (see above) and  $k_5/k_1 = 0.52$  at  $100^\circ\text{C}$  and  $0.72$  at  $200^\circ\text{C}$ .<sup>24</sup>  $\gamma$  may be computed by an iterative procedure using a rough initial value of  $k_2/k_3$  and  $k_{3a}/k_3 = 0.025$  at  $100^\circ\text{C}$  and  $0.048$  at  $200^\circ\text{C}$  obtained in experiments without added CO (see Results). The correction due to  $\beta$  was never  $> 22\%$ , and usually  $< 10\%$ . The correction due to  $\gamma$  was less than a few percent in most runs. However in a few runs, where  $[C_3H_6]/[CO]$  was large, the correction amounted to about 20%.

A plot of  $\Phi^i \{CO_2\}^{-1}$  vs.  $[C_3H_6]/[CO]$  should be linear with an intercept of  $k_1/2k_{1a}$  and a slope of  $\frac{k_1}{2k_{1a}k_2} \frac{k_3}{k_2}$ . This plot is shown in Figure 2 for  $100^\circ\text{C}$  and  $200^\circ\text{C}$ . The plots are linear with an intercept of  $1/2$  and a slope of  $37.5$  at  $100^\circ\text{C}$  and an intercept of  $1/2$  and a slope of  $27.4$  at  $200^\circ\text{C}$ . Consequently  $k_{1a}/k_1 = 1.0$  and  $k_3/k_2 = 75.0$  at  $100^\circ\text{C}$ . At  $200^\circ\text{C}$   $k_{1a}/k_1 = 1.0$  and  $k_3/k_2 = 55.0$ . The Arrhenius expression from the data at the two temperatures may be obtained to be  $k_3/k_2 = 293 \exp(+1000/RT)$ . An absolute value for  $k_3$  may be calculated from the known value of  $k_2$ , which has been the subject of many investigations. Baulch et al.<sup>16</sup> give  $k_2 = 7.0 \times 10^{-13} \exp(-1100/RT) \text{ cm}^3/\text{sec}$  based on a critical evaluation of all the determinations. The result is  $k_3 = 1.36 \times 10^{-11} \exp(-100/RT) \text{ cm}^3/\text{sec}$ . Thus at room temperature  $k_3 = 1.1 \times 10^{-11} \text{ cm}^3/\text{sec}$ . This value compares well the only other available value of  $1.7 \times 10^{-11} \text{ cm}^3/\text{sec}$  obtained by Morris, et al.<sup>13</sup> using a discharge-flow method with mass spectrometric detection.




Figure 2



# REFERENCES

1. W. D. McGrath and R. G. W. Norrish, Nature, 182, 235 (1958).
2. W. D. McGrath and R. G. W. Norrish, Proc. Roy. Soc., A254, 317 (1960).
3. N. Basco and R. G. W. Norrish, Proc. Roy. Soc., A260, 293 (1961).
4. D. Biedenkapp, L. G. Hartzhorn and E. J. Bair, Chem. Phys. Lettr., 5, 379 (1970).
5. G. Paraskevopoulos and R. J. Cvetanović, Chem. Phys. Lettr., 9, 603 (1971).
6. P. M. Scott and R. J. Cvetanović, J. Chem. Phys., 54, 1440 (1971).
7. E. Lissi and J. Heicklen, J. Photochem., 1, xxxx (1972).
8. S. J. Moss and K. R. Jennings, Trans. Faraday Soc., 64, 686 (1968).
9. R. Simonaitis and J. Heicklen, J. Chem. Phys., in press.
10. J. Heicklen, K. Westberg, and N. Cohen, "Chemical Reactions in Urban Atmospheres," Ed. C. S. Tuesday, Elsevier (1971) p 55.
11. W. E. Wilson and A. A. Westenberg, 11th Symposium (International) on Combustion (1967) p 1143.
12. N. R. Greiner, J. Chem. Phys., 53, 1284 (1970).
13. E. D. Morris, Jr., D. H. Stedman, and H. Niki, J. Am. Chem. Soc., 93, 3570 (1971).
14. L. J. Avramenko and R. V. Kolesnikova, Adv. Photochem., 2, 25 (1964).
15. F. P. DelGreco and F. Kaufman, Disc. Faraday Soc., 33, 128 (1962).
16. For a critical review see D. L. Baulch, D. D. Drysdale and A. C. Lloyd, "High Temperature Reaction Rate Data," Dept. of Phys. Chem., Leeds University, No. 1 (1968).
17. H. Yamazaki and R. J. Cvetanović, J. Chem. Phys., 39, 1902 (1963).
18. H. Yamazaki and R. J. Cvetanović, J. Chem. Phys., 40, 582 (1964).
19. H. Yamazaki and R. J. Cvetanović, J. Chem. Phys., 41, 3703 (1964).

20. R. J. Cvetanović, J. Chem. Phys., 43, 1850 (1965).
21. K. F. Preston and R. J. Cvetanović, J. Chem. Phys., 45, 2888 (1966).
22. P. M. Scott, K. F. Preston, R. J. Andersen and T. M. Quick, Canad. J. Chem., 49, 1808 (1971).
23. R. Simonaitis, R. I. Greenberg, and J. Heicklen, Intern. J. Chem. Kinetics, in press.
24. R. Simonaitis and J. Heicklen, Int. J. Chem. Kinetics, in press.
25. R. I. Greenberg, unpublished results at Penn State University (1970).
26. R. Simonaitis, J. Heicklen, M. M. McGuire and R. A. Bernheim, J. Phys. Chem., 75, 3205 (1971).