

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

X-691-73-329

PREPRINT

NASA TM X- 70511

PHOTOCHEMISTRY OF PLANETARY ATMOSPHERES

(NASA-TM-X-70511) PHOTOCHEMISTRY OF
PLANETARY ATMOSPHERES (NASA) 11 p HC
\$3.00

CSCI 03B

N74-10782

Unclas

G3/30 21593

L. J. STIEF

OCTOBER 1972



GSFC

GODDARD SPACE FLIGHT CENTER

GREENBELT, MARYLAND

X-691-73-329

PHOTOCHEMISTRY OF PLANETARY ATMOSPHERES*

L. J. Stief

Astrochemistry Branch (691)
Laboratory for Extraterrestrial Physics
NASA/Goddard Space Flight Center
Greenbelt, Maryland 20771

October 1973

*This is part of a paper presented at the NATO Advanced Study Institute on "Spectroscopy and Photochemistry in the Vacuum-ultraviolet," Valmorin, Quebec, Canada, August 1973.

GODDARD SPACE FLIGHT CENTER
Greenbelt, Maryland

PHOTOCHEMISTRY OF PLANETARY ATMOSPHERES

Photodissociation is the primary loss process for the major components of the atmospheres of planets. A fundamental problem is the explanation for the apparent stability of, for example, the CO_2 atmosphere on Mars. It now appears that the ultimate solution to this problem is more likely to depend on new knowledge concerning the dynamics of atomic and free radical reactions occurring subsequent to photodissociation rather than on additional details concerning the primary photochemical process. In the following sections, the atmospheric composition of Mars is presented and the applicability of laboratory data on CO_2 absorption cross sections and quantum yields of dissociation is discussed. This is followed by a summary and critical evaluation of the various mechanisms proposed for converting the photodissociation products CO and O_2 back to CO_2 .

INTRODUCTION

The scope of the subject the photochemistry of planetary atmospheres is obviously too broad to be adequately described in a limited presentation. I had initially intended to discuss two examples of photochemical problems relating to the atmospheres of planets, viz. the stability of the CO_2 atmosphere on Mars and the stability of the NH_3 and CH_4 in the atmosphere of Jupiter. However, the latter problem will not be discussed because the astronomical "facts" are not yet sufficiently firm and a choice between the two theories relating to this problem depends on these "facts" as well as laboratory data not presently available. The interested reader is referred to papers by McNesby¹ and Strobel² for details regarding the Jupiter problem. The present discussion will therefore be limited to the question of the stability of the Martian CO_2 atmosphere.

The composition of the lower atmosphere of Mars is shown in Table 1. The densities given are those at or near the surface where the temperature is about 150°K . There are some time variations observed as well as differences between the results of different observers, but the results for CO and O_2 are certainly good to within a factor of 2 and the CO_2 density is now known to an accuracy of $\pm 15\%$. The O_3 observations were made with the ultraviolet spectrometer aboard Mariner 7 and 9. No ozone is present in the equatorial region, but in the polar regions ozone appears in the fall, reaches a maximum in winter and decreases in the spring³. The inverse correlation of ozone with water vapor suggests that the presence of water suppresses the ozone formation.

The contents of Table 1 should be very disturbing to any laboratory photochemist who has used CO_2 as a chemical actinometer. It is well known that CO_2 is

Table 1
Mars: Atmospheric Composition



Molecule	Concentration (cm ⁻³)
CO ₂	1.5×10^{17}
O ₂	2.0×10^{14}
CO	1.2×10^{14}
H ₂ O	$\sim 10^{14}$
O ₃	$< 10^{10} \text{ a}$

- a. Estimated limit for planet outside the polar regions; ozone only observed over polar regions and reaches maximum in winter of $\sim 10^{11} \text{ cm}^{-3}$.



photodissociated in the laboratory and that CO is formed with a quantum yield ϕ equal to or close to unity. It may be readily calculated from the known flux of solar radiation, the absorption cross section for CO₂ as a function of wavelength and assuming $\phi = 1$ that the amount of CO now present in the atmosphere of Mars would be produced in only two years. Even if $\phi = 0.5$, the essential problem remains since the observed CO would then be produced in four years. One is thus forced to conclude that either the absorption cross sections or quantum yields of dissociation are many orders of magnitude smaller than inferred from laboratory data or that there is an efficient process for converting CO and O₂ back to CO₂. Julianne, Neuman and Krauss⁴ have drawn attention to the possible errors involved in using room temperature laboratory cross sections in modeling studies for CO₂ kinetics. Their theoretical calculations suggest that halving the temperature from 300° to 150°K produces only about a 10% decrease in CO₂ absorption for the region 1200-1750 Å. In addition, DeMore and Patapoff⁵ have shown that cross sections for the region 1700-2000 Å measured at 200°K are about one-half the room temperature values. Thus, while this effect should be considered for accurate modeling, it appears that there are no order of magnitude effects on absorption cross sections. The situation is similar for the primary quantum yield. While there may be evidence for ϕ as low as 0.5 (see for example the work of Inn⁶), order of magnitude effects appear to be definitely excluded. The task thus reduces to identifying the mechanism or mechanisms by which the photodissociation products CO and O₂ are converted back to CO₂.

The processes occurring in the upper and lower atmosphere leading to the formation of O₂ and CO may be summarized as follows:

Upper Atmosphere



Lower Atmosphere




Three body recombination of O and CO may be neglected based on the results of Slanger, Wood and Black⁷.

THE HO₂ MECHANISMS

There are a number of mechanisms which may be considered jointly since they all involve reaction and formation of the HO₂ free radical. The source of H is considered to be the photolysis of H₂O, known to be present in the atmosphere (~0.1% of CO₂). It is well to recall here the CO₂ photochemical equilibrium studies of Harteck et al.⁸ using the 1633 Å bromine line. It was found that with as little as 0.1% water vapor present (the same level as for Mars), complete recombination to CO₂ occurred (O₂/CO₂ < 0.005). While these experiments do not point to a specific mechanism, they clearly show that the presence of water is important and that under conditions comparable to that of the Martian atmosphere, there is little net CO₂ dissociation.

The HO₂ mechanism may be summarized by the following cycle of reactions:




The processes occurring in the upper and lower atmosphere leading to the formation of O_2 and CO may be summarized as follows:

Upper Atmosphere



Lower Atmosphere



Three body recombination of O and CO may be neglected based on the results of Slanger, Wood and Black⁷.

THE HO_2 MECHANISMS

There are a number of mechanisms which may be considered jointly since they all involve reaction and formation of the HO_2 free radical. The source of H is considered to be the photolysis of H_2O , known to be present in the atmosphere ($\sim 0.1\%$ of CO_2). It is well to recall here the CO_2 photochemical equilibrium studies of Harteck et al.⁸ using the 1633 Å bromine line. It was found that with as little as 0.1% water vapor present (the same level as for Mars), complete recombination to CO_2 occurred ($O_2/CO_2 < 0.005$). While these experiments do not point to a specific mechanism, they clearly show that the presence of water is important and that under conditions comparable to that of the Martian atmosphere, there is little net CO_2 dissociation.

The HO_2 mechanism may be summarized by the following cycle of reactions:





It is clear that the net effect of X is to convert less reactive HO_2 to OH which leads to recombination of CO and O_2 to CO_2 and regeneration of the chain carriers H and OH. We will briefly consider here mechanism for $\text{X} = \text{CO}, \text{NO}, \text{O}_3, \text{H}, \text{O}$ and HO_2 . Table 2 lists the rate constant for each $\text{HO}_2 + \text{X}$ reaction, the approximate steady state concentration required at or near the surface to compensate for the O_2 production rate. In the full model, many factors are included and concentrations are calculated as a function of altitude.

Table 2
Steady State Concentration Required by the Various HO_2 Mechanisms

X	k ($\text{cm}^3 \text{ s}^{-1}$)	(X) _{ss} (cm^{-3})	
		Req.	Obs.
CO	$< 10^{-20}$	10^{18}	10^{14}
NO	3×10^{-13}	10^{10}	—
O_3	3×10^{-15}	10^{12}	$< 10^{10}$
H	($\sim 10^{-11}$)	10^9	10^4 ^a
O	7×10^{-11}	10^8	10^8 ^a
HO_2	3×10^{-12}	10^9	—

a. Calculated from Mariner observations at 250 km for H and 135 km for O.
See ref. 9.

Despite early suggestions that reaction (4) should be fast^{8,10} and the very indirect determination by Westenberg and de Haas¹¹ that $k_4 \approx 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$, it is now relatively certain that the reaction



is negligibly slow near 300°K. This was first suggested by the high temperature

work of Baldwin¹² and subsequently verified by attempts to measure k_4 at 300°K.¹³⁻¹⁶ The best estimate comes from the isotopic labelling-product analysis experiment¹⁴ and this result of $k_4 < 10^{-20} \text{ cm}^3 \text{ molec}^{-1} \text{ sec}^{-1}$ would require a CO density that exceeds the observed value by 10^4 . Using the same isotopic labelling technique, an estimate has been made¹⁷ of the rate constant for the reaction¹⁵



As seen from Table 2, a concentration NO equal to $10^{10} \text{ molec cm}^{-3}$ would be required. While photolysis of NO_2 would return NO to the atmosphere, it should be emphasized that such levels of NO and NO_2 should be capable of detection by ultraviolet spectroscopy.

The $\text{HO}_2 + \text{O}_3$ reaction has been included since a good limit exists for O_3 concentration and both De More¹⁹ and Simonaitis and Heicklen²⁰ have recently measured the rate constant at 300°K. The results in Table 2 show that the required O_3 concentration exceeds the observed upper limit by a factor of 10^2 . Reaction with H should also be considered since atomic hydrogen is a known constituent of the atmosphere. Although the rate constant for the reaction



has not been measured, the estimated^{11,21} value of $k_6 = 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ sec}^{-1}$ leads to concentration of H many orders of magnitude too high.

Thus we are finally left with only two possibilities. One is the reaction²²



which is part of the complete model proposed by McElroy and Donahue²¹. The required O atom concentration is at least not inconsistent with observations; the model does however require large eddy diffusion to transport atomic oxygen from the upper to the lower atmosphere. The final reaction is HO_2 disproportionation²³



which is followed by photolysis of H_2O_2



This suggestion is due to Parkinson and Hunten²² and the large eddy diffusion requirement of the McElroy-Donahue model is avoided. These are some minor problems with both models, but it is not inconceivable that both could be important depending on the relative concentration of O and HO₂ in the lower atmosphere. These could be variable with time. At the present writing, these two models seem to offer the most hope for eventually understanding the stability of CO₂ in the atmosphere of Mars.

OZONE MECHANISM

This may be treated much more briefly. The proposed^{8,18} sequence of reactions is:



Again, the net result is to convert the less reactive O and O₂ to OH to achieve conversion of CO and O₂ back to CO₂. However, since there are laboratory measurements for k_{11} ²⁵ and k_1 ²⁶ and observational data for concentration of O₃, O₂ and CO₂ (see Table 1), we can readily calculate the O₃ concentration required for H + O₃ (11) to be competitive with H + O₂ + CO₂ (1).

$$k_{11}(\text{H})(\text{O}_3) = k_1(\text{H})(\text{O}_2)(\text{CO}_2)$$

$$\text{O}_3 = \frac{5 \times 10^{-32} \times 2 \times 10^{14} \times 1.5 \times 10^{17}}{2.6 \times 10^{-11}}$$

$$\text{O}_3 = 6 \times 10^{10} \text{ molec cm}^{-3}$$

This is well above the limit of O₃ < 10¹⁰ molec cm⁻³ from the Mariner observations for the equatorial regions and is within a factor of 2 of the maximum

values observed over the polar region in winter. In addition, for the $H + O_3$ mechanism to dominate over the HO_2 mechanism, we should assume $R_{11} = 10R_1$ which leads to O_3 concentrations of 6×10^{11} molec cm^{-3} . Clearly this mechanism is not important under the known condition of the Martian atmosphere.

IONIC MECHANISM

Finally, for completeness we consider the mechanism proposed by Parks²⁷ which is based on the interesting observation that CO_2 shows apparent stability under radiolysis. Essentially the idea is that CO_2 is converted by ultraviolet radiation to CO and O_2 and that the photochemical products are converted back to CO_2 by means of ion-molecule reactions involving positive ions produced from CO_2 . The mechanism requires positive ion concentrations of the order of 10^3 cm^{-3} and rate constants all of the order of 10^{-11} cm^{-3} molec $^{-1}$ sec $^{-1}$. Without going into the details of the full mechanism, we point out two serious problems. The first is that a laboratory measurement²⁸ of one of the key reactions of the mechanism



shows that the rate constant is much too slow ($k_{12} \leq 10^{-12}$ cm^3 molec $^{-1}$ sec $^{-1}$). The second problem, which applies to any ionic mechanism proposed, is that a means must be found to transport the positive ions from the upper atmosphere where they are formed to the very low atmosphere near the surface where CO and O_2 concentrations are many orders of magnitude larger.

SUMMARY

The persistence of the CO_2 atmosphere on Mars appears to require some mechanism which converts the photolysis products CO and O_2 back to CO_2 . At the present time, the mechanism most consistent with both astronomical observations and laboratory data involves the chain sequence: $OH + CO \rightarrow CO_2 + H$, $H + O_2 \rightarrow HO_2$ and $HO_2 + CO_2 \rightarrow HO_2 + CO_2$ and $HO_2 + O \rightarrow OH + O_2$ and/or $HO_2 + HO_2 \rightarrow H_2O_2$ $h\nu \rightarrow 2OH$. The chain carries H and OH are supplied by photolysis of H_2O which is present to the extent of 0.1% of the CO_2 .

REFERENCES

1. J. R. McNesby, *J. Atmos. Sci.* 26, 594 (1969); M. Lenzi, J. R. McNesby, A. Mele and C. N. Xuan, *J. Chem. Phys.* 57, 319 (1972).
2. D. F. Strobel, *J. Atmos. Sci.* 26, 906 (1969); *ibid.* 30, 489 (1973); *ibid.* (in press).
3. C. A. Barth, C. W. Hord, A. I. Stewart, A. L. Lane, M. L. Dick and G. P. Anderson, *Science* 179, 795 (1973).
4. P. S. Julienne, D. Neumann and M. Krauss, *J. Atmos. Sci.* 28, 833 (1971).
5. W. B. De More and M. Patapoff, *J. Geophys. Res.* 77, 6291 (1972).
6. E. C. Y. Inn, *J. Geophys. Res.* 77, 1991 (1972).
7. T. G. Slanger, B. J. Wood and G. Black, *J. Chem. Phys.* 57, 233 (1972).
8. P. Harteck, R. R. Reeves, Jr., B. A. Thompson and R. W. Waldron, *Tellus XVIII*, 192 (1966); *ibid.*, *J. Phys. Chem.* 70, 1637 (1966).
9. D. E. Anderson and C. W. Hord, *J. Geophys. Res.* 76, 6666 (1971); G. E. Thomas, *J. Atmos. Sci.* 28, 859 (1971).
10. S. W. Benson, "The Foundations of Chemical Kinetics," McGraw-Hill Book Co., Inc., New York, N.Y., 1960, p. 461.
11. A. A. Westenberg and N. de Haas, *J. Phys. Chem.* 76, 1586 (1972).
12. R. R. Baldwin, R. W. Walker and S. J. Webster, *Combust. Flames* 15, 167 (1970).
13. D. H. Volman and R. A. Gorse, *J. Phys. Chem.* 76, 3301 (1972).
14. D. D. Davis, W. A. Payne and L. J. Stief, *Science* 179, 280 (1973).
15. R. Simonaitis and J. Heicklen, *J. Phys. Chem.* 77, 1096 (1973).
16. D. Wyrsh, H. R. Wendt and H. E. Hunziker, preprints XI International Symposium on Free Radicals (1973).
17. W. A. Payne, L. J. Stief and D. D. Davis, *J. Amer. Chem. Soc.* 95, 0000 (1973).

18. M. B. McElroy and J. C. McConnell, *J. Atmos. Sci.* 28, 879 (1971).
19. W. B. De More, *Science* 180, 735 (1973).
20. R. Simonaitis and J. Heicklen, *J. Phys. Chem.* 77, 1932 (1973).
21. M. B. McElroy and T. M. Donahue, *Science* 177, 986 (1972).
22. C. J. Hochanadel, J. A. Ghormley and P. J. Ogren, *J. Chem. Phys.* 56, 4426 (1972).
23. T. T. Pankert and H. S. Johnston, *J. Chem. Phys.* 56, 2824 (1972).
24. T. D. Parkinson and D. M. Hunten, *J. Atmos. Sci.* 29, 1380 (1972).
25. L. F. Phillips and H. I. Schiff, *J. Chem. Phys.* 37, 1233 (1962).
26. M. J. Kurylo, *J. Phys. Chem.* 76, 3518 (1972); D. L. Baulch, D. D. Drysdale, D. G. Horne and A. C. Lloyd, "Evaluated Kinetic Data for High Temperature Reactions," Vol. 1, Butterworth & Co., London, 1972.
27. D. A. Parkes, *Nature* 241, 110 (1973).
28. L. W. Sieck, R. Gorden, Jr. and P. Ausloos, *Planet. Space Sci.* (in press).