

POSSIBLE ACCELERATION OF CHARGED PARTICLES  
THROUGH THE RECONNECTION OF MAGNETIC FIELD  
LINES IN INTERPLANETARY SPACE

E. H. Levy

Bartol Research Foundation of The Franklin Institute  
Swarthmore, Pennsylvania 19081

F. M. Ipavich and G. Gloeckler

University of Maryland, College Park, MD 20742

Technical Report # 74-116

June 1974



(NASA-CR-139387) POSSIBLE ACCELERATION OF  
CHARGED PARTICLES THROUGH THE RECONNECTION  
OF MAGNETIC FIELD LINES IN INTERPLANETARY  
SPACE (Maryland Univ.) 18 p HC \$4.00

N74-31318

CSCL 03B G3/30

Unclas  
46205

UNIVERSITY OF MARYLAND  
DEPARTMENT OF PHYSICS AND ASTRONOMY  
COLLEGE PARK, MARYLAND

Space Physics Group

**This is a preprint of research carried out at the University of Maryland. In order to promote the active exchange of research results, individuals and groups at your institution are encouraged to send their preprints to**

**PREPRINT LIBRARY  
DEPARTMENT OF PHYSICS AND ASTRONOMY  
UNIVERSITY OF MARYLAND  
COLLEGE PARK, MARYLAND  
20742  
U.S.A.**

POSSIBLE ACCELERATION OF CHARGED PARTICLES  
THROUGH THE RECONNECTION OF MAGNETIC FIELD  
LINES IN INTERPLANETARY SPACE\*

E. H. Levy

Bartol Research Foundation of The Franklin Institute  
Swarthmore, Pennsylvania 19081

F. M. Ipavich and G. Gloeckler

University of Maryland, College Park, MD 20742

Technical Report # 74-116

June 1974

\*Submitted to Geophysical Research Letters, June 1974.

Possible Acceleration of Charged Particles  
Through the Reconnection of Magnetic Field  
Lines in Interplanetary Space

E. H. LEVY

*Bartol Research Foundation of The Franklin Institute  
Swarthmore, Pennsylvania 19081*

F. M. IPAVICH and G. GLOECKLER

*University of Maryland, College Park 20742*

ABSTRACT

Prominent intensity spikes in the flux of protons and alphas with less than 0.5 MeV per charge have been observed in the region several hours behind an interplanetary shock front. The small spatial scale of these events and the high anisotropy of the particle flux suggest local acceleration. The spectra of the particles, which are cut off at equal energy per charge, suggest acceleration through an electric field. We examine the possibility that these events have their origin in active magnetic neutral sheets in the shocked solar wind.

It has been known for some years that large increases in the low to medium energy cosmic ray flux accompany interplanetary shock waves. Several authors have suggested that these delayed cosmic ray events are the result of particle acceleration associated with the propagation of the shock wave itself (Axford and Reid, 1962, 1963; Jokipii, 1966; Rao et al., 1967; Lanzerotti and Robbins, 1969; Armstrong et al., 1970; Ogilvie and Arens, 1971; Singer and Montgomery, 1971). Parker (1965) has noted that particle acceleration seems to be a ubiquitous property of the agitation associated with nonequilibrium plasmas as they relax to a more homogeneous state. However, the actual mechanism through which the particle acceleration takes place is not known. Indeed, the panoply of activities of a nonequilibrium plasma is vast and charged particle acceleration is likely to be produced, not by a unique mechanism, but rather by at least several, and perhaps many, mechanisms.

Gloeckler et al. (1974), hereinafter called Paper I) have reported observations of intensity increases of low energy protons and alphas associated with the interplanetary shock wave during 30 October to 1 November 1972. In particular they found a series of prominent but short-lived intensity increases — the so called "post-shock" spikes — several hours behind the shock front. In this letter we present more detailed observations on these post-shock spikes which are evident up to energies of about 0.5 MeV per charge. The peculiar features of these particle spikes, which are discussed in detail in this paper, lead us to

suggest that they are evidence of local cosmic ray acceleration in the solar wind and that the particles were observed closely adjacent to the acceleration region. We also explore, in a preliminary way, the possibility that these particles are accelerated through the electric field associated with magnetic field line reconnection in the shocked solar wind.

The observations were made using the University of Maryland experiment on the IMP 7 (Explorer 47) satellite between 1900 and 2100 UT on 31 October 1972 when the spacecraft was in interplanetary space. The instrument uniquely identifies low energy ( $\geq 100$  keV/charge) protons, alphas and heavier particles through a simultaneous measurement of the kinetic energy of an incoming particle and of its deflection in a known electrostatic field (Fan et al., 1971; also see Paper I).

Figure 1 shows the time variations of protons in the energy range 125 to 160 keV and of alpha particles in the range 95 to 135 keV/nucleon. The fluxes of both species exhibit a series of spikes, coincident in time and of several minutes duration. The counting rates between the spikes represent real particles and are not background (Paper I). These spikes, to which we are confining our attention, show the following features in common:

- a) The fluxes of protons and alpha particles, with energies greater than about 0.5 MeV per charge show no increase during the spikes (Figure 2, Paper I).
- b) The  $p/\alpha$  ratio, at equal energy per charge, is constant ( $\sim 20$ ) throughout the time when the spikes

occurred, but shows large changes in the spikes when evaluated at either equal velocity or equal rigidity (see Figure 3 in Paper I).

c) The particle anisotropy approaches 95% during the spikes (Ipavich, 1974).

d) The low energy flux of heavier ions ( $z > 6$ ) also exhibits spikes coincident with those in the proton and alpha fluxes; at higher energies the flux of heavy ions shows no increase and its behavior is consistent with the cutoff at about 0.5 MeV per charge displayed by the protons and alphas (D. Hovestadt, private communication, 1974).

e) There is no indication of an increase in the electron flux during these events; from the counting rate of our lowest energy electron channel we estimate that the intensity of electrons between 130 and 180 keV is at most a few percent of the proton flux at that energy.

Magnetic field data obtained with instruments on the same satellite (R. Lepping and N. Ness, private communication, 1974) permit the following further conclusions:

a) These events occurred several hours after the passage of an interplanetary shock wave in the region of high magnetic field;  $B \approx 15\gamma$ .

b) At the time that the spikes were observed the magnetic field displayed moderate fluctuations which, very

loosely, appear to have a wave-like character.

- c) Taking the Compton-Getting effect into account, the particles are found to be streaming generally along the magnetic field from an easterly, antisolar direction.
- d) The particle spikes do not appear to be confined to magnetic field lines which intersect the earth's bow shock.

It is interesting to note that the particle fluxes are accompanied by wave-like disturbances in the magnetic field. We suggest that these disturbances may be excited by the particle streams themselves. While a detailed discussion of this question is beyond the scope of the present paper we note briefly that a flux of particles is expected to drive hydromagnetic waves in a plasma when: 1) the streaming speed of the particles exceeds the Alfvén speed; 2) the energy density of the particles is comparable to the energy density of the magnetic field. In the present case 1) the streaming speed of the highly anisotropic beam of particles is of the order of a few times  $10^8$  cm sec<sup>-1</sup> which is significantly larger than the Alfvén speed; 2) the flux of particle energy in the beam is of the order of  $10^{11}$  eV cm<sup>-2</sup> sec<sup>-1</sup> (see below) so that the energy density due to the particles is of the order of 500 eV cm<sup>-3</sup>; by comparison the energy density of the 15γ magnetic field is about 600 eV cm<sup>-3</sup>.

The high particle anisotropy, together with the very well defined, beam-like nature of the observed particle fluxes, sug-



gests that the region of acceleration was nearby to the detector and that the particles were accelerated contemporaneously with their detection. In addition, the particle spectra — protons, helium nuclei and heavy nuclei — in which each species appears to be cut off at the same maximum energy per charge, suggest local acceleration through an electric field. The most promising candidate for the particle acceleration seems to be the electric field associated with the reconnection of magnetic field lines in the presence of resistive plasma instabilities. Although the magnetic field data (R. Lepping and N. Ness, private communication, 1974) show the presence of at least one and perhaps several magnetic neutral sheets associated with the region where the particles are accelerated, we cannot, unfortunately, report having caught one in the act of instability and active particle acceleration. However, in view of the importance of cosmic ray acceleration for astrophysics and in view of the theoretical uncertainties surrounding the question of magnetic field reconnection, it seems worthwhile to pursue the question further and to examine the consequences of assuming that these beams of energetic particles were accelerated in regions of magnetic field reconnection.

Consider the idealized neutral sheet, represented pictorially in Figure 2, having thickness  $d$  and longitudinal and transverse scales  $L$ . With a density of ions of about  $10 \text{ cm}^{-3}$ , the observed magnetic field of  $1.5 \times 10^{-4}$  gauss in the shocked solar wind plasma leads to an Alfvén speed,  $V_A$ , of about  $10^7 \text{ cm sec}^{-1}$ . The rate of energy production in phenomena for which magnetic field

reconnection is thought to play a central role suggests that the merging velocity,  $V_m$  associated with active neutral sheet regions is of the order of  $0.1 V_A$  (see the discussion in Parker, 1973). Then the electric field characteristic of the merging region is

$$E = 0.1 \frac{V_A B}{c} \quad (1)$$

Using the present values for  $V_A$  and  $B$  we find that  $E = 5 \times 10^{-9}$  stat-volt  $\text{cm}^{-1} = 2 \times 10^{-6}$  volt  $\text{cm}^{-1}$ . Neutral sheets in the solar wind are thought to have a coherence scale of about  $3 \times 10^{11}$  cm (McCracken and Ness, 1966; Bartley et al., 1966; Michel, 1967; Siscoe et al., 1968; McCracken et al., 1968) and Parker (1972) has already suggested that rapid reconnection of the interplanetary field may occur across such neutral sheets. Associating this scale of  $3 \times 10^{11}$  cm with the longitudinal scale  $L$ , we find that the maximum energy (per unit of charge) attained by a particle traversing the neutral sheet region is of the order of  $5 \times 10^5$  eV. This energy is very closely the maximum energy at which the spikes are apparent in the present observations. Furthermore the close agreement between the proton to alpha ratio at equal energy per charge, in the spikes, with the proton to alpha ratio in the quiet-time solar wind is in agreement with these ideas of local acceleration through an electric field.

It is instructive to compare the rate at which magnetic field energy enters the dissipation region with the estimated energy appearing in the particle fluxes. The energy density of

the  $1.5 \times 10^{-4}$  gauss magnetic field is about  $600 \text{ eV cm}^{-3}$ . This energy flows into the reconnection region at about  $10^6 \text{ cm sec}^{-1}$  over an area  $L^2$  on either side of the neutral sheet. Thus, when  $L \approx 3 \times 10^{11} \text{ cm}$ , magnetic field energy flows into the reconnection region at a rate of about  $10^{32} \text{ eV sec}^{-1}$ . In order to estimate the energy in the observed particle beam, it is necessary to know the spectrum at low energies (well below the 125 keV lower limit of our detectors). Now, in the spikes the spectrum varies as  $E^{-\gamma}$ , where  $\gamma$  is between 2 and 3 in the energy range above 125 keV. If the spectrum were to remain steep — say  $\gamma \sim 2-3$  — down to low energies (5 or 10 keV) then we estimate the energy flux in the observed beam to be about  $10^{11} \text{ eV cm}^{-2} \text{ sec}^{-1}$ . From the time structure of the particle spikes in Figure 1, together with the velocity of the shocked solar wind ( $\sim 7 \times 10^7 \text{ cm sec}^{-1}$ , H. Rosenbauer and H. Grünwald, 1974, private communication) we estimate that the profile of the particle beams has a small scale (thickness) of about  $10^{10} \text{ cm}$  and a large scale (width) about equal to  $L$  —  $3 \times 10^{11} \text{ cm}$ . The area of an observed beam is then about  $3 \times 10^{21} \text{ cm}^2$  and the particle beam requires about  $3 \times 10^{32} \text{ eV sec}^{-1}$  for its support. This power is of the order of, but somewhat larger than, the rate at which magnetic field energy is delivered to the reconnection region. Should the spectrum become substantially flatter below the range of our detectors then the estimated power required to sustain the beam of particles falls to between  $10^{31}$  and  $10^{32} \text{ eV sec}^{-1}$ . The energy balance seems to require that spectrum flattens at low energies. On the other hand, it should be remembered that the

simple model we have used here is presented only as a basis for discussion. It is not intended to be a quantitative representation of the acceleration region. The primary conclusion to be drawn from this discussion of the energy balance is: Should our suggestion that these particles are accelerated in interplanetary neutral sheets survive further scrutiny, then particle acceleration through reconnecting magnetic fields in a tenuous medium like the solar wind is an efficient process.

In the absence of collective effects, the electrical conductivity of a plasma is generally too high to permit rapid reconnection of magnetic fields (Parker, 1963). Thus it is thought that reconnection occurs primarily when conditions favor plasma instabilities which interrupt the flow of electrical current and raise the resistivity of the plasma. In the neutral sheet of Figure 2, the electrical current density,  $j(c\vec{v} \times \vec{B} = 4\pi j)$ , in the neutral layer is  $cB(2\pi d)^{-1}$ . If the density of electrons in the ambient plasma is  $\sim 10 \text{ cm}^{-3}$ , then the electron-ion relative drift speed is  $v_d \sim 10^{14} \text{ d}^{-1} \text{ cm sec}^{-1}$ . In the solar wind the electron temperature is greater than the proton temperature and, in addition, plasma turbulence tends to preferentially heat the electrons. Under these conditions the plasma is unstable to the growth of ion-sound waves when the electron-ion drift velocity exceeds the ion-sound speed —  $\sim 10^6 \text{ cm sec}^{-1}$ . The neutral sheet depicted in Figure 2 is therefore subject to plasma turbulence, driven by ion-sound waves when  $d \lesssim 10^8 \text{ cm}$ . (In the initial transient stage, however, it may be necessary for the current layer to be com-

pressed to less than  $10^7$  cm to get the plasma turbulence going.)  $10^8$  cm is less than the thickness of the observed layer of energetic particles, as is necessary on the basis of the suggested model.

Altogether the properties of the particle fluxes in the post-shock spikes agree well with the properties that would be expected if these particles were accelerated locally through electric fields associated with the reconnection of magnetic lines of force. In the framework of this model, the short-lived intensity spikes result from the convection, past the spacecraft, of magnetic field lines which connect with the acceleration region. On this basis, the observations suggest that particle acceleration through active magnetic field line merging is a feature of the solar wind magnetic field, at least in the disturbed region behind an interplanetary shock front.

We acknowledge the contribution of C.Y. Fan and D. Hovestadt as co-investigators on the University of Maryland IMP-7 experiment. We are also grateful to N. Ness and R. Lepping for the use of their IMP-7 magnetic field data and to H. Rosenbauer and H. Grünwald for providing us with the solar wind velocity data. This research was supported, in part, by the National Aeronautics and Space Administration under contract NAS-5-110<sup>63</sup> and Grant NGR 21-002-224 at the University of Maryland. At the Bartol Research Foundation this research was supported in part by the National Science Foundation.

## References

- Armstrong, T.P., S.M. Krimigis, and K.W. Behannon, Proton fluxes at 300 keV associated with propagating interplanetary shock waves, *J. Geophys. Res.* 75, 5980, 1970.
- Axford, W.I., and G.C. Reid, Polar cap absorption and the magnetic storm of February 11, 1958, *J. Geophys. Res.* 67, 1692, 1962.
- Axford, W.I., and G.C. Reid, Increases in intensity of solar cosmic rays before sudden commencements of geomagnetic storms, *J. Geophys. Res.* 68, 1793, 1963.
- Bartley, W.C., R.P. Bukata, K.G. McCracken, and U.R. Rao, Anisotropic cosmic ray fluxes of solar origin, *J. Geophys. Res.* 71, 3297, 1966.
- Gloeckler, G., F.M. Ipavich, C.Y. Fan, and D. Hovestadt, Post-shock spikes: A new feature of proton and alpha enhancements associated with an interplanetary shock wave, *Geophys. Res. Lett.* 1, 65, 1974.
- Ipavich, F.M., The Compton-Getting effect for low energy particles, to be published, 1974.
- Jokipii, J.R., A model of Fermi acceleration at shock fronts with an application to the earth's bow shock, *Astrophys. J.* 143, 961, 1966.
- Lanzerotti, L.J., and M.F. Robbins, Solar flare alpha to proton ratio changes following interplanetary disturbances, *Solar Physics* 10, 212, 1969.

- Lin, R.P., C.-I. Meng, and K.A. Anderson, 30-100 keV protons upstream from the earth's bow shock, *J. Geophys. Res.* 79, 489, 1974.
- McCracken, K.G., and N.F. Ness, The collimation of cosmic rays in the interplanetary magnetic field, *J. Geophys. Res.* 71, 3315, 1966.
- McCracken, K.G., U.R. Rao, and N.F. Ness, Interrelationship of cosmic ray anisotropies and the interplanetary magnetic field, *J. Geophys. Res.* 73, 4159, 1968.
- Michel, F.C., Model of solar wind structure, *J. Geophys. Res.* 72, 1917, 1967.
- Ogilvie, K.W., and J.F. Arens, Acceleration of protons by interplanetary shocks, *J. Geophys. Res.* 76, 13, 1971.
- Parker, E.N., The solar flare phenomenon and the theory of reconnection and annihilation of magnetic fields, *Astrophys. J. Suppl.* No 77, VIII, 177, 1963.
- Parker, E.N., Interplanetary origin of energetic particles, *Phys. Rev. Letters* 14, 55, 1965.
- Parker, E.N., Topological dissipation and the small scale fields in turbulent gases, *Astrophys. J.* 174, 499, 1972.
- Parker, E.N., The reconnection rate of magnetic fields, *Astrophys. J.* 180, 247, 1973.
- Rao, U.R., K.G. McCracken, and R.P. Bukata, Cosmic ray propagation processes, 2. The energetic storm particle event, *J. Geophys. Res.* 72, 4325, 1967.

- Singer, S.F., and M.D. Montgomery, Detailed directional and Temporal properties of solar energetic particles associated with propagating interplanetary shock waves, *J. Geophys. Res.* 76, 6628, 1971.
- Siscoe, G.L., L. Davis, Jr., P.J. Coleman, Jr., E.J. Smith, and D.E. Jones, Power spectra and discontinuities of the interplanetary magnetic field; Mariner 4, *J. Geophys. Res.* 73, 61, 1968.

#### Figure Captions

- Figure 1 Counting rates, integrated over 20 seconds, measured in the 125-160 keV proton channel and 95-136 keV/nucleon alpha particle channel. Particle fluxes ( $\text{Particles cm}^{-2} \text{ sec}^{-1} \text{ Sr}^{-1} (\text{keV/nucleon})^{-1}$ ) may be obtained by multiplying the counting rates by 1.4 in the case of the protons and 0.8 in the case of the alpha particles.
- Figure 2 An idealized representation of a magnetic neutral sheet. a) Crude general aspect of the magnetic field line configuration and length scales. b) Stylized representation of the magnetic and electric fields,  $B$  and  $E$  respectively, the current  $j$  and the merging speed  $V_m$ .






Figure 1





Figure 2