


## **General Disclaimer**


### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.


NG6-05-007-190-57



PLASMA PHYSICS GROUP



UNIVERSITY OF CALIFORNIA  
LOS ANGELES



N76-23124

(NASA-CR-147104) RELATIVISTIC ELECTRONS AND  
WHISTLERS IN JUPITER'S MAGNETOSPHERE  
(California Univ.) 34 p HC \$4.00 CSCI 03B

G3/91  
Unclas  
25303

RELATIVISTIC ELECTRONS  
AND  
WHISTLERS IN JUPITER'S MAGNETOSPHERE

D. D. Barbosa<sup>1</sup> and F. V. Coroniti<sup>1,2</sup>

PPG-260

April 1976

<sup>1</sup>Department of Physics  
<sup>2</sup>Department of Geophysics and Space Physics  
University of California  
Los Angeles, California 90024

## ABSTRACT

We have computed the path-integrated gain of parallel propagating whistlers driven unstable by an anisotropic distribution of relativistic electrons in the stable trapping region of Jupiter's inner magnetosphere. The requirement that a gain of 3 e-foldings of power balance the power lost by imperfect reflection along the flux tube sets a stably-trapped flux of electrons  $J^* = 4 \times 10^{10} L^{-4} \text{ cm}^{-2} \text{ sec}^{-1}$  which is close to the non-relativistic result. Comparison with measurements shows that observed fluxes are near the stably-trapped limit, which suggests that whistler wave intensities may be high enough to cause significant diffusion of electrons accounting for the observed reduction of phase space densities. A crude estimate of the wave intensity necessary to diffuse electrons on a radial diffusion time scale yields a magnetic field fluctuation intensity of  $I_B = 1.5 \cdot 10^{-18} (\Omega_{ce}/\omega)^2 L^{\alpha-2} \text{ watts m}^{-2} \text{ Hz}^{-1}$  as a lower limit.

Decreases in observed phase space densities of energetic electrons in Jupiter's inner magnetosphere have led to the suggestion that pitch angle scattering of the particles by wave microturbulence is an active loss mechanism (Fillius et al., 1975; McIlwain and Fillius, 1975; Baker and Van Allen, 1976). A likely suspect is the electron whistler wave or R mode, the theory of which when applied to earth's environment has been successful in explaining electron precipitation losses in the stable trapping region of the inner magnetosphere (Kennel and Petschek, 1966; Lyons et al., 1972).

Radio observations at decimeter wavelengths of the synchrotron emissions from Jupiter's radiation belts indicated long before Pioneer 10 that the electron distribution was highly anisotropic and that most of the energetic electrons were confined to the magnetic equator by a pancaked pitch angle distribution (Roberts and Komesaroff, 1965; Thorne, 1965). Pioneers 10 and 11 confirmed this expectation and also verified the theory that the immediate source of the radiation belt electrons was inward radial diffusion which, conserving the first and second particle adiabatic invariants, would flatten the pitch angle distribution. Since induced emission of whistlers is a consequence of anisotropic distribution, the hypothesis that whistler turbulence is responsible for observed losses is an attractive one both because of the ample growth rates possible and the theoretical simplicity of the instability.

The first hint that non-synchrotron associated losses were present at Jupiter was given by Stansberry and White (1974) before Pioneer 10. They set up a radial diffusion model for electrons to compute the strip-scan brightness and flux density spectrum of synchrotron emissions and found that an ad hoc loss process was needed at low L values for a reasonable fit with radio observations. Coroniti (1974), in deriving a comprehensive theoretical model of radiation belt electron fluxes, employed the stably-trapped limit concept for whistlers using the best pre-Pioneer 10 values for relevant parameters. The data from Pioneers 10 and 11 have narrowed the range of parameter space (in particular, the cold plasma density and anisotropy) available for theoretical models and the observed losses invite a reexamination of the relativistic electron-whistler interaction.

The purpose of this paper is to detail some of the consequences of relativistic electrons in stably-trapped equilibrium with parallel-propagating whistler waves. Approximate scaling laws for the stably-trapped electron flux and equilibrium wave intensity are derived. For simplicity of analysis and clarity of content, the major restrictions to our model are the following: 1) we treat the waves as generated locally and travelling strictly parallel to the ambient dipole magnetic field; 2) all resonant electrons are ultrarelativistic in that the total energy is proportional to particle momentum,  $E \approx pc$ .

In Section II the equatorial growth rate for whistlers is derived for a distribution modeled as  $f(\vec{p}) \propto p^{-(N+2)} \sin^M \theta$ .

The logarithmic gain for maximally-amplified waves is computed by an approximation to the path-integrated growth rate along a flux tube and the stably-trapped limit is defined as that which will produce 3 e-foldings of power. A similar model for the distribution was used by Liemohn (1967) to compute whistler amplification by relativistic electrons at earth. Schulz and Vampola (1975) considered an artificial radiation belt produced by the beta decay of nuclear-fission debris and computed the relativistic stably-trapped limit for a distribution with an angular distribution similar to ours but with an exponential energy dependence. Our procedure and results are consistent with the above authors. The stably-trapped limit of relativistic electron fluxes is found to compare closely with non-relativistic Kennel-Petechek theory. Comparison with measurements shows that observed fluxes lie near the limit, which lends support to the idea that whistlers are active. In Section III we briefly treat the aspects of quasilinear diffusion of relativistic electrons and estimate a level of wave intensity that will support diffusion (losses) on a time scale comparable to radial diffusion.

## II RELATIVISTIC ELECTRON-WHISTLER INTERACTION

### a. Equatorial Growth Rate

If we restrict our attention to whistlers propagating parallel to the local magnetic field  $\vec{B}_0 = B_0 \hat{z}$ , the dispersion relation for R modes with  $k_\perp = 0$  is (Lerche, 1969)

$$K(k, \omega) = 0 = -\frac{k^2 c^2}{\omega^2} + 1 + \sum \frac{2\pi e^2}{m\omega} \int d\vec{p} \frac{p_\perp}{\omega\gamma - kp_z/m + \Omega} \cdot \left\{ \frac{\partial f_0}{\partial p_\perp} + \frac{k}{m\omega\gamma} \left[ p_\perp \frac{\partial f_0}{\partial p_z} - p_z \frac{\partial f_0}{\partial p_\perp} \right] \right\} \quad (1)$$

The  $\sum$  represents a sum over particle species (species labels on all quantities are implicit),  $\gamma = [1 + p^2/m^2 c^2]^{1/2}$ ,  $m$  is the particle rest mass, and  $\Omega = \frac{qB_0}{mc}$ . If the particle distribution function  $f_0(\vec{p})$  is composed of a cold plasma background and a small relativistic component  $f$ , then defining  $t = p/mc$  and  $x = \cos \theta = \frac{\vec{p} \cdot \vec{B}_0}{pB_0}$ , equation (1) can be written in spherical coordinates, assuming azimuthal symmetry in  $\vec{p}$ -space, as

$$\frac{k^2 c^2}{\omega^2} = 1 + \frac{\omega_p^2}{(\Omega_{ci} + \omega)(\Omega_{ce} - \omega)} - \sum \frac{4\pi^2 e^2}{m\omega k c} \cdot \int_0^\infty t^2 dt \int_{-1}^1 \frac{(1-x^2) dx}{x - \frac{(\omega\gamma + \Omega)}{kct}} \left\{ \frac{\partial f}{\partial t} + \frac{\partial f}{\partial x} \left( \frac{kc}{\omega\gamma} - \frac{x}{t} \right) \right\} \quad (2)$$

where

$$\omega_p^2 = \frac{4\pi n_0 e^2}{m} \quad \text{and} \quad \Omega_{ce} = \left| \frac{eB_0}{mc} \right|$$



for the electrons. If the wave number has a small imaginary part,  $k = k_r + ik_i$ , we can approximate  $k_i = \frac{-K_i}{\partial K_r / \partial k}$  and integrating around the indented pole of the integrand for the electron contribution yields

$$k_i(\omega) = \frac{2\pi^3 e^2 m}{mk_c^3} \int_{t_0}^{\infty} dt \, t^2 (1-x^2) \left[ \frac{\partial f}{\partial t} + \frac{\partial f}{\partial x} \left( \frac{n}{\gamma} - \frac{x}{t} \right) \right] \Bigg|_{x = \frac{\omega \gamma - \Omega_{ce}}{kct}} \quad (3)$$

The integration is taken along the resonance contour  $x = \frac{\omega \gamma - \Omega_{ce}}{kct}$  and  $t_0$  represents the minimum momentum (magnitude) satisfying  $-1 = \frac{\omega \gamma - \Omega_{ce}}{kct_0}$ . The subscript on the real part of the wavenumber has been dropped [ $k_r \rightarrow k$ ] and

$$\frac{k_c^2}{\omega^2} = n^2 = 1 + \frac{\omega_p^2}{(\Omega_{ci} + \omega)(\Omega_{ce} - \omega)} \quad (4)$$

The resonance curve in momentum space is a hyperbola, plotted in Fig. 1, satisfying the relation

$$\left[ t_z + \frac{n\bar{\Omega}}{n^2 - 1} \right]^2 - \frac{t_1^2}{n^2 - 1} = \frac{\bar{\Omega}^2}{(n^2 - 1)^2} + \frac{1}{n^2 - 1} \quad (5)$$

with  $\bar{\Omega} = \frac{1}{\omega} = \frac{\Omega_{ce}}{\omega}$ . The minimum momentum is given by

$$t_0 = \frac{n\bar{\Omega} - \sqrt{\bar{\Omega}^2 + n^2 - 1}}{n^2 - 1} \quad (6)$$

The resonance curve crosses the  $t_1$  axis at the point  $t_1 = \sqrt{\bar{\Omega}^2 - 1}$  and asymptotes to the angle  $\phi = \sin^{-1} \frac{1}{n}$ . The passage to the non-relativistic regime requires  $n \gg 1$ ,  $t \ll 1$  and the resonance curve flattens to approximately a straight line. Equation (6)

then gives the familiar result  $-t_0 = t_R = \frac{V_R}{c}$  where  $\frac{V_R}{c} = \frac{1 - \bar{\Omega}}{n} = \frac{-\bar{\Omega}}{n}$  for  $\bar{\omega} \ll 1$  and  $E_R = \frac{1}{2} m V_R^2$  (Kennel and Petchek, 1966). Relativistic particles can interact with whistlers either by Doppler-shifting the wave frequency up to the local electron gyrofrequency or by lowering their gyrofrequency sufficiently by a mass increase, thus the resonance curve departs from a straight line. Induced emission or absorption of the whistler depends on the local derivatives of the distribution function along the resonance curve. The growth or damping of a mode depends on the net energy contribution to a wave from the particles. For a distribution such as  $f \propto \sin^M \theta$ , a given frequency may be unstable by equation (3) but for those particles with  $p_z = 0$ ,  $\left. \frac{\partial f}{\partial \theta} \right|_{\theta=\pi/2} = 0$  and consequently those particles are energized by the growing mode.

For application to energetic electrons in Jupiter's magnetosphere it is convenient to use the ultrarelativistic approximation where  $t \gg 1$ . In that case the resonance curve can be approximated by  $x \approx \frac{1}{n} [1 - \bar{\Omega}/t]$  valid so long as  $t_0 \approx \frac{\bar{\Omega}}{1+n} \gg 1$ . If the relativistic electron distribution obeys a power law in momenta (energy) we can model the distribution at the magnetic equator as  $f(t) = \frac{B \sin^M \theta}{t^{N+2}}$  where  $M$  and  $N$  are the pitch angle and spectral indices of the distribution and  $B$  is a normalization. A convenient normalization is in terms of the omnidirectional integral flux. For relativistic particles where  $\gamma = \sqrt{1+t^2} \approx t$  we have

$$f(t) = \frac{(N-1) \gamma^{N-1} J(\gamma v)}{4\pi c} \frac{2}{\sqrt{\pi}} \frac{\Gamma(\frac{M+3}{2})}{\Gamma(\frac{M+2}{2})} \frac{\sin^M \theta}{t^{N+2}} \quad (7)$$

such that  $J(>\gamma) = \int_{\gamma}^{\infty} d\gamma \int_{4\pi} d\Omega t^2 c f(\vec{t})$ . Since the growth rate depends on the distribution above  $t_0$  by (3), if we consider frequencies such that  $t_0$  lies in the domain where (7) is valid, we do not have to specify the behavior of the distribution at low energies. When (7) is inserted into the growth rate and the change of variable  $x = \frac{1}{n}[1 - \bar{n}/t]$  is made we have the following expression for the logarithmic gain scaled to a Jovian radius at the magnetic equator:

$$k_i R_J = \frac{\pi^2 e R_J}{B_0(L)} \frac{J(>\gamma)}{c} \left[ \frac{\gamma}{\bar{n}/n} \right]^{N-1} F(N, M, u) \quad (8)$$

where

$$F(N, M, u) = (N-1) \frac{1}{\sqrt{\pi}} \frac{\Gamma(\frac{M+3}{2})}{\Gamma(\frac{M+2}{2})} \int_{-1}^u dx (u-x)^{N-1} \cdot (1-x^2)^{\frac{M}{2}} [u(N+2)(1-x^2) + Mx(1-ux)] \quad (9)$$

REPRODUCIBILITY OF THE

and  $u = \frac{1}{n}$  is the normalized phase velocity of the wave. The growth rate in the relativistic approximation is dependent more fundamentally on the refractive index (actually phase velocity) rather than frequency.

If the quantity  $t_R = \bar{n}/n$  is considered as an effective resonant momentum, we can interpret equation (8) such that  $\frac{\pi^2 e R_J}{B_0(L)}$  is the electromagnetic coupling of the wave to the distribution,  $\frac{J(>\gamma)}{c} \left[ \frac{\gamma}{\bar{n}/n} \right]^{N-1}$  is the number "density" of particles with momenta  $t \geq t_R$ , and  $F(N, M, u)$  is the kinematical resonance integral over the curvature of the distribution. Figure 2 is a plot

of  $F(N, M, u)$  for  $N = 3$  and  $M = 0, 2, 4, 6, 8$ . Negative values of  $k_{\perp} R_J$  correspond to growth. At low frequencies,  $\bar{\omega} \rightarrow 0$ , the growth rate goes to zero falling off like  $\tau_R^{1-N}$ . At higher frequencies where  $u \sim 0.5$ , the growth rate turns over and goes to zero since at a high enough phase velocity as much energy is taken from the wave as is given, the resonance curve in Fig. 1 becoming increasingly concave.

If the cold plasma density is known the growth rate can be computed as a function of frequency through the transformation  $\frac{1}{u} = n(\omega)$ . A basic role of the cold plasma density is to determine a range of possible values for the refractive index:  $n_{\text{MAX}}(\omega = 0) \geq n(\omega) \geq n_{\text{MIN}}(\omega_p \text{ or } \frac{1}{2}\Omega_{ce})$ ,  $n_{\text{MIN}}$  occurring at the lesser of  $\omega_p$  or  $\frac{1}{2}\Omega_{ce}$ . This range specifies a phase velocity window which restricts the values of  $u$  in Fig. 2. For Jupiter, Frank et al. (1976) have reported cold proton densities, which have been used to plot Fig. 3. The cross-hatched region represents the accessible values of phase velocity at each  $L = R/R_J$  on the magnetic equator. By comparison with Fig. 2, using representative values of  $M = 4$  and  $N = 3$  for the pitch angle and spectral indices, there exists a range of unstable frequencies for  $1 \leq L \leq 12$ .

In the non-relativistic regime one can determine the marginally stable frequency and the corresponding resonant particle energy above which whistlers are unstable. For relativistic particles, since the resonance curve is hyperbolic, the marginal stability point is model-dependent and a function of  $M$  and  $N$ . We can estimate  $E_{M.S.}$  as follows. Choosing  $N = 3$  and

$M = 4$ , the marginally stable refractive  $n_{M.S.} \approx 2$ . If  $n_{MIN} \gg 1$ ,  $E_{M.S.}$  is non-relativistic and all relativistic electrons resonate with unstable whistlers. If  $n_{MIN} = 0(1)$ ,  $t_0 = t_{M.S.} = \frac{n_{M.S.}}{n_{M.S.} + 1}$ . Taking  $\bar{\omega} \ll 1$  gives  $t_{M.S.} = \frac{\Omega^2}{\omega_p^2} (n_{M.S.} - 1)$  and

$$E_{M.S.} = mc^2 \frac{\Omega^2}{\omega_{pc}^2} (n_{M.S.} - 1) = \frac{B^2}{4\pi n_0} \quad (10)$$

essentially the non-relativistic result.

### b) Path Integrated Growth

The actual logarithmic power gain of a ducted whistler requires the path-integrated quantity  $G = -2 \int_{s_1}^{s_2} k_1(s) ds$  along a flux tube. With certain approximations the integration can be done analytically and the result is good at least to the accuracy that fluxes are measured. We treat the cold plasma density as constant along the flux tube and require equal contributions to the gain from above and below the magnetic equator. As the wave convects along the tube from one hemisphere to the other, the phase velocity changes as demonstrated in Fig. 2 by either of the arrows A or B depending on the phase velocity (frequency) at the initial point. Liemohn (1967) found that waves which were locally damped at the magnetic equator could have a net positive gain because of greater growth contributions at higher latitudes. This situation is demonstrated by the arrow A for the curve  $M = 4$ . However, the stably-trapped flux is determined by the waves with maximal gain, which case

is represented by the arrow B; the situation carries the maximum of  $F(M, N, u)$  with the maximum flux of resonant particles at the equator. But from inspection of Fig. 2, if we choose  $N = 3$  and  $M = 4$  the excursion due to  $F(M, N, u)$  is small provided we consider phase velocities such that  $u(\omega) \lesssim .35$ . We thus approximate  $F(M, N, u)$  as constant in this frequency range and having the constant value  $F(N = 3, M = 4, u = 0) = -5/16$  (dotted lines) along the flux tube. The path integration then simply involves the scaling of the other factors in equation (8) along the flux tube. The coupling scales like  $(B_0/B)$  off the equator. For a  $\sin^M \theta$  distribution the flux scales like  $(B_0/B)^{M/2}$  (Roeberer, 1970). If  $n = 0(1)$ ,  $t_R = \bar{\Omega}/n$  scales like  $(B/B_0)$ . The growth rate along the tube is then approximately

$$k_i(s) = \left(\frac{B_0}{B}\right)^{\frac{M+2N}{2}} k_i(EQ) = \left(\frac{B_0}{B}\right)^5 k_i(EQ) \quad (11)$$

Then if  $G = -4 \int_0^{s_{\max}} k_i(s) ds$  and using the harmonic approximation  $B/B_0 = 1 + s^2/s_0^2$  for a dipole magnetic field where  $s_0^2 = \frac{2}{9} L^2 R_J^2$  we have

$$G = -4k_i(EQ) L R_J \int_0^{y_{\max}} dy (1 + y^2)^{\frac{-(M+2N)}{2}} \quad (12)$$

When  $y_{\max} = 1$  we are at the limit of the harmonic approximation, but since the integrand has decreased significantly, we extend the upper limit to infinity and for  $N = 3, M = 4$  the integration yields  $G = -4k_i R_J L(0.2)$ . Thus

$$G(N = 3, M = 4) \approx \frac{1}{4} \frac{\pi^2 e R_J L^4}{B_0} \frac{J(>\gamma)}{c} \left[ \frac{\gamma}{\bar{\Omega}/n} \right]^2 \quad (13)$$

We note that in scaling  $t_R \approx (\frac{B}{B_0})$  we required  $n = 0(1)$ . If the opposite extreme prevailed,  $n \gg 1$ , then  $t_R \sim (\frac{B}{B_0})^{3/2}$  and  $k_i(s) \approx (\frac{B_0}{B})^6 k_i(EQ)$ , with the integration providing  $G = -4k_i R_J L(0.18)$ , a slight difference. All quantities in equation (13) refer to the equator and (13) is valid for frequencies such that  $u(\omega) \lesssim 0.35$ .

Generalizing this procedure for arbitrary values of  $M$  and  $N$  and approximating  $F(N, M, u) \approx F(N, M, u = 0)$  we have

$$G \approx \sqrt{\frac{2}{9}}^{M(N-1)} \frac{\Gamma(\frac{N+1}{2}) \Gamma(\frac{M+3}{2}) \Gamma(\frac{M+2N-1}{2})}{\Gamma(\frac{M+2N}{2}) \Gamma(\frac{M+N+3}{2})} \frac{\pi^2 e R_J L^4}{B_0} \frac{J(>\gamma)}{c} \left[ \frac{\gamma}{\bar{\Omega}/n} \right]^{N-1} \quad (14)$$

We note that over the parameter range  $M = 2, 4, N = 2, 3, 4$ ,

$G$  is a relatively weak function of the indices, viz.

$G(N = 2, M = 4) \approx (1.27) G(N = 4, M = 2)$  at the extreme. This is due to the compensating effects of, say, larger growth rate ( $M$  increases) with smaller effective path length from equation (11). Therefore, over this parameter range we will take equation (13) as valid generally.

### c. Stably-Trapped Limit

Application of equation (13) to the concept of the stably-trapped limit (Kennel and Petchek, 1966) requires a knowledge of the power reflection coefficient,  $R$ , for the whistlers in the flux tube. If we treat the problem as strictly one-dimensional and define the volume emissivity  $\eta$  for parallel-propagating whistlers, the equation of radiative transfer  $\frac{\partial \bar{I}}{\partial s} = -2k_i(s)\bar{I} + \eta(s)$  yields for the intensity

$$I(s_{MAX}) = \frac{\int_{-s_{MAX}}^{s_{MAX}} ds \eta(s) \exp\left[-2 \int_s^{s_{MAX}} ds' k_i(s')\right]}{1 - Re^G} \quad (15)$$

It is clear that as  $Re^G \rightarrow 1$ , sufficiently high wave intensities will result that can relax the distribution on a time scale comparable to that of the particle source. No in situ wave measurements have been conducted at Jupiter, but if we assume that as in the Kennel-Petchek theory 5% of the wave energy is reflected, a gain of  $G = \ln(1/R) = 3$  is sufficient to maintain the stably trapped equilibrium. Equation (13) then gives a stably-trapped flux limit of

$$J^*( > t_R ) \approx 4 \times 10^{10} L^{-4} \frac{\text{electrons}}{\text{cm}^2 \text{ sec}} \quad (16)$$

The non-relativistic Kennel-Petschek result scaled to Jupiter is

$$J_{N-R}^*( > E_R ) = 7 \times 10^{10} L^{-4} \left( \frac{B_J R_E}{B_E R_J} \right) = 8.5 \times 10^{10} L^{-4} \quad (17)$$



which does not differ greatly from the relativistic result.

This fact is borne out by inspection of (8). The relativistic aspect is manifest only in the resonance integral  $F(N, M, u)$  and if the distribution is not too different topologically in the non-relativistic regime,  $F(N, M, u)$  should not vary a great deal. This can be further motivated by assuming a power law distribution in non-relativistic energy. If we take the distribution with  $\frac{1}{2}mv^2 = T$  as  $f(\vec{v}) = B \frac{\sin^M \theta}{T^{N+1}}$  where  $M$  and  $N$  are the pitch angle and spectral indices, a non-relativistic analysis yields the result equivalent to equation (8) for  $\bar{w} \ll 1$  of

$$k_i R_J = \frac{\pi^2 e R_J}{B_0} \frac{(T/T_R)^{N-1} J(>T)}{c} F_{N-R}(N, M, \bar{w}) \quad (18)$$

where

$$T_R = \frac{1}{2} m V_R^2 = \frac{1}{2} m \left( \frac{\bar{\Omega} c}{n} \right)^2$$

and

$$F_{N-R}(N, M, \bar{w}) = -M(N-1) \frac{2}{\sqrt{\pi}} \frac{\Gamma\left(\frac{M+3}{2}\right)}{\Gamma\left(\frac{M+2}{2}\right)} \left[ 1 - \frac{2\bar{w}}{M(1-\bar{w})} \right] \cdot \int_0^{\pi/2} d\theta \sin^{M+1} \theta \cos^{2N-1} \theta \quad (19)$$

If  $\bar{w} \ll 1$  and we neglect the second term in (19), then for  $N = 3$  and  $M = 4$ ,  $F_{N-R} = -\frac{1}{4}$  compared to  $F \approx -\frac{5}{16}$  taken previously.

Equation (16) describes a limit which should not be exceeded statistically by a flux of electrons at any energy. If the

flux  $J(>\nu)$  is known and obeys a power law, the distribution should depart from the power law and harden considerably at energies below the transition momentum given by

$$t_R = \gamma \left[ \frac{J(>\nu)}{J^*(>t_R)} \right]^{\frac{1}{N-1}} \quad (20)$$

since further extrapolation of the distribution by a power law to lower momenta would produce excessive amplification of the whistlers.

In Fig. 4 we have plotted  $J^*(>t_R)$  for comparison with observation. For  $L \leq 12$  measured electron spectra have spectral indices from 3 - 3.5 at high energies and pitch angle indices from 2 - 4 (Van Allen et al., 1974; Baker and Van Allen, 1976). We have taken the representative values of  $N = 3$  and  $M = 4$  throughout. The circles are equatorial electron fluxes of 5 Mev electrons from an empirical formula given by McIlwain and Fillius (1975); the diamonds are 5 Mev fluxes reported by Baker and Van Allen (1976) with error bars (D.N. Baker, private communication).

The theoretical uncertainty is more sensitive to the basic model we have chosen rather than to  $M$  and  $N$  values. We have taken the most efficient situation of parallel-propagation for which the growth rate is maximum (Kennel and Petschek, 1966); the finite  $k_{\perp}$  effects should raise  $J^*$ . A reflection coefficient of 0.5% rather than 5% raises  $J^*$  by a factor of 1.8. Another possibility is that centrifugal forces at large  $L$  confine the cold plasma to within  $\pm 1 R_J$  of the equator as suggested by Ioannidis and Brice (1971). The effective path length is then just  $2 R_J$

and for  $N = 3$ ,  $M = 4$  the modification to the limiting flux becomes  $J^* \sim 8 \times 10^9 L^{-3} \text{ cm}^{-2} \text{ sec}^{-1}$  which is also plotted in Fig. 4. Within the experimental and theoretical uncertainties involved, the data lie near the stably-trapped limit and suggest that whistlers may be active and that the transition energy is approximately 5 Mev at low  $L$  values.

### III DIFFUSION THEORY

#### a. Homogeneous Quasi-Linear Theory

If we consider a homogeneous plasma with parallel-propagating whistlers described by a one-dimensional electric field spectral density  $e(\omega)$  for waves travelling in just one direction, then

$$\left\langle \frac{E^2(\vec{x}, \tau)}{8\pi} \right\rangle = e_{TOT} = 2 \int_0^\infty d\omega e(\omega);$$

the factor of two includes the waves travelling in the opposite sense. The quasilinear diffusion equation for the resonant electrons can be written

$$\frac{\partial f(\vec{x}, \vec{v})}{\partial \tau} \Big|_{Q.L.} = - \frac{\partial}{\partial \vec{v}} \cdot \vec{J} \quad (21)$$

where

$$\begin{aligned} \vec{J} = & - \frac{4\pi^2 e^2}{m^2 c^2} \sum_{\omega_{\pm}} |v_g| \frac{e(\omega)}{|v_g - v_z|} \left\{ \frac{\partial f}{\partial t_{\perp}} \pm \frac{n}{\gamma} \left[ t_{\perp} \frac{\partial f}{\partial t_z} - t_z \frac{\partial f}{\partial t_{\perp}} \right] \right\} \\ & \left\{ \left( 1 \mp \frac{n t_z}{\gamma} \right) \hat{t}_{\perp} \pm \frac{n t_{\perp}}{\gamma} \hat{t}_z \right\} \end{aligned} \quad (22)$$

and  $\hat{t}_1, \hat{t}_2$  are unit vectors. Since a relativistic particle interacts with two waves going in opposite directions, the summation is over the frequencies satisfying  $\omega_{\pm} \left[ 1 \mp n(\omega_{\pm}) \frac{v_z}{c} \right] = \frac{\Omega_{ce}}{\gamma}$  and  $n(\omega_{\pm}) = \frac{|k|c}{\omega_{\pm}}$  for the R mode travelling in the  $\pm$  direction along  $B_0$ . The combination in the first bracket is proportional

to the incremental growth rate (see equation (1)) and has a sign,  $\sigma = +1$  ( $-1$ ), corresponding to the particles giving (taking) energy from the wave. The diffusion current is thus proportional to the spectral density and incremental growth rate and, if we consider the interaction with just an upward travelling wave, has the direction given by the angle  $\psi_+$  (see Fig. 5), where

$$\tan \psi_+ = \frac{J_z}{J_\perp} = \frac{nv \sin \theta}{c - nv \cos \theta}. \quad \text{For } v = c \text{ we have}$$

$$\psi_+ = \tan^{-1} \left[ \frac{n \sin \theta}{1 - n \cos \theta} \right] + \frac{\pi}{2}(1 + \sigma) \quad (23)$$

In general,  $\psi_+$  is directed such that those particles giving energy increase their pitch angle,  $\theta \rightarrow \pi$ , and those receiving energy decrease their pitch angle,  $\theta \rightarrow 0$ . If  $n \cos \theta \ll 1$ ,  $\psi_+ \approx \tan^{-1} [-\tan \theta] + \frac{\pi}{2}(1 + \sigma)$ , which describes pitch angle diffusion along approximately iso-energy surfaces, since  $t_z^2 + t_\perp^2 = \text{constant}$  leads to  $\frac{dt_z}{dt_\perp} = -\tan \theta$ . At  $\theta = \pi/2$   $\psi_+ = \tan^{-1} n$ , but these particles react equally to downgoing waves and the net current is in the  $\hat{t}_\perp$  direction. Thus  $90^\circ$  pitch angle particles should random-walk along the  $t_\perp$  axis to higher energies. The features of the diffusion are sketched qualitatively in Fig. 5. The locus of points where the incremental growth is zero defines a cone inside of which the diffusion is generally directed outwards. We note that non-relativistically, if we use the resonance condition  $1 - \frac{nv}{c} \cos \theta = \frac{1}{\gamma} \approx \bar{n}$ ,

$$\psi_+ = \tan^{-1} [(\bar{n} - 1) \tan \theta] + \frac{\pi}{2}(1 + \sigma) \quad (24)$$

and for low frequencies,  $\bar{\omega} \ll 1$ , pitch angle diffusion results.

### b. Radial Diffusion Particle Source

The immediate source of relativistic electrons in the inner Jovian magnetosphere has been identified as inward radial diffusion resulting from third adiabatic invariance violation (see, e.g., Simpson et al., 1974). As a source term the process can be written

$$\left. \frac{\partial f(\vec{x}, \vec{t})}{\partial \tau} \right)_{R.D.} = L^2 \frac{\partial}{\partial L} \left[ \frac{1}{L^2} D_{LL} \frac{\partial f}{\partial L} \right] \quad (25)$$

where the derivatives,  $\left. \frac{\partial}{\partial L} \right)_{M,J.}$  are taken at constant first and second invariant,  $M$  and  $J$ . If we neglect synchrotron losses and bounce average,  $\langle \rangle$ , equation (21), the evolution of the distribution under combined radial diffusion and quasilinear momentum-space diffusion by parallel-propagating whistlers is described by

$$\frac{\partial f(\vec{x}, \vec{t})}{\partial \tau} = \left. \frac{\partial f}{\partial \tau} \right)_{R.D.} + \langle \left. \frac{\partial f}{\partial \tau} \right)_{Q.L.} \rangle \quad (26)$$

If electric field intensities are small the first term dominates, but if the distribution is sufficiently unstable to whistlers and in a stably-trapped equilibrium, quasilinear diffusion must occur on a time scale comparable to that of radial diffusion. The solution of (26) in the steady state is formidable, but we can extract a crude approximation of the bounce-averaged  $\langle \mathcal{E}(\omega) \rangle$  in the stably-trapped regime. The radial diffusion coefficient

can be modeled as  $D_{LL} = D_0 L^\alpha$  where  $D_0 \sim 10^{-10} \text{ sec}^{-1}$  and  $\alpha \sim 4$  (Barbosa and Coroniti, 1975). Thus we estimate  $\tau_{R.D.} \sim \frac{1}{D_0 L^{\alpha-2}}$ . Writing (21) in spherical coordinates, we estimate the Q.L. relaxation time as  $\tau_{Q.L.} \sim \frac{m^2 c^2 t^2}{4\pi^2 e^2} \frac{|V_g - v_z|}{\langle V_g^2(\omega) \rangle}$ . The crudeness of the approximation is manifest in the neglect of the pitch angle dependence of  $\tau_{Q.L.}$  and also by the neglect of the derivatives in (22) and (25) which to some degree are compensatory. Equating characteristic times yields

$$\langle V_g^2(\omega) \rangle = \langle I_E(\omega) \rangle \sim \frac{m^2 c^2 t^2}{4\pi^2 e^2} |V_g - v_z| D_0 L^{\alpha-2} \quad (27)$$

If  $|V_g - v_z| \sim c$  and  $t = t_R = \bar{\Omega}/n$ , then letting  $2\pi\nu = \omega$  we have for the magnetic field intensity of modes travelling parallel to the field in one direction

$$\langle I_B(\nu) \rangle = 2\pi \langle n^2 I_E(\omega) \rangle \sim 1.5 \times 10^{-18} \bar{\Omega}^2 L^{\alpha-2} \frac{\text{WATTS}}{\text{m}^2 \text{Hz}} \quad (28)$$

$\langle I_B \rangle$  is a lower limit such that intensities much lower than (28) will not produce significant decreases in phase space densities evolving under pure radial diffusion.

We can estimate typical fluctuation field strengths from (28). If the bandwidth  $\Delta\nu \sim \nu \sim \frac{\Omega_{ce}}{2\pi}$ , then  $\frac{(\delta B)^2}{8\pi} \sim \frac{2}{V_g} \Delta\nu \langle I_B(\nu) \rangle$  and  $\delta B \sim .5 \text{ m}\gamma L^{\frac{1}{2}(\alpha-5)} = .5 \text{ m}\gamma L^{-\frac{1}{2}}$  for  $\alpha = 4$ . Such fields should be detectable by future spacecraft to Jupiter.

#### IV DISCUSSION

The evaluation of the limiting flux assumed whistlers were generated locally and did not propagate across L-shells. Without more detailed knowledge of the cold plasma distribution, ray path computations would be speculative. However, the fact that observed fluxes are lower than the limiting flux inside of  $L \leq 4$ , where phase space losses are still apparent, suggests that whistlers may propagate inward.

#### ACKNOWLEDGEMENT

We are indebted to M. Schulz for useful discussions concerning his work on relativistic electron fluxes in the earth's magnetosphere. This work was supported by NASA grant NGL 05-007-190-S7.



## REFERENCES

- Baker, D.N. and J.A. Van Allen, Energetic electrons in the Jovian magnetosphere, *J. Geophys. Res.*, 81, 617, 1976.
- Barbosa, D.D. and F.V. Coroniti, Lossy radial diffusion of relativistic Jovian electrons, *J. Geophys. Res.*, submitted, 1975.
- Coroniti, F.V., Energetic electrons in Jupiter's magnetosphere, *Astrophys. J., Suppl. Ser.*, 27, 261, 1974.
- Fillius, W., C. McIlwain, A. Mogro-Campero and G. Steinberg, Pitch angle scattering as an important loss mechanism for energetic electrons in the inner radiation belt of Jupiter, *J. Geophys. Res.*, submitted, 1975.
- Frank, L.A., K.L. Ackerson, J.H. Wolfe and J.D. Mihalov, Observations of plasmas in the Jovian magnetosphere, *J. Geophys. Res.*, 81, 457, 1976.
- Ioannidis, G. and N. Brice, Plasma densities in the Jovian magnetosphere: Plasma slingshot or Maxwell demon?, *Icarus*, 14, 360, 1971.
- Kennel, C.F. and H.E. Petchek, Limit on stably trapped particle fluxes, *J. Geophys. Res.*, 71, 1, 1966.
- Lerche, I., Wave phenomena in the interstellar plasma, *Advan. Plasma Phys.*, 2, 47, 1969.
- Liemoohn, H., Cyclotron-resonance amplification of VLF and ULF whistlers, *J. Geophys. Res.*, 72, 39, 1967.
- Lyons, L.R., R.M. Thorne and C.F. Kennel, Pitch-angle diffusion of radiation belt electrons within the plasmasphere, *J.*

- Geophys. Res., 77, 3455, 1972.
- McIlwain, C.E. and R.W. Fillius, Differential spectra and phase space densities of trapped electrons at Jupiter, J. Geophys. Res., 80, 1341, 1975
- Roberts, J.A. and M.M. Komesaroff, Observations of Jupiter's radio spectrum and polarization in the range from 6 cm to 100 cm, Icarus, 4, 127, 1965.
- Roederer, J.G., Dynamics of Geomagnetically Trapped Radiation, Springer-Verlag, Berlin, 1970.
- Schulz, M. and A.L. Vampola, Magnetospheric stability criteria for artificial electron belts, J. Geophys. Res., submitted, 1975.
- Simpson, J.A., D.C. Hamilton, R.B. McKibben, A. Mogro-Campero, K.R. Pyle and A.J. Tuzzolino, The protons and electrons trapped in the Jovian dipole magnetic field region and their interaction with Io, J. Geophys. Res., 79, 3522, 1974.
- Stansberry, K.G. and R.S. White, Jupiter's radiation belts, J. Geophys. Res., 79, 2331, 1974.
- Thorne, K.S., Dependence of Jupiter's decimeter radiation on the electron distribution in its Van Allen belts, Radio Sci., 69D, 1557, 1965.
- Van Allen, J.A., D.N. Baker, B.A. Randall and D.A. Sentman, The magnetosphere of Jupiter as observed with Pioneer 10 1. Instrument and principal findings, J. Geophys. Res., 79, 3559, 1974.

## FIGURE CAPTIONS

- Figure 1. Resonance curves plotted in normalized momentum space for  $R = \frac{\omega_p^2}{\omega_{ce}^2} = \frac{1}{4}$ . The solid line is the resonance with a whistler ( $\bar{n} = 10$ ,  $n = 1.94$ ) travelling along the field; the dashed line with a whistler ( $\bar{n} = 4$ ,  $n = 1.53$ ) travelling anti-parallel.
- Figure 2. Plot of the resonance function  $F(M, N, u)$  for  $M = 0, 2, 4, 6, 8$  and  $N = 3$ . Growth occurs for negative values of  $F(M, N, u)$  increasing with  $M$ . The arrows represent convective changes of  $F$  along the flux tube.
- Figure 3. Accessible phase velocities for whistlers using Frank et al. (1976) cold plasma observations.
- Figure 4. Plot of the limiting flux  $J^*(>t_R)$ . The circles are 5 Mev equatorial fluxes of McIlwain and Fillius (1975); the diamonds are 5 Mev fluxes of Baker and Van Allen (1976) with error bars. The dotted line is the modification to  $J^*$  from centrifugal effects.
- Figure 5. Qualitative view of diffusion in normalized momentum space arising from interaction with whistler propagating parallel to  $\vec{B}_0$  only. The small arrows give only the direction of the diffusion. The cone edge is defined by the integrand of (3) being zero for  $N = 3$  and  $M = 4$ .




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5



# REPRODUCIBILITY OF THE ORIGINAL PAGE IS POOR

## UCLA PLASMA PHYSICS GROUP REPORTS

- PG-1 Propagation of Ion Acoustic Waves Along Cylindrical Plasma Columns", A.Y. Wong (July 1965). *Phys. Fluids* 9, 1261 (1966).
- PG-2 Stability Limits for Longitudinal Waves in Ion Beam-Plasma Interaction", B.D. Fried and A.Y. Wong (August 1965). *Phys. Fluids* 9, 1084 (1966).
- PG-3 The Kinetic Equation for an Unstable Plasma in Parallel Electric and Magnetic Fields", B.D. Fried and S.L. Ossakow, (November 1965). *Phys. Fluids* 9, 2428 (1966).
- PG-4 Low-Frequency Spatial Response of a Collisionless Electron Plasma", B.D. Fried, A.N. Kaufman and D.L. Sachs (August 1965). *Phys. Fluids* 9, 292 (1966).
- PG-5 Effects of Collisions on Electrostatic Ion Cyclotron Waves", A.Y. Wong, D. Judd and F. Hal (December 1965). *Phys. Letters*, 21, 157 (1966).
- PG-6 Interaction Between Ion Beams and Plasmas", R. Rowberg, A.Y. Wong and J.M. Sellen (April 1966). *APS Bull.* 10, 1182 (1965).
- PG-7 Observation of Cyclotron Echoes from a Highly Ionized Plasma", D.E. Kaplan and R.M. Hill (May 1966). *Phys. Lett.* 21, 157 (1966).
- PG-8 Excitation and Damping of Drift Waves", A.Y. Wong and R. Rowberg (July 1966). *Proceedings of 1st Int. Conf. On Quiescent Plasmas*, Frascati, Rome, 1967. *Phys. Rev. Lett.* 18, 526 (1967).
- PG-9 The Guiding Center Approximation in Lowest Order", A. Bales, Jr. (Sept. 1966). *J. Plasma Phys.* 1, 305 (1967).
- PG-10 Plasma Streaming into a Magnetic Field", S.L. Ossakow (November 1966). Dissertation.
- PG-11 Cooperative Effects in Plasma Echo Phenomena", A.Y. Wong (March 1967). *Proc. of 1st Int. Conf. on Quiescent Plasmas*, Frascati, Rome, 1967.
- PG-12 A Quantum Mechanical Theory of Grid Excitation of Low Frequency Waves in a Plasma", G.L. Johnston (April 1967) *Dissertation*.
- PG-13 Linear and Nonlinear Diffusion of an Isolated Plasma Column", J. Newman (May 1967). *Dissertation*.
- PG-14 Two-Pole Approximation for the Plasma Dispersion Function", B.D. Fried, C.L. Hedrick and J. McCune (Aug. 1967). *Phys. Fluids* 11, 249, (1968).
- PG-15 Experimental Investigation of Electron Runaway Phenomena", J.S. McCrone (August 1967) *Dissertation*.
- PG-16 Parametric Coupling Between Drift Waves", F. Hal, R. Rowberg and A.Y. Wong (Oct. 1967). *Proc. of 2nd Int. Symp. on Fluctuations and Diffusion in Plasmas*, June, 1967.
- PG-17 Cyclotron Echoes from Drift Waves", A.Y. Wong (March 1968). *Phys. Rev. Lett.* 20, 310 (1968).
- PG-18 Ion Wave Echoes", D. Bales, H.B. Akers and A.Y. Wong (Nov. 1967). *Phys. Rev. Lett.* 20, 310 (1968).
- PG-19 Cyclotron Echoes in Plasmas", D. Judd (March 1968). *Dissertation*.
- PG-20 Test Particle Theory for Quantum Plasmas", M.E. Rensink (Oct. 1967). *Phys. Rev.* 164, 175 (1967).
- PG-21 Artificial Van Allen Belt", C.F. Kennel (Nov. 1967).
- PG-22 Landau Damping of Ion Acoustic Waves in a Cesium Plasma with Variable Electron-Ion Temperature Ratio", K.B. Ranganathan (Oct. 1967). *Dissertation*.
- PG-23 The Inhomogeneous Two-Stream Instability", G. Inzer (Sept. 1967).
- PG-24 Magnetic Turbulence in Shocks", C.F. Kennel and H.E. Petschek (Dec. 1967). In *Physics of the Magnetosphere*, R. Carovillano, J.F. McCloy, and H. R. Hudson, eds. 485-515, D. Reidel, Dordrecht, Holland, 1968.
- PG-25 Small Amplitude Waves in High  $\beta$  Plasmas", V. Formisano and C.F. Kennel (Feb. 1968). *J. Plasma Phys.* 3, 55 (1969).
- PG-26 Low  $\beta$  Plasma Penetration Across a Magnetic Field", B.D. Fried and S. Ossakow (Mar. 1968). *Phys. Fluids* 12, 702 (1969).
- PG-27 Annual Status Report", Feb. 1, 1967-Jan. 31, 1968. Principal Investigators, A. Bales, Jr., B.D. Fried, C.F. Kennel.
- PG-28 The Theorist's Magnetosphere", C.F. Kennel (April 1968).
- PG-29 Electromagnetic Pitch Angle Instabilities in Space", C.F. Kennel and F.L. Scarf (April 1968) in *Plasma Waves in Space and in the Laboratory*, ed. J.O. Thomas and B.J. Landmark, Edinburgh U. Press, Edinburgh, Vol. 11, 1969.
- PG-30 Electromagnetic Echoes in Collisionless Plasmas", A.Y. Wong (April 1968). *Phys. Fluids* 12, 866 (1969).
- PG-31 Parametric Excitation of Drift Waves in a Resistive Plasma", G. Weyl and M. Goldman (June 1968). *Phys. Fluids* 12, 1097 (1969).
- PG-32 Parametric Excitation from Thermal Fluctuations at Plasma Drift Wave Frequencies", A.Y. Wong, M.V. Goldman, F. Hal, R. Rowberg (May 1968). *Phys. Rev. Letters* 21, 518 (1968).
- PG-33 Current Decay in a Streaming Plasma Due to Weak Turbulence", S.L. Ossakow and B.D. Fried (June 1968). *Phys. Fluids* 11, 274 (1968).
- PG-34 Temperature Gradient Instabilities in Axisymmetric Systems", C.S. Liu (Aug. 1968). *Phys. Fluids* 12, 1489 (1969).
- PG-35 Electron Cyclotron Echo Phenomena in a Hot Collisionless Plasma", D. Judd (Aug. 1968). *Dissertation*.
- PG-36 Transverse Plasma Wave Echoes", B.D. Fried and C. Olson (Oct. 1968). *Phys. Rev.* 180, 214 (1969).
- PG-37 Also presented at APS Plasma Physics Div. Annual Meeting, Nov. 1968.
- PG-38 Low Frequency Interchange Instabilities of the Ring Current Belt", C.S. Liu (Jan. 1969). *J. Geophys. Res.* 75, 3789 (1970).
- PG-39 Drift Waves in the Linear Regime", R.E. Rowberg and A.Y. Wong (Feb. 1969). *Phys. Fluids* 13, 661 (1970).
- PG-40 Parametric Mode-Mode Coupling Between Drift Waves in Plasmas", F. Hal and A.Y. Wong (Sept. 1970).
- PG-41 Nonlinear Mode-Mode Coupling Between Drift Waves in Plasmas", F. Hal and A.Y. Wong (Sept. 1970).
- PG-42 Ion-Burst Excited by a Grid in a Plasma", H. Ikeshi and R.J. Taylor (Feb. 1969). *J. Appl. Phys.* 41, 738 (1970).
- PG-43 Measurements of Diffusion in Velocity Space from Ion-Ion Collisions", A. Wong and D. Baker (Mar. 1969). *Phys. Rev.* 180, 1 (1969).
- PG-44 Nonlinear Excitation in the Ionosphere", A.Y. Wong (Mar. 1969).
- PG-45 Observation of 1st order Ion Energy Distribution in Ion Acoustic Waves", H. Ikeshi, R. Taylor, *Phys. Rev. Lett.* 22, 923, (1969).
- PG-46 A New Representation for the Conductivity Tensor of a Collisionless Plasma in a Magnetic Field", B.D. Fried and C. Hedrick (Mar. 1969). *Zeitschrift für Geophysik*, U. of Chicago Press, (1969).
- PG-47 Direct Measurements of Linear Growth Rates and Nonlinear Saturation Coefficients", A.Y. Wong and F. Hal (April 1969). *Phys. Rev. Lett.* 23, 165 (1969).
- PG-48 Electron Precipitation Pulsations", F. Coroniti and C.F. Kennel (April 1969). *J. Geophys. Res.* 75, 1279 (1970).
- PG-49 Auroral Microprecipitation Instability", F. Coroniti and C.F. Kennel (May 1969). *J. Geophys. Res.* 75, 1863 (1970).
- PG-50 Effect of Fokker-Planck Collisions on Plasma Wave Echoes", G. Johnston (June 1969). *Phys. Fluids* 13, 116 (1970).
- PG-51 Linear and Nonlinear Theory of Grid Excitation of Low Frequency Waves in a Plasma", G. Johnston (July 1969).
- PG-52 Theory of Stability of Large Amplitude Periodic (MCK) Waves in Collisionless Plasmas", M.V. Goldman (June 1969). *Phys. Fluids* 13, 1281 (1970).
- PG-53 Observation of Strong Ion Wave-Wave Interaction", R. Taylor and H. Ikeshi (Aug. 1969). *Phys. Fluids* 13, 1802 (1970).
- PG-54 Optical Mixing in a Magnetospheric Plasma", G. Weyl (Aug. 1969). *Phys. Fluids* 13, 1863 (1970).
- PG-55 Trapped Particles and Echoes", A.Y. Wong and R. Taylor (Oct. 1969). *Phys. Rev. Lett.* 23, 958 (1969).
- PG-56 Formation and Interaction of Ion-Acoustic Solitons", H. Ikeshi, R.J. Taylor and D.R. Baker, *Phys. Rev. Lett.* 25, 11 (1970).
- PG-57 Observation of Collisionless Electrostatic Shocks", R. Taylor, D. Baker and H. Ikeshi, *Phys. Rev. Lett.* 25, 206 (1970).
- PG-58 Turbulent Loss of Ring Current Protons", J.M. Cornwall, F.V. Coroniti and R.M. Thorne (June 1970). *J. Geophys. Res.* 75, 4699 (1970).
- PG-59 Efficient Modulation Coupling Between Electron and Ion Resonances in Magnetospheric Plasmas", A. Wong, D.R. Baker, H. Booth (Dec. 1969). *Phys. Rev. Lett.* 24, 804 (1970).
- PG-60 Interaction of Quasi-Transverse and Quasi-Longitudinal Waves in an Inhomogeneous Vlasov Plasma", C.L. Hedrick (Jan. 1970). *Dissertation*.
- PG-61 Observation of Strong Ion Acoustic Wave-Wave Interaction", R.J. Taylor and H. Ikeshi (Jan. 1970).
- PG-62 Perturbed Ion Distributions in Ion Waves and Echoes", H. Ikeshi and R.J. Taylor (Jan. 1970). *Phys. Fluids* 13, 2348 (1970).
- PG-63 Propagation of Ion Cyclotron Harmonic Wave", E.R. Ault and H. Ikeshi (Nov. 1970). *Phys. Fluids* 13, 2076 (1970).
- PG-64 The Analytic and Asymptotic Properties of the Plasma Dispersion Function", A. Bales, Jr., & G. Johnston (Feb. 1970).
- PG-65 Effect of Ion-Ion Collision and Ion Wave Turbulence on the Ion Wave Echo", D. Baker (June 1970).
- PG-66 Dispersion Discontinuities of Strong Collisionless Shocks", F.V. Coroniti (Mar. 1970). *J. Plasma Phys.* 4, 26, (1970).
- PG-67 An Ion Cyclotron Instability", E.S. Weibel (Apr. 1970). *Dissertation*.
- PG-68 Turbulence Structure of Finite- $\beta$  Perpendicular Fast Shocks", F.V. Coroniti (Apr. 1970). *J. Geophys. Res.* 75, 7007 (1970).
- PG-69 Strengthening of Ion Acoustic Waves and Formation of Collisionless Electrostatic Shocks", R. Taylor (Apr. 1970).
- PG-70 A Method of Studying Trapped Particles Behavior in Magnetic Geometries", C.S. Liu and A.Y. Wong (Apr. 1970).
- PG-71 Phys. Rev. Lett. 25, 1702 (1970).
- PG-72 A Note on the Differential Equation  $g'' + a^2g = 0$ ", E.S. Weibel (Apr. 1970).
- PG-73 Plasma response to a Step Electric Field Greater than the Critical Runaway Field, With and Without an External Applied Magnetic Field", J.E. Robin (June 1970). *Dissertation*.
- PG-74 The UC Mathematical On-Line Systems as a Tool for Teaching Physics", B.D. Fried and R.B. White (Aug. 1970). *Proc. of Comp. Graphics*, 134-154, 1970.
- PG-75 High Frequency Hall Current Instability", F. Lee, C.F. Kennel, J.M. Kindel (Aug. 1970). *Radio Sci.* 6, 209 (1971).
- PG-76 Laminar Wave Train Structure of Collisionless Magnetic Slow Shocks", F.V. Coroniti (Sept. 1970).
- PG-77 Field-Aligned Current Instabilities in the Topside Ionosphere", J.M. Kindel and C.F. Kennel (Aug. 1970). *J. Geophys. Res.* 76, 3005 (1971).
- PG-78 Spatial Cyclotron Damping", C. Olson (Sept. 1970). *Phys. Fluids* 13, 160 (1972).
- PG-79 Electromagnetic Plasma Wave Propagation Along a Magnetic Field, C. Olson (Sept. 1970). *Dissertation*.
- PG-80 Electron Plasma Waves and Free-Streaming Electron Bursts", H. Ikeshi, F.J. Barrett, R.B. White and A.Y. Wong (Nov. 1970). *Phys. Fluids* 14, 1997 (1971).
- PG-81 Relativistic Electron Precipitation During Magnetic Storm Main Phase", R.M. Thorne and C.F. Kennel (Nov. 1970). *J. Geophys. Res.* 76, 4446 (1971).
- PG-82 A Unified Theory of SAR-Arc Formation at the Plasmapause", J.M. Cornwall, F.V. Coroniti and R.M. Thorne (Nov. 1970). *J. Geophys. Res.* 76, 4428 (1971).
- PG-83 Nonlinear Collisionless Interaction Between Electron and Ion Modes in Inhomogeneous Magnetospheric Plasmas", H. Booth (Dec. 1970). *Dissertation*.
- PG-84 Observation of Parametrically Excited Ion Acoustic Waves", R. Stenzel (Mar. 1971).
- PG-85 Remote Double Resonance Coupling of Radar Energy to Ionospheric Irregularities", C.F. Kennel (Jan. 1971). *Comm. Astro & Space Phys.* 3, 2, 87 (1971).
- PG-86 Ion Acoustic Waves in a Multi-Ion Plasma", B.D. Fried, R.B. White, T. Saenz (Jan. 1971). *Phys. Fluids* 14, 2388 (1971).
- PG-87 Current-Driven Electrostatic and Electromagnetic Ion Cyclotron Instabilities", D.H. Forslund, C.F. Kennel, J.M. Kindel (Feb. 1971).
- PG-88 Locating the Magnetospheric Ring Current", C.F. Kennel and R. Thorne (Mar. 1961). *Comm. Astrophys. & Space Phys.* C3, 115 (1971).
- PG-89 Ion Acoustic Instabilities Due to Ions Streaming Across Magnetic Field", P.J. Barrett, R.J. Taylor (Mar. 1971).
- PG-90 Evolution of Turbulent Electronic Shocks", A.Y. Wong and R. Means (July 1971). *Phys. Rev. Lett.* 27, 973 (1971).
- PG-91 Density Step Production of Large Amplitude Collisionless Electrostatic Shocks and Solitons", D.B. Cohen (June 1971).
- PG-92 Turbulent Resistivity, Diffusion and Heating", B.D. Fried, C.F. Kennel, E. McKenzie, F.V. Coroniti, J.M. Kindel, R. Stenzel, R.J. Taylor, R.B. White, A.Y. Wong, W. Bernstein, J.M. Sellen, D. Forslund & R.A. Sagdeev (June 1971). *Proc. 4th Conf. Plasma Phys. & Cont. Nucl. Fusion Res.* Madison, Wis., IAEA-CN-20/E-4, Plasma Phys. & Cont. Nucl. Fusion Res. 11, Vienna (1971).
- PG-93 Nonlinear Evolution and Saturation of an Unstable Electrostatic Wave", B.D. Fried, C.S. Liu, R.M. Means & R.Z. Sagdeev (Aug. 1971).
- PG-94 Cross-Field Current-Driven Ion Acoustic Instability", P.J. Barrett, B.D. Fried, C.F. Kennel, J.M. Sellen & R.J. Taylor (Dec. 1971). *Phys. Rev. Lett.* 28, 337 (1972).
- PG-95 3-D Velocity Space Diffusion in Beam-Plasma Interaction Without Magnetic Field", P.J. Barrett, D. Gresslin & A.Y. Wong (Sept. 1971). *Proc. 3rd Int. Conf. on Quies. Plas.* 291, Elsinore, Calif., 1971.
- PG-96 Diverse Auroral Oval Plasma Density and Conductivity Enhancements Due to Magnetospheric Electron Precipitation", C.F. Kennel & M.H. Rees (Sept. 1971). *J. Geophys. Res.* 77, 2294 (1972).
- PG-97 Collisionless Wave-Particle Interactions Perpendicular to the Magnetic Field", A.Y. Wong, D.L. Jassby (Sept. 1971). *Phys. Rev. Lett.* 28, 41 (1972).
- PG-98 Magnetospheric Substorms", F.V. Coroniti & C.F. Kennel (Sept. 1971). *Cos. Plas. Phys.* *Proc. of Conf. Cos. Plas. Phys. held at Euro Space Res. Inst.* Frascati, Italy, Sept. 1971. *Plenum Press*, 1972, ed. Karl Schindler.
- PG-99 Magnetopause Motions, DP-2, and the Growth Phase of Magnetospheric Substorms", F.V. Coroniti & C.F. Kennel (Sept. 1971).
- PG-100 Structure of Ion Acoustic Solitons and Shock Waves in a Two-Component Plasma", R.B. White, B.D. Fried & F.V. Coroniti (Sept. 1971). *Phys. Fluids* 15, 1484 (1972).
- PG-101 Solar Wind Interaction with Lunar Magnetic Field", G. Siscoe (Meteorology Dept.) & D. Goldstein (JPL) (Nov. 1971).
- PG-102 Changes in Magnetospheric Configuration During Substorm Growth Phase", F.V. Coroniti & C.F. Kennel (Nov. 1971). *J. Geophys. Res.* 77, 3361 (1972).
- PG-103 Trip Report-1971. Elev. Conference on Plasma Theory & Visits to Lebedev & Burchakov Institutes", B.D. Fried (Oct. 1971).
- PG-104 Pitch-Angle Diffusion of Radiation Belt Electrons Within the Plasmasphere", L.R. Lyons, R.M. Thorne, C.F. Kennel (Jan. 1972). *J. Geophys. Res.* 77, 3455 (1972).
- PG-105 Remote Feedback Stabilization of a High- $\beta$  Plasma", F.F. Chen, D. Jassby & M. Maric (Dec. 1971). *Phys. Fluids* 15, 1864 (1972).
- PG-106 Remote Plasma Control, Heuristic Measurements of Electron Distribution and Trapped Particles by Nonlinear Electromagnetic Interaction", A.Y. Wong, F.F. Chen, H. Booth, D.L. Jassby, R. Stenzel, D. Baker & C.S. Liu (June 1971). *J. Plasma Phys. & Cont. Nucl. Fusion Res.* 1, 335 (1971).
- PG-107 Computational and Experimental Plasma Physics for Theoreticians", B.D. Fried (Jan. 1972). *Adv. Plas. Phys.* 5, 211 (1974).
- PG-108 Threshold and Saturation of the Parametric Decay Instability", R. Stenzel & A.Y. Wong (Nov. 1971). *Phys. Rev. Lett.* 28, 274 (1972).
- PG-109 Laser Amplification in an Inhomogeneous Plasma", R. White (Jan. 1972).
- PG-110 External Production and Control of Electrojet Irregularities", E. Lee, P.R. Kav & C.F. Kennel (Jan. 1972). *J. Geophys. Res.* 77, 4197 (1972).
- PG-111 Ion Heating Via Turbulent Ion Acoustic Waves", R.J. Taylor & F.V. Coroniti (Feb. 1972). *Phys. Rev. Lett.* 29, 34 (1972).
- PG-112 Polarization of the Auroral Electrojet", F.V. Coroniti & C.F. Kennel (Feb. 1972). *J. Geophys. Res.* 77, 2035 (1972).
- PG-113 Mode Coupling and Wave Particle Interactions for Unstable Ion Acoustic Waves", F. Martin & B.D. Fried (Feb. 1972). *Phys. Fluids* 15, 2275 (1972).
- PG-114 Parallel Magnetic Multi-pole Confinement of a Magnetic Field-Free Plasma", R. Limpacher (Mar. 1972). *Dissertation*.
- PG-115 Turbulence in Electrostatic Collisionless Shock Waves", R.M. Means (Apr. 1972). *Dissertation*.
- PG-116 Large Diameter, Quiescent Plasma in a Magnetospheric Field", E. Ault (Apr. 1972). *Dissertation*.
- PG-117 Parasitic Pitch-Angle Diffusion of Radiation Belt Particles by Ion-Cyclotron waves", L.R. Lyons & R.M. Thorne (Mar. 1972). *J. Geophys. Res.* 77, 5008 (1973).
- PG-118 A New Role for Infrared Lasers", F.F. Chen (May 1972). *Comm. Plasma Phys. & Cont. Nucl. Fusion* 1, 81 (1972).
- PG-119 Electrostatic Instability of Ring Current Protons Beyond the Plasmapause During Injection Events", F.V. Coroniti, R.W. Fredericks & R. White (May 1972). *J. Geophys. Res.* 77, 6243 (1972).
- PG-120 Magnetospheric Structure of the Outer Planets", C.F. Kennel (May 1972). *Space Sci. Rev.* 14, 511 (1973).
- PG-121 Measurement of Transverse and Longitudinal Heat Flow in a Laser-Heated, Magnetically Confined Arc Plasma", S. B. Fav (June 1972). *Dissertation*. Cond. version by S.W. Fav, F. Chen & D. Jassby, *Phys. Rev. Lett.* 42a, 261 (1972).

- PPG-122 Plasmaspheric Hiss", R.N. Thorne, E.J. Smith, R.E. Burton, R.B. Nisler (July 1972) *J. Geophys. Res.* 78, 1581 (1973).
- PPG-123 Magnetospheric Electrons", F.V. Coroniti & R.N. Thorne (July 1972) *Ann. Rev. Earth & Plan. Sci.* 1, 107 (1973).
- PPG-124 Calculation of Reflection and Transmission Coefficients for a Class of One-Dimensional Wave Propagation Problems in Inhomogeneous Media", A. Baños (Sept 1972) *J. Math. Phys.* 14, 963 (1973).
- PPG-125 Electromagnetic Wave Functions for Parabolic Plasma Density Profiles", A. Baños & D. Keller (Sept 1972) *Phys. Fluids* 17, 2275 (1974).
- PPG-126 Amplification of Electromagnetic Waves in Overdense Plasmas", F.F. Chen & R.B. White (Sept 1972) *J. Plas. Phys.* 16, 565 (1974).
- PPG-127 Abstracts presented at An Phys Soc Div of Plas Phys Ann Meeting, Monterey (Nov 13-16, 1972).
- PPG-128 Can the Ionosphere Regulate Magnetospheric Convection", F.V. Coroniti & C.F. Kennel (Oct 1972) *J. Geophys. Res.* 78, 2837 (1973).
- PPG-129 Nonlinear Stabilization of Oscillating Two-Stream Instability", K. Nishikawa, Y.C. Lee & P.E. Kaw (Oct 1972) *Phys. Fluids* 16, 1380 (1973).
- PPG-130 Drift Waves in Finite  $\beta$  Plasmas", H.S. Chance (Oct 1972) Thesis.
- PPG-131 Wave Packet Formulation of Nonlinear Plasma Kinetics", K. Nishikawa & B.D. Fried (Oct 1972) *Phys. Fluids* 16, 1521 (1973).
- PPG-132 Electron Cyclotron Drift Instability Experiment", B.H. Ripin & R.L. Stenzel (Oct 1972) *Phys. Rev. Lett.* 30, 45 (1973).
- PPG-133 Resonant Excitation of Electrostatic Modes with Electromagnetic Waves", G. Schmidt (Oct 1973) *Phys. Fluids* 16, 1676 (1973).
- PPG-134 Energetic Ion Beam Source and Free-Stream Beam Diagnostic Techniques", R.L. Stenzel & B.H. Ripin (Nov 1972) *Rev. Sci. Instr.* 44, 617 (1973).
- PPG-135 Electron Plasma Waves in an Unbounded Uniform Magnetoplasma", R.L. Stenzel (Nov 1972) *Phys. Fluids* 16, 565 (1973).
- PPG-136 Convective Amplification of Type I Irregularities in the Equatorial Electrojet", E. Lee & C.F. Kennel (Nov 1972) *J. Geophys. Res.* 78, 4619 (1973).
- PPG-137 Effects of Propagation Parallel to the Magnetic Field on the Type I Electrojet Irregularity Instability", E. Lee & C.F. Kennel (Nov 1972) *Plan. & Space Sci.* 21, 1339 (1973).
- PPG-138 Analog Computer Simulation of Parametric Instabilities", R.L. Stenzel (Nov 1972) Thesis.
- PPG-139 Theory of Double Resonance Parametric Excitation in Plasmas", D. Arnush, B.D. Fried, C.F. Kennel, K. Nishikawa & A.Y. Hong (Nov 1972) *Phys. Fluids* 16, 2270 (1973).
- PPG-140 Filamentation and Trapping of Electromagnetic Radiation in Plasmas", P. Kaw, G. Schmidt & T. Wilcox (Dec 1972) *Phys. Fluids* 16, 1522 (1973).
- PPG-141 Finite  $\beta$  Drift Wave Instability", H.S. Chance, F.V. Coroniti & C.F. Kennel (Jan 1973) *J. Geophys. Res.* 78, 7521 (1973).
- PPG-142 The Formation of Ion Acoustic Shocks", R.B. White, B.D. Fried & F.V. Coroniti (Jan 1973) *Phys. Fluids* 17, 211 (1974).
- PPG-143 Experiments on Parametric Instabilities", A.Y. Hong (Mar 1973) Thesis.
- PPG-144 On Cosmic Ray Generation by Pulsars", C.F. Kennel, G. Schmidt & T. Wilcox (Mar 1973) *Phys. Rev. Lett.* 31, 1364 (1973).
- PPG-145 On the Marginally Stable Saturation Spectrum of Unstable Type I Equatorial Electrojet Irregularities", E. Lee, C.F. Kennel & F.V. Coroniti (Apr 1973) *J. Geophys. Res.* 78, 249 (1974).
- PPG-146 Spatial Growth Properties of Parametric & Backscattering Plasma Instabilities", B.D. Fried, R. Gould & G. Schmidt (Apr 1973) *J. Plas. Phys.* 16, 121 (1975).
- PPG-147 Evolution of BIC-Like Modes with Trapped Electrons", A.Y. Hong, B.H. Ripin & B. Ripin (Apr 1973) *Phys. Rev. Lett.* 30, 1299 (1973).
- PPG-148 Stabilization of Ion Acoustic Waves by Electron Trapping", N. Albrecht (Apr 1973) *Phys. Fluids* 17, 206 (1974).
- PPG-149 Turbulence in Electrostatic Ion Acoustic Shocks", R.W. Hoens, F.V. Coroniti, A.Y. Hong & R.B. White (May 1973) *Phys. Fluids* 16, 2304 (1973).
- PPG-150 Theory of Dielectric Function in a Magnetized Plasma", Y.C. Lee & C.S. Liu (June 1973) Thesis.
- PPG-151 Physical Interpretation of the Oscillatory Two-Stream Instability", A.Y. Hong & G. Schmidt (June 1973) Thesis.
- PPG-152 Relativistic Particle Motion in Nonuniform Electromagnetic Waves", G. Schmidt & T. Wilcox (June 1973) *Phys. Rev. Lett.* 31, 1380 (1973).
- PPG-153 The Ring Current and Magnetic Storms", F.V. Coroniti (July 1973) *Radio Sci.* 8, 1007 (1973).
- PPG-154 Energetic Electrons in Jupiter's Magnetosphere", F.V. Coroniti (July 1973) *Phys. Fluids* 17, 261 (1974).
- PPG-155 Stably Trapped Proton Fluxes in the Jovian Magnetosphere", F.V. Coroniti, C.F. Kennel & R.N. Thorne (July 1973) *Astrophys. J.* 185, 385 (1974).
- PPG-156 Absolute Raman Scattering Instabilities in an Inhomogeneous Plasma", J.F. Drake & Y.C. Lee (July 1973) *Phys. Rev. Lett.* 31, 1197 (1973).
- PPG-157 Growth & Saturation of the Absolute Electron Cyclotron Drift Instability", R.L. Stenzel & B.H. Ripin (July 1973) *Phys. Rev. Lett.* 31, 1545 (1973).
- PPG-158 Parametric Instabilities of Electromagnetic Waves in Plasmas", J. Drake, P.E. Kaw, Y.C. Lee, G. Schmidt, C.S. Liu & M.N. Rosenbluth (July 1973) *Phys. Fluids* 17, 778 (1974).
- PPG-159 Nonlinear Optics of Plasmas", F.F. Chen (Aug 1973) *Survey Lect. Int. Cong. Waves & Instab. Plas.* Innsbruck Austria, 1973. Institute for Theoretical Physics, Innsbruck pp. C1-C19.
- PPG-160 Physical Mechanisms for Laser & Plasma Heating - Parametric Instabilities", F.F. Chen (Aug 1973) in *Laser Interaction and Related Plasma Phenomena*, ed. W.J. Schwartz & W. More, Vol. 3A, p. 291-313 (Plenum Press 1974).
- PPG-161 Trip Report - 6th Euro Conf. on Fusion & Plas. Phys., July 30-Aug. 4, 1973, Moscow", B.D. Fried (Aug 1973).
- PPG-162 Abstracts presented at Phil Meeting of Am Phys Soc, Div. of Plas. Phys., Oct 31-Nov 3, 1973.
- PPG-163 Enhancement of Plasma DC Currents by Intense AC Fields", A.T. Lin & J.M. Dawson, Oct 1973. *Phys. Fluids* 17, 987 (1974).
- PPG-164 Temporal Electrostatic Instabilities in Inhomogeneous Plasmas", Y.C. Lee & P.E. Kaw (Nov 1973) *Phys. Rev. Lett.* 32, 135 (1974).
- PPG-165 Nonlinear Schrödinger Equation Model of Oscillating Two-Stream Instability", G.J. Morales, Y.C. Lee & R.B. White, (Dec 73) *Phys. Rev. Lett.* 32, 457 (1974).
- PPG-166 Backscattering Decay Processes in Electron Beam-Plasma Interactions Including Ion Dynamics", B.H. Ripin, A.Y. Hong & B.H. Ripin, (Dec 1973) *Phys. Rev. Lett.* 32, 406 (1974).
- PPG-167 Conversion of Electromagnetic Waves to Electrostatic Waves in Inhomogeneous Plasmas", R. Stenzel, A.Y. Hong & H.C. Kim, (Dec 1973) *Phys. Rev. Lett.* 32, 451 (1974).
- PPG-168 Langmuir Wave Turbulence - Condensation & Collapse", Y.C. Lee, C.S. Liu & K. Nishikawa (Jan 1974) *Comm. Plas. Phys. & Cont. Phys.*
- PPG-169 The Consequences of Micropulsations on Geomagnetically Trapped Particles", R.N. Thorne (Jan 1974) *Rev. Space Sci.* 16, 443 (1974).
- PPG-170 Linear Wave Conversion in Inhomogeneous Plasmas", D.L. Kelley & A. Baños (Mar 1974) Thesis.
- PPG-171 The Cause of Storm After Effects in the Middle Latitude D-Region Ionosphere", W.N. Spjeldvik & R.N. Thorne (Mar 1974) *J. Atmos. & Terres. Phys.*
- PPG-172 Application of an Electromagnetic Particle Simulation Code to the Generation of Electromagnetic Radiation", A.T. Lin, J.M. Dawson & H. Oshida (Mar 1974) *Phys. Fluids* 17, 1995 (1974).
- PPG-173 The Ponderomotive Force Exerted on a Plasma by an Infrared Laser Beam", M. Murbic (Mar 1974) Dissertation.
- PPG-174 What we have Learned from the Magnetosphere", C.F. Kennel, Comm. Astrophys. & Space Phys. 6, 71 (1975).
- PPG-175 Observation of the Ponderomotive Force & Oscillating Two-Stream Instability", H.C. Kim, R. Stenzel & A.Y. Hong, (Apr 1974).
- PPG-176 Electron Beam Plasma Interaction Including Ion Dynamics", B.H. Ripin (June 1974) Thesis.
- PPG-177 Linear Conversion & Parametric Instabilities in a Non-Uniform Plasma", H.C. Kim, R. Stenzel & A.Y. Hong, (June 1974) Thesis.
- PPG-178 Stimulated Compton Scattering of Electromagnetic Waves in Plasma", A.T. Lin & J.M. Dawson (June 1974) *Phys. Fluids* 18, 201 (1975).
- PPG-179 Equatorial Spread F - Low Frequency Modes in a Collisional Plasma", M.N. Hudson (July 1974) Dissertation.
- PPG-180 Effect of the Ponderomotive Force in the Interaction of a Capacitor RF Field with a Nonuniform Plasma", G.J. Morales & Y.C. Lee (July 1974) *Phys. Rev. Lett.* 33, 1016 (1974).
- PPG-181 Deduction of Ionospheric Tidal Winds by Dynamo Simulation", J.P. Schieldge & S.V. Venkateswaran (Aug 1974) Thesis.
- PPG-182 Response of the Middle Latitude D-Region to Geomagnetic Storms", W. Spjeldvik & R.N. Thorne (Aug 1974) *J. Atmos. & Terres. Phys.*
- PPG-183 Production of Negative Ions & Generation of Intense Neutral Beams", A.Y. Hong, J.M. Dawson & W. Gekelman (Aug 1974) *App. Phys. Lett.* 25, 579 (1974).
- PPG-184 Development of Cavities & Trapping of RF Fields", H.C. Kim, R. Stenzel & A.Y. Hong (Aug 1974) *Phys. Rev. Lett.* 33, 886 (1974).
- PPG-185 Albuquerque Abstracts: Papers presented at Albuquerque Meeting of Am Phys Soc Div. of Plas. Phys., Oct 28-31, 1974).
- PPG-186 Localized Quasi-Stationary Plasma Modes in One, Two & Three Dimensions", J.Z. Wilcox & T.J. Wilcox (Sept 1974) *Phys. Rev. Lett.*
- PPG-187 Denouement of Jovian Radiation Belt Theory", F.V. Coroniti (Sept 1974) *Proc. Conf. Magnetospheres of Earth & Jupiter*, Frascati, Italy (May 28-June 1, 1974).
- PPG-188 Is Jupiter's Magnetosphere Like a Pulsar's or Earth's?", C.F. Kennel & F.V. Coroniti (Sept 1974) *Ibid.*
- PPG-189 Parametric Instability of the Sheath Plasma Resonance", R. Stenzel, H.C. Kim & A.Y. Hong (Sept 1974) *Bull. Am. Phys. Soc.* (Oct 1974), *Rad. Sci.* 10, 485 (1974).
- PPG-190 Effect of Localized Electric Fields on Evolution of the Velocity Distribution Function", G.J. Morales & Y.C. Lee (Sept 74) *Phys. Rev. Lett.* 33, 1534 (1974).
- PPG-191 Parametric Instabilities in Plasma", J.M. Dawson & A.T. Lin (Sept 1974) Thesis.
- PPG-192 Surmac - A Large Surface Magnetic Confinement Device", A.Y. Hong (Sept 1974) Thesis.
- PPG-193 Extraction of Energy from High Intensity Ion Beams", A.T. Forrester (Sept 1974) Presented at 2nd Symp. on Ion Sources and Formation of Ion Beams - Berkeley, Calif. (22-25 Oct 1974).
- PPG-194 A Large Quasistatic Magnetized Plasma for Wave Studies", W. Gekelman & R.L. Stenzel (Dec 1974) *Rev. Sci. Instr.* 46, 1586 (1975).
- PPG-195 Electrostatic Waves Near the Lower Hybrid Frequency", R. Stenzel & W. Gekelman (Dec 1974) *Phys. Rev. Lett.* 33, 2075 (1975).
- PPG-196 A Corrugated Micro-Cyclotron Frequency Direct Conversion System (Coax-Cyfer)", A.T. Forrester, J. Busnardo-Yeto & J.T. Crow (Oct 1974) *IEEE Transactions on Plasma Science*
- PPG-197 The Study of Comparative Magnetospheres: The Future of Space Physics", F.V. Coroniti & C.F. Kennel (Oct 1974) Presented at NASA Study Group On "Outlook for Space", Goddard Space Flight Center (Sept 10, 1974).
- PPG-198 Application of the Fokker-Planck Numerical Method to Anisotropic & Energy-Dependent Electron Precipitation", W. Spjeldvik (Oct 1974) Thesis.
- PPG-199 Self-focusing & Filamentation of Laser Light in Plasmas", Y.C. Lee, C.S. Liu, H.H. Chen & K. Nishikawa (Dec 1974) *Proceedings of IAEA Sixth Conf. on Plas. Phys.* Tokyo (Nov 1974).
- PPG-200 Stimulated Brillouin Backscatter in the Equatorial Electrojet", D.D. Barbosa & C.F. Kennel (Nov 1974) Thesis.
- PPG-201 The Electromagnetic Interchange Mode in a Partially Ionized Collisional Plasma", M.N. Hudson & C.F. Kennel (Dec 1974) *J. Plas. Phys.* 18, 135 (1975).
- PPG-202 The Collisional Drift Mode in a Partially Ionized Plasma", M.N. Hudson & C.F. Kennel (Dec 1974) *J. Plas. Phys.* 18, 135 (1975).
- PPG-203 High Density Constraint on the Entropy Instability", M.N. Hudson & C.F. Kennel (Dec 1974) Thesis.
- PPG-204 Excitation of Zero-Frequency Instabilities & Spatial Collapse of Beam Driven Plasma Waves", A.Y. Hong & B.H. Ripin (Dec 74) *Phys. Rev. Lett.* 34, 1499 (1975).
- PPG-205 A Recursive Numerical Method to Solve the Pure Pitch Angle Diffusion Equation, A Technical Report", W.N. Spjeldvik (Dec 1974) Thesis.
- PPG-206 The Equilibrium Radiation Belt Electron Pitch Angle Distribution & its Dependence on the Radial Diffusive Source", W.N. Spjeldvik (Jan 1975) Thesis.
- PPG-207 Optimization of Plasma Confinement with Permanent Magnet Multipoles", K.N. Leung, T.E. Saneck & A. Lam
- PPG-208 Annular Electron Transport & Lower-Hybrid Wave Damping", C. Chu, J.M. Dawson & H. Oshida (1975) *Phys. Fluids* 18, 1762 (1975).
- PPG-209 Plasma Confinement by a Picket-Fence", K.N. Leung, N. Hershkowitz & T. Rostovsky (Jan 1975) Thesis.
- PPG-210 Secular Mode Coupling & Nonlinear Drag: A Theory of Plasma Turbulence with Application to the Electron Beam-Plasma Instability", J. Drake (Feb 1975) Dissertation.
- PPG-211 Nonlinear Generation of Intense Localized Electric Fields in Plasmas", G.J. Morales & Y.C. Lee (Feb 1975) Thesis.
- PPG-212 Plasma Electron Heating by Injection of Low Energy Electrons", N. Hershkowitz & K.N. Leung (Feb 1975) Thesis.
- PPG-213 Ion Confinement by Electrostatic Potential Well in Magnetic Multiple Device", Y. Sakamura, B. Quon & A.Y. Hong (Feb 1975) *Phys. Lett.* 52A, 85 (1975).
- PPG-214 Scattering of Electromagnetic Waves into Plasma Oscillations via Plasma Particles", A.T. Lin & J.M. Dawson *Phys. Fluids* 18, 1542 (1975).
- PPG-215 Surface Magnetic Confinement", A.Y. Hong, Y. Sakamura & B. Quon (Mar 1975) *Phys. Rev. Lett.* 35, 1156 (1975).
- PPG-216 Experimental Facilities of Plasma Physics Research Laboratories at UCLA (Mar 1975) Thesis.
- PPG-217 A Modified Surmac Configuration", A.T. Forrester (Mar 1975) Thesis.
- PPG-218 Some New Ideas on Hot Hood Burners", J.M. Dawson & A.T. Lin (Apr 1975) Thesis.
- PPG-219 Plasma Heating at Frequencies Near the Lower Hybrid", C. Chu, J.M. Dawson & H. Oshida (Apr 1975) Thesis.
- PPG-220 Unstable High-Frequency Electrostatic Modes Induced by a Loss Cone Distribution of Electrons", B.D. Fried & D. Balmain
- PPG-221 ICRH Heating of a Magnetized Plasma and Associated Longitudinal Cooling", J. Busnardo-Yeto, J. Dawson, T. Sakamura & T. Lin
- PPG-222 Plasma Confinement by Localized Magnetic Cusps", K. Leung (May 1975) Thesis.
- PPG-223 The Generation of Spiky Turbulence", G.J. Morales & Y.C. Lee (May 1975) Thesis.
- PPG-224 A Multipole Antenna-Single Grid Extraction Ion Source", A.T. Forrester, A.T. Crow, N.A. Hassie & D.M. Goshel
- PPG-225 Q-Mechanism - by Robert K. Mottel" A book review by F.F. Chen.
- PPG-226 Flux Leakage Through a Low  $\beta$  Line Cusp", N. Hershkowitz, K.N. Leung, T. Rostovsky, W. Gekelman & R.L. Stenzel
- PPG-227 Localized Fields & Density Perturbations in Nonlinear Lower-Hybrid Waves", W. Gekelman & R.L. Stenzel *Phys. Rev. Lett.* 35, 1708 (1975).
- PPG-228 Effect of Mirroring on Convective Transport in Plasmas", T. Sakamura & J.M. Dawson. *Phys. Rev. Lett.* 36, 315 (1976).
- PPG-229 Parametric Excitation of Ion Fluctuations in the Relativistic Beam-Plasma Interaction", H. Schamel, Y.C. Lee, & G.J. Morales (June 1975) Thesis.
- PPG-230 UCLA Plasma Physics & Controlled Fusion Research, Presented for ERDA Program Review (July 18, 1975) B. Fried, J. Dawson, A. Hong, T. Forrester & F. Chen
- PPG-231 The Nonlinear Filamentation of Lower-Hybrid Cusps", G.J. Morales & Y.C. Lee. *Phys. Rev. Lett.* 35
- PPG-232 Fusion Reactor with Picket-Fence Walls", N. Hershkowitz & J. Dawson.
- PPG-233 Shock Formation in Large Amplitude Waves Due to Relativistic Electron Mass Variation", J.F. Drake, Y.C. Lee, & K. Nishikawa, *Phys. Rev. Lett.* 36, 196 (1976).
- PPG-234 Characteristics of a Large Volume RF Grid Discharge Plasma", B. Schumacher, N. Hershkowitz & K.R. Mackenzie
- PPG-235 Surface Magnetic Confinement: a Review", A.Y. Hong (July 1975) Reports.
- PPG-237 St. Petersburg Abstract: Papers to be presented at St. Petersburg Meeting of Am Phys Soc Div. of Plas. Phys. (10-13 Nov 1975)
- PPG-238 Parametric Instabilities in Strongly Relativistic, Plane Polarized Electromagnetic Waves", J. Drake, Y.C. Lee & N.L. Tsintsadze, *Phys. Rev. Lett.* 36, 31 (1976).

- PPG-239 "Detection of Brillouin Backscattering in Underdense Plasmas", J. Turechek & F. F. Chen.
- PPG-240 "Adiabatic Invariance & Charged Particle Confinement in a Geometric Mirror", T. K. Samec, Y. C. Lee & B. D. Fried.  
Phys Rev Let 35, 1763 (1975).
- PPG-241 "Magnetic Fields for Surface Containment of Plasmas", A. T. Forrester & J. Busnardo-Neto. J Ap Phys
- PPG-242 "Electrostatic Parametric Instabilities Arising from Relativistic Electron Mass Oscillations", A. T. Lin & N. L. Tsintsadze.  
Phys Fluids
- PPG-243 "Relativistic Nonlinear Plasma Waves in a Magnetic Field", C. F. Kennel & R. Pellat (Oct 1975) J Plas Phys
- PPG-244 "Flux Limits on Cosmic Acceleration by Strong Spherical Pulsar Waves", E. Asseo, C. F. Kennel & R. Pellat.  
Astron & Astrophys 44, No 1, 31 (1975).
- PPG-245 "On the Detection of Magnetospheric Radio Bursts from Uranus & Neptune", C. F. Kennel & J. E. Maggs (Nov 1975).
- PPG-246 "Parametric Instabilities with Finite Wavelength Pump", B. D. Fried, T. Ikemura, K. Nishikawa, & G. Schmidt.
- PPG-247 "The Scattering of Cosmic Rays by Magnetic Bubbles", R. F. Flewelling & F. V. Coroniti. Astrophys. Jnl. (May 1976)
- PPG-248 "Plasma Simulation on the CHI Microprocessor System", T. Kamimura, J. M. Dawson, B. Rosen, G. J. Culler, R. D. Levee & G. Bail.
- PPG-249 "Nonlinear Interactions of Focused Resonance Cone Fields with Plasmas", R. Stenzel & W. Cekelman.
- PPG-250 "The Effect of Pump Cutoff on Parametric Instabilities in an Inhomogeneous Plasma", J. VanDam & Y. C. Lee.
- PPG-251 "Parametric Excitation of Ion Density Fluctuations in the Relativistic Beam-Plasma Interaction", H. Schamel, Y. C. Lee, & G. J. Morales. (Jan 1976) Phys Fluids
- PPG-252 "The Spiky Turbulence Generated by a Propagating Electrostatic Wave of Finite Spatial Extent", G. J. Morales & Y. C. Lee.  
(Jan 1976) Phys Fluids
- PPG-253 "Departures from Theory of the Experimental Line Profile of Helium I  $4471.5\text{\AA}$ ", J. Turechek. (Jan 1976)
- PPG-254 "Maintenance of the Middle Latitude Nocturnal D-Layer by Energetic Electron Precipitation", W. Spjeldvik & R. Thorne  
(Jan 1976)
- PPG-255 "Free Boundaries for Plasmas in Surface Magnetic Field Configurations, B. D. Fried, J. W. VanDam & Y. C. Lee.  
Feb. 1976 Submitted to Phys. of Fluids.
- PPG-256 "Soliton-Like Structures in Plasmas, Y. C. Lee & G. J. Morales. Mar. 1976.
- PPG-257 "Magnetic Field Reconnection in a Collisionless Plasma, F. V. Coroniti & A. Eviatar. Mar. 1976 Submitted to Astrophys. Jnl.
- PPG-258 "The Earth's Magnetosphere, F. V. Coroniti. Accepted by Encyclopedia of Physics.
- PPG-259 "Lossy Radial Diffusion of Relativistic Jovian Electrons, D. D. Barbosa & F. V. Coroniti. Mar. 1976 Accepted by Jnl. Geophys. Res.
- PPG-260 "Relativistic Electrons & Whistlers in Jupiter's Magnetosphere, D. D. Barbosa & F. V. Coroniti. April 1976  
Submitted to Jnl. Geophys. Res.

REPRODUCIBILITY OF THIS  
ORIGINAL PAGE IS POOR