

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NITROUS OXIDE IN FRESH WATER SYSTEMS:  
AN ESTIMATE FOR THE YIELD OF ATMOSPHERIC N<sub>2</sub>O  
ASSOCIATED WITH DISPOSAL OF HUMAN WASTE

(NASA-CR-153978) NITROUS OXIDE IN FRESH  
WATER SYSTEMS: AN ESTIMATE FOR THE YIELD OF  
ATMOSPHERIC N<sub>2</sub>O ASSOCIATED WITH DISPOSAL OF  
HUMAN WASTE (Harvard Univ.) 40 p HC A03/MF  
A01

N77-30642

Unclas  
42029

CSCL 13B G3/45

W. A. Kaplan  
J. W. Elkins  
C. E. Kolb\*  
M. B. McElroy  
S. C. Wofsy  
A. P. Durán



Center for Earth and Planetary Physics, Harvard University,  
Cambridge, Mass. 02138

\* Primary affiliation: Center for Chemical and Environmental  
Physics, Aerodyne Research, Inc., Bedford Research Park, Bedford  
Mass. 01730

August 1977

## Abstract

The  $N_2O$  content of waters in the Potomac and Merrimack Rivers was measured on a number of occasions over the period April to July 1977. The concentrations of dissolved  $N_2O$  exceed those which would apply in equilibrium with air by factors ranging from about 46 in the Potomac to 1.2 in the Merrimack. Highest concentrations of dissolved  $N_2O$  were associated with sewage discharges from the vicinity of Washington, D. C. and analysis indicates a relatively high yield, 1.3 to 11%, for prompt conversion of waste nitrogen to  $N_2O$ . The yield could be even higher if bubbles originating in sediments should contain as little as 0.3%  $N_2O$ . Measurements of dissolved  $N_2O$  in fresh water ponds near Boston demonstrate that aquatic systems may provide both strong sources and sinks for atmospheric  $N_2O$ .

## 1. Introduction

Oxides of nitrogen play an important role in stratospheric chemistry. Ozone in the natural environment is thought to be removed mainly by catalytic reactions involving NO and NO<sub>2</sub> (Crutzen, 1970; Johnston, 1971). Nitric oxide is formed in the stratosphere by reaction of O(<sup>1</sup>D) with N<sub>2</sub>O (Crutzen, 1971; McElroy and McConnell, 1971; Nicolet and Vergison, 1971). Nitrous oxide is believed to originate as a by-product of microbiological activity at and near the earth's surface (Bates and Witherspoon, 1952; Bates and Hays, 1967), and concern has arisen in recent years that a variety of human activities might lead to a significant increase in the net global source of atmospheric N<sub>2</sub>O (Crutzen, 1974; McElroy, 1974; Pierotti and Rasmussen, 1976; Weiss and Craig, 1976).

Recent interest has focussed on perturbations associated with agriculture. Quantitative assessment is hampered however by major gaps in our understanding of the global nitrogen cycle, and by specific difficulties which arise in the description of the human influence.

One possible approach to the problem has been outlined by McElroy (1976) and by McElroy et al. (1977). Under conditions appropriate to North American agriculture, approximately 70% of nitrogen applied as chemical fertilizer may be recovered in crops. Approximately 6% of crop nitrogen is consumed directly by the human population. A much larger fraction, 67%, is fed

to animals. The remainder, 27%, is lost to a variety of factors including processing waste, spoilage, fire, insects, and plant pathogens. The human population obtains 30% of its nitrogen requirement directly from plants. The remainder is supplied indirectly through the animal food chain as meat, eggs, and dairy products.

Human waste per person in the developed world amounts to some  $5.4 \times 10^3$  grams of fixed nitrogen per year (National Academy of Sciences, 1972), equivalent to some  $1.14 \times 10^6$  tons N per year for the human population of the United States. The comparable figure for the world population is estimated at  $1.5 \times 10^7$  tons N per year, using data from a variety of sources (Deevey, 1970; National Academy of Sciences, 1977). Animals in the United States excrete an estimated  $3.7 \times 10^6$  tons N per year. By way of comparison, we might note that photochemical processes in the stratosphere represent a sink for  $N_2O$  of magnitude  $10^7$  tons N per year. It is clearly important that one should define the magnitude of the anthropogenic contribution to atmospheric  $N_2O$ .

This paper offers a preliminary account of results obtained in a series of measurements intended to explore the distribution of dissolved  $N_2O$  in a variety of aquatic systems. The paper includes results for two rather different river systems which receive human wastes, the Merrimack in Massachusetts and the Potomac near Washington, D. C. The Merrimack provides an example of a fast moving stream with few tributaries, draining

a large industrial area with a population of about 500,000 persons. The Potomac is relatively sluggish at sites sampled here, receiving wastes from a concentrated population in excess of 2 million. In addition, data are presented for a number of shallow fresh water ponds in Massachusetts.

## 2. Experimental procedures

Waters deeper than a meter were sampled by hydrocast using 1.7 liter Niskin bottles. Each bottle was equipped with a self-reversing thermometer triggered by the Niskin messenger unit. A waiting period between 7 and 10 minutes was observed at depth in order to allow the thermometer and sample bottle to equilibrate with their environment.

After retrieval of bottles, bubble-free water samples were removed by slow gravity feed to permit analyses for salinity, pH, dissolved oxygen and nitrous oxide. Dissolved oxygen, salinity and pH were measured immediately using a Yellow Springs Instruments Model 57 oxygen meter, a Yellow Springs Instruments Model 33 conductivity/salinity meter, and an Orion Model 407A pH meter respectively.

Samples for  $N_2O$  determination were transferred from the Niskin bottles to 300 ml BOD bottles. Microbial activity was inhibited by addition of 1 ml of a saturated  $HgCl_2$  solution. Samples were cooled to a temperature between 5 and 10°C until analyzed.

Measurements of  $N_2O$  were usually performed within 1 to 2 days of sample acquisition. Analyses were performed using a Perkin-Elmer 3920B electron capture gas chromatograph equipped with an 8' by 1/8" OD column of Porapak Q held at a temperature of 40°C. The  $^{63}Ni$  detector was operated at a temperature of 350°C with a standing current of  $3.5 \times 10^{-9}$

amps. This procedure allowed excellent separation of  $N_2O$  from possible contaminating gases as illustrated in Figure 1. We verified experimentally that samples treated with  $HgCl_2$  exhibit no significant gain or loss of  $N_2O$  during storage intervals employed here (Elkins et al., 1977).

Nitrous oxide was removed from water samples using the static gas partitioning method described by McAuliffe (1969, 1971). A 100 ml glass syringe with a locking input orifice (Precision Sampling Corporation) was flushed with  $N_2O$ -free helium. A 50 ml sample of water was displaced from the BOD bottle into the empty syringe again using helium. This procedure was devised in order to minimize contamination by outside air. Finally 50 ml of helium was introduced into the syringe and the syringe barrel was isolated, leaving a 1:1 volume ratio of helium and water. The syringe was placed on a wrist action shaker for 10 minutes to equilibrate the two phases. Gas from the syringe was then passed through the sample loop of the gas chromatograph and a 1 ml sample was injected for analysis. Residual helium was released and 50 ml of fresh helium were added. This process was repeated 2 or 3 times. A plot of  $\log N_2O$  peak area versus number of equilibrations yields a straight line. The y intercept of this line is a function of the initial  $N_2O$  concentration, as discussed by McAuliffe (1969, 1971).

Air samples obtained at the same time as the water samples were analyzed for atmospheric  $N_2O$ . Samples were acquired using



a 12V DC portable Speedaire compressor and were stored in 1 liter Whitey 304 stainless steel cylinders. Sample cylinder pressure delivered air to the chromatograph through a six port Carle Model 5518 valve equipped with a 1 ml sample loop. Column and detector parameters were identical to those used for water measurements. Two or three injections of synthetic air standards with known  $N_2O$  concentration (209, 326 or 420 ppb) were measured prior to analysis of each unknown. Standards were obtained from Scott Environmental Technology Incorporated, who certified their  $N_2O$  concentrations to an accuracy of  $\pm 2\%$ . An absolute pressure gauge was used to achieve equal sample loop pressures of unknowns and standards. Chromatographic peaks from both air and water samples were integrated using a Hewlett Packard 3380 A electronic integrator. Replicate analyses of air and water samples were used to establish overall precisions of 1% and 3% respectively.

### 3. Results

Figure 2 summarizes the  $N_2O$  data obtained for the Merrimack River on April 13, 1977 and June 29, 1977.

ly. Samples were acquired in the order indicated by the station numbers, 1 through 14. The initial measurements were taken when the flow rate in the river was high,  $466 \text{ m}^3 \text{ sec}^{-1}$ , corresponding to a linear flow speed of about  $50 \text{ cm sec}^{-1}$ . Flow rates were smaller during the June sampling period,  $131 \text{ m}^3 \text{ sec}^{-1}$ , corresponding to a flow speed of about  $14 \text{ cm sec}^{-1}$ . Stations 9 and 14 were occupied at approximately the same position and give a direct indication of temporal variability of dissolved  $N_2O$ .

Results for  $N_2O$  in Figure 2 are quoted as saturation ratios for dissolved gas with respect to the measured air concentration of  $294 \pm 3$  ppbv. Solubility data were taken from Markham and Kobe (1941). Fresh water sections of the river show at most small spatial variations in the concentration of dissolved  $N_2O$  for either measurement period. Other measured properties of the water, pH,  $O_2$  and temperature, are also spatially uniform. Concentrations of dissolved  $N_2O$  are spatially variable near the mouth of the river, downstream from station 4, consistent with a model in which fresh waters of high  $N_2O$  content are admixed with salt waters containing relatively little  $N_2O$ , as illustrated in Figure 3. Average properties of the river as measured for all 14 stations are summarized in Table 1.

Selected profiles for dissolved  $N_2O$  in the Potomac River are shown in Figures 4 and 5. Measurements were taken over the time period July 16, 1977 to July 23, 1977. Exceedingly high concentrations of  $N_2O$  were observed in waters near and downstream from the Alexandria and Blue Plains Wastewater Treatment Plants. As indicated in Figure 5, the water at station 12 was supersaturated in  $N_2O$  with respect to ambient air by a factor in excess of 40 on July 16, 1977. The concentration of  $N_2O$  in surface waters at station 11, closer to the Treatment Plants, was larger than that at station 12 by a factor of 1.2. The influence of the Treatment Plants extends approximately 25 km downstream, as may be seen from the data in Figure 5.

Lowest concentrations of  $N_2O$ , approximately 150% of saturation with respect to ambient air, were measured upstream of the Treatment Plants at stations 4, 5 and 6. The surface concentration of  $N_2O$  at station 16, near Quantico, Virginia, was somewhat higher than that at stations 4-6, about 200% of saturation with respect to ambient air. In mid-stream where the Potomac River enters Chesapeake Bay, the surface concentration of  $N_2O$  had a value equivalent to a supersaturation of 164% with respect to ambient air on July 18, 1977. At Harpers Ferry, 60 miles northwest of Washington, D. C. the surface concentration of  $N_2O$  was equivalent to 119% of saturation on July 17, 1977.

A significant, factor of 1.8, change in the concentration of dissolved  $N_2O$  was detected in waters near the Wastewater treatment plants between July 16 and July 23, 1977. The change

may be attributed to the influence of strong winds associated with the passage of a storm through the Washington area on July 21 and July 22, 1977, as indicated in Figure 6. The rate for gas exchange across an air-water interface is known to depend in a fairly sensitive fashion on wind speed (Broecker and Peng, 1974; Emerson, 1975), and we shall return to this matter later. The Blue Plains Plant releases approximately  $2 \times 10^4$  kg of N per day with trivial variation in outflow rate over the period covered by these measurements. The nitrogen source from the Alexandria Plant is approximately 10% of that from Blue Plains.

Dissolved oxygen concentrations showed substantial undersaturations ( $\sim 40\%$ ) near the bottom of the Potomac River. Surface waters were generally close to equilibrium, or slightly supersaturated, with respect to atmospheric oxygen.

Measurements for a variety of shallow ponds in the Boston area are summarized in Table 2. The data from Concord Town Forest are perhaps most intriguing, in that they illustrate the remarkable spatial heterogeneity of  $N_2O$  in fresh water systems. Measurements were carried out in two ponds separated by an embankment approximately 5 meters wide by about 0.5 meters high. The larger pond (Fairylane Pond) is approximately 60 meters long by 30 meters wide and is drained from its lower end through a small dam. It is choked with a variety of aquatic plants at its upper reach. Stations 1-4 are situated along the mid section of the pond beginning at its upper end. Surface concentrations of

$N_2O$  range from 650% of saturation with respect to ambient air in the upper region, to about 150% of saturation in clear waters near the center of the pond. Surface waters of the pond are super-saturated with respect to atmospheric  $O_2$  by about a factor of 2, reflecting the influence of photosynthesis. The vertical profile for  $N_2O$  at station 2 suggests that the source for  $N_2O$  may occur at depth, and that it may be largest in areas of highest biological productivity. In contrast the adjacent Concord bog which has no surface outlet, appears to act as a vigorous sink for  $N_2O$ . Waters there were undersaturated in  $N_2O$  with respect to ambient air by factors which varied between about 2 and 5. Water samples from this system were incubated for five days under a variety of conditions in the laboratory. Samples treated with  $HgCl_2$  showed at most small changes in the concentration of dissolved  $N_2O$  over this period. The concentration of  $N_2O$  in untreated samples dropped below detectable limits, about  $10 \text{ ng l}^{-1}$ . This behavior was observed for both light and dark incubations. Oxygen was also consumed in untreated samples. Its concentration was reduced to  $0.4 \text{ mg l}^{-1}$  and  $0.25 \text{ mg l}^{-1}$  for light and dark incubations respectively.

The data for Great Meadows National Wildlife Refuge were taken from a system of two ponds each about 200 meters square, separated by a dike. The upper pond, fed with treated sewage from the town of Concord, exhibited patches of floating algal growth apparently absent from the lower pond. Dissolved oxygen concentrations were relatively high in the upper pond. The

$N_2O$  concentration was relatively low however, though samples were not taken close to the sewage outflow region. The data suggest that the upper pond may be slightly undersaturated in  $N_2O$ , although the discrepancy could be attributed to experimental error.

#### 4. Discussion

The data described above may be used to place quantitative limits on the yield for  $N_2O$  associated with disposal of human waste in rivers. The results for the Potomac are especially relevant in this regard, in that the Blue Plains and Alexandria Treatment Plants may be considered point sources for anthropogenic nitrogen. The discharges have a major influence on the river, providing as much as 20% of the total stream flow over the period covered by our measurements. In addition, the temporal variability observed for  $N_2O$  allows us to make a direct estimate for the magnitude of the  $N_2O$  flux to the atmosphere.

Figure 7 illustrates the spatial distribution of  $N_2O$  as measured in the river near the treatment plants between July 16, 1977 and July 23, 1977. The  $N_2O$  plume associated with the plants is relatively extensive. Flow conditions in the stream were such that a period of approximately 40 days would be required for waters to traverse a distance of 25 km in the interval prior to July 23, 1977. The change in  $N_2O$  detected between July 16 and July 23 might be attributed to dilution associated with the higher flow rates which occurred on July 23 after the storm. The time and space scales exhibited in Figure 7 would not appear, however, to support this conjecture. It seems more probable that the variation in  $N_2O$  should be attributed to a change in the rate for gas exchange across the river surface associated with

passage of the storm on July 22. Future measurements may be expected to resolve possible ambiguities about the mechanism for temporal variations of  $N_2O$ .

Gas exchange across an air-water interface is frequently described using a simple diffusion model (Broecker and Peng, 1974), in which gas flux is given by

$$\phi = \frac{D}{Z} (C_L - C_E) ,$$

where  $D$  is a molecular diffusion coefficient,  $Z$  is an empirical parameter which defines the thickness of a hypothetical molecular boundary layer,  $C_L$  is the observed concentration of gas in the liquid phase, and  $C_E$  is the concentration of dissolved gas which should apply if the liquid were in equilibrium with the atmosphere. The parameter  $Z$  may be estimated using an empirical relationship between  $Z$  and wind speed developed by Emerson (1975) from studies of radon and  $CO_2$  in a variety of aquatic systems. Following his data, we derive a value for  $Z$  of approximately  $75 \mu$  for the period 0900 July 21 to 0600 July 23 (see Figure 6). The river has a mean depth of about 4 meters over the spatial domain of interest here. A film thickness of  $75 \mu$  would imply therefore a residence time for dissolved gas of about 2 days.

Other factors, in addition to those included by the simple wind-driven model, could contribute to the rate for gas exchange across the air-water interface in the Potomac. We might note that the river carries a significant traffic in moderate-sized power



boats. Four boats from the Wilson Line, for example, have an underwater cross section of approximately  $10 \text{ m}^2$ . These boats pass through the waters sampled here on an average of 6 times per day, displacing as much as 1% of the total water mass per day, at least 7% of the river encountered during the sampling interval, or 40% of the river during the 40 day residence time of water in the polluted area. It would appear that boat traffic could have a significant influence on gas exchange, at least during periods of relatively low wind speed and summer flow conditions.

The observed variation in  $\text{N}_2\text{O}$  may be used to place empirical limits on the rate for  $\text{N}_2\text{O}$  exchange. It appears that the  $\text{N}_2\text{O}$  content of the river downstream from Blue Plains decreased by approximately  $580 \times 10^3 \text{ g (N)}$  over the period July 16 to July 23. If we assume that the storm represented the dominant environmental change in this interval, and ignore sources of  $\text{N}_2\text{O}$  during the storm, we may place a lower bound on the  $\text{N}_2\text{O}$  release rate equivalent to  $290 \times 10^3 \text{ g N day}^{-1}$ , or 1.3% of the total nitrogen released from the wastewater treatment facilities over this period. Alternatively, we might suppose that the waters were entirely depleted of excess  $\text{N}_2\text{O}$  due to the influence of the storm, and that the concentration of disequilibrium  $\text{N}_2\text{O}$  observed on July 23, 6 hours after storm passage, indicates the recovery rate of the water system. This assumption would imply an upper limit to the prompt source strength for  $\text{N}_2\text{O}$  of magnitude  $2460 \times 10^3 \text{ g N day}^{-1}$ , or a yield of about 11% for conversion of wastewater N to  $\text{N}_2\text{O}$ . A source strength for  $\text{N}_2\text{O}$  of  $290 \times 10^3 \text{ g}$

$\text{N day}^{-1}$  could be rationalized on the basis of the diffusive exchange model if the film thickness,  $Z$ , were taken to have an average value of about  $200 \mu$  during the storm. A source strength as high as  $2460 \times 10^3 \text{ g N day}^{-1}$  would require a film thickness of about  $30 \mu$ .

We should caution that the efficiency for prompt conversion of waste nitrogen to  $\text{N}_2\text{O}$  could be even larger than values quoted here. The measured profiles for dissolved  $\text{N}_2\text{O}$  suggest that the gas may originate at or below the sediment-water interface. Bubbles were observed to reach the surface at station 11 at a rate of approximately  $3.5 \text{ ft}^{-2} \text{ min}^{-1}$ . These bubbles had an average diameter of about  $\frac{1}{4}$ ". If we assume that the bubbles originate in the sediment and are composed primarily of  $\text{CH}_4$ , reasonable assumptions based on dissolved  $\text{CH}_4$  data reported by Swinnerton, Linnenbom and Cheek (1967) (see also Hammond et al. (1975)), we may estimate a  $\text{CH}_4$  source strength of magnitude  $2 \times 10^8 \text{ g C day}^{-1}$  over the polluted stretch of the Potomac. Bubbles containing a ratio of  $\text{N}_2\text{O}$  to  $\text{CH}_4$  as small as 0.3% could provide an additional source for atmospheric  $\text{N}_2\text{O}$  which might exceed even the largest value derived above.

Flow rates in the Merrimack River are much higher than those measured for the Potomac. It is unlikely that transport of gas across the air-water interface in the Merrimack should be described by the thin film model given above for the relatively placid Potomac. A model outlined by Fortescue and Pearson (1967) provides an alternate description for gas exchange across the boundary of a turbulent liquid.

The flux is given in this case by

$$\phi = 1.46 \left( \frac{Du}{L} \right)^{1/2} (C_L - C_E)$$

where  $u$  is the root mean square speed for the river,  $L$  is the average depth of the liquid, and other symbols are as described above. The turbulent model suggests a lifetime for dissolved gas in the Merrimack of about a day for conditions encountered during the initial measurements on April 13, 1977, and is approximately twice as long for the measurements taken in June.

Flow rates in the Merrimack are such that the water would be expected to travel a distance in excess of 40 km over the one day residence time inferred here for dissolved  $N_2O$  in April, or 25 km for conditions encountered in June. We may note in this context that the Merrimack River is dammed near Lawrence and that it subsequently traverses a 1 km stretch of rapids located about 46 km upstream from the mouth of the river. One would not expect, therefore, any direct contribution to the excess concentrations of  $N_2O$  measured near the river mouth due to export of  $N_2O$  from regions of the river above Lawrence. Indeed, the residence time for dissolved  $N_2O$  estimated above, when combined with measured flow speeds for the river, suggest that the excess  $N_2O$  measured in April must be generated in close proximity to our sampling stations, in any event within an upstream displacement of 40 km. Supporting this view, we may note that the concentration of excess dissolved  $N_2O$  at Nashua, 80 km from the river

mouth is approximately half that measured in the estuary. Nitrous oxide should be a conservative property of the river over the 15 km interval sampled in April. A similar conclusion holds for the measurements made in June.

The measurements in April may be used to obtain a direct estimate for the quantity of  $N_2O$  released to the ocean by the Merrimack in this period. We find in this manner that the river exported a quantity of  $N_2O$  equivalent to  $2 \times 10^4$  g N day<sup>-1</sup> at that time.

It would seem reasonable to suppose that a major fraction of the excess  $N_2O$  measured in the Merrimack should reflect denitrification of N associated with diffuse sources of human waste. If the efficiency for conversion to  $N_2O$  were similar to that observed for the Potomac, the excess concentrations of  $N_2O$  in the Merrimack could be attributed to prompt denitrification of waste from a population of between 10,000 and 100,000 persons, with the lower value corresponding to a conversion efficiency of 11%. The river over its bottom 40 km drains an area occupied by a total population of about 180,000 people. The fraction of this population which exhausts its human waste directly into the Merrimack is somewhat uncertain. Moreover, a significant fraction of waste nitrogen entering the river is likely to be exported directly to the ocean as  $NO_3^-$  during the high flow conditions encountered during our measurements (U. W. Geological Survey, unpublished). Within the accuracy of the present analysis we may conclude that the April data for  $N_2O$  in the Merrimack are consistent with the range of yield factors

inferred earlier from the Potomac for prompt conversion of human waste nitrogen to  $N_2O$ , and a similar analysis may be made for the Merrimack data taken in June.

## 5. Concluding remarks

The data presented here allow us to estimate the magnitude of  $N_2O$  source associated with one component of the anthropogenically perturbed nitrogen cycle. If yields for  $N_2O$  inferred from the Potomac are representative, human wastes could provide a global source for  $N_2O$  of between  $2 \times 10^5$  and  $1.6 \times 10^6$  tons N per year, or between 2% and 15% of the rate at which the gas is consumed photochemically today in the stratosphere. We should note that these estimates reflect prompt sources for  $N_2O$ . They do not account for  $N_2O$  formed on a delayed basis following export of nitrogenous material. They omit also possible sources of atmospheric  $N_2O$  associated with bubble transport of gas from polluted sediments.

Extrapolation to global scale is hazardous, however, for several reasons. Results for the Potomac reflect sources of  $N_2O$  associated with one particular method for waste disposal at one particular time. Results from the Merrimack are consistent with a relatively high prompt yield for  $N_2O$ , but our understanding of the Merrimack system is not sufficiently complete to allow more definitive conclusions.

Animal wastes contain additional sources of nitrogen which exceed those for the human population of the developed world by more than a factor of 3. We are unable from the present study to offer any quantitative estimate for the corresponding yield of  $N_2O$  from this source. It could however be significant.

Our understanding of processes affecting atmospheric  $N_2O$  remains deficient. The limited data on rivers and ponds described here indicate that biological processes may offer both sources and sinks for the gas. Further observational data are clearly needed in order to quantify the overall budget of atmospheric  $N_2O$ .

### Acknowledgement

This work was supported by the National Science Foundation and the National Aeronautics and Space Administration under contracts #NSF-ENV76-24239 and #NAS-NASW-2952 respectively, to Harvard University.

We wish to acknowledge help offered by E. Jones of the Blue Plains Sewage Treatment Plant, W. White of the United States Geological Survey, L. Clark of the Environmental Protection Agency, and G. Goldberg from Aerodyne Research, Inc.

We are indebted to C. Martin of the University of Massachusetts Marine Biological Station and J. W. Elkins, Sr. for hospitality during critical phases of this study.



## References

- Bates, D. R. and P. B. Hays, 1967. "Atmospheric Nitrous Oxide", Planet. Space Sci. 15, 189.
- Bates, D. R. and A. E. Witherspoon, 1952. "The photo-chemistry of some minor constituents of the Earth's atmosphere", Mon. Not. Roy. Astron. Soc. 112, 101.
- Broecker, W. S. and T. H. Peng, 1974. "Gas exchange rates between air and sea", Tellus 26, 21.
- Crutzen, P. J., 1970. "The influence of nitrogen oxides on the atmospheric ozone content", Quart. J. Roy. Meteor. Soc. 96, 320.
- Crutzen, P. J., 1971. "Ozone production rates in an oxygen-hydrogen-nitrogen atmosphere", J. Geophys. Res. 76, 7311.
- Crutzen, P. J., 1974. "Estimates of possible variations in total ozone due to natural causes and human activities", Ambio 3, 201.
- Deevey, E. S., Jr., 1970. Mineral cycles in The Biosphere, W. H. Freeman and Co., San Francisco, 81.

- Elkins, J. W.; W. Kaplan, C. E. Kolb, M. B. McElroy, and S. C. Wofsy, 1977. "Determination of spring 1977 water column and marine air nitrous oxide concentrations in the New York Bight", Aerodyne Res. Inc., Tech. Rep. #ARI-RR-109, June 1977, Bedford, Mass.
- Emerson, S., 1975. "Gas exchange rates in small Canadian Shield lakes ", Limnology and Oceanography 20, 754.
- Fortescue, G. E. and J. R. A. Pearson, 1967. "On gas absorption into a turbulent liquid", Chem. Eng. Sci. 22, 1163.
- Hammond, D. E., H. J. Simpson, and G. Mathieu, 1975. "Methane and Radon-222 as tracers for mechanisms of exchange across the sediment-water interface in the Hudson River estuary, in marine chemistry in the coastal environment", Amer. Chem. Soc. Symp. Ser. 18, T. M. Church, Ed., American Chemical Society, 119.
- Johnston, H. S., 1971. "Reduction of stratospheric ozone by nitrogen oxide catalysts from supersonic transport exhaust", Science 173, 517.
- Markham, A. E. and K. A. Kobe, 1941. "Solubility of carbon dioxide and nitrous oxide in salt solution". J. Amer. Chem. Soc. 63, 449.

McAuliffe, C., 1969. "Determination of dissolved hydrocarbons in subsurface brines", Chem. Geol. 4, 225.

McAuliffe, C., 1971. "G. C. determination of solutes by multiple phase equilibrium", Chem. Tech. 46 (Jan. 71)

McElroy, M. B., 1974. Testimony presented to the Committee on Interstate and Foreign Commerce, U. S. House of Representatives, Washington D. C. 11 December 1974.

McElroy, M. B., 1975. "Chemical processes in the solar system: A kinetic perspective ", MTP International Review of Science, Series Two, Volume 9, 127. D. R. Herschback, Ed. Butterworths, London.

McElroy, M. B. and J. C. McConnell, 1971 "Nitrous oxide: A natural source of stratospheric NO", J. Atmos. Sci. 28, 1095.

McElroy, M. B., S. C. Wofsy and Y. L. Yung, 1977. "The nitrogen cycle: Perturbations due to man and their impact on atmospheric  $N_2O$  and  $O_3$ . Phil. Trans. Roy. Soc., B, 277, 159.

National Academy of Sciences, 1972. "Accumulation of nitrate", National Academy of Science, Washington, D. C. 106 pp.

National Academy of Science, 1977. "World food and nutrition study, the potential contributions of research", National Academy of Science, Washington, D. C., 192 pp.

Nicolet, M. and A. Vergison, 1971. "L'oxyde azoteux dans la stratosphere", Aeron. Acta. 90, 1.

Pierotti, D. and R. A. Rasmussen, 1976. "Combustion as a source of nitrous oxide in the atmosphere", Geophys. Res. Lett. 3, 615.

Swinnerton, J. W., V. J. Linnenbom, and C. H. Cheek, 1967. "Distribution of methane and carbon monoxide between atmosphere and natural waters", Envi. Sci. Tech. 3, 836.

Weiss, R. F. and H. Craig, 1976. "Production of atmospheric nitrous oxide by combustion", Geophys. Res. Lett. 3, 751.

Table I

## Merrimack River Station Average Data

April 13, 1977      Mean air concentration  $295 \pm 5$  ppbv  $N_2O$       Mean River Flow  $466 \text{ m}^3 \text{ sec}^{-1}$

Station #	Dissolved $N_2O$ ( $\mu\text{g m}^{-3}$ )	Excess $N_2O$ ( $\text{mg m}^{-3}$ )	pH (range)	$O_2$ (ml/l)	$O_2$ Saturation ratio (%) (range)	Salinity (‰) (range)	Total # Depths Sampled	Temp ( $^{\circ}\text{C}$ ) (range)	Remarks
1	0.65	0.14	8.0-8.2	6.31	64-97%	24.8 (16.1-28.0)	17.8 (5)	5.0 (3.4-6.8)	River plume to 3m with high $N_2O$ . River temp. $7.5-8.5^{\circ}\text{C}$ , ocean $3.5^{\circ}\text{C}$ .
2	0.684	0.15	7.97-8.16	5.57	55-83	24.5 (17.8-27.8)	10.7 (5)	5.0 (3.7-5.9)	
3	0.822	0.28	7.80-7.98	6.06	71-83	12.7 (11.2-16.8)	16.2 (5)	6.6 (6.4-6.7)	
4	0.876	0.34	7.4-7.7	6.01	69-81	7.5 (4.5-8.9)	16.2 (5)	7.6 (7.5-7.8)	
5	0.984	0.42	7.0-7.6	6.32	70-84	0	4.9 (4)	7.65 (7.6-7.75)	Incubated samples (1 week) grew in $0.07 \mu\text{g/l}$ $N_2O$ (average of 3).
6	1.05	0.49	6.9-7.0	6.37	71-82	0	6.1 (4)	7.73 (7.6-7.8)	
7	1.05	0.49	6.9-7.0	6.30	73-79	0	4.6 (3)	8.0 (7.9-8.1)	
8	1.07	0.52	6.8	6.45	65-73	0	9.8 (5)	8.3 (8.1-8.8)	
9	1.04	0.49	6.9	6.66	72-76	0	8.2 (5)	8.25 (8.2-8.3)	River plume above 3.5 m.
10	1.03	0.47	6.9	6.0	70-76	0	6.1 (4)	8.2 (8.1-8.4)	
11	0.63	0.09	8.1-8.2	5.2	64-73	28.0 (25-30)	11.6 (5)	4.2 (3.3-6.9)	

June 29, 1977      Mean air concentration  $291 \pm 4$  ppbv  $N_2O$       Mean River Flow  $131 \text{ m}^3 \text{ sec}^{-1}$

12	1.50	1.17	7.5-8.1	6.22	105-107%	0	3.0 (2)	24.0	Approx. same location as Station 9 of 4/13/77.
13	1.40	1.09	8.2-8.3	7.0	120	0	2.0 (2)	24.5	
14	1.44	1.12	7.8-8.0	6.87	115-120	0	8.5 (5)	24.2	

Table 2

## Concord Town Forest: Concord Bog (6/9/77)

Location	Time	$\text{N}_2\text{O}$ (mg m <sup>-3</sup> )	% $\text{N}_2\text{O}$ Saturation	°C	$\text{O}_2$ (ml l <sup>-1</sup> )	% $\text{O}_2$ Saturation	pH
I	6/9/77	0.152	36.5	15.5	6.9	98.9	6.5
II	6/9/77	0.214	51.4	15.5	6.9	98.9	6.5
III	6/16/77	0.205	54.3	19.9	6.2	103	6.7
IV	6/16/77	0.078	20	19.2	6.5	100	6.7
V	6/16/77	0.227	57.2	18.5	3.9	59.6	6.5

Table 2 (cont.)

Concord Town Forest: Fairyland Pond (6/9/77)

 $f_{\text{N}_2\text{O}} = 285 \text{ ppb}$ 

Location	$\text{N}_2\text{O} \text{ (mg m}^{-3}\text{)}$	$\% \text{ N}_2\text{O Saturation}$	$^{\circ}\text{C}$	$\text{O}_2 \text{ (ml l}^{-1}\text{)}$	$\% \text{ O}_2 \text{ Saturation}$	pH
1, < 8"	2.59	650	16.8	12.3	181	6.7
2, surface	0.958	225	14.9	11.8	167	6.6
2, 2.5'	1.249	292	14.7	13	183	6.6
3, surface	0.624	156	16.7	13.1	192	6.5
3, 2.5'	0.645	159	16.4	11.7	171	6.6
4, surface	0.753	181	15.5	12.2	175	6.8
4, 2'	0.775	197	17.2	11.9	177	6.8

Table 2 (cont.)

## Great Meadow National Wildlife Refuge

6/9/77

Station	N <sub>2</sub> O (mg m <sup>-3</sup> )	% N <sub>2</sub> O Saturation	°C	O <sub>2</sub> (ml l <sup>-1</sup> )	% O <sub>2</sub> Saturation	pH
Algal mat I	0.367	99.1	19.0	12.6	194	6.9
Algal mat II	0.387	104	18.9	18.0	277	7.4
III	0.353	95.3	19	8.4	129	6.6
IV	0.338	90.7	18.8	10.1	149	6.8
V	0.348	93.3	18.8	10.1	149	6.8



### Figure Captions

Figure 1. Chromatograms showing response of the ECD to  $O_2$  and  $N_2O$ . Chromatogram A is an air sample containing  $295 \pm 3$  ppbv  $N_2O$ . Chromatograms B and C are successive extractions of  $N_2O$  from a water sample using the technique of McAuliffe (1969). This water sample contained  $1.404 \mu g N_2O/liter$ .

Figure 2. Merrimack River  $N_2O$  profiles are shown for April 13, 1977 and June 29, 1977. Data are given as saturation ratios of  $N_2O$  with respect to air containing  $294 \pm 3$  ppbv  $N_2O$ . The distance between stations 3 and 9 is 12 km. Stations 14 and 9 are located at the same site. The extent of tidal influence in the river is approximately 10 km from the mouth of the river.


Figure 3. Dissolved  $N_2O$  is plotted against salinity in a 12 km segment of the Merrimack River for the data from April 13, 1977.


Figure 4. Locations of sampling stations 1-16 are illustrated for the Potomac River between Washington, D. C. and Quantico, Va.

Figure 5. Selected depth profiles of  $N_2O$  in the Potomac are presented in terms of supersaturation with respect to ambient air. Station numbers refer to sites given in Figure 4.

Figure 6. Surface wind data from Washington National Airport are given for the period July 21-23, 1977. The passage of a cold front is indicated by the arrow. Effective film thicknesses  $Z$  (Broecker and Peng, 1974) are shown which correspond to the various wind speeds. If the wind data were extrapolated to 10 cm elevation (cf. Emerson, 1975), the film thicknesses would be increased by about a factor of 2.

Figure 7. Mean concentrations of  $N_2O$  ( $mg\ m^{-3}$ ) are plotted versus distance along the Potomac from the D. C. boundary (Chain Bridge), for various times during the week 16-23 July 1977. Stations where only surface data were acquired are marked by an asterisk (\*). The location of the Blue Plains and Alexandria sewage discharges are indicated by the arrows.





2

