


CR 151616



"AS-BUILT" DESIGN SPECIFICATION JAN 24 1978  
FOR THE  
DIGITAL DERIVATION OF  
DAILY AND MONTHLY DATA BASES  
FROM SYNOPTIC OBSERVATIONS  
OF TEMPERATURE AND PRECIPITATION FOR  
THE PEOPLE'S REPUBLIC OF CHINA

Job Order 73-763  
AD 63-1347-3763-54

Prepared By  
Lockheed Electronics Company, Inc.  
Systems and Services Division  
Houston, Texas  
Contract NAS 9-15200

For  
EARTH OBSERVATIONS DIVISION  
SPACE AND LIFE SCIENCES DIRECTORATE



*National Aeronautics and Space Administration*  
**LYNDON B. JOHNSON SPACE CENTER**  
*Houston, Texas*

December 1977

LEC- 11680

(NASA-CR-151616) AS-BUILT DESIGN N78-18624  
SPECIFICATION FOR THE DIGITAL DERIVATION OF  
DAILY AND MONTHLY DATA BASES FROM SYNOPTIC  
OBSERVATIONS OF TEMPERATURE AND  
PRECIPITATION FOR THE (Lockheed Electronics G3/47 05431  
HC A06/HF A01  
Unclas

DOCUMENTATION DISTRIBUTION LIST

System: CCEA/Yield Estimation System (YES)

Title: "As-Built" Design Specification For The Digital Derivation of  
Daily and Monthly Data Bases From Synoptic Observations of  
Temperature and Precipitation For The People's Republic of China

Job Order: 74-963

Responsible Organization: Applications Software Section, 333-6311

JSC

SF/L. F. Childs  
SF2/R. L. Eason  
SF4/V. S. Whitehead  
TF3/T. E. Barnett  
R. G. Stuff

NOAA/J. Hill

T. Phillips  
N. Strommen  
S. Laduc  
C. Sakamoto

USDA/J. Rogers

LEC

C09/D. E. Phinney  
B. L. Carroll  
C42/J. M. Everette  
J. A. Wilkinson  
B. H. Jeun  
R. L. Davenport  
G. L. Barger

Job Order File

Technical Publications (5) ✓

Data Research and Control (3)

[ ]

"AS-BUILT" DESIGN SPECIFICATION  
FOR THE  
DIGITAL DERIVATION OF  
DAILY AND MONTHLY DATA BASES  
FROM SYNOPTIC OBSERVATIONS  
OF TEMPERATURE AND PRECIPITATION FOR  
THE PEOPLE'S REPUBLIC OF CHINA

Job Order 73-763

AD 63-1347-3763-54

Prepared By

B. H. Jeun

G. L. Barger

APPROVED BY

*for* *James A. Wilkinson*  
Philip L. Krumm, Supervisor  
Applications Software Section

*B. L. Carroll*  
B. L. Carroll, Manager  
LACIE Development and Evaluation Department

Prepared By

Lockheed Electronics Company, Inc.

For

Earth Observations Division

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
LYNDON B. JOHNSON SPACE CENTER  
HOUSTON, TEXAS

December 1977

LEC- 11680

## CONTENTS

| SECTION                                           | PAGE |
|---------------------------------------------------|------|
| 1. SCOPE.....                                     | 1-1  |
| 2. APPLICABLE DOCUMENTS.....                      | 2-1  |
| 3. SYSTEM DESCRIPTION.....                        | 3-1  |
| 3.1 HARDWARE DESCRIPTION.....                     | 3-4  |
| 3.2 SOFTWARE DESCRIPTION.....                     | 3-4  |
| 3.2.1 THE DERIVATION OF INPUT VARIABLES.....      | 3-4  |
| 3.2.2 THE COMPUTATION ALGORITHMS.....             | 3-5  |
| 3.2.3 THE EVALUATION OF OUTPUT VALIDITY.....      | 3-7  |
| 4. OPERATIONS.....                                | 4-1  |
| 4.1 THE LEAST SQUARES INTERPOLATION.....          | 4-1  |
| 4.2 THE MODIFIED MAX-MIN PROGRAM.....             | 4-1  |
| 4.3 THE MONTHLY DATA BASES ALGORITHM.....         | 4-2  |
| 4.4 THE ANALYSIS OF VARIANCE PROCEDURE.....       | 4-3  |
| 5. ACKNOWLEDGEMENT.....                           | 5-1  |
| APPENDIX                                          |      |
| A. THE LEAST SQUARES INTERPOLATION ALGORITHM..... | A-1  |
| B. THE MONTHLY DATA BASES ALGORITHM.....          | B-1  |
| C. THE ANALYSIS OF VARIANCE PROCEDURE.....        | C-1  |

## 1. SCOPE

As an integral part of the Large Area Crop Inventory Experiment (LACIE) which is sponsored by the Department of Agriculture (USDA), the National Oceanic and Atmospheric Administration (NOAA) and the National Aeronautics and Space Administration (NASA), continued building of daily and monthly data bases to test various advanced wheat yield models is an active objective and goal. The People's Republic of China (PRC) is one of the largest agricultural countries in the world. Winter and spring wheat is one of its major crops; therefore, a knowledge of China's crop production is very important to the LACIE Project. The China data base consists of meteorological information in the form of daily precipitation and maximum and minimum temperatures and their monthly totals and averages, respectively, for about 95 weather stations covering the entire winter and spring wheat production area of China over the period 1965 through 1975. The daily and monthly meteorological information was built up from synoptic (4 or 8 times daily) weather observations available on magnetic tape from the National Climatic Center (N.C.C.) in Asheville, North Carolina.

There are three major sets of computer algorithms for building the China data base, that is: 1) the least squares interpolation; 2) modified version of the max-min program; and 3) the AOV procedure.

The least squares interpolation algorithm is used to generate input variables; the modified version of the max-min program is used to generate daily and monthly precipitation and max-min temperatures; the analysis of variance algorithm is used to carry out a simple statistical hypothesis test on the digital estimated data. The China data base is considered complete and unique. It may be the only one of this kind of meteorological data existing outside, or inside the People's Republic of China.

## 2. APPLICABLE DOCUMENTS

Action Documentation (AD) 63-1347-3763-54, China Data Base,  
June 28, 1977

AD 63-1557-4963-22, YES Data Base Maintenance, Oct 17, 1977.

Job Order 73-763

"As-Built" Design Specification for Historical Daily Data Bases  
for Testing Advanced Models (JSC-12891; LEC-10572), April 1977.

### 3. System Description

#### CHINA SYNOPTIC DATA SOURCE

A total of 32 tapes were obtained by the Center for Climatic and Environmental Assessment (CCEA) from the National Climatic Center (NCC) in Asheville, North Carolina, covering World Meteorological Organization (WMO) blocks, 53, 54, 57 and 58 in China for the years 1965 through 1975.

Data were contained in 3 tapes per year except for 1968 which has only 2 tapes. These tapes, containing synoptic data (i.e., 4-8 observations per day) for all reporting weather stations within these blocks, are recorded in NCC deck format 9685.

#### CHINA DATA BASE DESIGN

A total of 95 WMO weather stations providing uniform coverage of the entire winter and spring wheat production area of China (see map on page 3.3), was selected on the basis of relative reliability and consistency of record. Daily precipitation and max-min temperatures are the major meteorological variables generated for the 95 representative weather stations covering the entire winter and spring wheat production area of China over the period from 1965 to 1975. Daily data were stored in the magnetic tapes in the following format:

| <u>Character String<br/>(per record)</u> | <u>Content</u>             |
|------------------------------------------|----------------------------|
| 1 - 6                                    | Station number             |
| 7 - 8                                    | Year                       |
| 9 - 10                                   | Month                      |
| 11 - 12                                  | Day                        |
| 13                                       | Blank                      |
| 14 - 16                                  | Daily Maximum temperature* |
| 17 - 19                                  | Daily Minimum temperature* |
| 20 - 22                                  | Blank                      |
| 23 - 26                                  | Precipitation **           |

\* Temperatures are in whole degrees Fahrenheit with a leading minus sign if negative; missing temperature observations are denoted by 999.


\*\* Precipitation is in hundredths of inches with no decimal points; missing precipitation is indicated by 9999. The above format is known as the Modified Kansas Format (as opposed to NCC Format).

Monthly data were generated by a separate computer program; output is available in hard copies containing the following information:

1. Station
2. Year
3. Month
4. Monthly average maximum temperature
5. Monthly average minimum temperature
6. Monthly total precipitation
7. Total number of observations
8. Number of missing observations
9. Total annual precipitation

This section describes the system software, Appendices A, B, and C contain detail flowcharts for subroutines used.

# THE WMO WEATHER STATIONS IN CHINA



### 3.1 HARDWARE DESCRIPTION

All computer algorithms described in this document were written in standard FORTRAN language and will operate on any IMB 360/370 system with FORTRAN compiler and with 1600 BPI magnetic drive. They will also operate on mini-computer systems such as PDP 11/50 with FORTRAN compiler and 800 BPI with minor change.

### 3.2 SOFTWARE DESCRIPTION

#### 3.2.1 THE DERIVATION OF INPUT VARIABLES

The least squares interpolation computer algorithm was designed to generate a complete set of monthly temperature ranges. The monthly average daily temperature ranges for each weather station are needed as input variables to the max-min program to generate the daily maximum and minimum temperatures for each month. The program consists of one main program and seven subroutines i.e., SUM, COR, LSTSQ, MEQSOL, GRAPH, STANDV AND MULTR, and one IBM built-in library routine, PLTSCT, which is used to plot the scatter diagram for the input data; the above subroutines can be found in computer literature and journals.

#### SUBROUTINE SUM

This subroutine is used to calculate the total of input observations such as  $X$ ,  $X^2$  and  $Y$  and  $Y^2$ .

#### SUBROUTINE COR

This subroutine is used to calculate the correlation coefficient ( $r$ ) of the input data, the correlation coefficient is used as a criterion to determine whether a straight line or a 2nd degree curve fits a given set of data; that is, if  $|r| > 0.7$ , a straight line should be fitted; otherwise, a higher degree curve is required.

#### SUBROUTINE LSTSQ

This subroutine is used to determine the best fitting equation by establishing the proper matrix equation  $AX = B$

For example, if a linear least squares equation is needed then,

$$A = \begin{pmatrix} N & \sum x_i \\ \sum x_i & \sum x_i^2 \end{pmatrix} ; \quad X = \begin{pmatrix} C_1 \\ C_0 \end{pmatrix} ; \quad B = \begin{pmatrix} \sum y_i \\ \sum x_i y_i \end{pmatrix}$$

where  $C_1$  are the coefficients of the least squares equation. For higher order least squares polynomials, the elements of different matrices will be changed automatically by this subroutine.

#### SUBROUTINE MEQSOL

This subroutine is used to solve the matrix equation  $AX = B$  which has been determined by the subroutine LSTSQ.

ORIGINAL PAGE IS  
OF POOR QUALITY

#### SUBROUTINE GRAPH

This subroutine is used to plot the least squares equation in the X - Y Plane.

#### SUBROUTINE MULTR

This subroutine is used to calculate the multiple correlation coefficient (RMULT) which serves as the criterion to determine if a 2nd degree or higher order least squares equation is needed.

#### SUBROUTINE STANDV

This subroutine is used to calculate the standard deviation of a given set of data points; this subroutine is needed in the subroutine MULTR.

### 3.2.2. THE COMPUTATION ALGORITHMS

There are two major computer algorithms for building the China data bases. One for generating the daily data and the other for generating monthly data. All programs are written in standard FORTRAN language.

The computer program for daily data is a modified version of the max-min program which is basic to the non-linear statistical model developed by a research team at Kansas State University led by Dr. Arlin Feyerherm, Professor of Statistics. Complete information about the max-min program can be found in the "As-Built Design Specification for Historical Daily Data Bases for Testing Advanced Models", prepared by Buddy H. Jeun and K. Williams and published by the National Aeronautics and Space Administration (NASA), Lyndon B. Johnson Space Center, Houston, Texas 77058, April 1977.

In generating daily precipitation for the PRC, the precipitation amounts reported at synoptic hours 0600, 0900, 1200, and 1800 (GMT) were totaled. (When a similar application was carried out for the USSR only 0600 and 1800 (GMT) observations were summed). The reporting conventions differ region-to-region and must be verified locally by summing the data by day, month, and year. The computer algorithm that generates the monthly meteorological data, i.e., average maximum and minimum temperature and total precipitation, consists of one main routine and five subroutines, which are SORTMD, DMEAN, TOTAL, ANP and PBAR, and one built-in function, i.e., EXTERNAL SIGN, which is used to distinguish if the observation is blank or zero, since blank denotes a missing observation for maximum and minimum temperatures and for precipitation, the missing observation is denoted by 9999.

#### SUBROUTINE SORTMD

This subroutine is used to sort the daily observations by month; in doing so, for every station, the maximum and minimum of temperatures and precipitation for every month are sorted out.

#### SUBROUTINE DMEAN

This subroutine is used to calculate the average maximum and minimum temperatures for each month. At the same time, it also keeps track of the number of missing temperatures by the argument ICOUNT and KOUNT. In this subroutine, ICOUNT represents the number of missing observations for the maximum temperature and KOUNT represents the number of missing observations for the minimum temperature.

#### SUBROUTINE TOTAL

This subroutine is used to calculate the sum of a given set of observations. Input arguments are N, DZ and output arguments are SUM and KNT, where N is the number of observations. DZ is an array of N observations. SUM is the total value of N observations, KNT is the number of missing observations.

#### SUBROUTINE PBAR

This subroutine is used to calculate the monthly total of precipitation to each station. Input argument PC is an array of N observations of precipitation. Output argument PAV is the monthly total of precipitation.

#### SUBROUTINE ANP

This subroutine is used to calculate the annual precipitation to each weather station. Input arguments are STATN, YEAR, N and VP; output arguments are BP and the missing observations are represented by ICOUNT.

#### 3.2.3. THE EVALUATION OF OUTPUT VALIDITY

ORIGINAL PAGE IS  
OF POOR QUALITY


The analysis of variance.

This analysis of variance computer algorithm was designed to test the reliability of the monthly data derived from the modified version of the max-min program. A few weather stations were selected for a study comparing the estimated means with PRC published means obtained from the USAF through the National Climatic Center at Asheville, North Carolina and other reliable records from the Hong Kong Royal Observatory and Chinese educational institutions obtained through personal correspondence. The six stations for which comparative monthly temperature data are available are shown on the map on page 3-9. They are confined to the more humid eastern portions of China but they do give a good temperature range north-south across the major wheat-producing area. The tables and graphs (page 3-10 through 3-21) include individual monthly average temperatures (EST) as summarized from the computed daily

max-min values alongside the most reliable check data (PUB) available. Whether the check data are true internal published amounts or "usually reliable" figures is not known. An analysis of variance (page 3-22 through 3-24) indicates that the estimated mean monthly temperatures are not significantly different from the available check data. The null hypothesis of the analysis of variance is tenable.

For precipitation, only four stations were supplied for comparison (pages 3-25 through 3-29). The WMO station numbers are the same as in daily source tapes but it is suspected the pairs of records compared may come from different, but proximate, rain gauges. Nevertheless, the monthly and annual precipitation totals compare favorably.

# SELECTED WMO WEATHER STATIONS



ORIGINAL PAGE IS  
OF POOR QUALITY.

PEOPLE'S REPUBLIC OF CHINA  
中华人民共和国


## MONTHLY AVERAGE TEMPERATURE

CHI-NAN

#54823

| MONTH | 1965  |      | 1966  |      | 1967  |      | 1968  |      | 1969  |      | 1970  |      |
|-------|-------|------|-------|------|-------|------|-------|------|-------|------|-------|------|
|       | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  |
| JAN   | -0.38 | -0.1 | -0.63 | 0.3  | -2.70 | -1.1 | -3.27 | -1.3 | -4.66 | -3.4 | -3.86 | -1.8 |
| FEB   | 2.70  | 2.7  | 3.56  | 4.2  | -0.18 | 0.4  | -3.16 | -1.3 | -4.86 | -3.2 | 0.35  | 2.7  |
| MAR   | 8.20  | 7.8  | 6.54  | 7.6  | 7.25  | 8.6  | 7.43  | 9.4  | 3.37  | 5.2  | 3.21  | 5.2  |
| APR   | 11.88 | 12.7 | 13.84 | 15.1 | 14.50 | 16.1 | 14.84 | 16.7 | 11.81 | 14.0 | 13.04 | 15.1 |
| MAY   | 22.48 | 22.9 | 21.88 | 22.7 | 23.58 | 25.5 | 21.29 | 22.3 | 19.68 | 21.4 | 19.42 | 21.5 |
| JUN   | 26.00 | 27.1 | 24.50 | 26.4 | 25.60 | 27.3 | 26.83 | 28.4 | 23.76 | 26.0 | 22.80 | 24.8 |
| JUL   | 27.23 | 27.7 | 27.15 | 27.8 | 27.48 | 27.8 | 25.56 | 28.7 | 25.60 | 27.5 | 24.82 | 26.4 |
| AUG   | 24.89 | 25.5 | 27.15 | 28.1 | 27.48 | 28.1 | 25.56 | 26.5 | 25.60 | 26.9 | 24.82 | 25.9 |
| SEP   | 23.50 | 23.3 | 19.47 | 20.6 | 18.98 | 20.2 | 21.46 | 22.8 | 19.92 | 21.2 | 19.42 | 21.2 |
| OCT   | 20.12 | 17.6 | 14.79 | 15.9 | 14.66 | 16.2 | 12.69 | 14.2 | 14.58 | 16.4 | 14.31 | 16.3 |
| NOV   | 9.14  | 9.0  | 8.33  | 9.4  | 4.46  | 6.1  | 7.27  | 8.8  | 4.84  | 6.7  | 5.50  | 7.2  |
| DEC   | 1.20  | 1.2  | -2.10 | -1.0 | -4.16 | -3.0 | 1.42  | 3.2  | -1.66 | 0.3  | 0.46  | 1.5  |

ORIGINAL PAGE IS  
OF POOR QUALITY




MONTHLY AVERAGE TEMPERATURE

TA-LIEN

#54662

| MONTH | 1965  |      | 1966  |      | 1967  |      | 1968  |      | 1969  |      | 1970  |      |
|-------|-------|------|-------|------|-------|------|-------|------|-------|------|-------|------|
|       | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  |
| JAN   | -5.60 | -4.7 | -6.20 | -4.5 | -6.80 | -5.2 | -5.16 | -5.8 | -8.90 | -7.1 | -7.35 | -6.0 |
| FEB   | -2.96 | -2.3 | -3.58 | -1.6 | -4.95 | -4.0 | -5.46 | -6.6 | -9.28 | -7.4 | -4.17 | -2.1 |
| MAR   | 1.76  | 2.3  | -0.33 | 1.2  | 1.52  | 3.4  | 4.29  | 3.1  | -1.82 | -0.2 | -2.68 | -0.8 |
| APR   | 7.03  | 7.7  | 5.87  | 7.9  | 7.00  | 9.0  | 10.81 | 9.8  | 6.35  | 8.4  | 7.93  | 9.7  |
| MAY   | 14.83 | 15.7 | 14.26 | 15.9 | 14.29 | 16.6 | 15.10 | 14.0 | 13.02 | 15.0 | 14.31 | 16.5 |
| JUN   | 19.76 | 20.1 | 17.48 | 18.0 | 17.40 | 19.4 | 20.34 | 19.2 | 17.44 | 19.2 | 16.25 | 18.3 |
| JUL   | 22.00 | 23.1 | 21.71 | 22.6 | 21.90 | 23.5 | 22.26 | 23.4 | 21.37 | 22.5 | 20.67 | 21.6 |
| AUG   | 22.13 | 22.6 | 23.34 | 24.6 | 24.00 | 25.3 | 22.60 | 24.1 | 22.34 | 24.0 | 22.33 | 23.8 |
| SEP   | 20.69 | 20.7 | 17.27 | 18.5 | 17.11 | 18.7 | 19.50 | 20.7 | 18.75 | 19.8 | 18.50 | 20.1 |
| OCT   | 16.19 | 15.5 | 13.23 | 14.2 | 13.36 | 14.7 | 10.68 | 12.3 | 11.83 | 14.0 | 13.03 | 14.3 |
| NOV   | 5.53  | 6.0  | 5.47  | 6.9  | 2.38  | 3.9  | 5.72  | 7.1  | 3.70  | 4.8  | 4.22  | 5.9  |
| DEC   | -2.77 | -2.1 | -5.50 | -4.0 | -6.80 | -5.4 | -1.65 | 0.4  | -3.71 | -1.9 | -2.25 | -0.6 |

ORIGINAL PAGE IS  
OF POOR QUALITY




## MONTHLY AVERAGE TEMPERATURE

NANKING

#58238

| MONTH | 1965  |      | 1966  |      | 1967  |      | 1968  |      | 1969  |      | 1970  |      |
|-------|-------|------|-------|------|-------|------|-------|------|-------|------|-------|------|
|       | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  |
| JAN   | 3.41  | 3.8  | 2.36  | 2.9  | 0.96  | 1.4  | 0.35  | 1.5  | -0.15 | 1.3  | -0.68 | 0.8  |
| FEB   | 4.37  | 5.1  | 4.90  | 5.9  | 2.88  | 3.8  | 0.76  | 1.8  | -1.56 | 0.3  | 4.02  | 5.2  |
| MAR   | 7.58  | 7.9  | 8.81  | 9.7  | 8.04  | 8.8  | 8.73  | 9.7  | 5.57  | 7.1  | 4.60  | 5.9  |
| APR   | 12.10 | 12.7 | 13.84 | 14.6 | 13.33 | 14.5 | 16.00 | 15.1 | 13.69 | 14.6 | 12.89 | 14.0 |
| MAY   | 20.31 | 20.9 | 18.61 | 19.8 | 20.04 | 20.7 | 20.78 | 19.8 | 20.10 | 21.3 | 18.63 | 20.0 |
| JUN   | 23.41 | 23.6 | 23.51 | 24.8 | 25.51 | 26.0 | 25.67 | 24.9 | 22.10 | 23.7 | 22.01 | 23.3 |
| JUL   | 27.99 | 28.4 | 27.86 | 28.9 | 27.36 | 28.2 | 27.20 | 26.8 | 25.12 | 26.6 | 26.55 | 27.2 |
| AUG   | 25.26 | 25.9 | 28.56 | 29.6 | 29.49 | 30.5 | 26.43 | 27.4 | 26.98 | 28.4 | 27.39 | 28.2 |
| SEP   | 21.52 | 22.0 | 19.96 | 21.1 | 21.73 | 23.0 | 21.98 | 23.1 | 22.96 | 24.1 | 21.39 | 22.6 |
| OCT   | 16.91 | 17.0 | 15.76 | 17.0 | 14.95 | 16.4 | 14.55 | 15.4 | 16.21 | 17.1 | 16.15 | 17.4 |
| NOV   | 11.31 | 11.9 | 9.18  | 10.4 | 8.31  | 9.8  | 11.22 | 11.9 | 6.79  | 8.3  | 8.86  | 10.0 |
| DEC   | 2.21  | 3.4  | 2.26  | 3.0  | -0.56 | -0.1 | 5.92  | 7.5  | 2.00  | 3.2  | 3.32  | 4.7  |




MONTHLY AVERAGE TEMPERATURE

CHANG-CHUN

#54161

| MONTH | 1965   |       | 1966   |       | 1967   |       | 1968   |       | 1969   |       | 1970   |       |
|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|-------|
|       | EST    | PUB   | EST    | PUB   | EST    | PUB   | EST    | PUB   | EST    | PUB   | EST    | PUB   |
| JAN   | -15.49 | -15.9 | -16.87 | -16.9 | -14.27 | -14.4 | -18.43 | -17.8 | -17.72 | -17.8 | -20.73 | -20.5 |
| FEB   | -13.61 | -14.0 | -10.24 | -9.8  | -10.92 | -11.2 | -16.43 | -16.4 | -15.95 | -15.8 | -14.20 | -13.4 |
| MAR   | -4.25  | -4.5  | -5.22  | -4.8  | -1.72  | -1.3  | -2.00  | -1.8  | -4.36  | -4.7  | -8.15  | -8.1  |
| APR   | 5.22   | 4.7   | 6.26   | 6.0   | 7.48   | 7.6   | 10.15  | 9.5   | 6.37   | 6.0   | 8.10   | 7.9   |
| MAY   | 15.29  | 14.9  | 15.78  | 16.1  | 17.68  | 17.4  | 15.11  | 15.3  | 12.93  | 13.3  | 15.66  | 15.9  |
| JUN   | 21.51  | 21.3  | 20.40  | 20.3  | 19.00  | 18.9  | 19.11  | 18.9  | 18.02  | 17.9  | 22.39  | 21.9  |
| JUL   | 22.88  | 22.3  | 22.52  | 22.4  | 23.24  | 22.9  | 24.36  | 23.7  | 22.75  | 22.6  | 22.13  | 21.7  |
| AUG   | 20.83  | 20.4  | 21.42  | 21.5  | 23.25  | 23.4  | 21.01  | 21.2  | 20.61  | 20.5  | 22.28  | 22.1  |
| SEP   | 16.25  | 15.2  | 13.94  | 13.8  | 15.90  | 15.2  | 15.07  | 14.9  | 14.44  | 14.4  | 15.06  | 15.2  |
| OCT   | 8.28   | 7.9   | 8.38   | 8.7   | 9.27   | 8.7   | 6.12   | 6.2   | 6.65   | 7.1   | 7.22   | 7.5   |
| NOV   | -4.30  | -4.1  | -4.08  | -3.2  | -6.73  | -6.1  | -3.98  | -3.3  | -6.93  | -6.6  | -2.72  | -2.4  |
| DEC   | -14.81 | -15.3 | -15.95 | -15.9 | -18.30 | -17.9 | -10.04 | -9.2  | -16.81 | -16.1 | -12.42 | -12.6 |




# MONTHLY AVERAGE TEMPERATURE

PEKING

#54511

| MONTH | 1965  |      | 1966  |      | 1967  |      | 1968  |      | 1969  |      | 1970  |      |
|-------|-------|------|-------|------|-------|------|-------|------|-------|------|-------|------|
|       | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  |
| JAN   | -7.06 | -4.9 | -6.50 | -5.7 | -6.08 | -4.8 | -6.96 | -5.6 | -8.22 | -6.4 | -7.04 | -5.6 |
| FEB   | -1.41 | -1.7 | -2.43 | -2.5 | -4.79 | -3.3 | -6.37 | -4.9 | -7.15 | -5.1 | -4.03 | -2.0 |
| MAR   | 4.50  | 4.4  | 1.70  | 3.4  | 3.63  | 4.8  | 4.38  | 5.7  | -0.15 | 2.1  | -0.10 | 2.0  |
| APR   | 10.65 | 10.9 | 10.51 | 12.6 | 10.45 | 12.8 | 12.11 | 14.2 | 10.34 | 12.9 | 11.31 | 13.6 |
| MAY   | 20.35 | 20.8 | 17.79 | 19.7 | 20.63 | 22.1 | 17.61 | 20.3 | 16.96 | 19.3 | 16.59 | 19.7 |
| JUN   | 23.98 | 24.6 | 22.28 | 23.8 | 21.16 | 23.7 | 23.06 | 25.5 | 20.92 | 23.3 | 20.12 | 22.8 |
| JUL   | 25.87 | 26.9 | 24.02 | 25.6 | 23.63 | 25.3 | 25.47 | 26.9 | 23.13 | 24.6 | 23.16 | 24.8 |
| AUG   | 24.19 | 24.8 | 23.94 | 25.0 | 23.72 | 25.1 | 22.68 | 24.2 | 22.12 | 24.2 | 22.51 | 27.8 |
| SEP   | 20.83 | 20.2 | 17.35 | 18.7 | 16.67 | 18.3 | 17.99 | 19.7 | 17.56 | 19.6 | 17.69 | 18.9 |
| OCT   | 16.20 | 14.1 | 11.42 | 12.8 | 11.44 | 13.0 | 9.83  | 11.2 | 10.19 | 12.3 | 11.31 | 13.0 |
| NOV   | 5.42  | 4.6  | 2.50  | 3.7  | 0.49  | 2.0  | 2.86  | 3.9  | 1.38  | 2.8  | 1.39  | 3.0  |
| DEC   | -4.10 | -3.9 | -5.70 | -4.7 | -7.59 | -6.0 | -3.25 | -1.7 | -5.09 | -3.4 | -4.40 | -3.1 |




## MONTHLY AVERAGE TEMPERATURE

SHANGHAI

#58367

| MONTH | 1965  |      | 1966  |      | 1967  |      | 1968  |      | 1969  |      | 1970  |      |
|-------|-------|------|-------|------|-------|------|-------|------|-------|------|-------|------|
|       | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  | EST   | PUB  |
| JAN   | 4.56  | 5.1  | 3.62  | 4.3  | 1.42  | 2.0  | 1.85  | 2.8  | 2.10  | 3.6  | 0.81  | 2.0  |
| FEB   | 4.98  | 5.7  | 5.55  | 6.6  | 3.50  | 4.0  | 0.34  | 1.8  | 0.93  | 2.6  | 4.80  | 6.0  |
| MAR   | 7.34  | 7.7  | 9.25  | 10.2 | 8.08  | 8.7  | 7.94  | 8.8  | 5.37  | 6.8  | 4.87  | 6.0  |
| APR   | 11.10 | 12.2 | 13.36 | 13.8 | 12.88 | 13.6 | 14.11 | 13.7 | 12.91 | 14.2 | 11.67 | 13.1 |
| MAY   | 19.06 | 19.5 | 17.64 | 19.0 | 18.18 | 19.3 | 18.98 | 18.7 | 18.55 | 19.6 | 17.70 | 18.7 |
| JUN   | 21.29 | 22.2 | 22.49 | 23.6 | 23.91 | 24.3 | 22.90 | 22.6 | 21.31 | 22.5 | 21.19 | 22.1 |
| JUL   | 27.28 | 28.1 | 26.09 | 27.1 | 27.03 | 28.0 | 27.07 | 26.7 | 25.68 | 26.7 | 26.73 | 27.1 |
| AUG   | 26.46 | 26.9 | 27.44 | 28.6 | 28.53 | 29.5 | 26.37 | 27.5 | 27.10 | 28.0 | 26.89 | 28.1 |
| SEP   | 21.96 | 22.7 | 21.10 | 22.3 | 23.14 | 24.0 | 22.57 | 23.7 | 23.96 | 25.0 | 23.85 | 24.7 |
| OCT   | 18.09 | 18.0 | 16.66 | 18.1 | 17.10 | 18.4 | 16.08 | 17.4 | 16.42 | 18.0 | 17.55 | 18.5 |
| NOV   | 13.31 | 13.9 | 11.13 | 12.6 | 11.04 | 12.5 | 12.19 | 13.3 | 9.26  | 10.3 | 11.22 | 11.8 |
| DEC   | 4.26  | 4.9  | 5.19  | 6.1  | -0.04 | 1.0  | 7.88  | 9.5  | 3.93  | 4.9  | 5.87  | 6.9  |

ORIGINAL PAGE IS  
OF POOR QUALITY



# ANALYSIS OF VARIANCE

| SOURCE OF VARIATIONS | DEGREES OF FREEDOM | SUM OF SQUARES | MEAN SQUARE       | F - RATIO     |
|----------------------|--------------------|----------------|-------------------|---------------|
| BETWEEN GROUPS       | $p - 1$            | BSS            | $BMS = BSS/(p-1)$ |               |
| WITHIN GROUPS        | $N - p$            | WSS            | $WMS = WSS/(N-p)$ | $F = BMS/WMS$ |
| TOTAL                | $N - 1$            | TSS            |                   |               |

WHERE  $p$  IS THE # OF GROUPS  
 $N$  IS THE # OF OBSERVATIONS  
 $R_i$  IS THE # OF OBSERVATION PER GROUP

$$TSS = \sum_i \sum_j X_{ij}^2 - (\sum_i \sum_j X_{ij})^2 / N$$

$$BSS = \sum_i (\sum_j X_{ij})^2 / R_i - (\sum_i \sum_j X_{ij})^2 / N$$

$$WSS = \sum_i \sum_j X_{ij}^2 - \sum_i (\sum_j X_{ij})^2 / R_i$$

$i, j = 1, 2, 3, \dots, N$

ORIGINAL PAGE IS  
OF POOR QUALITY

ORIGINAL PAGE IS  
OF POOR QUALITY

| LOCKHEED     | TA-LIEN(#54662)<br>U.S. AIR FORCE | HONG KONG ROYAL |
|--------------|-----------------------------------|-----------------|
| 21.80        | 23.34                             | 25.000          |
| 26.67        | 27.86                             | 25.000          |
| 35.18        | 36.14                             | 36.000          |
| 45.60        | 45.86                             | 48.000          |
| 58.69        | 60.26                             | 59.000          |
| 67.58        | 68.18                             | 68.000          |
| 71.61        | 73.58                             | 73.000          |
| 71.48        | 72.68                             | 76.000          |
| 69.24        | 69.24                             | 69.000          |
| 61.15        | 59.90                             | 58.000          |
| 41.95        | 42.80                             | 43.000          |
| 27.00        | 28.00                             | 30.000          |
| SUM= 1815.79 | SUMSQ= 103413.06                  |                 |

SUM OF GROUP( 1) IS 597.95  
 SUM OF GROUP( 2) IS 607.84  
 SUM OF GROUP( 3) IS 610.00

| SOURCE OF VARIATION | ANALYSIS OF VARIANCE |               |             | F RATIO |
|---------------------|----------------------|---------------|-------------|---------|
|                     | DEGREE OF FREEDOM    | SUM OF SQUARE | MEAN SQUARE |         |
| BETWEEN GROUP       | 2                    | 6.938         | 3.469       |         |
| WITHIN GROUPS       | 33                   | 11820.375     | 358.193     | 0.010   |
| TOTAL               | 35                   | 11827.313     |             |         |

SINCE  $FCAL < FTAB$ , SO ACCEPT THE HYPOTHESE

|      | LOCKHEED | PEKING(#54511)<br>U.S.AIR FORCE | HONG KONG ROYAL |
|------|----------|---------------------------------|-----------------|
|      | 19.28    | 23.18                           | 23.000          |
|      | 29.44    | 28.94                           | 29.000          |
|      | 40.45    | 39.92                           | 41.000          |
|      | 51.17    | 51.62                           | 57.000          |
|      | 68.64    | 69.44                           | 68.000          |
|      | 75.17    | 76.28                           | 76.000          |
|      | 78.58    | 80.42                           | 79.000          |
|      | 75.54    | 76.64                           | 76.000          |
|      | 70.50    | 68.36                           | 68.000          |
|      | 61.18    | 57.36                           | 54.000          |
|      | 41.77    | 40.28                           | 38.000          |
|      | 24.65    | 24.38                           | 27.000          |
| SUM= | 1909.19  | SUMSQ= 116026.31                |                 |

SUM OF GROUP( 1) IS 636.37  
 SUM OF GROUP( 2) IS 636.82  
 SUM OF GROUP( 3) IS 636.00

| SOURCE OF VARIATION | ANALYSIS OF VARIANCE |               |             | F RATIO |
|---------------------|----------------------|---------------|-------------|---------|
|                     | DEGREE OF FREEDOM    | SUM OF SQUARE | MEAN SQUARE |         |
| BETWEEN GROUP       | 2                    | 0.125         | 0.063       | 0.000   |
| WITHIN GROUPS       | 33                   | 14776.313     | 447.767     |         |
| TOTAL               | 35                   | 14776.438     |             |         |

SINCE  $FCAL < FTAB$ , SO ACCEPT THE HYPOTHESE

ORIGINAL PAGE IS  
OF POOR QUALITY

# MONTHLY PRECIPITATION

(In Millimeters)

| MONTH       | 1967             |       | 1968              |       | 1969              |       | 1970              |        |
|-------------|------------------|-------|-------------------|-------|-------------------|-------|-------------------|--------|
|             | PEKING<br>#54511 |       | CHI-NAN<br>#54823 |       | TA-LIEN<br>#54662 |       | NANKING<br>#58238 |        |
|             | EST              | PUB   | EST               | PUB   | EST               | PUB   | EST               | PUB    |
| JAN         | 2.0              | 2.8   | 5.8               | 7.3   | 0.0               | 0.7   | 15.7              | 9.3    |
| FEB         | 1.7              | 2.4   | 1.7               | 1.5   | 5.3               | 3.4   | 66.0              | 84.1   |
| MAR         | 10.6             | 7.7   | 5.8               | 3.4   | 27.1              | 16.0  | 50.8              | 66.6   |
| APR         | 29.7             | 26.8  | 41.4              | 45.1  | 80.2              | 70.3  | 115.8             | 120.0  |
| MAY         | 30.2             | 17.4  | 19.3              | 22.1  | 42.4              | 32.0  | 132.0             | 155.5  |
| JUN         | 104.6            | 129.7 | 10.4              | 10.0  | 26.1              | 51.4  | 87.1              | 91.5   |
| JUL         | 149.1            | 173.5 | 34.5              | 27.9  | 105.4             | 87.8  | 176.2             | 218.2  |
| AUG         | 177.1            | 180.4 | 25.1              | 61.3  | 101.1             | 137.0 | 198.8             | 155.2  |
| SEP         | 27.4             | 26.0  | 8.8               | 10.2  | 15.7              | 38.7  | 119.3             | 154.7  |
| OCT         | 37.8             | 19.7  | 64.5              | 68.9  | 10.4              | 7.9   | 74.4              | 71.1   |
| NOV         | 7.5              | 7.0   | 37.5              | 37.5  | 15.4              | 32.1  | 12.9              | 13.7   |
| DEC         | 0.0              | 0.0   | 40.6              | 25.5  | 5.6               | 4.6   | 26.4              | 14.7   |
| ANNUAL (mm) | 577.7            | 593.4 | 295.4             | 320.7 | 434.7             | 481.9 | 1075.4            | 1154.6 |
| TOTAL (in)  | 22.7             | 23.3  | 11.6              | 12.6  | 17.1              | 18.0  | 42.0              | 45.0   |

MONTHLY PRECIPITATION IN MM

DE KING  
\*54511  
北京市

PUBLISHED

ESTIMATED

1 2 3 4 5 6 7 8 9 10 11 12

MONTHS  
< 1967 >

ORIGINAL PAGE IS  
OF POOR QUALITY

Print

MONTHLY PRECIPITATION IN MM


CHI-NAN  
\* 54823  
济南市


PUBLISHED  
PRECIPITATION


ESTIMATED  
PRECIPITATION

1 2 3 4 5 6 7 8 9 10 11 12

MONTHS  
< 1968 >







#### 4. OPERATIONS

##### 4.1 LEAST SQUARES INTERPOLATION

This program was designed to work on more than one data set which is denoted by a parameter NS.

Each data set contains XP and YP up to 100 observations as input data; in each card, punch XP in columns 2-6 and YP in columns 7-11, with one decimal point at column 5 and column 10 for XP and YP; respectively.

At the end of each data set punch a trailer card with -99.0000.0 in columns 2-11.

Example of a program setup:

```
// JOB CARD
//SS EXEC FCALCMGO, PARM.FORM = 'MAP'
//FORT*SYSIN DD *
SOURCE PROGRAM OF LEAST SQUARES INTERPOLATION
//GO*SYSIN DD *
DATA CARDS
/*
//
```

ORIGINAL PAGE IS  
OF POOR QUALITY.

##### 4.2 THE MODIFIED MAX-MIN PROGRAM

The daily estimates of maximum and minimum air temperatures and the sum of precipitation measurements reported each day are obtained according to procedures described in "As-Built Design Specifications for Historical Daily Data Bases for Testing Advanced Models", AD 63-1347-4963-08 (JSC-12891, LEC-10572, April 1977).

### 4.3 THE MONTHLY DATA BASES ALGORITHM

This program needs no input data from cards; all its input variables are read in from the magnetic tape.

Daily data are stored in tapes by years, each file contains one year's daily data, each of 85-95 stations.

Example of a program setup:

```
// JOB CARD
/* SETUP      TAPE = 1
//SS EXEC FTG1CLG, PARM.FORT = 'MAP'
// FORT.SYSIN      DD      *
        SOURCE PROGRAM OF THE MONTHLY DATA BASES ALGORITHM
//LKED.SYSIN      DD      *
        INCLUDE BMDSUB (BMDZ992)
//LKED.BMDSUB      DD      DSN=SYS2.BMD.LOAD, DISP = SHR
//GO.FT 09F001      DD      DSN=CH68A,
// UNIT=TAPE, VOL=(, RETAIN, SER = X06121),
// DCB=(RECFM=FB, LBECL=26, BLKSIZE=7800, DEN=4),
// LABEL=(44,,,IN),
// DISP=OLD, SHR)
//GO.FT10F001      DD      DSN=CH68B
// UNIT = TAPE, VOL= SER = X06121,
// DCB= (RECFM = FB, LBECL = 26, BLKSIZE=7800, DEN=4),
// LABEL = (45,,,IN)
// DISP = (OLD, KEKP)
//GO.SYSIN      DD
/*
//
```

#### 4.4 THE ANALYSIS OF VARIANCE ALGORITHM

Twelve input cards (one for each month) are required for running the program. Each card contains three estimates of the monthly mean temperature based on the KSU algorithm output, the "published" values obtained through NCC from ETAC for years available and period of record means obtained from various Chinese publications. These values are located as follows:

Columns 1-5 contains KSU estimates

" 7-11 " ETAC record

" 13-18 " period of record means

and each field contains two digits, i.e., one decimal point punched in columns 3, 9, and 15, for KSU, ETAC and period of record means.

Example of a program setup:

```
// JOB CARD
//SS EXEC FTG1CLG, PARM. FORT = 'MAP'
//FORT=SYSIN DD *
        SOURCE PROGRAM OF AOV ALGORITHM
//GO=SYSIN DD *
        DATA CARD

/*
//
```

## 5. ACKNOWLEDGEMENT:


The contribution of the University of Missouri-Columbia, in furnishing computer time for the derivation of these data bases is gratefully acknowledged. Without the cooperation of Dr. Wayne L. Decker, Chairman of the Department of Atmospheric Sciences, University of Missouri-Columbia, this data source would have been much later in making its appearance.


## APPENDIX A

### TABLE OF CONTENTS


|                                                                                  | PAGE |
|----------------------------------------------------------------------------------|------|
| LEAST SQUARES INTERPOLATION PROGRAM FLOWCHART                                    | A- 1 |
| LEAST SQUARES INTERPOLATION PROGRAM LISTING                                      | A-14 |
| SUBROUTINE SUM FLOWCHART                                                         | A-18 |
| SUBROUTINE SUM LISTING                                                           | A-19 |
| SUBROUTINE LSTSQ FLOWCHART                                                       | A-20 |
| SUBROUTINE LSTSQ LISTING                                                         | A-23 |
| SUBROUTINE MEQSOL FLOWCHART                                                      | A-24 |
| SUBROUTINE MEQSOL LISTING                                                        | A-28 |
| SUBROUTINE MULTR FLOWCHART                                                       | A-29 |
| SUBROUTINE MULTR LISTING                                                         | A-31 |
| SUBROUTINE GRAPH FLOWCHART                                                       | A-32 |
| SUBROUTINE GRAPH LISTING                                                         | A-34 |
| SUBROUTINE COR FLOWCHART                                                         | A-35 |
| SUBROUTINE COR LISTING                                                           | A-36 |
| SUBROUTINE STANDV FLOWCHART                                                      | A-37 |
| SUBROUTINE STANDV LISTING                                                        | A-39 |
| <br>EXAMPLE OF PROGRAM OUTPUT FROM THE LEAST SQUARES<br>INTERPOLATIONS ALGORITHM |      |
| A 2nd DEGREE CURVE                                                               | A-40 |
| A 3rdDEGREE CURVE                                                                | A-41 |

# LEAST SQUARES INTERPOLATION







ORIGINAL PAGE IS  
OF POOR QUALITY



ORIGINAL PAGE IS  
OF POOR QUALITY




ORIGINAL PAGE IS  
OF POOR QUALITY







ORIGINAL PAGE IS  
OF POOR QUALITY












ORIGINAL PAGE IS  
OF POOR QUALITY





ORIGINAL PAGE IS  
OF POOR QUALITY



```

*****
* LEAST SQ INTERPOLATION *
*           BY           *
*        BUDDY H JEUN    *
*****

```

IN THIS PROGRAM, ALL YOU HAVE TO DO IS TO FIT A SET OF DATA PTS TO THE COMPUTER, AND THE COMPUTER WILL RETURN A BEST FIT LEAST SQ EQUATION FOR YOUR DATA PTS, AND THAT RESULTING EQUATION IS INDEED THE BEST FIT IN THE STATISTICAL SENSE. AND THE RESULTING EQUATION IS USE FOR INTERPOLATION.

THIS PROGRAM CONSISTING OF ONE MAIN PROGRAM AND SEVEN SUBROUTINE SUBPROGRAMS, THAT IS SUBROUTINE SUM, COR, LSTSQ, MEQSOL, AND GRAPH AND STANDV. MULTR  
 SUBROUTINE SUM IS USE TO FIND ALL SUMS FOR X AND Y  
 SUBROUTINE COR IS USE TO FIND THE COR. COEFF  
 SUBROUTINE LSTSQ IS USE TO FIND THE N DEG EQUATION  
 SUBROUTINE MEQSOL IS USE TO SOLVE THE N\*N SYSTEM OF EQUATION  
 SUBROUTINE GRAPH IS USE TO OUTPUT THE DATA IN THE FORM OF A STRAIGHT LINE OR A CURVE  
 SUBROUTINE STANDV IS USE TO CALCULATE THE STANDARD DEVIATION OF A GIVEN SET OF DATA PTS  
 SUBROUTINE RMULT IS USE TO FIND THE MULTIPLE CORRELATION COEFF  
 THE CRITERIA TO DETERMINE WHICH ONE IS THE BEST FIT IS ACCORDING TO THE VALUE OF THE MULTIPLE CORRELATION COEFF  
 THE HIGHEST VALUE OF MULTIPLE R IS CONSIDER AS THE BEST FIT

```

0001 REAL XP(100),YP(100),XL(200),YL(200),RES(100)
0002 COMMON/AREA1/SX,SY,SXSQ,SYSQ,SXY/AREA2/C(25),A(25,25),B(25)
0003 REAL XSQ(100),XCUB(100)
0004 NS=28
0005 NS IS THE NUMBER OF THE DATA SET TO BE FIT INTO THE REGRESSION LINE
0006 K=1
0007 9 WRITE(6,20)K
0008 20 FORMAT('1',52X,'THE INPUT DAT #',I3,' IS:',/60X,'X2',10X,'YP'//
0009      #'+',59X,3('--'),8X,3('--'))
0010 DO 111 I=1,100
0011 READ(5,3)XP(I),YP(I)
0012 3 FORMAT(1X,2F5.1)
0013 IF (XP(I).EQ.-99.0) GO TO 222
0014 WRITE(6,66)XP(I),YP(I)
0015 66 FORMAT(56X,F7.2,5X,F7.2)
0016 111 CONTINUE
0017 222 NE=I-1
0018 DO 50 I=1,NE
0019 XSQ(I)=XP(I)**2
0020 XCUB(I)=XP(I)**3
0021 50 CONTINUE
0022 EN=FLOAT(NE)
0023 CALL SUM(NE,XP,YP)
0024 WRITE(6,7)
0025 7 FORMAT(///20X,'THE RESULT OF CALCULATIONS ARE:',/'+',20X,27('_''))
0026 WRITE(6,6) NE

```

ORIGINAL PAGE IS  
OF POOR QUALITY

```

0025      6 FORMAT(20X,'N=',I3)
0026      WRITE(6,8)SX,SY,SXSQ,YSQ,SXY
0027      8 FORMAT(20X,'SX=',F10.2/20X,'SY=',F10.2/20X,'SXSQ=',F10.2/20X,
+ 'YSQ=',F10.2/20X,'SXY=',F10.2)
0028      CALL COR(NE,R)
0029      WRITE(6,77)R
0030      77 FORMAT(/50X,'THE COR. COEFF IS R=',F5.2//)
0031      IF(ABS(R).LT.0.7) GO TO 444
C      SINCE X AND Y ARE HIGHLY CORRELATED, SO FIT A STRAIGHT LINE
0032      ANEW=SXSQ*SY-SX*SXY
0033      BNEW=EN*SXY-SX*SY
0034      DENEW=EN*SXSQ-SX**2
0035      A1=ANEW/DENEW
0036      B1=BNEW/DENEW
0037      WRITE(6,33)A1,B1
0038      33 FORMAT(/50X,'THE LINEAR REG LINE IS: '/50X,'Y=',F8.2,2X,'+',F8.2,
+ '*X'//)
0039      WRITE(6,21)
0040      XMIN=1.0
0041      XMAX=26
0042      DALTAX=1.0
0043      X=XMIN
0044      666 Y=A1+B1*X
0045      WRITE(6,22)X,Y
0046      CALL GRAPH(Y)
0047      X=X+DALTAX
0048      IF(X-XMAX)666,666,317
C      FIND A HIGHER DEGREE EQUATION BY THE LEAST SQ METHOD
0049      444 NC=3
0050      CALL PLTSCT(XP,YP,XL,YL,NE,100,2,
+ 36HLEAST SQ INTERPOLATION PLOT
+ ,36HTHE MONTHS
+ ,36HTHE MONTHLY TEMP RANGE
+)
0051      CALL LSTSQ(XP,YP,NE,NC)
0052      CALL MULTR(NE,XP,YP,RXY)
0053      CALL MULTR(NE,XSQ,YP,RXSQY)
0054      C1=C(1)
0055      C2=C(2)
0056      C3=C(3)
C      AFER CALL THE 2ND DEGREE POLYN
0057      CALL STANDV(NE,XP,SDX)
0058      CALL STANDV(NE,YP,SDY)
0059      CALL STANDV(NE,XSQ,SDXSQ)
C      B(I)=C(I)*SD(I)/SDY
0060      B1=(C2*SDX)/SDY
0061      B2=(C3*SDXSQ)/SDY
C      THE MULTIPLE R FOR 2ND DEGREE IS:
0062      S2=ABS(RXY*B1+RXSQY*B2)
0063      R2=SQRT(S2)
0064      NC=4
0065      CALL LSTSQ(XP,YP,NE,NC)
0066      CALL MULTR(NE,XCUB,YP,RXCUBY)
C      AFTER CALL THE 3RD DEGRE POLYN;
0067      CALL STANDV(NE,XCUB,SDXCUB)
0068      B1=(C(2)*SDX)/SDY

```

```

0069      B2=(C(3)*SDXSQ)/SDY
0070      B3=(C(4)*SDXCUB)/SDY
          C      THE MULTIPLE R FOR THE 3DEGREE IS:
0071      S3=(RXY*B1+RXSQY*B2+RXCUBY*B3)
0072      R3=SQRT(S3)
0073      IF(R2.GT.R3) GO TO 500
0074      CALL PLTSCT(XP,YP,XL,YL,NE,100,3,
+ 36HLEAST SQ INTERPOLATION PLOT
+ ,36HTHE MONTHS
+ ,36HTHE MONTHLY TEMP RANGE
+)
0075      CALL LSTSQ(XP,YP,NE,NC)
0076      WRITE(6,25)(I,C(I),I=1,NC)
0077      25  FORMAT(5X,'C(',I1,') =',F15.5)
0078      WRITE(6,11)
0079      11  FORMAT(/50X,'THE LEAST SQ EQUATION IS :'/)
0080      WRITE(6,12)(C(I),I=1,NC)
0081      12  FORMAT(50X,'Y=',F7.2,'+',F7.2,'X','+',F7.2,'X**2','+',F7.2,'X**3')
          C      HEADING
0082      WRITE(6,21)
0083      21  FORMAT('1',45X,'THE GRAPH OF THE BEST FIT LEAST SQ EQUATION IS:/'
+ 4X,'X',5X,'Y',28X,78(' '),Y-AXIS'/
+ ' ',6(' '),2X,6(' '))
0084      XMIN=1.0
0085      XMAX=26
0086      DALTAX=1.0
0087      X=XMIN
0088      201 Y=C(1)+C(2)*X+C(3)*X**2+C(4)*X**3
0089      CALL GRAPH(Y)
0090      WRITE(6,22)X,Y
0091      22  FORMAT(1X,F6.2,F8.2)
0092      X=X+DALTAX
0093      IF(X-XMAX)201,201,202
0094      202 WRITE(6,318)
0095      GO TO 505
0096      500 WRITE(6,319)C1,C2,C3
0097      319 FORMAT(50X,'Y=',F7.2,'+',F7.2,'X','+',F7.2,'X**2')
0098      XMIN=0.0
0099      XMAX=26.0
0100      DALTAX=1.0
0101      X=XMIN
0102      404 Y=C1+C2*X+C3*X**2
0103      WRITE(6,78)X,Y
0104      78  FORMAT(60X,2F10.2)
0105      X=X+DALTAX
0106      IF(X-XMAX)404,404,505
0107      505 WRITE(6,55)R2,R3
0108      55  FORMAT(/20X,'THE MULTIPLE R FOR THE 2ND DEGREE POLYN IS',F10.2/
+ 20X,'THE MULTIPLE R FOR THE 3RD DEGREE POLYN IS',F10.2//)
          IF(K.EQ.NS) GO TO 690
0109      K=K+1
0110      GO TO 9
0111      317 WRITE(6,318)
0112      318 FORMAT(38X,'X-AXIS')
0113      IF(K.EQ.NS) GO TO 690
0114      K=K+1
0115

```

A-16

ORIGINAL PAGE IS  
OF POOR QUALITY

FORTRAN IV G1 RELEASE 2.0

MAIN

DATE = 77321


00/54/57

PAG

0116  
0117  
0118


GO TO 9  
690 STOP  
END


# SUBROUTINE SUM




```
0001      SUBROUTINE SUM(NE,X,Y)
0002      COMMON/AREA1/SX,SY,SXSQ,SYSQ,SXY
      C
      C      IN THIS SUBROUTINE SUM, THE INPUT DATA ARE NE,X,Y AND THE OUTPUT
      C      ARGUMENTS ARE SX,SY,SXSQ,SYSQ,SXY;ALL VALUES FOR THE SUMS ARE
      C      CALCULATED IN THIS SUBROUTINE.
      C
0003      DIMENSION X(100),Y(100)
0004      SX=0.0
0005      SY=0.0
0006      SXSQ=0.0
0007      SYSQ=0.0
0008      SXY=0.0
0009      DO 10 I=1,NE
0010      SX=SX+X(I)
0011      SY=SY+Y(I)
0012      SXSQ=SXSQ+X(I)**2
0013      SYSQ=SYSQ+Y(I)**2
0014      SXY=SXY+X(I)*Y(I)
0015      10 CONTINUE
0016      RETURN
0017      END
```

# SUBROUTINE LSTSQ







```


0001      SUBROUTINE LSTSQ(X,Y,N,M)
0002      COMMON/AREA2/C(25),A(25,25),B(25)


      C
      C
      C      LEAST SQUARE CURVE FITTING
      C      TO FIT N GIVEN POINTS X(N),Y(N) BY A POLYNOMIAL OF
      C      M-1 DEGREE, SUM C(I)*X**(I-1) FOR I=1 TO I=M
      C      C(M) IS THE OUTPUT COEFF ARRAY
      C      NEEDS SUBROUTINE MEQSOL TO SOLVE FOR AC=B


0003      REAL X(100),Y(100)
0004      DO 7 I=1,M
0005      B(I)=0.0
0006      DO 5 K=1,N
0007      IF((X(K).EQ.0.0).AND.(I.EQ.1))X(K)=1.0
0008      5 B(I)=B(I)+X(K)**(I-1)*Y(K)
0009      DO 7 J=1,M
0010      A(I,J)=0.0
0011      DO 7 K=1,N
0012      IF((X(K).EQ.0.0).AND.((I+J).EQ.2))X(K)=1.0
0013      A(I,J)=A(I,J)+X(K)**(I+J-2)
0014      7 CONTINUE
0015      CALL MEQSOL(M)
0016      RETURN
0017      END


```

# SUBROUTINE MEASOL









0001 SUBROUTINE MEQSOL(N)  
0002 COMMON/AREA2/X(25),A(25,25),B(25)

C  
C  
C  
C  
C  
C


THIS SUBROUTINE MEQSOL IS USE TO SOLVE THE MATRIX EQUATION  $AX=B$  OF N, IN THIS PARTICULAR CASE, N IS 4, AND THE RESULT IS RETURNED TO THE MAIN PROGRAM BY THE LABEL COMMON X(25). IF HIGHER ORDER MATRIX EQUATION IS NEEDED TO BE SOLVE, THEN THE VALUE OF N HAVE TO BE CHANGE


0003 INTEGER M(25)  
0004 DO 9 I=1,N  
0005 M(I)=1  
0006 AMAX=A(I,1)  
0007 DO 2 J=2,N  
0008 IF(ABS(A(I,J))-ABS(AMAX)) 2,2,1  
0009 1 AMAX=A(I,J)  
0010 M(I)=J  
0011 2 CONTINUE  
0012 IF(AMAX) 3,98,3  
0013 DO 4 J=1,N  
0014 4 A(I,J)=A(I,J)/AMAX  
0015 B(I)=B(I)/AMAX  
0016 DO 9 IP=1,N  
0017 IF(IP-1)5,9,5  
0018 5 MMM=M(I)  
0019 ZMULT=A(IP,MMM)  
0020 DO 8 J=1,N  
0021 IF(J-MMM)7,6,7  
0022 6 A(IP,J)=0.0  
0023 GO TO 8  
0024 7 A(IP,J)=A(IP,J)-ZMULT\*A(I,J)  
0025 8 CONTINUE  
0026 B(IP)=B(IP)-ZMULT\*B(I)  
0027 9 CONTINUE  
0028 DO 95 I=1,N  
0029 95 X(M(I))=B(I)  
0030 RETURN  
0031 98 WRITE(6,99)  
0032 99 FORMAT(12H NO SOLUTION)  
0033 RETURN  
0034 END

A-28

ORIGINAL PAGE IS  
OF POOR QUALITY

# SUBROUTINE MULTR






```


0001      SUBROUTINE MULTR(N,X,Y,RMULT)
          C      THIS SUBROUTINE MULTR IS USE TO CALCULATE THE MULTIPLE
          C      CORRELATION COEFF R AND DENOTED THAT BY THE OUTPUT PARAMETER
          C      MULTR AND THE FOMULA IS  $R = \sqrt{\text{SUME OF } R(I,Y)*B(J)}$ ,
          C      WHERE B(J) IS THE STANDARDIZE REGRESSION COEFFAND THE
          C      UNSTANDARDIZE REGRESSION COEFF IS  $B = SQ(J)/SD(Y)*B(J)$  ....
0002      REAL X(100),Y(100)
0003      EN=N
0004      SX=0.0
0005      SY=0.0
0006      SXY=0.0
0007      SXSQ=0.0
0008      SYSQ=0.0
0009      DO 100 I=1,N
0010      SX=SX+X(I)
0011      SY=SY+Y(I)
0012      SXY=SXY+X(I)*Y(I)
0013      SXSQ=SXSQ+X(I)**2
0014      SYSQ=SYSQ+Y(I)**2
0015      100 CONTINUE
0016      A=(EN*SXY-(SX)*(SY))
0017      B=SQRT(EN*SXSQ-(SX)**2)
0018      C=SQRT(EN*SYSQ-(SY)**2)
0019      RMULT=(A/(B*C))
0020      RETURN
0021      END

```

ORIGINAL PAGE IS  
OF POOR QUALITY

# SUBROUTINE GRAPH






```
0001      SUBROUTINE GRAPH(Y)
0002      DIMENSION IP(81)
0003      DATA IASTR/'*'/,IBLNK/' '/
0004      J=ABS(Y)
0005      IF(J-81)500,500,600
0006      DO 333 I=1,81
0007          IP(I)=IBLNK
0008      333 CONTINUE
0009      IP(J)=IASTR
0010      WRITE(6,51) IP
0011      51 FORMAT('+',38X,'1',81A1)
0012      600 RETURN
0013      END
```


ORIGINAL PAGE IS  
OF POOR QUALITY


# SUBROUTINE COR



```
0001      SUBROUTINE COR(NE,R)
0002      COMMON/ARFA1/SX,SY,SXSQ,SYSQ,SXY
      C
      C      IN THIS SUBROUTINE COR,THE INPUT ARGUMENTS ARE NE,SX,SY,SXSQ,SYSQ,
      C      SXY; AND THE OUPUT ARGUMENT IS R;THE COR. COEFF IS RETURN TO THE
      C      MAIN PROGRAM BY R.
0003      EN=FLOAT(NE)
0004      RNEW=EN*SXY-SX*SY
0005      DENEWX=EN*SXSQ-SX**2
0006      DENEWY=EN*SYSQ-SY**2
0007      R=RNEW/SQRT(DENEWX*DENEWY)
0008      RETURN
0009      END
```

# SUBROUTINE STANDV





```
0001      SUBROUTINE STANDV(N,Z,STDV)
      C      THIS SUBROUTINE IS USE TO CALCULATE THE STANDARD DEVIATION
      C      OF A GIVEN SET OF DATA PTSAND DENOTED THAT BY THE OUTPUT
      C      PARAMETER STDV
      REAL Z(100)
0002      EN=N
0003      SZ=0.0
0004      SZSQ=0.0
0005      DO 200 I=1,N
0006      SZ=SZ+Z(I)
0007      SZSQ=SZSQ+Z(I)**2
0008      CONTINUE
0009      A=(EN*SZSQ-SZ**2)
0010      B=A/(EN*(EN-1.0))
0011      STDV=SQRT(B)
0012      RETURN
0013      END
0014
```

ORIGINAL PAGE IS  
OF POOR QUALITY

# LEAST SQ INTERPOLATION PLOT


THE MONTHLY TEMP RANGE

( X 10\*\* 0 )

31.547 I  
31.241 I  
30.935 I  
30.629 I  
30.322 I  
30.016 I  
29.710 I  
29.404 I  
29.098 I  
28.792 I  
28.486 I  
28.180 I  
27.873 I  
27.567 I  
27.261 I  
26.955 I  
26.649 I  
26.343 I  
26.037 I  
25.731 I  
25.424 I  
25.118 I  
24.812 I  
24.506 I  
24.200 I  
23.894 I  
23.588 I  
23.282 I  
22.975 I  
22.669 I  
22.363 I  
22.057 I  
21.751 I  
21.445 I  
21.139 I  
20.833 I  
20.527 I  
20.220 I  
19.914 I  
19.608 I  
19.302 I  
18.996 I  
18.690 I  
18.384 I  
18.078 I  
17.771 I  
17.465 I  
17.159 I  
16.853 I  
16.547 I

1.000 3.000 5.000 7.000 9.000 11.000 13.000 15.000 17.000  
THE MONTHS  
( X 10\*\* 0 )

A-40



# LEAST SQ INTERPOLATION PLOT

THE MONTHLY TEMP RANGE

( X.10\*\* 0 )

25.398 I  
 25.194 I  
 24.990 I  
 24.786 I  
 24.582 I  
 24.378 I  
 24.173 I  
 23.969 I  
 23.765 I  
 23.561 I  
 23.357 I  
 23.153 I  
 22.949 I  
 22.745 IO  
 22.541 I  
 22.337 I  
 22.133 I  
 21.929 I  
 21.724 I  
 21.520 I  
 21.316 I  
 21.112 I  
 20.908 I  
 20.704 I  
 20.500 I  
 20.296 I  
 20.092 I  
 19.888 I  
 19.684 I+  
 19.480 I+  
 19.275 I  
 19.071 I  
 18.867 I++  
 18.663 IO+  
 18.459 IO+  
 18.255 I++  
 18.051 I+  
 17.847 I++  
 17.643 I++  
 17.439 I++++  
 17.235 I++++  
 17.031 I  
 16.827 I  
 16.622 I  
 16.418 I  
 16.214 I  
 16.010 I  
 15.806 I  
 15.602 I  
 15.398 IO

ORIGINAL PAGE IS  
 OF POOR QUALITY

1.000 3.000 5.000 7.000 9.000 11.000 13.000 15.000 17.000

THE MONTHS


( X 10\*\* 0 )


## APPENDIX B

### TABLE OF CONTENTS


|                                                | Page |
|------------------------------------------------|------|
| Monthly Data Bases Program Flowchart . . . . . | B-1  |
| Monthly Data Bases Program Listing . . . . .   | B-6  |
| Subroutine SORTMD Flowchart. . . . .           | B-7  |
| Subroutine SORTMD Listing. . . . .             | B-12 |
| Subroutine DMEAN Flowchart . . . . .           | B-16 |
| Subroutine DMEAN Listing . . . . .             | B-18 |
| Subroutine TOTAL Flowchart . . . . .           | B-19 |
| Subroutine TOTAL Listing . . . . .             | B-20 |
| Subroutine ANP Flowchart . . . . .             | B-21 |
| Subroutine ANP Listing . . . . .               | B-23 |
| Subroutine PBAR Flowchart. . . . .             | B-24 |
| Subroutine PBAR Listing. . . . .               | B-26 |


*ESTIMATION OF MONTHLY MAX AND MIN TEMPERATURES  
AND PRECIPITATION FOR THE ENTIRE WHEAT PRODUCTION  
AREA OF THE PEOPLE'S REPUBLIC OF CHINA*






ORIGINAL PAGE IS  
OF POOR QUALITY





ORIGINAL PAGE IS  
OF POOR QUALITY



END of the MAIN PROGRAM

```


0001      EXTERNAL SIGN
0002      INTEGER VS (600),VY(600),VN(600),VD(600),VDMAX(600),VDMIN(600),
      @VP(600),STATN,YEAR,MONTH,DAY,DMAX,DMIN,PRECIP
0003      INTEGER N1(6),N2(2),N3(2),N4(3),N5(3),N6(3),N7(3)
0004      INTEGER STN1,YR1,BF
0005      REAL RMAX,RMIN
0006      DO 500 IN=9,10
0007      READ(IN,20)N1,N2,N3,N4,N5,N6,N7
0008      20 FORMAT(6A1,2A1,2A1,3A1,1X,3A1,3A1,3X,3A1)
0009      10 N=1
0010      50 READ(IN,40,END=499)STATN,YEAR,MONTH,DAY,DMAX,DMIN,PRECIP,RMAX,
      @RMIN
0011      VS(N)=STATN
0012      40 FORMAT(I6,3I2,1X,2I3,3X,I4,T14,2F3.0)
0013      VY(N)=YEAR
0014      VN(N)=MONTH
0015      VD(N)=DAY
0016      VP(N)=PRECIP
0017      IF((DMAX.EQ.C).AND.(SIGN(1.0,RMAX).EQ.-1.0))DMAX=999
0018      VDMAX(N)=DMAX
0019      IF((DMIN.EQ.C).AND.(SIGN(1.0,RMIN).EQ.-1.0))DMIN=999
0020      VDMIN(N)=DMIN
0021      M1=VN(N)
0022      M2=VD(N)
0023      M3=VS(N)
0024      M4=VS(1)
0025      IF(((M1.EQ.12).AND.(M2.EQ.31)).OR.((M3.NE.M4))) GO TO 200
0026      N=N+1
0027      GO TO 50
0028      200 CALL      SORTND(VS,VY,VN,VD,VDMAX,VDMIN ,N,STN1,YR1,VP,PIBAR)
0029      CALL ANP(STN1,YR1,VP,BF,N)
      CLEAR ALL STORAGE FOR NEXT STATION
0030      DO 300 I=1,N
0031      VS(I)=C
0032      VY(I)=C
0033      VN(I)=C
0034      VD(I)=C
0035      VP(I)=C
0036      300 CONTINUE
0037      GO TO 10
0038      499 REWIND IN
0039      500 CONTINUE
0040      STOP
0041      CEBLG ELBCHK
0042      END


```





B-6





ORIGINAL PAGE IS  
 OF POOR QUALITY


# Subroutine SORTMD











```

      C      THIS SUBROUTINE IS TO SORTED THE 12 MONTHS OBSERVATIONS FOR
      C      EACH STATION
0001      SUBROUTINE SORTMD(VS,VY,VM,VD,VDMAX,VDMIN,N,STN1,YR1,VP,P1BAR)
0002      REAL BM1,BM2,BM3,BM4,BM5,BM6,BM7,BM8,BM9,BM10,BM11,BM12,
      +BN1,BN2,BN3,BN4,BN5,BN6,BN7,BN8,BN9,BN10,BN11,BN12
0003      INTEGER DX1(50),DX2(50),DX3(50),DX4(50),DX5(50),DX6(50),DX7(50),
      +DX8(50),DX9(50),DX10(50),DX11(50),DX12(50),DN1(50),DN2(50),
      +DN3(50),DN4(50),DN5(50),DN6(50),DN7(50),DN8(50),DN9(50),DN10(50),
      +DN11(50),DN12(50),I1,I2,I3,I4,I5,I6,I7,I8,I9,I10,I11,I12,YR1,YR2,
      +YR3,YR4,YR5,YR6,YR7,YR8,YR9,YR10,YR11,YR12,STN1,STN2,STN3,STN4,
      +STN5,STN6,STN7,STN8,STN9,STN10,STN11,STN12
0004      INTEGER VS(600),VY(600),VM(600),VD(600),VDMAX(600),VDMIN(600),
      +JAN,FEB,MAR,APR,MAY,JUN,JUL,AUG,SEP,OCT,NOV,DEC,D1(50),D2(50),
      +D3(50),D4(50),D5(50),D6(50),D7(50),D8(50),D9(50),D10(50),D11(50),
      +D12(50),P1(50),P2(50),P3(50),P4(50),P5(50),P6(50),P7(50),P8(50),
      +P9(50),P10(50),P11(50),P12(50),VP(400),P1BAR
0005      INTEGER P1BAR,P2BAR,P3BAR,P4BAR,P5BAR,P6BAR,P7BAR,P8BAR,
      +P9BAR,P10BAR,P11BAR,P12BAR
0006      I1=0
0007      I2=0
0008      I3=0
0009      I4=0
0010      I5=0
0011      I6=0
0012      I7=0
0013      I8=0
0014      I9=0
0015      I10=0
0016      I11=0
0017      I12=0
0018      DO 100 I=1,N
0019      K=VM(I)
0020      GO TO(1,2,3,4,5,6,7,8,9,10,11,12),K
0021      1 I1=I1+1
0022      P1(I1)=VP(I)
0023      DX1(I1)=VDMAX(I)
0024      DN1(I1)=VDMIN(I)
0025      D1(I1)=VD(I)
0026      JAN=VM(I)
0027      YR1=VY(I)
0028      STN1=VS(I)
0029      GO TO 100
0030      2 I2=I2+1
0031      P2(I2)=VP(I)
0032      DX2(I2)=VDMAX(I)
0033      DN2(I2)=VDMIN(I)
0034      D2(I2)=VD(I)
0035      FEB=VM(I)
0036      YR2=VY(I)
0037      STN2=VS(I)
0038      GO TO 100
0039      3 I3=I3+1
0040      P3(I3)=VP(I)
0041      DX3(I3)=VDMAX(I)
0042      DN3(I3)=VDMIN(I)
0043      D3(I3)=VD(I)

```

B-12

ORIGINAL PAGE IS  
OF POOR QUALITY

B-13

```
0044      MAR=VM(I)
0045      YR3=VY(I)
0046      STN3=VS(I)
0047      GO TO 100
0048  4 I4=I4+1
0049      P4(I4)=VP(I)
0050      DX4(I4)=VDMAX(I)
0051      DN4(I4)=VDMIN(I)
0052      D4(I4)=VD(I)
0053      APR=VM(I)
0054      YR4=VY(I)
0055      STN4=VS(I)
0056      GO TO 100
0057  5 I5=I5+1
0058      P5(I5)=VP(I)
0059      DX5(I5)=VDMAX(I)
0060      DN5(I5)=VDMIN(I)
0061      D5(I5)=VD(I)
0062      MAY=VM(I)
0063      YR5=VY(I)
0064      STN5=VS(I)
0065      GO TO 100
0066  6 I6=I6+1
0067      P6(I6)=VP(I)
0068      DX6(I6)=VDMAX(I)
0069      DN6(I6)=VDMIN(I)
0070      D6(I6)=VD(I)
0071      JUN=VM(I)
0072      YR6=VY(I)
0073      STN6=VS(I)
0074      GO TO 100
0075  7 I7=I7+1
0076      P7(I7)=VP(I)
0077      DX7(I7)=VDMAX(I)
0078      DN7(I7)=VDMIN(I)
0079      D7(I7)=VD(I)
0080      JUL=VM(I)
0081      YR7=VY(I)
0082      STN7=VS(I)
0083      GO TO 100
0084  8 I8=I8+1
0085      P8(I8)=VP(I)
0086      DX8(I8)=VDMAX(I)
0087      DN8(I8)=VDMIN(I)
0088      AUG=VM(I)
0089      YR8=VY(I)
0090      D8(I8)=VD(I)
0091      STN8=VS(I)
0092      GO TO 100
0093  9 I9=I9+1
0094      P9(I9)=VP(I)
0095      DX9(I9)=VDMAX(I)
0096      DN9(I9)=VDMIN(I)
0097      D9(I9)=VD(I)
0098      SEP=VM(I)
0099      YR9=VY(I)
```

```

0100      STN9=VS(I)
0101      GO TO 100
0102      10 I10=I10+1
0103      P10(I10)=VP(I)
0104      DX10(I10)=VDMA>(I)
0105      DN10(I10)=VDMIN(I)
0106      O10(I10)=VD(I)
0107      OCT=VM(I)
0108      YR10=VY(I)
0109      STN10=VS(I)
0110      GO TO 100
0111      11 I11=I11+1
0112      P11(I11)=VP(I)
0113      DX11(I11)=VDMA>(I)
0114      DN11(I11)=VDMIN(I)
0115      O11(I11)=VD(I)
0116      NOV=VM(I)
0117      YR11=VY(I)
0118      STN11=VS(I)
0119      GO TO 100
0120      12 I12=I12+1
0121      P12(I12)=VP(I)
0122      DX12(I12)=VDMA>(I)
0123      DN12(I12)=VDMIN(I)
0124      O12(I12)=VD(I)
0125      DEC=VM(I)
0126      YR12=VY(I)
0127      STN12=VS(I)
0128      100 CONTINUE
0129      WRITE(6,50)
0130      50 FORMAT(20X,'THE MONTHLY MAX,AND MIN TEMP TABLE'/
@20X,'MONTH',5X,'MAX TEMP',10X,'MIN TEMP',10X,'PRECIP'/'
@20X,52(' '))
0131      CALL DMEAN(DX1,DN1,D1,JAN,YR1,STN1,I1,BN1,BN1)
0132      CALL DMEAN(DX2,DN2,D2,FEB,YR2,STN2,I2,BN2,BN2)
0133      CALL DMEAN(DX3,DN3,D3,MAR,YR3,STN3,I3,BN3,BN3)
0134      CALL DMEAN(DX4,DN4,D4,APR,YR4,STN4,I4,BN4,BN4)
0135      CALL DMEAN(DX5,DN5,D5,MAY,YR5,STN5,I5,BN5,BN5)
0136      CALL DMEAN(DX6,DN6,D6,JUN,YR6,STN6,I6,BN6,BN6)
0137      CALL DMEAN(DX7,DN7,D7,JUL,YR7,STN7,I7,BN7,BN7)
0138      CALL DMEAN(DX8,DN8,D8,AUG,YR8,STN8,I8,BN8,BN8)
0139      CALL DMEAN(DX9,DN9,D9,SEP,YR9,STN9,I9,BN9,BN9)
0140      CALL DMEAN(DX10,DN10,D10,OCT,YR10,STN10,I10,BN10,BN10)
0141      CALL DMEAN(DX11,DN11,D11,NOV,YR11,STN11,I11,BN11,BN11)
0142      CALL DMEAN(DX12,DN12,D12,DEC,YR12,STN12,I12,BN12,BN12)
0143      CALL PEAR(P1,I1,P1BAR)
0144      CALL PEAR(P2,I2,P2BAR)
0145      CALL PEAR(P3,I3,P3BAR)
0146      CALL PEAR(P4,I4,P4BAR)
0147      CALL PEAR(P5,I5,P5BAR)
0148      CALL PEAR(P6,I6,P6BAR)
0149      CALL PEAR(P7,I7,P7BAR)
0150      CALL PEAR(P8,I8,P8BAR)
0151      CALL PEAR(P9,I9,P9BAR)
0152      CALL PEAR(P10,I10,P10BAR)
0153      CALL PEAR(P11,I11,P11BAR)


```

B-14


ORIGINAL PAGE IS  
OF POOR QUALITY

```
0154      CALL PEAR(P12,I12,F12BAR)
0155      WRITE(6,52)STN1,YR1
0156      52 FORMAT(3X,'STATION #',I5,' IN 19',I2)
0157      WRITE(6,51) JAN,BM1,BN1,F1BAR,
        @FEB,BM2,BN2,P2BAR,
        @MAR,BM3,BN3,P3BAR,
        @APR,BM4,BN4,P4BAR,
        @MAY,BM5,BN5,P5BAR,
        @JUN,BM6,BN6,P6BAR,
        @JUL,BM7,BN7,P7BAR,
        @AUG,BM8,BN8,P8BAR,
        @SEP,BM9,BN9,P9BAR,
        @OCT,BM10,BN10,P10BAR,
        @NOV,BM11,BN11,P11BAR,
        @DEC,BM12,BN12,P12BAR
0158      51 FORMAT(2CX,I4,F12.2,10X,F10.2,10X,I8)
0159      RETURN
0160      DEBLC SLBCHK
0161      END
```

# SUBROUTINE MEAN




C-2




```
      C      THIS SUBROUTINE IS USE TO CALCULATE THE MONTHLY MAX,MIN TEMP
0001      SUBROUTINE DMEAN(DX,DN,D,MCN,YR,STN,N,MAXBAR,MINBAR)
0002      INTEGER DX(50),DN(50),MCN,YR,STN
0003      INTEGER D(50)
0004      REAL MAXBAR,MINBAR
0005      CALL TOTAL(N,SMA),DX,ICCLNT)
0006      EN=N-ICCLNT
0007      IF(EN.EQ.0.0)EN=1.0
0008      MAXBAR=SMA/EN
0009      CALL TOTAL(N,SMIN,DN,KCLNT)
0010      EM=N-KCLNT
0011      IF(EM.EQ.0.0)EM=1.0
0012      MINBAR=SMIN/EM
0013      RETURN
0014      DEBLC SLBCHK
0015      END
```


# SUBROUTINE TOTAL



```
0001      SUBROUTINE TOTAL(N,SLM,DZ,KNT)
0002      INTEGER DZ(50)
0003      KNT=C
0004      SLM=C.C
0005      DO 100 K=1,N
0006      IF(DZ(K).EQ.555)GO TO 200
0007      SLM=SLM+DZ(K)
0008      GO TO 100
0009      200 KNT=KNT+1
0010      100 CONTINUE
0011      RETURN
0012      DEBLG SLBCHK
0013      END
```


# SUBROUTINE AND






```
0001      C      THIS SUBROUTINE IS USE TO FIND THE ANN PRECIP CF EACH STATION
0002      SLBROTIME ANP(STATN, YEAR, VP, BP, N)
0003      INTEGER STATN, YEAR, VP(600), BP
0004      L=0
0005      ICOUNT=0
0006      DO 100 I=1, N
0007      IF(VP(I).EQ.5555) GO TO 200
0008      L=L+VP(I)
0009      GO TO 100
0010 200 ICOUNT=ICOUNT+1
0011 100 CONTINUE
0012      BP=L
0013      WRITE(6,30)N, ICOUNT, STATN, YEAR, BP
0014 30 FORMAT(3CX, 'TOTAL# OF OBSERVATIONS=', I8, /
0015      @3CX, '#OF MISSING OBSERVATIONS=', I8, /
0016      @3CX, 'FOR STATION', I8, ' IN THE YEAR ', I4,
0017      @' THE ANNUAL PRECIP IS ', I6)
0018      RETURN
0019      DEBLC SLBCHK
0020      END
```

# SUBROUTINE PBAR





ORIGINAL PAGE IS  
OF POOR QUALITY


```
0001      SUBROUTINE PBAR(PC,N,PAV)
0002      INTEGER PC(50),PAV
0003      IS=0
0004      DO 10 I=1,N
0005      IF(PC(I).EQ.9999) GC TC 10
0006      IS=IS+PC(I)
0007      10 CONTINUE
0008      EN=N
0009      PAV=IS
0010      RETURN
0011      DEBLG SLBCHK
0012      END
```


APPENDIX C  
TABLE OF CONTENTS

|                                                  | Page |
|--------------------------------------------------|------|
| Analysis of Variance Program Flowchart . . . . . | C-1  |
| Analysis of Variance Program Listing . . . . .   | C-8  |


ORIGINAL PAGE IS  
OF POOR QUALITY

# THE ANALYSIS OF VARIANCE








ORIGINAL PAGE IS  
OF POOR QUALITY




ORIGINAL PAGE IS  
OF POOR QUALITY







ORIGINAL PAGE IS  
OF POOR QUALITY



COMPILER OPTIONS - NAME= MAIN,OPT=00,LINECNT=55,SIZE=0000K,  
SOURCE,EBCDIC,NOLIST,NODECK,LOAD,MAP,NOEDIT,ID,XREF

```

C      *****
C      ***** THE AOV ANALYSIS *****
C      ***** HY *****
C      ***** BUDDY H JEUN *****
C      *****
ISN 0002 REAL X(100,100),T(100),G(100)
ISN 0003 INTEGER R(100)
ISN 0004 FTAB=6.93
ISN 0005 R(1)=12
ISN 0006 R(2)=12
ISN 0007 R(3)=12
ISN 0008 M=12
ISN 0009 N=3
ISN 0010 NG=3
ISN 0011 NO=0
ISN 0012 DO 100 I=1,NG
ISN 0013 NO=NO+R(I)
ISN 0014 100 CONTINUE
ISN 0015 DO 22 I=1,M
ISN 0016 READ(5,10)(X(I,J),J=1,N)
ISN 0017 10 FORMAT(3(F5.2,1X))
ISN 0018 22 CONTINUE
ISN 0019 WRITE(6,98)
ISN 0020 98 FORMAT('1',59X,'TA-LIEN(#54662)'/
@43X,'LOCKHEED',5X,'U.S.AIR FORCE',5X,'HONG KONG ROYAL'//
@43X,46(' '))
ISN 0021 DO 11 I=1,M
ISN 0022 WRITE(6,9)(X(I,J),J=1,N)
ISN 0023 9 FORMAT(43X,F10.2,5X,F12.2,5X,F13.3)
ISN 0024 11 CONTINUE
ISN 0025 SUM=0.0
ISN 0026 SUMSQ=0.0
ISN 0027 DO 200 J=1,NG
ISN 0028 M=R(J)
ISN 0029 DO 200 I=1,M
ISN 0030 SUM=SUM+X(I,J)
ISN 0031 SUMSQ=SUMSQ+X(I,J)**2
ISN 0032 200 CONTINUE
ISN 0033 WRITE(6,105) SUM,SUMSQ
ISN 0034 105 FORMAT(20X,'SUM=',F10.2,2X,'SUMSQ=',F10.2//)
C      NO IS THE # OF OBSERVATIONS
C      NG IS THE # OF GROUP
ISN 0035 ENO=NO
ISN 0036 SQSUM=SUM**2
ISN 0037 TSS=(SUMSQ-SQSUM/ENO)
C      TSS IS THE TOTAL SUM SQ OF OBSERVATIONS
ISN 0038 DO 300 J=1,NG
ISN 0039 T(J)=0.0
ISN 0040 L=R(J)
ISN 0041 DO 300 I=1,L
ISN 0042 T(J)=T(J)+X(I,J)

```

C-8

ORIGINAL PAGE IS  
OF POOR QUALITY

```

C-7
ISN 0043      300 CONTINUE
ISN 0044      DO 12 I=1,NG
ISN 0045      WRITE(6,106)I,T(I)
ISN 0046      106 FORMAT(20X,'SUM OF GROUP(' ,I3,' ) IS ',F10.2/)
ISN 0047      12 CONTINUE
ISN 0048      SUMGSQ=0.0
ISN 0049      DO 400 I=1 ,NG
ISN 0050      F=R(I)
ISN 0051      SUMGSQ=SUMGSQ+T(I)**2/E
ISN 0052      400 CONTINUE
ISN 0053      BSS=(SUMGSQ-SQSUM/ENO)
ISN 0054      C   RSS IS THE SUMSQ BETWEEN GROUP
ISN 0055      WSS=TSS-BSS
ISN 0056      C   NGM1=NG-1
ISN 0057      C   NGM1 IS THE DEGREE OF FREEDOM FOR THE WITHIN THE GROUP
ISN 0058      C   NOMNG=NO-NG
ISN 0059      C   NOMNG IS THE DEGREE OF FREEDOM FOR WITHIN THE GROUP
ISN 0060      C   NOM1=NO-1
ISN 0061      C   NOM1 IS THE DEGREE OF FREEDOM FOT THE TOTAL OBSERVATIONS
ISN 0062      BMS=BSS/FLOAT(NGM1)
ISN 0063      WMS=WSS/FLOAT(NOMNG)
ISN 0064      FCAL=BMS/WMS
ISN 0065      WRITE(6,20)
ISN 0066      20 FORMAT (55X,'ANALYSIS OF VARIANCE'//
ISN 0067      +50X,'DEGREE OF ',7X,'SUM OF '/
ISN 0068      +26X,'SOURCE OF VARIATION',5X,'FREEDOM',10X,'SQUARE',5X,..
ISN 0069      +1MEAN SQUARE',5X,'F RATIO'/
ISN 0070      +26X,75(' '))//
ISN 0071      WRITE(6,Z1)NGM1,HSS,BMS,
ISN 0072      +NOMNG,WSS,WMS,FCAL, .....
ISN 0073      +NOM1,TSS
ISN 0074      21 FORMAT(25X,'BETWEEN GROUP',12X,I3,9X,F13.3,5X,F10.3/
ISN 0075      +25X,'WITHIN GROUPS',12X,I3,9X,F13.3,5X,F10.3,5X,F10.3/
ISN 0076      +25X,'TOTAL',20X,I3,9X,F13.3/
ISN 0077      +25X,77(' '))////////
ISN 0078      IF(FCAL.GE.FTAB) GO TO 101
ISN 0079      WRITE(6,23)
ISN 0080      23 FORMAT(30X,'SINCE FCAL<FTAB,SO ACCEPT THE HYPOTHESE!')
ISN 0081      GO TO 600
ISN 0082      101 WRITE(6,24)
ISN 0083      24 FORMAT(30X,'SINCE FCAL>FTAB,SO REJECT THE HYPOTHESE!')
ISN 0084      600 STOP
ISN 0085      END

```