

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.



Technical Memorandum 79599

Interplanetary Magnetic Holes: Theory


L. F. Burlaga

J. F. Lemaire

July 1978

National Aeronautics and
Space Administration

Goddard Space Flight Center
Greenbelt, Maryland 20771



(NASA-TM-79599) INTERPLANETARY MAGNETIC
HOLES: THEORY (NASA) 24 p HC A02/MF A01

N78-30023

CSCI 038

Unclas
G3/90 29031

**INTERPLANETARY MAGNETIC HOLES:
THEORY**

**L. F. Burlaga
NASA/Goddard Space Flight Center
Laboratory for Extraterrestrial Physics
Greenbelt, MD 20771**

**ORIGINAL PAGE IS
OF POOR QUALITY**

**J. F. Lemaire
Institut D'Aeronomie Spatiale De Belgique
3 Avenue Circulaire
B-1180 Brussels
Belgium**

TO BE SUBMITTED TO: Journal of Geophysical Research

ABSTRACT

Magnetic holes in the interplanetary medium are explained as stationary, non-propagating, equilibrium structures in which there are field-aligned enhancements of the plasma density and/or temperature. Magnetic anti-holes are considered to be associated with depressions in the plasma pressure. In this model, the observed changes in the magnetic field intensity and direction are due to diamagnetic currents that are carried by ions which drift in a sheath as the result of gradients in the magnetic field and in the plasma pressure within the sheath. The thickness of the sheaths that we consider is approximately a few ion Larmor radii. An electric field is normal to the magnetic field in the sheath. Solutions of Vlasov's equation and Maxwell's equations are presented which account for several types of magnetic holes, including "null-sheets", that have been observed.

I. INTRODUCTION

In an analysis of high resolution (12.5 samples/s) magnetic field measurements obtained by Explorer 43 in the period March 18 to April 9, 1971, Turner et al. (1977) found that most low-field intensity regions ($B \leq 1\gamma$) were isolated and very thin ($\approx 10^4$ km or $\approx 100 R_L$, where R_L is the proton Larmor radius). They called these regions magnetic holes. Observations of such structures with two spacecraft (Fitzenreiter and Burlaga, 1977) show that the extent of magnetic holes is more than $\approx 10^5$ km. Magnetic holes form a class of current sheets.

Three types of magnetic holes were identified by Turner et al. (1977) and analyzed by Fitzenreiter and Burlaga (1977): 1) holes with no change in the magnetic field direction across the current sheet; 2) holes with a reversal in magnetic field direction, but no rotation of the field vector across the current sheet; and 3) holes in which there is a rotation of the magnetic field vector across the current sheet. Turner et al. also exhibited current sheets in which the magnetic field intensity increased with no change in direction ("anti-holes"). The purpose of this paper is to apply the theory presented previously by Lemaire and Burlaga (1976) to explain the three types of magnetic holes and to explain anti-holes.

For the calculations, we make the following assumptions: 1) magnetic holes are stationary inhomogeneities convected with the solar wind speed. 2) They are bounded by thin current sheaths and the radius of curvature of these sheaths is very large compared to the thickness. As a consequence, we consider the geometry of the sheath to be planar, with a normal in the \hat{z} direction. 3) The physical variables are functions only of z . 4) The magnetic field, $\underline{B}(z)$, is normal to the \hat{z} direction as for tangential

discontinuities. 5) In the frame of reference moving with the magnetic hole there is an electric field $\underline{E}(z)$ along \hat{z} i.e., $E_x = E_y = 0$: this implies that there is no net plasma flow across the boundary layer. 6) The plasma is quasi-neutral and consists of electrons and protons. Multi-ionic plasmas have been considered but will not be discussed in this paper. 7) The plasma is collisionless. 8) The diamagnetic currents are carried only by the protons (the electron velocity distribution is assumed to be isotropic). 9) The plasma is in thermal equilibrium on both sides of the current sheath and the bulk velocities at $z = \pm\infty$ are both equal to zero in a frame of reference comoving with the solar wind speed.

The basic equations are Maxwell's and Vlasov's equations.

We obtained solutions of these equations which describe magnetic holes using the method described by Lemaire and Burlaga (1976). In all of the examples discussed below, we assume that the ambient magnetic intensity and particle density on each side of the current sheath are 5γ (nT) and 5 protons or electrons/cm³ respectively. For simplicity we also assume that the temperatures of the electrons and ions are the same on both sides of the layer at $z = \pm\infty$: $T_e = T_i = 7.5 \times 10^4$ K; calculations with $T_e \neq T_i$ and with different temperatures on both sides have also been made but are not discussed in this paper. This complicates the structure of the layer but does not alter the basic physical effects, which are our principal interest for the present article. The following sections present solutions to the problems that we have posed above, and they include heuristic discussions, based on particle orbit theory, aimed at identifying the basic physical processes in magnetic holes.

II. MAGNETIC HOLES WITH NO CHANGE IN \hat{B} : $\Delta\omega = 0$

Let us consider magnetic holes in which the direction of magnetic field remains along \hat{x} : $\underline{B} = B(z) \hat{x}$. In particular, consider the case where $B = 1\gamma$ (nT) in the middle of the hole, together with the other boundary conditions discussed in Section I. A self-consistent solution of Vlasov's equation and Maxwell's equations, obtained by the method of Lemaire and Burlaga (1976) is shown in Figure 1.

The current in Figure 1 reaches a maximum at the two points where $|\frac{dB}{dz}|$ is largest. This follows directly from

$$\mu_0 \underline{J} = \nabla \times \underline{B} = \hat{y} \frac{dB}{dz} = \hat{y} \frac{dB}{dz}.$$

Note that \underline{J} flows normal to \underline{B} and \hat{z} in this current sheet. $|\underline{J}|$ is zero in the middle of the magnetic hole because there $\frac{dB}{dz}$ goes to zero as J_y changes from negative values to positive values.

In the first orbit approximation \underline{J} is actually the sum of two currents, the gradient drift current

$$J_D = - \frac{W_{\perp}}{B^2} \frac{dB}{dz}$$

(where $W_{\perp} = \sum \frac{1}{2} m v_{\perp}^2 = Nk (T_{i\perp} + T_{e\perp})$ which results from the gradient drift of protons in the inhomogeneous field, and the magnetization current

$$J_M = - \frac{d}{dz} \left(\frac{W_{\perp}}{B} \right).$$

(There is no current due to \underline{E} , since electrons and protons drift in the same direction and with the same speed due to $\underline{E} \times \underline{B}$.) Setting $\mu_0 (\underline{J}_D + \underline{J}_M) = \nabla \times \underline{B}$ gives

$$\frac{B^2}{2\mu_0} + Nk (T_{i\perp} + T_{e\perp}) = \text{constant}, \quad (1)$$

which shows that the pressure is constant across the current sheet, in agreement with our assumption of equilibrium. The increase in N coincident with the decrease in B shown in Figure 1 is a consequence of the constant pressure condition, (1), since for the case considered the kinetic temperature is nearly constant across the sheath.

In the example illustrated in Figure 1, we assumed that the kinetic pressure enhancement was due to a larger thermal proton and electron density inside the magnetic hole. But a similar structure would be obtained when the excess of particle pressure is due to a localized beam of suprathermal electrons or ions spiraling in an interplanetary magnetic flux tube. High intensity electrons spikes or protons bursts events sporadically detected in the interplanetary medium can produce such an excess kinetic pressure in an otherwise uniform solar wind plasma. Correlative studies would however be necessary to verify if some of the observed magnetic holes can be associated with these suprathermal particle beams.

The electric field E_z in Figure 1 preserves local quasi-neutrality in the plasma. Its value is largest where $|\frac{dN}{dz}|$ is largest. This can be understood heuristically as follows. The electron velocity distribution is assumed to be isotropic. This implies that the bulk velocity of the electrons is zero and that their pressure tensor is isotropic with $p_e = NkT_e$. The momentum equation for the electrons is therefore given by

$$\frac{dp_e}{dz} = -N|e|E_z = kT_e \frac{dN}{dz} + N \frac{dkT_e}{dz}$$

which shows that the charge separation electric field is indeed proportional to $|dN/dz|$ when $dT_e/dz \approx 0$; this is the case for the solution in Figure 1. If the temperature changed and N were everywhere constant, the same argument would apply and the $|E_z|$ needed to maintain equilibrium in this case would be greatest where $|dT/dz|$ was greatest. In general both $N(z)$ and $T(z)$ can vary and they need not be in phase; in this case $E_z(z)$ could be complicated, but the physical processes would be basically the same as in the simpler cases just described.

The first inflection point in $N(z)$ in Figure 1 is indicated by a dashed line; note the small accumulation of positive charge at a distance approximately R_{L2} (the proton Larmor radius on side 2) to the right of this point and the small accumulation of negative charge at $\approx R_{L1}$ to the left of this point. Since $B_1 > B_2$, $R_{L1} < R_{L2}$ and the peak of negative excess charge is closer to the inflection point of N than the peak of positive excess charge. This accounts for the asymmetry in the $E_z(z)$ profile in Figure 1.

A charge separation $\Delta n/N$ is required to support E_z . Poisson's equation gives $|e| \Delta n = \frac{dE_z}{dz}$ or $\Delta n \propto -\frac{d^2N}{dz^2}$, indicating that there will be an accumulation of negative charge in the region where E_z is decreasing most rapidly (i.e., where the dN/dz is increasing with z) and an accumulation of positive charge where E_z is increasing most rapidly (i.e., where dN/dz is decreasing with z). Such electrostatic double layers generally appear at the surface of a magnetic hole having an excess kinetic plasma pressure. The excess of positive charges at the edges of a dense plasma region results from the larger gyroradius of the ions compared to the gyroradius of the electrons. Indeed, the positive charges are deflected at a larger distance from the interior of the density (or pressure) enhancement than

the lighter negative charges. On the contrary an excess of negative charges is expected to accumulate at the edge of a plasma depression. Indeed, because of their small mass-to-charge ratio, the electrons tend to be reflected before the ions. This builds up a negative surface charge on the low density side and a positive charge accumulation toward the high density side (see Figure 2a).

Willis (1971, 1975) has discussed the charge separation electric field in a Ferraro-type magnetopause boundary layer, assuming that there is no plasma in the magnetosphere (i.e., on the low density side). In this extreme case when the density is strictly equal to zero outside the plasma region, only the positive charge accumulation is present at the edge of the enhancement (see Figure 2b). This "single-layer" structure is also described by the kinetic theory of Lemaire and Burlaga (1976) when the plasma density is forced to become equal to zero at $z = +\infty$.

ORIGINAL PAGE IS
OF POOR QUALITY

III. HOLES WITH A REVERSAL IN FIELD DIRECTION WITHOUT ROTATION ($\Delta\omega = 180^\circ$)

Let us again consider the case where $\underline{B} = B(z) \hat{x}$, but now we allow $B(z)$ to vary monotonically from $B_1 > 0$ to $B_2 < 0$. Clearly, $B(z)$ must go through zero at some point in the magnetic hole in this case. Note that "neutral sheets" or "null-sheets" ideally correspond to this special case of magnetic field reversal without rotation. As before, let us take $B_1 = 5\gamma$, $B_2 = -5\gamma$, $N_e = N_p = 5/\text{cm}^3$ and $T_i = T_e = \text{constant} = 7.5 \times 10^{40} \text{K}$ on each side of the current sheet.

Figure 3 shows one possible solution of Vlasov's equation and Maxwell's equations for the case considered. One sees that there is a single current peak, which is maximum in the center of the diamagnetic current layer, where $|B| = 0$. The density rises to a maximum at the center of the hole, as required by the pressure balance condition for equilibrium. The charge separation electric field is maximum in each of the two regions where N changes most rapidly, as required to maintain approximate local quasi-neutrality of the plasma. Each electric field region is maintained by a concentration of positive charge on one side of the maximum of $|E|$ and a concentration of negative charge on the other side; note that the positive charge concentrations overlap. Although the appearance of the magnetic hole in this case is rather different from that in Section II, the basic physical processes are the same in both cases.

Figure 4 shows another solution of the Vlasov-Maxwell equations, again for $\underline{B} = B(z) \hat{x}$ and for the same boundary conditions that were used in the preceding example. Although Figures 3 and 4 are solutions for the same boundary value problem, the solution in Figure 4 is rather

different from that in Figure 3. This illustrates the well-known point that boundary conditions do not specify a unique solution to the Vlasov-Maxwell equations. The principal difference in this case is that Figure 4 shows a double layer current distribution while Figure 3 shows a single peaked current distribution. The current distribution in Figure 4 resembles that in Figure 1 in being double-peaked, and it resembles that in Figure 3 in having a non-zero current where B is a minimum.

IV. MAGNETIC HOLES WITH A ROTATION OF \underline{B}

Consider the case in which \underline{B} rotates 180° from one side of a magnetic hole to the other, and consider the same boundary conditions that were used in our previous examples. A solution of the Vlasov-Maxwell equations obtained by our numerical program is shown in Figure 5. The magnetic field intensity profile in this case differs qualitatively from those considered above in that there is now a secondary maximum in B where in previous examples B was a minimum. The reason for the secondary maximum is basically geometrical, as can be seen by considering the rotation of \underline{B} in the current sheet (see the insert in Figure 5). When \underline{B} rotates in a plane, the current has a component parallel to the local magnetic field direction as well as a component perpendicular to \underline{B} :

$$\mu_0 \underline{J} = \hat{y} \frac{dB_x}{dz} - \hat{x} \frac{dB_y}{dz} = (\hat{z} \times \hat{B}) \frac{dB}{dz} + \underline{B} \frac{d\omega}{dz}$$

where $B_x = B \sin \omega$ and $B_y = B \cos \omega$. Therefore if the magnetic field rotates without changing intensity (i.e., $dB/dz = 0$) the diamagnetic current \underline{J} is field aligned and proportional to the rate of rotation $d\omega/dz$. In Figure 5, the peaks in \underline{J} occur near the points where the direction, ω , changes most rapidly (and where $\frac{dB}{dz}$ happens to be zero, as can be seen in the insert in Figure 5). \underline{J} is also non-zero at the secondary maximum of B even though $\frac{dB}{dz} = 0$ there, because the direction of \underline{B} is changing ($\frac{d\omega}{dz} \neq 0$) at that point.

The density profile in Figure 5 is anticorrelated with $B(z)$ as required by the constant pressure condition. The electric field is again extremal at the extrema of dN/dz for the same reasons given

earlier. The charge separation profile shows a positive and negative peak beside each extremum of E_z , as required to maintain E_z , and the charge peaks are separated from the peaks in E_z by $\approx 1 R_L$.

V. ANTIHOLES

ORIGINAL PAGE IS
OF POOR QUALITY

Figure 6 shows a solution of the Vlasov-Maxwell's equations for a linear "anti-hole" for the same boundary conditions that we have used in previous examples. By definition, B is enhanced in some region with dimensions on the order of several ion Larmor radii or more. Since $\underline{B} = B \hat{z}$, $|\underline{J}|$ is maximum where $\frac{dB}{dz}$ is extremum. We have assumed $T = \text{constant}$, so N is anticorrelated with B as required by the pressure balance condition. E_z is again extremum where $\frac{dN}{dz}$ is extremum, and each peak in E is maintained by a charge separation over a distance $\approx \text{few } R_L$. Thus, this type of anti-hole is physically equivalent to the type of magnetic hole described in Figure 1.

VI. CONCLUSIONS

Magnetic holes are considered here as kinetic scale plasma inhomogeneities convected with the solar wind speed. These localized density and/or temperature enhancements are assumed to be sheet-like structures which are stationary in the solar wind flow. The plasma in the magnetic hole is separated from the background solar wind by sheaths whose thicknesses are only a few proton gyroradii. Diamagnetic currents flow in these sheaths. In the kinetic model presented here to describe the structure of these sheaths, we have assumed that the radius of curvature of the surface is much larger than the thickness of the current sheath itself. Neglecting Coulomb collisions and other irreversible processes we have found, by using the models of Lemaire and Burlaga (1976), a variety of magnetic field structures similar to those observed in solar wind magnetic holes and anti-holes. Even null sheets, in which the magnetic field reverses with $|\vec{B}|$ going strictly through zero, can be modeled by the theory presented above (e.g., case III).

There are a number of other factors that might be considered in constructing more refined models of magnetic holes. Coulomb collisions although infrequent would tend to broaden the holes. In some cases the ion drift speed might exceed the Alfvén speed and cause local instabilities. Variations in both ion density and temperature should be considered.

The effects of alpha particle drifts and curvature of the current sheet might also be significant in some cases. These factors are essentially complications, however. The aim of this paper has been to explain the basic mechanisms

ORIGINAL PAGE IS
OF POOR QUALITY

responsible for magnetic holes. It will be appropriate to consider the other factors when high resolution plasma observations are available which give $n(z)$, $T(z)$, etc., in the magnetic holes.

ACKNOWLEDGMENTS

One of us (J. L.) was supported by the University of Maryland and wishes to thank N. F. Ness and K. W. Ogilvie for their hospitality and support at the Laboratory for Extraterrestrial Physics.

ORIGINAL PAGE IS
OF POOR QUALITY

REFERENCES

- Fitzenreiter, R. and L. F. Burlaga, Structure of Current Sheets in
Magnetic Holes at 1 AU, submitted to J. Geophys. Res., 1977.
- Lemaire, J. and Burlaga, L. F., Diamagnetic boundary layers: A kinetic
model, Astrophys. and Space Sci., 45, 303-325, 1976.
- Turner, J. M., Burlaga, L. F., Ness, N. F., and Lemaire, J., Magnetic
holes in the solar wind, J. Geophys. Res., 82, 1921-1924, 1977.
- Willis, D. M., Structure of the magnetopause, Rev. Geophys. Space Phys.,
9, 953-985, 1971.
- Willis, D. M., Microstructure of the magnetopause, Geophys. J. R.
Astro Soc., 41, 355-389, 1975.

FIGURE CAPTIONS

- Figure 1 Model of an isothermal magnetic hole in which $|\underline{B}|$ decreases to 1 without changing direction and then increases symmetrically to its initial value.
- Figure 2 An illustration of the charge accumulation which produces an electric field in a magnetic hole and which determines the thickness of a magnetic hole. Case a) shows a low density on one side and a high density on the other side of a sheath. Case b) shows zero density on one side of the sheath.
- Figure 3 Model of an isothermal magnetic hole in which the direction of \underline{B} reverses without a rotation of \hat{B} .
- Figure 4 Another isothermal magnetic hole in which the direction of \underline{B} reverses without a rotation of \hat{B} . Although the boundary conditions are the same as for the hole in Figure 3, the structures of the two holes are very different, illustrating the non-uniqueness of solutions of Vlasov's equation.
- Figure 5 Model of an isothermal magnetic hole in which the direction of \underline{B} changes by a rotation of \hat{B} through 180° . In this case there is a component of current along \underline{B} due to the rotation as well as a component of current normal to \underline{B} due to the gradients.
- Figure 6 Model of an isothermal anti-hole. Compare with the hole in Figure 1.




Figure 1





Figure 2

$$\Delta\omega = 180^\circ$$

$$T_i = T_e = 7.5 \times 10^4 \text{ }^\circ\text{K}$$




Figure 3

$$\Delta\omega = 180^\circ$$

$$T_i = T_e = 7.5 \times 10^4 \text{ }^\circ\text{K}$$




Figure 4

ORIGINAL PAGE IS
POOR QUALITY

$$\Delta\omega = 180^\circ$$

$$T_i = T_e = 7.5 \times 10^4 \text{ }^\circ\text{K}$$




Figure 5




Figure 6

ORIGINAL PAGE IS
 OF POOR QUALITY

1. Report No. 79599	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Interplanetary Magnetic Holes: Theory		5. Report Date July 1978	
		6. Performing Organization Code	
7. Author(s) L. F. Burlaga and J. F. Lemaire		8. Performing Organization Report No.	
9. Performing Organization Name and Address NASA/GSFC Laboratory for Extraterrestrial Physics Interplanetary Physics Branch, Code 692 Greenbelt, MD 20771		10. Work Unit No.	
		11. Contract or Grant No.	
		13. Type of Report and Period Covered Technical Memorandum	
12. Sponsoring Agency Name and Address		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract <p>Magnetic holes in the interplanetary medium are explained as stationary, non-propagating, equilibrium structures in which there are field-aligned enhancements of the plasma density and/or temperature. Magnetic anti-holes are considered to be associated with depressions in the plasma pressure. In this model, the observed changes in the magnetic field intensity and direction are due to diamagnetic currents that are carried by ions which drift in a sheath as the result of gradients in the magnetic field and in the plasma pressure within the sheath. The thickness of the sheaths that we consider is approximately a few ion Larmor radii. An electric field is normal to the magnetic field in the sheath. Solutions of Vlasov's equation and Maxwell's equations are presented which account for several types of magnetic holes, including "null-sheets", that have been observed.</p>			
17. Key Words (Selected by Author(s)) Interplanetary Plasma, current sheets		18. Distribution Statement	
19. Security Classif. (of this report) U	20. Security Classif. (of this page) U	21. No. of Pages 23	22. Price*