

N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED
IN THE INTEREST OF MAKING AVAILABLE AS MUCH
INFORMATION AS POSSIBLE

JPL PUBLICATION 80-23

(NASA-CR-163284) SUBMILLIMETER, MILLIMETER,
AND MICROWAVE SPECTRAL LINE CATALOGUE (Jet
Propulsion Lab.) 141 p HC A07/MF A01

N80-26408

CSCCL 07D

Unclas

GJ/25

23615

Submillimeter, Millimeter, and Microwave Spectral Line Catalogue

R. L. Poynter
H. M. Pickett

June 1, 1980

National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute Of Technology
Pasadena, California



TECHNICAL REPORT STANDARD TITLE PAGE

1. Report No. JPL Pub. 80-23	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Submillimeter, Millimeter, and Microwave Spectral Line Catalogue		5. Report Date June 1, 1980	
		6. Performing Organization Code	
7. Author(s) R.L. Poynter and H.M. Pickett		8. Performing Organization Report No.	
9. Performing Organization Name and Address JET PROPULSION LABORATORY California Institute of Technology 4800 Oak Grove Drive Pasadena, California 91103		10. Work Unit No.	
		11. Contract or Grant No. NAS 7-100	
		13. Type of Report and Period Covered JPL Publication	
12. Sponsoring Agency Name and Address NATIONAL AERONAUTICS AND SPACE ADMINISTRATION Washington, D.C. 20546		14. Sponsoring Agency Code	
15. Supplementary Notes			
<p>16. Abstract</p> <p>This report describes a computer accessible catalogue of submillimeter, millimeter, and microwave spectral lines in the frequency range between 0 and 3000 GHz (i.e., wavelengths longer than 100 m). The catalogue can be used as a planning guide or as an aid in the identification and analysis of observed spectral lines. The information listed for each spectral line includes the frequency and its estimated error, the intensity, lower state energy, and quantum number assignment.</p> <p>The catalogue has been constructed by using theoretical least squares fits of published spectral lines to accepted molecular models. The associated predictions and their estimated errors are based upon the resultant fitted parameters and their covariances. Future versions of this catalogue will add more atoms and molecules and update the present listings (122 species) as new data appear.</p> <p>The catalogue is available from the authors as a magnetic tape recorded in card images and as a set of microfiche records.</p>			
17. Key Words (Selected by Author(s)) Physical Chemistry Geophysics Atomic and Molecular Physics Astronomy		18. Distribution Statement Unclassified - Unlimited	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 142	22. Price

JPL PUBLICATION 80-23

Submillimeter, Millimeter, and Microwave Spectral Line Catalogue

R. L. Poynter
H. M. Pickett

June 1, 1980

National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute Of Technology
Pasadena, California

The research described in this publication was carried out by the Jet Propulsion Laboratory, California Institute of Technology, under NASA Contract No. NAS7-100.

ABSTRACT

This report describes a computer accessible catalogue of submillimeter, millimeter, and microwave spectral lines in the frequency range between 0 and 3000 GHz (i.e., wavelengths longer than 100 μm). The catalogue can be used as a planning guide or as an aid in the identification and analysis of observed spectral lines. The information listed for each spectral line includes the frequency and its estimated error, the intensity, lower state energy, and quantum number assignment.

The catalogue has been constructed by using theoretical least squares fits of published spectral lines to accepted molecular models. The associated predictions and their estimated errors are based upon the resultant fitted parameters and their covariances. Future versions of this catalogue will add more atoms and molecules and update the present listings (122 species) as new data appear.

The catalogue is available from the authors as a magnetic tape recorded in card images and as a set of microfiche records.

"Page missing from available version"

CONTENTS

I.	Introduction	1
II.	Format of data tape	2
III.	Intensity units and conversions	3
IV.	General comments on precision	5
V.	Format of quantum numbers	7
VI.	Documentation by species (listed in order of species tag)	11

Tables

V-1.	Standard quantum number formats.	9
V-2.	Special-case quantum number formats.	10

I. Introduction

This report describes a computer accessible catalogue of submillimeter, millimeter, and microwave spectral lines in the frequency range between 0 and 3000 GHz (i.e. wavelengths longer than $100\mu\text{m}$). The catalogue is intended to be used as a guide in the planning of spectral line observations and as a reference which can facilitate identification and analysis of observed spectral lines. The selection of lines for the catalogue is based on the projected needs of astronomers and atmospheric scientists.

The catalogue is constructed using theoretical least squares fits and predictions based on spectral lines, mostly obtained from the literature. In subsequent versions of the catalogue, more molecules will be added and existing molecular listings will be updated as new data appears.

The catalogue is available as a magnetic data tape recorded in card images, with one card image per spectral line. The format of the data is given in Section II. Section III gives conversions between different measures of spectral line intensity. General comments on the precision of the spectral line positions and intensities are given in Section IV, while species specific comments are reserved for Section VI. Section V gives the format of quantum numbers as they appear in the catalogue. Documentation for each molecular or atomic species is listed in Section VI in order of the "species tag". This tag is a six digit number in which the three most significant digits represent the mass number of the molecule or atom and the last three digits are an accession number for the given mass. Usually there is a separate tag for each vibration-electronic state of a particular molecule.

II. Format of Data Tape

The catalogue data tape is composed of 80 character card images, with one card image per spectral line. The format of each card image is given below:

FREQ , ERR , LGINT, DR , ELOW , GUP, TAG, QNFORM, QN
(F13.4, F8.4 , F8.4 , I2 , F10.4 , I3 , I7 , I4 , I2I2, I1)

- FREQ: Frequency of the line in MHz.
- ERR: Estimated or experimental error of FREQ in MHz.
- LGINT: Base 10 logarithm of the integrated intensity in units of $\text{nm}^2 \text{MHz}$ at 300K. (See Section III for conversions to other units.)
- DR: Degrees of freedom in the rotational partition function (0 for atoms, 2 for linear molecules, 3 for non-linear molecules).
- ELOW: Lower state energy in cm^{-1} .
- GUP: Upper state degeneracy.
- TAG: Species tag or molecular identifier.
A negative value flags that the line frequency has been measured in the laboratory. The absolute value of TAG is then the species tag and ERR is the reported experimental error. The three most significant digits of the species tag are coded as the mass number of the species as explained above.
- QNFORM: Identifies the format of the quantum numbers given in the field QN. These quantum number formats are given in Section V.
- QN: Quantum numbers for the upper and lower states coded according to QNFORM.

III. Intensity Units and Conversions

The units of intensity given in the catalogue, $\text{nm}^2 \text{ MHz}$, are based on the integral of the absorption cross section over the spectral line-shape. The value of the intensity is calculated for 300K and is directly comparable with the common infrared intensity unit of $\text{cm}^{-1} / (\text{molecule}/\text{cm}^2)$. The latter is obtainable by dividing the catalogue intensity by $2.9979 \times 10^{+18}$.

The line intensity in the catalogue I_{ba} , is obtained from

$$I_{ba} = \frac{8\pi^3}{3hc} \nu_{ba} x_{S_{ba}} \mu_x^2 \left(e^{-E''/kT} - e^{-E'/kT} \right) / Q_{rs} \quad (1)$$

in which ν_{ba} is the line frequency, $x_{S_{ba}}$ is the line strength, μ_x is the dipole moment along the molecular axis x , E'' and E' are the lower and upper state energies respectively, and Q_{rs} is the rotation-spin partition function (using the same zero of energy as E' and E''). Care must be taken to assure that $x_{S_{ba}}$ and Q_{rs} are determined with same state degeneracies. Note that with this definition the intensities are defined with respect to the total concentration of the vibration electronic state of the species.

The temperature dependence of I_{ba} can be obtained from the value of I_{ba} at $T=300\text{K}$ by the use of Eq. (1) once the temperature dependence of Q_{rs} is known. For linear molecules Q_{rs} is proportional to T in the limit where the energy spacings are small compared to kT . For non-linear molecules Q_{rs} is proportional to $T^{3/2}$ in the same limit.

Peak intensities of collision broadened lines can be obtained from I_{ba} from the relation

$$a_{\text{max}} = \frac{I_{ba}}{\Delta\nu} \left(\frac{300\text{K}}{T} \right) \times 102.46 \quad (2)$$

in which $\Delta\nu$ is the half-width at half-height in MHz at 1 torr partial pressure of absorber, I_{ba} is in units of $\text{nm}^2 \text{ MHz}$, and α_{max} is in units of cm^{-1} . The corresponding value of α_{max} in the thermal Doppler limit is

$$\alpha_{\text{max}} = \frac{I_{ba}}{\nu_{ba}} p \left(\frac{T}{300\text{K}} \right)^{-3/2} \sqrt{\frac{m}{28}} \times 1.2898 \times 10^8 \quad (3)$$

in which p is the partial pressure of absorber, ν_{ba} is the line frequency in MHz, and m is the mass of the absorber.

The absorption cross-section of an interstellar absorber integrated over a 1 km/s velocity interval is

$$\sigma_{ba} = \frac{I_{ba}}{\nu_{ba}} \times 2.9979 \times 10^{-9} \text{ cm}^2. \quad (4)$$

The inverse of σ_{ba} is the column density per unit optical depth in the same 1 km/s velocity interval.

The average spontaneous emission rate from the upper states into the lower states is

$$A_{ba} = I_{ba} \nu_{ba} \frac{Q_{rs}}{g'} \left(e^{-E''/kT} - e^{-E'/kT} \right)^{-1} \times 2.7964 \times 10^{-16} \text{ sec}^{-1} \quad (5)$$

in which g' is the degeneracy of the upper state. The value of g' is listed as part of the spectral line information in the catalogue, and the value of Q_{rs} at 300K is listed in Section VI.

IV. General Comments on Precision

The expected errors of the frequency as listed in the catalog are usually based on a propagation of errors estimated from a least squares fit of the observed frequencies to a model Hamiltonian, using the following equation:

$$\epsilon_n^2 = \sum_{k,\ell} \frac{\partial v_n}{\partial p_k} \frac{\partial v_n}{\partial p_\ell} V_{k\ell} \quad (6)$$

in which ϵ_n is the estimated error of frequency v_n and $V_{k\ell}$ is an element of the least squares variance - covariance matrix for the parameters p_k . This variance - covariance matrix is determined from the observed lines by

$$(V^{-1})_{k\ell} = \sum_m \frac{\partial v_m}{\partial p_k} \frac{\partial v_m}{\partial p_\ell} \epsilon_m^{-2} \quad (7)$$

in which the summation over m is over the experimental lines using experimental uncertainties, ϵ_m . The diagonal elements of V are the squares of the parameter uncertainties and the off-diagonal elements of V are products of the parameter uncertainties and correlation coefficients.

Experimental uncertainties in the literature vary from 1.6 σ estimates to 3 σ estimates and are more usually "guess-timates". Many times the statistical nature of the uncertainties are not stated. Therefore, the expected errors in predicted lines will usually reflect this ambiguity in laboratory uncertainties through Eq (6) and (7). In some cases, the quality of the least squares fit of the parameters to the experimental lines can be a guide to the statistical nature of the experimental uncertainties. Whenever possible, the expected errors in the catalog will reflect an expected 95% confidence interval based on the model used to fit the data. However, the errors can be different from this design goal by factors of three just due to the quality of input error estimates. Lines with an expected error greater than 1 GHz have been dropped from the catalogue.

The expected errors can only be computed relative to the model used. There are at least two ways the model can be "wrong" for the predicted frequencies. First, higher-order centrifugal distortion terms may no longer be negligible for the predicted frequencies. This effect will generally be important for lines of higher J or K than the laboratory-determined data set. In a sense, the predictions are then a form of extrapolation rather than interpolation and are therefore more suspect. A second factor leading to discrepancies in the predicted frequencies comes from "resonances". These resonances come from a near overlap of energy states that are coupled by elements of the Hamiltonian matrix. Poor predictability comes when these elements are neglected in the model or are treated inadequately by some form of perturbation theory. Such a neglect of coupling elements is always necessary at some level in any practical calculation. A major contributing problem is that often the existing data set is not sensitive to the parameters which are needed to characterize the resonance.

Precision in the intensity estimates is generally less critical than precision in the frequency. Contributing to intensity uncertainty are errors in the dipole moment, errors in the line strength χS_{ba} and errors in the rotation-spin partition function. (The vibration-electronic partition function does not enter directly because the catalogued intensities are defined on the basis of concentrations of the given vibration-electronic state.) Dipole moment errors come directly from the experimental determination and indirectly from the J dependence of the dipole moment due to centrifugal mixing of vibrational states. Line strength errors can come from deficiencies in the model Hamiltonian and are particularly severe when resonances have been inadequately accounted for. Partition function errors are relatively benign but can become significant if the classical formulae are used at low temperatures for small molecules. With the exception of unanticipated resonances and poorly determined dipole moments, worst case errors in the intensity will generally be at the 1% level or lower.

V. Format of Quantum Numbers

The quantum numbers of the catalogued lines are given in the field QN according to a format QNFORM, as discussed in Section II. In this section the identities of the quantum numbers will be described for each distinct format. The field QN will be regarded as being divided into 13 integers. The first 12 integers are two digits long and the last integer is a single digit (a FORTRAN 12I2, I1 format).

The spectroscopic nomenclature followed is the standard one in which upper state quantities are denoted by a single prime and lower state quantities are denoted by a double prime.

In addition,

L	electron orbital angular momentum
R	rotational angular momentum
$N = R+L$	
S	electron spin
$J = N+S$	total electron and rotational angular momentum
I	nuclear spin
$F_1 = I_1+J$	intermediate F value for two nuclear spins I_1 and I_2
$F = I_2 + F_1$	(for two non-zero nuclear spin)
$= I + J$	(for one non-zero nuclear spin)
Ω	projection of J on the molecular axis of a linear molecule
K	projection of J on the symmetry axis of a symmetric rotor
p	parity under inversion (1 = gerade, -1 = ungerade)
v	vibrational quantum number
K_{-1}	prolate K quantum number for an asymmetric rotor
K_{+1}	oblate K quantum number for an asymmetric rotor

In all cases, half integer quantum numbers will be written into the QN field after rounding up to the next integer. For example, $J = 1/2$ will be entered in the catalogue as a 1. The format specification QNFORM is designed so that the correct half-integer value can be recovered unambiguously.

The format integer QNFORM is allocated four digits. The most significant digit is zero for the standard formats described in Table V-1 and is non-zero for special-case formats which will be allocated as the need arises. Of the remaining three digits, the most significant is used to describe the type of species involved and the least significant digits are used to specify the nature of the integers in the field. The formats are shown in Tables V-1 and V-2.

Table V-1: Standard Quantum Number Formats

QNF	QN	Remarks
A. Atoms		
010	$J' + \frac{1}{2}, J'' + \frac{1}{2}$	$I = 0, J$ half integer
020	J', J''	$I = 0, J$ integer
011	$J' + \frac{1}{2}, J'' + \frac{1}{2}, F' + \frac{1}{2}, F'' + \frac{1}{2}$	J and F half integer
012	$J' + \frac{1}{2}, J'' + \frac{1}{2}, F', F''$	J half integer, F integer
021	$J', J'', F' + \frac{1}{2}, F'' + \frac{1}{2}$	J integer, F half integer
022	J', J'', F', F''	J and F integer
B. Linear molecules and Diatomic molecules		
1XY	$J', J'', x', x'', y', y''$	1Σ state
2XY	$J' + \frac{1}{2}, N', J'' + \frac{1}{2}, N'', x', x'', y', y''$	2Σ state
3XY	$J', N', J'', N'', x', x'', y', y''$	3Σ state
4XY	$J', P', J'', P'', x', x'', y', y''$	1Π state
5XY	$J' + \frac{1}{2}, \Omega', J'' + \frac{1}{2}, P', J'' + \frac{1}{2}, \Omega'' + \frac{1}{2}, P'', x', x'', y', y''$	2Π state
6XY	$J', \Omega', P', J'', \Omega'', P'', x', y', y''$	3Π state
7XY	$J', \Omega', P', J'', \Omega'', P'', x', x'', y', y''$	1Δ state
C. Symmetric Rotors (3-fold symmetry or higher, $S = 0$)		
8XY	$J', K', J'', K'', x', x'', y', y''$	
D. Asymmetric Rotors		
9XY	$J', K'_{-1}, K'_{+1}, J'', K''_{-1}, K''_{+1}, x', x'', y', y''$	closed shell molecule
95Y	$N', K'_{-1}, K'_{+1}, N'', K''_{-1}, K''_{+1}, J' + \frac{1}{2}, J'' + \frac{1}{2}, y', y''$	$S = \frac{1}{2}$ open shell radical
96Y	$N', K'_{-1}, K'_{+1}, N'', K''_{-1}, K''_{+1}, J', J'', y', y''$	$S = 1$ triplet state
$X = 0:$	$x = 0$	$Y = 0:$
$X = 1:$	$x = F + \frac{1}{2}$	$Y = 1:$
$X = 2:$	$x = F$	$Y = 2:$
$X = 4:$	$x = v$	
		(Y must be zero)
		(F half integer)
		(F integer)
		vibrational state label (further specified in the species description)
		' signifies upper energy state, " signifies lower energy state

ORIGINAL PAGE IS
OF POOR QUALITY

Table V-2: Special-Case Quantum Number Formats

(Formats will be provided in later supplements.)

VI. Documentation by Species

The documentation for each molecular or atomic species appears in order of "species tag." On the following page, immediately preceding the documentation, is an "Index of Species." This index lists all the species documentation contained in this publication, ordered by "species tag."

INDEX OF SPECIES

1001 H-Atom	41003 CH3C-13-CH	60001 OCS
2001 D-Atom	41004 C-13-H3CCH	60002 SiS
3001 HD	41005 CH3CCD	61001 OC-13-S
12001 C-Atom	41006 CH2DCCH	61002 Si-29-S
13001 C-13-Atom	42001 CH3CN-15	62001 OCS-34
14001 N-Atom	42002 CH2CO	62002 O-18-CS
14002 N-Atom-D-State	43001 CHDCO	62003 Si-30-S
16001 O-Atom	44001 CS	62004 SiS-34
17001 HO	44002 SiO	64001 S2
17002 NH3	44003 CH3CHO-A	64002 SO2
17003 CH3D	44004 N2O	75001 HCCCCCN
18001 DO	45001 C-13-S	76001 HCCCCC-13-N
18002 N-15-H3	45002 Si-29-O	76002 HCCCC-13-CN
18003 H2O	45003 NH2CHO	76003 HCCC-13-CCN
19001 HO-18	46001 CS-34	76004 HCC-13-CCCN
19002 HDO	46002 Si-30-O	76005 HC-13-CCCN
27001 HCN	46003 H2CS	76006 HCCCCCN-15
27002 HNC	46004 C2H5OH	76007 DCCCCCN
28001 CO	46005 HCOOH	80001 HBr-79
28002 HC-13-N	47001 H2C-13-S	82001 HBr-81
28003 HCN-15	47002 HC-13-OOH	95001 Br-79-0
28004 DCN	47003 DCOOH	97001 Br-81-0
28005 HNC-13	47004 HCOOD	
28006 HN-15-C	48001 SO	
28007 DNC	48002 SOv1	
29001 C-13-O	48003 H2CS-34	
29002 HCO+	48004 O3	
29003 CH2NH	50001 S-34-0	
30001 CO-18	50002 SO-18	
30002 HC-13-O+	50003 O3-SYM-0-18	
30003 DCO+	50004 O3-ASYM-0-18	
30004 H2CO	51001 HCCCN	
30005 C-13-H2NH	51002 C1O	
30006 CH2N-15-H	52001 HCCC-13-N	
30007 CH2ND	52002 HCC-13-CN	
30008 NO	52003 HC-13-CCN	
31001 HCO-18+	52004 HCCCN-15	
31002 H2C-13-O	52005 DCCCN	
32001 O2	53001 C2H3CN	
32002 O2v1	53002 C1-37-O	
32003 CH3OH	54001 CH2CHC-13-N	
32004 H2CO-18	54002 CH2C-13-HCN	
34001 O-18-O	54003 C-13-H2CHCN	
34002 H2S	54004 CH2CDCN	
34003 PH3	55001 C2H5CN	
35001 HDS	56001 CH3CH2C-13-N	
36001 HCl	56002 CH3C-13-H2CN	
38001 HCl-37	56003 C-13-H3CH2CN	
40001 CH3CCH	56004 C2H5CN-15	
41001 CH3CN	56005 CH2DCH2CN-S	
41002 CH3CC-13-H	56006 CH2DCH2CN-A	

Species Tag: 1001	Page 1 of 1	Version: 1	Date: Oct, 1979
Species Name: Atomic Hydrogen $2S_{1/2}$ ground state		Author: H. M. Pickett	
		Min. Int. = 8×10^{-10} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4.0		J Max. = ($\frac{1}{2}$)	Lines Listed = 1
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The atomic hydrogen line at 21 cm has been measured to a precision of 0.001 Hz (L. Essen, R. W. Donaldson, M. J. Bangham, and E. G. Hope, 1971, Nature 229, 110). The average spontaneous emission lifetime of the three F = 1 states is 2.876×10^{-15} sec⁻¹.

Species Tag: 2001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Atomic Hydrogen Deuterium Isotope $2S_{1/2}$ ground state		Author: H. M. Pickett	
		Min. Int. = 5×10^{-11} nm ² MHz	
Q (Spin-Rot., T = 300K) = 6.0		J Max. = ($\frac{1}{2}$)	Lines Listed = 1
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The atomic deuterium line at 327 MHz has been measured to an accuracy of 0.002 Hz (D. J. Wineland and N. F. Ramsey, 1972, Phys. Rev. A 5, 821). The average spontaneous emission lifetime for the four $F = 3/2$ states is 4.695×10^{-17} sec⁻¹.

Species Tag: 3001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: HD Mono-deuterated molecular hydrogen		Author: H. M. Pickett	
		Min. Int. = 10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 5.016		J Max. = (1)	Lines Listed = 1
Dipole Moments/Debye: $\mu_a = 5.85 \times 10^{-4}$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1339100. C =			

The dipole moment is from M. Trefler and H. P. Gush, 1968, Phys. Rev. Letters, 20, 703. The rotational constant is from R. A. Durie and G. Herzberg, 1960, Can. J. Phys. 38, 806. The frequency and its error limits were obtained from the difference of the S₁ (0) and R₁ (1) transitions listed by A. McKellar, 1974, Can. J. Phys. 52, 1144.

Species Tag: 12001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Atomic Carbon ³ P ground state		Author: H. M. Pickett	
		Min. Int. = 10 ⁻⁵ nm ² MHz	
Q (Spin-Rot., T = 300K) = 7.8330		J Max. = (2)	Lines Listed = 2
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The lines of atomic carbon have been measured by Laser Magnetic Resonance (R. J. Saykally and K. M. Evenson, Symp. on Molecular Spectroscopy, Columbus, OH, 1979). Intensities were calculated using the free electron g factor. The partition function was calculated from the following table:

<u>J</u>	<u>Energy</u>	<u>Degeneracy</u>
0	0.0 cm ⁻¹	1
1	16.4168	3
2	43.4138	5

Species Tag: 13001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Atomic Carbon ¹³ C isotope		Author: H. M. Pickett	
		Min. Int. = 10 ⁻¹⁶ nm ² MHz	
Q (Spin-Rot., T = 300K) = 15.666		J Max. = (2)	Lines Listed = 7
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The fine structure intervals of ¹²C were multiplied by 0.999992967 to account for the isotope effect on the Rydberg constant. The hyperfine intervals of G. Wolber, H. Figger, R. A. Haberstroh and S. Penselin, 1969 (Physics Letters A29, 461) were then used to compute the line frequencies. Intensities were calculated using the free electron g value.

Species Tag: 14001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Atomic Nitrogen $^4S_{3/2}$ ground state		Author: H. M. Pickett	
		Min. Int. = 10^{-13} nm ² MHz	
Q (Spin-Rot., T = 300K) = 12.0		J Max. = (3/2)	Lines Listed = 2
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The hyperfine transitions of atomic nitrogen at 15.6 and 26.1 MHz were obtained from J. M. Hirsch, G. H. Zimmerman, III, D. J. Larson, and N. F. Ramsey, 1977, Phys. Rev. A 16, 484. Intensities were calculated using the experimental g_j value.

Species Tag: 14002	Page 1 of 1	Version: 1	Date: Nov. 1979
Species Name: Atomic Nitrogen 2D excited state		Author: H. M. Pickett	
		Min. Int. = 10^{-8} nm ² MHz	
Q (Spin-Rot., T = 300K) = 29.510		J Max. = (5/2)	Lines Listed = 6
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The $^2D_{5/2}$ state is 19223 cm^{-1} above the ground $^4S_{3/2}$ state. (C. E. Moore, 1949, Atomic Energy Levels I, U. S. N. Bureau of Stds.). The $^2D_{3/2} - ^2D_{5/2}$ inverted fine structure transition at 260 GHz has a predicted uncertainty of ± 600 MHz, but the hyperfine structure intervals are uncertain by less than 1 MHz. (H. E. Radford and K. M. Evenson, 1968, Phys. Rev. 168, 70). The average spontaneous emission rate from the 2D states to the 4S states is $1.06 \times 10^{-6} \text{ sec}^{-1}$. In contrast, the spontaneous emission rate from $^2D_{3/2} F = 5/2$ to $^2D_{5/2} F = 7/2$ is $3.15 \times 10^{-9} \text{ sec}^{-1}$.

Species Tag: 16001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Atomic Oxygen ³ P ground state		Author: H. M. Pickett	
		Min. Int. = 10 ⁻⁴ nm ² MHz	
Q (Spin-Rot., T = 300K) = 7.4943		J Max. = (2)	Lines Listed = 1
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = C =			

The 68 cm⁻¹ J = 1-2 line of oxygen has been measured by Laser Magnetic Resonance (R. J. Saykally and K. M. Evenson, Symposium on Molec. Specty., Columbus, 1979). Intensities were calculated using the free electron g factor. The partition function was calculated from the following table:

<u>J</u>	<u>Energy</u>	<u>Degeneracy</u>
0	227.0195	1
1	68.7165	3
2	0.0	5

Species Tag: 17001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydroxyl Radical OH ground $2\Pi_{3/2}$ and $2\Pi_{1/2}$ states		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 10^{-13} nm ² MHz	
Q (Spin-Rot., T = 300K) = 83.328		J Max. = 19.5	Lines Listed = 125
Dipole Moments/Debye: μ_a = 1.667 μ_b = μ_c =			
Rot. Const./MHz: A = B = 556140.9 C =			

The calculational method and microwave data are reported in R. A. Beaudet and R. L. Poynter, 1978, J. Phys. Chem. Ref. Data 7, 311. The partition function was calculated by a sum over states to J = 11.5. Other values are:

T	Q_{sr}
150K	40.230
70	21.800
30	16.354
10	15.696

Only the J = 5/2 - 3/2 and J = 3/2 - 1/2 submillimeter transitions are included in the catalogue.

Species Tag: 17002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Ammonia NH ₃ ground inversion states		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 10 ⁻¹⁷ nm ² MHz	
Q (Spin-Rot., T = 300K) = 581.4432		J Max. = 20	Lines Listed = 235
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ $\mu_c = 1.476$			
Rot. Const./MHz: A = B B = 298115.37 C = 185092.			

The calculational method and microwave data are reported in R. L. Poynter and R. K. Kakar, 1975, Ap. J. Suppl. 29, 87. The vibrational state designations in the catalogue are $v = 0$ for the lower inversion state and $v = 1$ for the upper state.

Species Tag: 17003	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Deutero Methane CH ₃ D		Author: R. L. Poynter	
		Min. Int. = 10 ⁻⁸ cm ² MHz	
Q (Spin-Rot., T = 300K) = 402.166	J Max. = 13	Lines Listed = 80	
Dipole Moments/Debye: $\mu_a = 5.6 \times 10^{-3}$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 157412. B = 116325.308 C = B			

The J = 0 → 1 line has been measured by H. M. Pickett and E. A. Cohen, 1980, Ap. J. Letters. Predicted lines are based upon the constants given by C. Chakerian and G. Guelachvilli, 1980, J. Mol. Spect., in press. The dipole moment has been measured by I. Ozier, W. Ho, and G. Birnbaum, 1969, J. Chem. Phys. 51, 4873, and by S. C. Wofsey, J. S. Muentzer, and W. Klemperer, (1970), J. Chem. Phys. 53, 4005.

Species Tag: 18001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Deuterated hydroxyl		Author: R.L. Poynter & H.M. Pickett	
radical: OD $^2_{\Pi}$ ground states		Min. Int. = 10^{-13} nm ² MHz	
Q (Spin-Rot., T = 300K) = 215.558		J Max. = 19.5	Lines Listed = 235
Dipole Moments/Debye: $\mu_a = 1.653$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 296312.0 C =			

The calculational method and microwave data are reported in R. A. Beudet and R. L. Poynter, 1978, J. Phys. Chem. Ref. Data 7, 311. The partition function was calculated using the relation $Q=6 Q_{rot} [1 + \exp(-188.112/T)]$, in which Q_{rot} is the rotational partition function for a diatomic molecule. No submillimeter lines are currently listed in the catalogue for OD.

Species Tag: 18002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Ammonia - ^{15}N $^{15}\text{NH}_3$ ground inversion. states		Author: R. L. Poynter	
		Min. Int. = 10^{-11} nm ² MHz	
Q (Spin-Rot., T = 300K) = 581.44		J Max. = 14	Lines Listed = 106
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ $\mu_c = 1.476$			
Rot. Const./MHz: A = B B = 298115.37 C = 185092.			

The lines used in the fit were reported by E. Schnabel, T. Törring, and W. Wilke, (1965) Z. Phys. 188, 167.

The most recent measurements of H Sasada, 1980, J. Mol. Spect., in press, were received too late to include in the catalogue.

The dipole moment was assumed to be the same as $^{14}\text{NH}_3$. The computational method is the same as that used for $^{14}\text{NH}_3$.

Species Tag: 18003	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Water		Author: R. L. Poynter	
H ₂ O		Min. Int. = 10 ⁻⁹ nm ² MHz	
Q (Spin-Rot., T = 300K) = 177.06		J Max. = 13	Lines Listed = 151
Dipole Moments/Debye: $\mu_a = 0$ $\mu_b = 1.855$ $\mu_c = 0$			
Rot. Const./MHz: A = 835840.29 B = 435351.72 C = 278138.7			

The data on H₂O are from F. C. DeLucia, P. Helminger, R. Cook, and W. Gordy (1972), Phys. Rev. A, 5, 487, R. T. Hall and J. M. Dowling (1967), J. Chem. Phys. 47, 2454, and R. T. Hall and J. M. Dowling, (1970), J. Chem. Phys. 52, 1161. The dipole moment is from S. Clough, Y. Beers, G. P. Klein, and L. Rothman (1973), J. Chem. Phys. 59, 3125. Other references are given in F.C. DeLucia, P. Helminger, and W. H. Kirchhoff, (1974), J. Phys. Chem. Ref. Data, 3, 211.

Due to model problems and an incomplete least squares fitting program for this special case, no least squares fit was done for this molecule. Rather, the line frequencies were calculated in double precision using the set of rotational constants determined by De Lucia, et al. (1972, 1974) with a Hamiltonian that included up through tenth order distortion terms. The lower frequency lines that are computed from this Hamiltonian are in reasonable agreement with De Lucia, et al., but the higher frequency, higher J lines show deviations, possibly due to round off problems. The error estimates were set to a fixed 160 MHz for all the calculated lines, since our present program does not give the information necessary to evaluate the predicted errors.

Further work will be done on this problem. The present list is therefore an interim set of results.

Species Tag: 19001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydroxyl radical Oxygen - 18 isotope ^{18}OH		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 10^{-13} cm^2MHz	
Q (Spin-Rot., T = 300K) = 83.882		J Max. = 20	Lines Listed = 113
Dipole Moments/Debye: $\mu_a = 1.667$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 552470. C =			

The calculational method and microwave data are reported in R. A. Beudet and R. L. Poynter, 1978, J. Phys. Chem. Ref. Data 7, 311. The partition function was calculated using the ^{16}OH value and the ratio of the B values. No pure rotational lines are currently listed in the catalogue.

Species Tag: 19002	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Water Single Deuterium Isotope HDO		Author: R. L. Poynter	
		Min. Int. = 10^{-13} nm ² MHz	
Q (Spin-Rot., T = 300K) = 146.05	J Max. = 13	Lines Listed = 221	
Dipole Moments/Debye: $\mu_a = 0.657$ $\mu_b = 1.732$ $\mu_c = 0.$			
Rot. Const./MHz: A = 701931.5 B = 272912.6 C = 192055.2			

The observed lines are from F. C. De Lucia, R. L. Cook, P. Helminger, and W. Gordy (1971), J. Chem. Phys. 55, 5334, and J. W. Fleming and M. J. Gibson, (1976), J. Mol. Spect. 62, 326. The dipole moment components are from S. A. Clough, Y. Beers, G. P. Klein, and L. S. Rothman, (1973), J. Chem. Phys. 59, 2254.

Further references can be found in F. C. De Lucia, P. Helminger, and W. H. Kirchhoff, (1974), J. Phys. and Chem. Ref. Data 3, 211.

Due to model problems and an incomplete least squares fitting program for this special case, no least squares fit was done for this molecule. Rather, the line frequencies were calculated in double precision using the set of rotational constants determined by De Lucia, et al. (1971, 1974) with a Hamiltonian that included up through tenth order distortion terms. The lower frequency lines that are computed from this Hamiltonian are in reasonable agreement with De Lucia, et al., but the higher frequency, higher J lines show deviations, possibly due to round off problems. The error estimates were set to a fixed 160 MHz for all the calculated lines, since our present program does not give the information necessary to evaluate the predicted errors.

Further work will be done on this problem. The present list is therefore an interim set of results.

Species Tag: 27001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Cyanide HCN		Author: R. L. Poynter	
		Min. Int. = 10^{-4} nm ² MHz	
Q (Spin-Rot., T = 300K) = 424.154		J Max. = 34	Lines Listed = 45
Dipole Moments/Debye: $\mu_a = 2.984$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 44315.975 C =			

The observed transitions are from F. C. De Lucia and W. Gordy (1969), Phys. Rev. 187, 58. The dipole moment is from G. Tomasevich, (1970), Thesis, Harvard Univ.

Species Tag: 27002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Isocyanide HNC		Author: R. L. Poynter	
		Min. Int. = 10^{-3} nm ² MHz	
Q (Spin-Rot., T = 300K) = 138.223		J Max. = (34)	Lines Listed = 34
Dipole Moments/Debye: $\mu_a = 2.699$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 45332.00 C =			

The observed lines are from R. A. Creswell, E. F. Pearson, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 221, and E. F. Pearson, R. A. Creswell, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 1394.

Species Tag: 28001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbon Monoxide CO		Author: R. L. Poynter	
		Min. Int. = 10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 108.787		J Max. = (26)	Lines Listed = 26
Dipole Moments/Debye: $\mu_a = 0.1098$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 57635.97 C =			

The experimental measurements were reported by B. Rosenblum, A. H. Nethercot, Jr., and C. H. Townes, 1958, Phys. Rev. 109, 400; W. Gordy and M. J. Cowan, 1957, Bull. Am. Phys. Soc. 2, 212; and by P. Helminger, F. C. De Lucia and W. Gordy, 1970, Phys. Rev. Lett. 25, 1397.

The dipole moment was measured by J. S. Muentner, 1975, J. Mol. Spect. 55, 490.

Species Tag: 28002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Cyanide Carbon - 13 isotope H ¹³ CN		Author: R. L. Poynter	
		Min. Int. = 10 ⁻⁴ nm ² MHz	
Q (Spin-Rot., T = 300K) = 435.385	J Max. = 34	Lines Listed = 45	
Dipole Moments/Debye: $\mu_a = 2.984$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 43170.137 C =			

The measured lines are from E. F. Pearson, R. A. Creswell, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 1394.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 28003	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Cyanide Nitrogen - 15 isotope HC ¹⁵ N		Author: R. L. Poynter	
		Min. Int. = 3×10^{-3} cm ² MHz	
Q (Spin-Rot., T = 300K) = 435.204	J Max. = 35	Lines Listed = 35	
Dipole Moments/Debye: $\mu_a = 2.984$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 43027.69 C =			

The measured lines are from E. F. Pearson, R. A. Creswell, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 1394.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 28004	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Cyanide Deuterium isotope DCN		Author: R. L. Poynter	
		Min. Int. = 7×10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 518.916		J Max. = 41	Lines Listed = 54
Dipole Moments/Debye: $\mu_a = 2.984$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 36207.46 C =			

The observed transitions are from F. C. De Lucia and W. Gordy (1969), Phys. Rev. 187, 58. The dipole moment is from G. Tomasevich, (1970), Thesis, Harvard Univ.

Species Tag: 28005	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Isocyanide Carbon - 13 isotope HN ¹³ C		Author: R. L. Poynter	
		Min. Int. = 2×10^{-3} cm ² MHz	
Q (Spin-Rot., T = 300K) = 143.880		J Max. = (35)	Lines Listed = 35
Dipole Moments/Debye: $\mu_a = 2.699$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 43545.61 C =			

The observed lines are from R. A. Creswell, E. F. Pearson, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 221, and E. F. Pearson, R. A. Creswell, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 1394.

Species Tag: 28006	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Isocyanide Nitrogen - 15 isotope $\text{H } ^{15}\text{NC}$		Author: R. L. Poynter	
		Min. Int. = 2×10^{-3} nm ² MHz	
Q (Spin-Rot., T = 300K) = 141.013		J Max. = (33)	Lines Listed = 33
Dipole Moments/Deby: $\mu_a = 2.699$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 44433.04 C =			

The observed lines are from R. A. Creswell, E. F. Pearson, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 221, and E. F. Pearson, R. A. Creswell, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 1394.

Species Tag: 28007	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Hydrogen Isocyanide Deuterium isotope DNC		Author: R. L. Poynter	
Q (Spin-Rot., T = 300K) = 164.169		Min. Int. = 10^{-3}	$\text{nm}^2 \text{MHz}$
		J Max. = (39)	Lines Listed = 39
Dipole Moments/Debye: $\mu_a = 2.699$ $\mu_b =$ $\mu_c =$			
Rot. Const./ MHz: A = B = 38152.995 C =			

The observed lines are from R. A. Creswell, E. F. Pearson, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 221, and E. F. Pearson, R. A. Creswell, M. Winnewisser and G. Winnewisser, (1976), Z. Naturforsch. 31a, 1394.

Species Tag: 29001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbon monoxide Carbon - 13 isotope ^{13}CO		Author: R. L. Poynter	
		Min. Int. = 8×10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 113.776		J Max. = (27)	Lines Listed = 27
Dipole Moments/Debye: $\mu_a = 0.110$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 55101.02 C =			

The experimental measurement was reported by B. Rosenblum, A. H. Nethercot, Jr. and C. H. Townes, 1958, Phys. Rev. 109, 400.

The dipole moment was measured by W. L. Meerts, F. H. De Leeuw, and A. Dymanus, 1977, Chem. Phys. 22, 319.

Species Tag: 29002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Formyl radical cation		Author: R. L. Poynter	
HCO ⁺		Min. Int. = 10 ⁻³ nm ² MHz	
Q (Spin-Rot., T = 300K) = 140.504		J Max. = (33)	Lines Listed = 33
Dipole Moments/Debye: $\mu_a = 3.30$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 44594.46 C =			

Only one line has been observed in the laboratory, by R. C. Woods, T. A. Dixon, R. J. Saykally, and P. G. Szanto, (1975), Phys. Rev. Lett. 35, 1269.

The dipole moment was estimated theoretically in the same paper.

Species Tag: 29003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Methylenimine CH ₂ NH		Author: R. L. Poynter Min. Int. = 6×10^{-11} nm ² MHz	
Q (Spin-Rot., T = 300K) = 5893.		J Max. = 40	Lines Listed = 4769
Dipole Moments/Debye: $\mu_a = 1.325$ $\mu_b = 1.530$ $\mu_c = 0.$			
Rot. Const./MHz: A = 196211.045 B = 34532.395 C = 29352.232			

The data set and computational method are referenced in W. H. Kirchhoff, D. R. Johnson, and F. J. Lovas, 1973, J. Phys. Chem. Ref. Data 2, 1.

Species Tag: 30001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbon Monoxide oxygen - 18 isotope $C^{18}O$		Author: R. L. Poynter	
		Min. Int. = 8×10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 114.2096		J Max. = (27)	Lines Listed = 27
Dipole Moments/Debye: $\mu_a = 0.110$ $\mu_b =$ $\mu_c = 28$			
Rot. Const./MHz: A = B = 54891.425 C =			

The experimental measurement was reported by B. Rosenblum, A. H. Nethercot, Jr. and C. H. Townes, 1958, Phys. Rev. 109, 400.

The dipole moment was reported by W. L. Meerts, F. H. De Leeuw, and A. Dymanus, 1977, Chem. Phys. 22, 319.

Species Tag: 30002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Formyl radical cation carbon - 13 isotope H^{13}CO^+		Author: R. L. Poynter	
		Min. Int. = 3×10^{-3} nm^2MHz	
Q (Spin-Rot., T = 300K) = 144.437		J Max. = (34)	Lines Listed = 34
Dipole Moments/Debye: $\mu_a = 3.30$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 43377.36 C = '			

Only one line has been observed in the laboratory, by R. C. Woods, R. J. Saykally, T. A. Dixon, P. G. Szanto, and T. Anderson, (1976) 31st Symposium on Molecular Spectroscopy, Columbus, Ohio.

Species Tag: 30003	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Formyl radical cation Deuterium isotope DCO ⁺		Author: R. L. Poynter	
		Min. Int. = 2×10^{-3} nm ² MHz	
Q (Spin-Rot., T = 300K) = 173.872		J Max. = (42)	Lines Listed = 41
Dipole Moments/Debye: $\mu_a = 3.30$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 36019.76 C =			

Only one line has been observed in the laboratory, by R. C. Woods, R. J. Saykally, T. A. Dixon, P. G. Szanto, and T. Anderson, (1976) 31st Symposium on Molecular Spectroscopy, Columbus, Ohio.

Species Tag: 30004	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Formaldehyde H ₂ CO		Author: R. L. Poynter	
		Min. Int. = 10 ⁻²⁰ nm ² MHz	
Q (Spin-Rot., T = 300K) = 2876.7		J Max. = 40	Lines Listed = 611
Dipole Moments/Debye: $\mu_a = 2.331$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 281925.97 B = 38836.582 C = 34001.6729			

The experimental data were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- J. K. Bragg and A. H. Sharbaugh, 1949, Phys. Rev. 75, 1774.
 J. S. Chardon and D. Guichon, 1977, J. Phys. (Paris), 38, 113; 1973, J. Phys. (Paris), 34, 791.
 F. Y. Chu, S. M. Freund, J. W. C. Johns, and T. Oka, 1973, J. Mol. Spect. 48, 328.
 D. Dangoisse, E. Willemot, and J. Bellet, 1978, J. Mol. Spect. 71, 414.
 D. R. Johnson, R. Lovas, and W. H. Kirchhoff, 1972, J. Phys. Chem. Ref. Data, 1, 1011.
 A. F. Krupnov, L. I. Gershtein, V. G. Shustrov, and V. V. Polyakov, 1970, Opt. Spect. (USSR), 28, 257.
 R. B. Lawrence and M. W. P. Strandberg, 1949, Phys. Rev. 75, 1774.
 T. Oka, H. Hirakawa, and K. Shimoda, 1960, J. Phys. Soc. Japan, 15, 2265.
 T. Oka, T. Takagi, Y. Morino, 1964, J. Mol. Spect. 14, 27.

The dipole moment was measured by K. Kondo and T. Oka, 1960, J. Phys. Soc. Japan, 15, 307.

Species Tag: 30005	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Methylenimine Carbon - 13 isotope $^{13}\text{CH}_2\text{NH}$		Author: R. L. Poynter	
		Min. Int. = 10^{-8} nm ² MHz	
Q (Spin-Rot., T = 300K) = 2012.45		J Max. = 10	Lines Listed = 439
Dipole Moments/Debye: $\mu_a = 1.325$ $\mu_b = 1.530$ $\mu_c = 0.$			
Rot. Const./MHz: A = 196195.217 B = 33747.872 C = 28707.047			

The computational method is the same as that used on the parent species, CH_2NH . The data are from R. Pearson, Jr. and F. J. Lovas, 1977, J. Chem. Phys. 66, 4149. The dipole moment has been assumed the same as for the parent species.

Species Tag: 30006	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Methylenimine Nitrogen - 15 isotope $\text{CH}_2 \text{ } ^{15}\text{NH}$		Author: R. L. Poynter	
		Min. Int. = 10^{-8} nm ² MHz	
Q (Spin-Rot., T = 300K) = 2015.792		J Max. = 10	Lines Listed = 440
Dipole Moments/Debye: $\mu_a = 1.325$ $\mu_b = 1.530$ $\mu_c = 0.$			
Rot. Const./MHz: A = 195738.07 B = 33736.100 C = 28688.614			

The computational method is the same as that used on the parent species, CH_2NH . The data are from R. Pearson, Jr. and F. J. Lovas, 1977, J. Chem. Phys. 66, 4149. The dipole moment has been assumed the same as for the parent species.

Species Tag: 30007	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Methylenimine Deuterium isotope on nitrogen atom CH ₂ ND		Author: R. L. Poynter	
		Min. Int. = 8×10^{-10} nm ² MHz	
Q (Spin-Rot., T = 300K) = 7181.07		J Max. = 15	Lines Listed = 1820
Dipole Moments/Debye: $\mu_a = 1.325$ $\mu_b = 1.530$ $\mu_c = 0.$			
Rot. Const./MHz: A = 157673.877 B = 32069.366 C = 26563.849			

The computational method is the same as that used on the parent species, CH₂NH. The data are from R. Pearson, Jr. and F. J. Lovas, 1977, J. Chem. Phys. 66, 4149. The dipole moment has been assumed the same as for the parent species.

Species Tag: 30008	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Nitric Oxide ground 2π states NO		Author: H. M. Pickett	
		Min. Int. = 10^{-20} nm ² MHz	
Q (Spin-Rot., T = 300K) = 1159.464		J Max. = 35.5	Lines Listed = 1909
Dipole Moments/Debye: $\mu_a = 0.15872$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = B = 50849.06 C =			

The spectrum of NO was fit to the fine structure Hamiltonian described by C. Amiot, R. Bacis and G. Guelachvili, 1978, Can. J. Phys. 56, 251, along with the hyperfine Hamiltonian described by W. L. Meerts, 1976, Chem. Phys. 14, 421. The radio frequency lines are from W. L. Meerts and A. Dymanus, 1972, J. Mol. Spectroscopy. 44, 320. The millimeter lines were measured at JPL by H. M. Pickett and E. A. Cohen. The partition function was determined by a sum over states to $J=71/2$. Other values are:

<u>T</u>	<u>Q</u>
10	30.394
30	80.168
60	162.551
100	296.509
150	492.323

Species Tag: 31001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Formyl radical cation Oxygen - 18 isotope HC ¹⁸ O ⁺		Author: R. L. Poynter	
		Min. Int. = 3×10^{-3} ^{cm} ² MHz	
Q (Spin-Rot., T = 300K) = 145.427		J Max. = (34)	Lines Listed = 34
Dipole Moments/Debye: μ_a = μ_b = 3.30 μ_c =			
Rot. Const./MHz: A = B = 43081.217 C =			

Only one line has been observed in the laboratory, by R. C. Woods, R. J. Saykally, T. A. Dixon, P. G. Szanto, and T. Anderson, (1976) 31st Symposium on Molecular Spectroscopy, Columbus, Ohio.

Species Tag: 31002	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Formaldehyde Carbon - 13 isotope $H_2^{13}CO$		Author: R. L. Poynter	
		Min. Int. = 5×10^{-10} nm ² MHz	
Q (Spin-Rot., T = 300K) = 2949.7		J Max. = 40	Lines Listed = 601
Dipole Moments/Debye: $\mu_a = 2.331$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 281930.85 B = 37811.92 C = 33213.19			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- D. Dangoisse, E. Willemot, and J. Bellet, 1978, J. Mol. Spect. 71, 414.
D. R. Johnson, F. Lovas, and W. H. Kirchhoff, 1972, J. Phys. Chem. Ref. Data, 1, 1011.
R. B. Lawrence and M. W. P. Strandberg, 1951, Phys. Rev. 83, 363.
R. Nerf, 1972, Ap. J., 174, 467.
T. Oka, H. Hirakawa, and K. Shimoda, 1960, J. Phys. Soc. Japan, 15, 2265.
T. Oka, T. Takagi, and Y. Morino, 1964, J. Mol. Spect. 14, 27.
K. D. Tucker, G. R. Tomasevich, and P. Thaddeus, 1972, Ap. J. 174, 463, 1971, Ap. 169, 429.

The dipole moment was assumed to be the same as the parent $H_2^{12}CO$.

Species Tag: 32001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Molecular Oxygen $^{16}\text{O}_2$ v = 0 state of ground $^3\Sigma_g^-$ electronic state		Author: H. M. Pickett	
		Min. Int. = 10^{-13} cm ² MHz	
Q (Spin-Rot., T = 300K) = 218.529		J Max. = (50)	Lines Listed = 74
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = 43099.795 C =			

The measurements and calculational method are from T. Amano and E. Hirota, 1974, J. Mol. Spectroscopy 53, 346. The Raman lines of O_2 (M. Loete and H. Berger, 1977, J. Mol. Spectroscopy 68, 317) were used with the millimeter wavelength measurements and the submillimeter-line of W. Steinbach and W. Gordy (1973, Phys. Rev A 8, 1953) in a combined fit of the v = 0 and v = 1 transitions. The intensities of the magnetic dipole transitions have been calculated using the g values obtained from laser magnetic resonance (D. M. Evenson and M. Misushima, 1972, Phys. Rev. A 6, 2197).

The value of Q is determined by a sum over states to J = 60. Other values are:

<u>T</u>	<u>Q</u>
10	8.306
30	22.615
60	44.326
100	73.327
150	109.605

Species Tag: 32002	Page 1 of 1	Version: 1	Date: Oct, 1979
Species Name: Molecular Oxygen $^{16}\text{O}_2$ v = 1 state of the ground $^3\Sigma_g^-$ state		Author: H. M. Pickett	
Q (Spin-Rot., T = 300K) = 220.918		J Max. = (40)	Lines Listed = 63
Dipole Moments/Debye: $\mu_a =$		$\mu_b =$ magnetic	$\mu_c =$
Rot. Const./MHz: A =		B = 42626.9	C =

The calculations are described for the ground state (Species 32001). The vibrationally excited state, v = 1, is $1556.38 \pm 0.01 \text{ cm}^{-1}$ above the ground state (M. Loete and H. Berger, 1977, J. Molec. Spectry. 68, 317). Intensities were calculated using the ground state g values.

The value of Q was determined by a sum over states to J = 60. Other values are:

<u>T</u>	<u>Q</u>
10	8.348
30	22.819
60	44.772
100	74.097
150	110.779

Species Tag: 32003	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Methyl Alcohol (Methanol) CH ₃ OH lowest A, E ₁ , and E ₂ vibrational states		Author: H.M. Pickett & R.L. Poynter	
		Min. Int. = 10 ⁻¹⁰ nm ² MHz	
Q (Spin-Rot., T = 300K) = 6414.2516	J Max. = 12	Lines Listed = 142	
Dipole Moments/Debye: $\mu_a = 0.885$ $\mu_b = 1.440$ $\mu_c = 0$			
Rot. Const./MHz: A = 127,484. B = 24,679.98 C = 23769.70			

The lines below 200 GHz are reported by R. M. Lees, F. J. Lovas, W. H. Kirckhoff, and D. R. Johnson, 1973, J. Phys. Chem. Ref. Data, 2, 205. Additional lines are referenced in F. J. Lovas, L. E. Snyder, and D. R. Johnson, 1979, Ap. J. Suppl., 41.

The partition function is twice the value for a rigid rotor with the rotational constants given above. The quantum numbers for methanol are listed in the 840 format even though methanol is not a symmetric rotor. The v quantum numbers in the catalogue are:

<u>v</u>	<u>species</u>	<u>asymmetric rotor correspondence</u>
1	A+	upper K state for K even, lower for K odd
2	A-	upper K state for K odd, lower for K even
3	E ₁	no direct correspondence
4	E ₂	no direct correspondence

Species Tag: 32004	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Formaldehyde Oxygen - 18 isotope $\text{H}_2\text{C}^{18}\text{O}$		Author: R. L. Poynter	
		Min. Int. = 10^{-9} nm^2MHz	
Q (Spin-Rot., T = 300K) = 3017.226		J Max. = 20	Lines Listed = 449
Dipole Moments/Debye: $\mu_a = 2.331$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 281993.0 B = 36903.573 C = 32514.740			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- D. Dangoisse, E. Willemot, and J. Bellet, 1978, J. Mol. Spect. 71, 414.
D. R. Johnson, F. Lovas, and W. K. Kirchhoff, 1972, J. Phys. Chem. Ref. Data, 1, 1011.
R. B. Lawrence and M. W. P. Strandberg, 1951, Phys. Rev. 83, 363.
T. Oka, H. Hirakawa, and K. Shimoda, 1960, J. Phys. Soc. Japan, 15, 2265.
T. Oka, T. Takagi, and Y. Morino, 1964, J. Mol. Spect. 14, 27.
K. D. Tucker, G. R. Tomasevich, and P. Thaddeus, 1972, Ap. J. 174, 463, 1971, Ap. J., 169, 429.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 34001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: $18_0 16_0$		Author: H. M. Pickett	
Molecular Oxygen - 0^{18} species ground state		Min. Int. = 2.5×10^{-11} nm ² MHz	
Q (Spin-Rot., T = 300K) = 462.32		J Max. = (42)	Lines Listed = 132
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = 40708. C =			

The measurements and calculational method are from T. Amano and E. Hirota, 1974, J. Mol. Spectroscopy 53, 346. Intensities of Magnetic dipole transitions have been calculated using the g values for the $^{16}O_2$ species (K. M. Evenson and M. Mizushima, 1972, Phys. Rev. A 6, 2197). The electric dipole moment is expected to be $\sim 5 \times 10^{-5}$ Debye, based on the arguments of P. R. Bunker, 1973, J. Mol. Spectroscopy 46, 119. Therefore, no electric dipole transitions have been included in the catalogue. At high J the error estimates are likely to be optimistic since the expected errors in the assumed centrifugal distortion constants are not included.

The value of Q is determined by a sum over states to J = 60. Other values are:

<u>T</u>	<u>Q</u>
10K	17.13
30	47.46
60	93.44
100	154.85
150	231.67

Species Tag: 34002	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Hydrogen sulfide		Author: R. L. Poynter	
H ₂ S		Min. Int. = 10 ⁻⁸ nm ² MHz	
Q (Spin-Rot., T = 300K) = 512.21		J Max. = 10	Lines Listed = 213
Dipole Moments/Debye: $\mu_a = 0.$ $\mu_b = 0.974$ $\mu_c = 0.$			
Rot. Const./MHz: A = 310182.24 B = 270884.05 C = 141705.880			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

P. Helminger, R. L. Cook, and F. C. De Lucia, 1972, J. Chem. Phys. 56, 4581.
P. Helminger, F. C. De Lucia, and W. H. Kirchhoff, 1973, J. Phys. Chem. Ref. Data; 2, 213,

The dipole moment was reported by C. Huiszoon and A. Dymanus, 1965, Physica, 31, 1049.

Species Tag: 34003	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Phosphine PH_3		Author: R. L. Poynter	
		Min. Int. = 10^{-18} nm ² MHz	
Q (Spin-Rot., T = 300K) = 608.55		J Max. = (19)	Lines Listed = 126
Dipole Moments/Debye: $\mu_a = *8.3 \times 10^{-5}$ $\mu_b =$ $\mu_c = 0.574$			
Rot. Const./MHz: A = B B = 33480.15 C = 117488.85			

*centrifugally induced

The measured lines are taken from:

D. Helms and W. Gordy (1977), J. Mol. Spect., 66, 206, F. Y. Chu and T. Oka, (1974), J. Chem. Phys. 60, 4612, P. B. Davies, R. M. Newman, S. C. Wofsy, and W. Klemperer, (1971), J. Chem. Phys. 55, 3564.

The J = 1 → 2 lines measured by A. F. Krupnov, A. A. Melnikov, and V. A. Skvortsov (1979) Opt. Spectrosc. (USSR) 46 (5), 570, came to our attention too late to include in this analysis.

The dipole moment measured by P. B. Davies, et al., above, is $0.57395 \pm 0.0003D$.

Species Tag: 35001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Hydrogen Sulfide Deuterium Isotope HDS		Author: R. L. Poynter	
		Min. Int. = 4×10^{-12} cm^2MHz	
Q (Spin-Rot., T = 300K) = 430.900	J Max. = 20	Lines Listed = 1138	
Dipole Moments/Debye: $\mu_a = 0.470$ $\mu_b = 0.974$ $\mu_c = 0.$			
Rot. Const./MHz: A = 292351.302 B = 147861.801 C = 96704.120			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 40, 125. G. Steenbeckliens, quoted in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was reported by R. E. Hillger and M. W. P. Strandberg, 1951, Phys. Rev. 83, 575.

Species Tag: 36001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Hydrogen Chloride H ³⁵ Cl		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 5×10^{-2} cm ² MHz	
Q (Spin-Rot., T = 300K) = 81.232		J. Max. = 4	Lines Listed = 17
Dipole Moments/Debye: $\mu_a = 1.109$ $\mu_b = 0$ $\mu_c = 0$			
Rot. Const./MHz: A = B = 312989.3 C =			

The observed lines were measured by F. C. De Lucia, P. Helminger, and W. Gordy, 1971, Phys. Rev., A 3, 1849.

The dipole moment was measured by F. A. deLeeuw and A. Dymanus, 1971, Symposium on Molec. Spect., Columbus, Ohio.

Species Tag: 38001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Hydrogen Chloride Chlorine - 37 isotope $H^{37}Cl$		Author: R. L. Poynter & H. L. Pickett	
		Min. Int. = 5×10^{-2} cm ² MHz	
Q (Spin-Rot., T = 300K) = 81.352	J Max. = 4	Lines Listed = 17	
Dipole Moments/Debye: $\mu_a = 1.109$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 312519.12 C =			

The observed lines were measured by F. C. De Lucia, P. Helminger, and W. Gordy, 1971, Phys. Rev., A 3, 1849.

The dipole moment was measured by F. A. DeLeeuw and A. Dymanus, 1971, Symposium on Molec. Spect., Columbus, Ohio.

Species Tag: 40001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Propyne CH ₃ CCH		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} cm ² MHz for J > 14	
Q (Spin-Rot., T = 300K) = 5428.80	J Max. = 80	Lines Listed = 813	
Dipole Moments/Debye: $\mu_a = 0.750$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158590. B = 8545.860 C = B			

The experimental measurements were obtained from A. Dubrille, D. Boucher, J. Burie, and J. Demaison, 1978, J. Mol. Spect. 72, 158. The A moment was estimated from the structure.

The dipole moment was measured by J. S. Muenter and V. W. Laurie, 1966, J. Chem. Phys. 45, 855.

Species Tag: 41001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Acetonitrile CH ₃ CN ground state		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} cm ² MHz for J > 14	
Q (Spin-Rot., T = 300K) = 15145.		J Max. = 82	Lines Listed = 1441
Dipole Moments/Debye: $\mu_a = 3.919$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158290. B = 9198.8993 C = B			

The experimental measurements were taken from D. Boucher, J. Burie, J. Demaison, A. Dubrille, J. Legrand, and B. Segard, 1977, J. Mol. Spect. 64, 290; A. Bauer and S. Maes, 1969, J. Phys. 30, 169; S. G. Kukolich, D. J. Ruben, J. H. S. Wang, and J. R. Williams, 1973, J. Chem. Phys. 58, 3155.

The dipole moment was reported by P. A. Steiner and W. Gordy, 1966, J. Mol. Spect. 21, 291.

Species Tag: 41002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Propyne, Carbon - 13 isotope on atom 1 $\text{CH}_3 \text{C}^{13}\text{CH}$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} cm^2/MHz for J > 14	
Q (Spin-Rot., T = 300K) = 5596.11		J Max. = 80	Lines Listed = 822
Dipole Moments/Debye: $\mu_a = 0.750$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158590. B = 8290.259 C = B			

The experimental measurements were taken from D. Boucher, J. Burie, J. Demaison, A. Dubrille, J. Legrand, and B. Segard, 1977, J. Mol. Spect. 64, 290; A. Bauer and S. Maes, 1969, J. Phys. 30, 169; S. G. Kukolich, D. J. Ruben, J. H. S. Wang, and J. R. Williams, 1973, J. Chem. Phys. 58, 3155.

The dipole moment was reported by P. A. Steiner and W. Gordy, 1966, J. Mol. Spect. 21, 291.

Species Tag: 41003	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Propyne, Carbon - 13 isotope on atom 2 $\text{CH}_3^{13}\text{CCH}$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} nm ² MHz for J > 14	
Q (Spin-Rot., T = 300K) = 5431.06	J Max. = 79	Lines Listed = 813	
Dipole Moments/Debye: $\mu_a = 0.750$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158590. B = 8542.304 C = B			

The experimental measurements were taken from D. Boucher, J. Burie, J. Demaison, A. Dubrille, J. Legrand, and B. Segard, 1977, J. Mol. Spect. 64, 290; A. Bauer and S. Maes, 1969, J. Phys. 30, 169; S. G. Kukolich, D. J. Ruben, J. H. S. Wang, and J. R. Williams, 1973, J. Chem. Phys. 58, 3155.

The dipole moment was reported by P. A. Steiner and W. Gordy, 1966, J. Mol. Spect. 21, 291.

Species Tag: 41004	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Propyne, carbon - 13 isotope on atom 3 $H_3^{13}CCCH$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} nm ² MHz for J > 14	
Q (Spin-Rot., T = 300K) = 5580.65		J Max. = 80	Lines Listed = 821
Dipole Moments/Debye: $\mu_a = 0.750$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158590. B = 81313.24 C = B			

The experimental measurements were taken from D. Boucher, J. Burie, J. Demaison, A. Dubrille, J. Legrand, and B. Segard, 1977, J. Mol. Spect. 64, 290; A. Bauer and S. Maes, 1969, J. Phys, 30, S. G. Kukolich, D. J. Ruben, J. H. S. Wang, and J. R. Williams, 1973, J. Chem. Phys. 58, 3155.

The dipole moment was reported by P. A. Steiner and W. Gordy, 1966, J. Mol. Spect. 21, 291.

Species Tag: 41005	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Propyne, Deuterium isotope on carbon atom 1 CH ₃ CCD		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} nm ² MHz for J > 14	
Q (Spin-Rot., T = 300K) = 5956.732		J Max. = 82	Lines Listed = 800
Dipole Moments/Debye: $\mu_a = 0.750$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158590.0 B = 7788.158 C = B			

The experimental measurements are from J. S. Muentner and V. W. Laurie, 1966, J. Chem. Phys. 45, 855; L. F. Thomas, E. I. Sherrard, and J. Sheridan, 1955, Trans. Far. Soc. 51, 619.

The dipole moment was assumed to be the same as the parent species for this calculation. An experimental value, 0.770D (avg.) corrected for the new OCS standard, has been reported by Muentner and Laurie (above). This result came to our attention too late to include in this version.

Species Tag: 41006	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Propyne Deuterium isotope on methyl carbon atom CH ₂ D CCH		Author: R. L. Poynter	
		Min. Int. = 2×10^{-10} nm ² MHz	
Q (Spin-Rot., T = 300K) = 9982.44		J Max. = 15	Lines Listed = 223
Dipole Moments/Debye: $\mu_a = 0.750$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 117744.370 B = 8155.784 C = 8025.577			

The experimental measurements are from L. F. Thomas, E. I. Sherrard, and J. Sheridan, 1955, Trans. Far. Soc. 54, 619.

The dipole moment was assumed to be the same as the parent species for this calculation.

Species Tag: 42001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Acetonitrile Nitrogen - 15 isotope $\text{CH}_3 \text{C}^{15}\text{N}$		Author: R. L. Poynter	
		Min. Int. = 2×10^{-5} nm^2MHz for J > 14	
Q (Spin-Rot., T = 300K) = 15145		J Max. = 90	Lines Listed = 2755
Dipole Moments/Debye: $\mu_a = 3.919$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = 158290. B = 9198.8993 C = B			

The experimental measurements were obtained from A. Bauer and S. Maes, 1969, J. de Phys. 30, 169; J. Demaison, A. Dubrulle, D. Boucher, J. Burie, 1969, J. Mol. Spect. 76, 1.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 42002	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ketene H_2CCO		Author: R. L. Poynter	
		Min. Int. = 3×10^{-15} nm ² MHz	
Q (Spin-Rot., T = 300K) = 10322.9		J Max. = 29	Lines Listed = 835
Dipole Moments/Debye: $\mu_a = 1.422$ $\mu_b = 0$ $\mu_c = 0$			
Rot. Const./MHz: A = 282473 B = 10293.80 C = 9916.38			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- B. Fabricant, D. Krieger, and J. S. Muentex, 1977, J. Chem. Phys. 67, 1576.
 J. W. C. Johns, J. M. R. Stone, and G. Winnewisser, 1972, J. Mol. Spect. 42, 523.
 H. R. Johnson and M. W. P. Strandberg, 1952, J. Chem. Phys. 20, 687.

The dipole moment was reported by Johnson (above) and by B. Fabricant, et al. (above).

Species Tag: 43001	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ketene Deuterium isotope HDCCO		Author: R. L. Poynter	
		Min. Int. = 10^{-23} nm ² MHz	
Q (Spin-Rot., T = 300K) = 6682.68	J Max. = 29	Lines Listed = 886	
Dipole Moments/Debye: $\mu_a = 1.422$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 194313.0 B = 9647.396 C = 9174.975			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

L. Nemes and M. Winnewisser, 1976, Z. Naturforsch. 31a, 272.

The dipole moment was assumed the same as the parent species.

Species Tag: 44001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbon Monosulfide ground state CS		Author: R. L. Poynter	
		Min. Int. = 4×10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 255.515		J Max. = (60)	Lines Listed = 60
Dipole Moments/Debye: $\mu_a = 1.957$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 24495.554 C =			

The experimental data were taken from the following papers:

R. Kewley, K. V. L. N. Sastry, M. Winnewisser, and W. Gordy, 1963, J. Chem. Phys. 39, 2856.

R. C. Mockler and G. R. Bird, 1955, Phys. Rev. 98, 1837.

The dipole moment is from G. Winnewisser and R. L. Cook, 1968, J. Mol. Spect. 28, 266.

Species Tag: 44002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monoxide $^{28}\text{Si O}$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 287.232	J Max. = 66	Lines Listed = 66	
Dipole Moments/Debye: $\mu_a = 3.098$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 21787.46 C =			

The experimental data were taken from the following papers:

E. Tiemann, 1974, J. Phys. Chem. Ref. Data, T. Törring, 1968, Z. Naturforsch. 23a, 777.

The dipole moment was reported by J. W. Raymonda, J. S. Muentner, and W. A. Klemperer, 1970, J. Chem. Phys. 52, 3458.

A more recent set of measurements, discovered too late to include in this edition, has been given by E. L. Manson, W. W. Clark, F. C. De Lucia, and W. Gordy, 1977, Phys. Rev. A15, 223. The catalogue predictions are within three standard deviations of their measurements.

Species Tag: 44003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Acetaldehyde ground vibrational A state CH ₃ CHO		Author: R. L. Poynter	
		Min. Int. = 6×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 12112.28		J Max. = 20	Lines Listed = 1347
Dipole Moments/Debye: $\mu_a = 2.550$ $\mu_b = 0.870$ $\mu_c = 0.$			
Rot. Const./MHz: A = 56609.474 B = 10162.766 C = 9100.412			

The data set used is referenced by A. Bauder, F. J. Lovas and D. R. Johnson 1976, J. Phys. Chem. Ref. Data 5, 53. The lines were fit to a rigid rotor with centrifugal distortion.

Species Tag: 44004	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Nitrous Oxide N_2O		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 2×10^{-7} nm^2MHz	
Q (Spin-Rot., T = 300K) =	J Max. = 61	Lines Listed = 61	
Dipole Moments/Debye: $\mu_a = .1608$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 12561.637 C =			

The experimental data were taken from the summary given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was measured by L. H. Scharpen, J. S. Muenter, and V. W. Laurie, 1970, J. Chem. Phys. 53, 2513.

Species Tag: 45001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbon Monosulfide Carbon - 13 isotope ^{13}CS		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} cm^2MHz	
Q (Spin-Rot., T = 300K) = 270.652		J Max. = (62)	Lines Listed = 62
Dipole Moments/Debye: $\mu_a = 1.957$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 23123.807 C =			

The experimental measurement was reported by R. C. Mockler and G. R. Bird, 1955, Phys. Rev. 98, 1837.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 45002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monoxide silicon - 29 isotope $^{29}\text{Si O}$		Author: R. L. Poynter	
		Min. Int. = 4×10^{-5} cm^2MHz	
Q (Spin-Rot., T = 300K) = 290.878		J Max. = 66	Lines Listed = 66
Dipole Moments/Debye: $\mu_a = 3.098$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 21514.07 C =			

The experimental data were taken from the following papers:

E. Tiemann, 1974, J. Phys. Chem. Ref. Data, T. Törring, 1968, Z. Naturforsch. 23a, 777.

The dipole moment was reported by J. W. Raymonda, J. S. Muentzer, and W. A. Klemperer, 1970, J. Chem. Phys. 52, 3458.

A more recent set of measurements, discovered too late to include in this edition, has been given by E. L. Manson, W. W. Clark, F. C. De Lucia, and W. Gordy, 1977, Phys. Rev. A15, 223. The catalogue predictions are within three standard deviations of their measurements.

Species Tag: 45003	Page: 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Formamide NH ₂ CHO		Author: R. L. Poynter	
		Min. Int. = 10 ⁻¹¹ nm ² MHz	
Q (Spin-Rot., 1 - 300K) - 29155.75		J Max. = 30	Lines Listed = 3246
Dipole Moments/Debye: μ_a 3.616		μ_b = 0.852	μ_c = 0.
Rot. Const./MHz: A = 72716.945		B = 11373.453	C = 9833.903

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- C. C. Costain and J. M. Dowling, 1960, J. Chem. Phys. 32, 290.
- D. R. Johnson, F. J. Lovas, and W. H. Kirchhoff, 1972, J. Phys. Chem. Ref. Data, 1, 1011.
- W. H. Kirchhoff and D. R. Johnson, 1973, J. Mol. Spect. 45, 159.
- S. G. Kukolich and A. C. Nelson, 1971, Chem. Phys. Lett. 11, 383.
- R. J. Kurland and E. B. Wilson, Jr., 1957, J. Chem. Phys. 27, 585.

The dipole moment was reported in the paper by Kurland and Wilson.

Species Tag: 46001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbon Monosulfide Sulfur - 34 isotope $C^{34}S$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 259.665	J Max. = (61)	Lines Listed = 61	
Dipole Moments/Debye: $\mu_a = 1.957$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 24103.550 C =			

The experimental measurement was reported by R. C. Mockler and G. R. Bird, 1955, Phys. Rev. 98, 1837.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 46002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monoxide Silicon - 30 isotope $^{30}\text{Si O}$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 294.358		J Max. = 67	Lines Listed = 67
Dipole Moments/Debye: $\mu_a = 3.098$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 21259.48 C =			

The experimental data were taken from the following papers:

E. Tiemann, 1974, J. Phys. Chem. Ref. Data, T. Törring, 1968, Z. Naturforsch. 23a, 777.

The dipole moment was reported by J. W. Raymonda, J. S. Muentner, and W. A. Klemperer, 1970, J. Chem. Ref. Data, T. Törring, 1968, Z. Naturforsch.

A more recent set of measurements, discovered too late to include in this edition, has been given by E. L. Manson, W. W. Clark, F. C. De Lucia, and W. Gordy, 1977, Phys. Rev. A15, 223. The catalogue predictions are within three standard deviations of their measurements.

Species Tag: 46003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Thioformaldehyde H_2CS		Author: R. L. Poynter	
		Min. Int. = 8×10^{-10} $nm^2 MHz$	
Q (Spin-Rot., T = 300K) = 5984.646		J Max. = 27	Lines Listed = 517
Dipole Moments/Debye: $\mu_a = 1.649$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 291291.641 B = 17699.628 C = 16651.830			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

D. R. Johnson, F. J. Lovas, and W. H. Kirchhoff, 1972, J. Phys. Chem. Ref. Data 1, 1011.

Other references were taken from this paper.

The dipole moment was taken from B. Fabricant, D. Krieger, and J. S. Muentzer, 1977, J. Chem. Phys. 67, 1576.

Species Tag: 46004	Page 1 of 1	Version: 1	Date: Jan 1980
Species Name: Ethyl Alcohol ground trans state C_2H_5OH		Author: R. L. Poynter	
		Min. Int. = 3×10^{-10} nm ² MHz	
Q (Spin-Rot., T = 300K) = 17010.5	J Max. = 20	Lines Listed = 1236	
Dipole Moments/Debye: $\mu_a = 0.046$ $\mu_b = 1.438$ $\mu_c = 0.$			
Rot. Const./MHz: A = 34891.75 B = 9350.635 C = 8135.236			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

J. Michielson-Effinger, 1969, J. de Phys. 30, 333.

Y. Sasada, M. Takano, and T. Satoh, 1971, J. Mol. Spect. 38, 33.

M. Takano, Y. Sasada, and T. Satoh, 1968, J. Mol. Spect. 26, 157.

Additional measurements were made by E. A. Cohen, 1979, private communication.

The dipole moment was reported by Takano, et al, above.

Species Tag: 46005	Page 1 of 1	Version: 1	Date: Jan, 1980
Species Name: Formic Acid ground trans state HCOOH		Author: R. L. Poynter	
		Min. Int. = 6×10^{-10} nm ² MHz	
Q (Spin-Rot., T = 300K) = 8884.2	J Max. = 20	Lines Listed = 1888	
Dipole Moments/Debye: $\mu_a = 1.396$ $\mu_b = 0.260$ $\mu_c = 0.$			
Rot. Const./MHz: A = 77512.25 B = 12055.11 C = 10416.12			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- J. Bellet, C. Samson, G. Steenbeckeliers, and R. Wertheimer, 1971, J. Mol. Structure 9, 49.
P. G. Lerner, B. P. Dailey, and J. P. Friend, 1957, J. Chem. Phys. 26, 680.
R. Trambarulo, A. Clark, and C. Hearn, 1958, J. Chem. Phys. 28, 736.

The dipole moment measured by H. Kim, R. Keller, and W. D. Gwinn, 1962, J. Chem. Phys. 37, 2748, was adjusted to the new OCS standard, (J. S. Muentzer, 1968, J. Chem. Phys. 48, 4544).

Species Tag: 47001	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Thioformaldehyde		Author: R. L. Poynter	
Carbon - 13 isotope $H_2^{13}CS$		Min. Int. = 2×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 6219.87	J Max. = 10	Lines Listed = 110	
Dipole Moments/Debye: $\mu_a = 1.649$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 291660.0 B = 16998.342 C = 16030.791			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

D. R. Johnson, F. J. Lovas, and W. H. Kirchhoff, 1972, J. Phys. Chem. Ref. Data 1, 1011.

Other references were taken from this paper.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 47002	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Formic Acid trans state Carbon - 13 isotcpe H ¹³ COOH		Author: R. L. Poynter	
		Min. Int. = 10 ⁻¹⁰ nm ² MHz	
Q (Spin-Rot., T = 300K) = 8896.76	J Max. = 20	Lines Listed = 1194	
Dipole Moments/Debye: $\mu_a = 1.396$ $\mu_b = 0.260$ $\mu_c = 0.$			
Rot. Const./MHz: A = 77580.494 B = 12053.567 C = 10378.997			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- J. Bellet, A. Deldalle, C. Samson, G. Steenbeckeliers, and R. Wertheimer, 1971, J. Mol. Structure, 9, 65.
R. G. Lerner, B. P. Dailey, and J. P. Friend, 1957, J. Chem. Phys. 26, 680.

The dipole moment was assumed the same as the parent species, tag #46005.

Species Tag: 47003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Formic Acid Deuterium isotope on C-atom DCOOH		Author: R. L. Poynter	
		Min. Int. = 2×10^{-9} cm ² MHz	
Q (Spin-Rot., T = 300K) = 10531.19	J Max. = 20	Lines Listed = 628	
Dipole Moments/Debye: $\mu_a = 1.396$ $\mu_b = 0.260$ $\mu_c = 0.$			
Rot. Const./MHz: A = 57709.33 B = 12055.971 C = 9955.609			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

J. Bellet, A. Deldalle, C. Samson, G. Steenbackeliers, and R. Wertheimer, 1971, J. Mol. Structure, 9, 65.

R. G. Lerner, B. P. Dailey, and J. P. Friend, 1957, J. Chem. Phys. 26, 680.

The dipole moment was assumed the same as the parent species, tag #46005.

Species Tag: 47004	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Formic Acid Deuterium isotope on O-atom HCOOD		Author: R. L. Poynter	
		Min. Int. = 10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 9954.87	J Max. = 20	Lines Listed = 612	
Dipole Moments/Debye: $\mu_a = 1.396$ $\mu_b = 0.260$ $\mu_c = 1.$			
Rot. Const./MHz: A = 66100.14 B = 11762.577 C = 9969.943			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

J. Bellet, A. Deldalle, C. Samson, G. Steenbeckeliers, and R. Wertheimer, 1971, J. Mol. Structure, 9, 65.

R. G. Lerner, B. P. Dailey, and J. P. Friend, 1957, J. Chem. Phys. 26, 680.

The dipole moment was assumed the same as the parent species, tag #46005.

Species Tag: 48001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Sulfur monoxide SO $^3\Sigma$ ground state		Author: H. M. Pickett	
		Min. Int. = 10^{-6} cm^2MHz	
Q (Spin-Rot., T = 300K) = 850.099		J Max. = 50	Lines Listed = 330
Dipole Moments/Debye: $\mu_a = 1.55$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 21523.02 C =			

The calculational method is described in T. Amano and E. Hirota, 1974, J. Molec. Spectroscopy 53, 346. The data used is from W. W. Clark and F. C. DeLucia, 1976, J. Molec. Spectroscopy 60, 332. Both electric dipole and magnetic dipole transitions are listed.

The partition function was obtained by a sum over states to J=50. Other values are

<u>I</u>	<u>Q</u>
10K	17.284
30	69.382
60	154.400
100	269.677
150	414.502

Species Tag: 48002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Sulfur Monoxide $v = 1$ vibrationally excited state of the ground $^3\Sigma$ electronic state		Author: H. M. Pickett	
		Min. Int. = 4×10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 856.726		J Max. = 50	Lines Listed = 261
Dipole Moments/Debye: $\mu_a = 1.55$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 21351.0 C =			

The calculational method is described in T. Amano and E. Hirota, 1974, J. Molec. Spectroscopy 53, 346. The data are from T. Amano, E. Hirota, and Y. Morino, 1967, J. Phys. Soc. Japan 22, 399. All centrifugal distortion constants were fixed to their ground state values. Magnetic and electric dipole transitions are included.

This state is 1111.5 cm^{-1} above $v = 0$ (G. Herzberg, 1950, Spectra of Diatomic Molecules, Van Nostrand, New York). The partition function is determined by a sum over states to $J = 50$. Other values are:

<u>T</u>	<u>Q</u>
10K	17.341
30	69.781
60	155.452
100	271.644
150	417.628

Species Tag: 48003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Thioformaldehyde		Author: R. L. Poynter	
Sulfur - 34 isotope $H_2C^{34}S$		Min. Int. = 3×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 6084.57		J Max. = 10	Lines Listed = 111
Dipole Moments/Debye: $\mu_a = 1.647$ $\mu_b = 0.$ $\mu_c = 0.$			
Rot. Const./MHz: A = 291660. B = 17387.949 C = 16376.922			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

D. R. Johnson, F. J. Lovas, and W. H. Kirchhoff, 1972, J. Phys. Chem. Ref. Data 1, 1011.

Other references were taken from this paper.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 48004	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ozone ground state $^{16}\text{O}_3$		Author: R. L. Poynter and H. M. Pickett	
		Min. Int. = 3×10^{-10} nm^2MHz	
Q (Spin-Rot., T = 300K) = 3378.28		J Max. = 10	Lines Listed = 1465
Dipole Moments/Debye: $\mu_a = 0.$ $\mu_b = 0.5324$ $\mu_c = 0.$			
Rot. Const./MHz: A = 106535.234 B = 13349.0901 C = 11834.5235			

The catalog of the ozone lines is based on the work of M.J.C. Depannemaecker, B. Duterage, and M. J. Bellet, 1977, J. Quant. Spect. Radiat. Transfer 17, 519. Additional lines are cited in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data 7, 1445.

Species Tag: 50001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Sulfur Monoxide Sulfur 34 isotope $^3\Sigma$ ground state ^{34}SO		Author: H. M. Pickett	
		Min. Int. = 10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 866.870		J Max. = 50	Lines Listed = 280
Dipole Moments/Debye: $\mu_a = 1.55$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 21102.72 C =			

The calculational method is described in T. Amano and E. Hirota, 1974, J. Molec. Spectroscopy 53, 346. The data used is reported in E. Tiemann, 1974, J. Phys. Chem. Ref. Data 3, 259. Both electric dipole and magnetic dipole transitions are listed in the catalogue.

The partition function was determined by a sum over states. Other values are:

<u>T</u>	<u>Q</u>
10K	17.554
30	70.654
60	157.354
100	274.922
150	422.629

Species Tag: 50002	Page 1 of 1	Version: 1	Date: Dec, 1979
Species Name: Sulfur Monoxide Oxygen 18 isotope		Author: H. M. Pickett	
		Min. Int. = 10^{-5} nm ² MHz	
Q (Spin-Rot., T = 300K) = 917.393		J Max. = 50	Lines Listed = 179
Dipole Moments/Debye: $\mu_a = 1.55$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 19929. C =			

The calculational method is described in T. Amano and E. Hirota, 1974, J. Molec. Spectroscopy 53, 346. The value of μ was fixed at -156.51 MHz and $\mu(1)$ was set to zero. Both electric dipole and magnetic dipole transitions are listed in the catalogue. The data used is listed in E. Tiemann, 1974, J. Phys. Chem. Ref. Data. 3, 259.

The partition function was determined by a sum over states. Other values are:

<u>T</u>	<u>Q</u>
10K	18.370
30	74.493
60	166.257
100	290.728
150	447.118

C-2

Species Tag: 50003	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Symmetric ^{18}O Ozone ground state $^{16}\text{O}-^{18}\text{O}-^{16}\text{O}$		Author: H. M. Pickett & R. L. Poynter	
		Min. Int. = 10^{-9} nm^2MHz	
Q (Spin-Rot., T = 300K) = 3525.63		J Max. = 40	Lines Listed = 1568
Dipole Moments/Debye: $\mu_a = 0.$ $\mu_b = 0.5324$ $\mu_c = 0.$			
Rot. Const./MHz: A = 98646.687 B = 13352.732 C = 11731.767			

The catalog of symmetric ^{18}O ozone is based on the work of J. Depannemaecker and J. Bellet (1977, J. Molo Specty. 66, 106). The dipole moment used is the $^{16}\text{O}_3$ value.

Species Tag: 50004	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Asymmetric ^{18}O Ozone $^{18}\text{O}-^{16}\text{O}-^{16}\text{O}$		Author: H. M. Pickett & R. L. Poynter	
		Min. Int. = 10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 7213.74		J Max. = 40	Lines Listed = 4036
Dipole Moments/Debye: $\mu_a = 0.0068$ $\mu_b = 0.5324$ $\mu_c = 0.$			
Rot. Const./MHz: A = 104573.161 B = 12591.533 C = 11212.5057			

The catalog of asymmetric ^{18}O ozone is based on the work of J. Depamemaecker and J. Bellet (1977, J. Mol. Specty. 66, 106). The dipole moment used is the $^{16}\text{O}_3$ value rotated to the inertial axes of asymmetric ^{18}O ozone (based on the equilibrium structure).

Species Tag: 51001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanoacetylene HCCCN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} cm^2MHz	
Q (Spin-Rot., T = 300K) = 4123.26		J Max. = 99	Lines Listed = 154
Dipole Moments/Debye: $\mu_a = 3.599$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 4549.059 C =			

The set of experimental lines used in the calculation was obtained from the references in W. J. Lafferty and F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 441.

The dipole moment was remeasured by the above authors as 3.724D. An old value, 3.599D, was used in the present calculations, so the intensities should be multiplied by a factor of 1.071.

Species Tag: 51002	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Chlorine Monoxide 2π states ^{35}ClO $v = 0$		Author: H. M. Pickett	
		Min. Int. = 10^{-10} nm^2MHz	
Q (Spin-Rot., T = 300K) = 3289.29		J Max. = 83	Lines Listed = 2610
Dipole Moments/Debye: $\mu_a = 1.239$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 18602.86 C =			

The experimental lines below 200 GHz are from R. K. Kakar, E. A. Cohen, and M. Geller, 1978, J. Mol. Spec. 70, 243. Lines above 200 GHz were measured at JPL by E. A. Cohen and H. M. Pickett. The lines were fit to a fine structure Hamiltonian which included p_D and q_D terms and centrifugal distortion on B to sixth power in J. The hyperfine terms in the Hamiltonian included all off-diagonal matrix elements for a, b, c, d, η_1 , η_2 , and η_3 . The partition function was determined by a sum over states to F = 49 for both the $\Omega = 1/2$ and $\Omega = 3/2$ states. Other values are:

<u>T</u>	<u>Q</u>
150	1424.27
100	921.36
60	553.46
30	284.09
10	105.25

Species Tag: 52001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanoacetylene carbon - 13 isotope on atom 1 H^{13}CCN		Author: R. L. Poynter	
Q (Spin-Rot., T = 300K) = 4140.42		J Max. = 99	Min. Int. = 2×10^{-7} nm ² MHz
Dipole Moments/Debye: $\mu_a = 3.599$		$\mu_b =$	$\mu_c =$
Rot. Const./MHz: A =		B = 4530.198	C =

The set of experimental lines used in the calculation was obtained from the references in W. J. Lafferty and F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 441.

The dipole moment was remeasured by the above authors as 3.724D. An old value, 3.599D, was used in the present calculations, so the intensities should be multiplied by a factor of 1.071.

Species Tag: 52002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanoacetylene Carbon - 13 isotope on atom 2 HC ¹³ CCN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4140.82	J Max. = 99	Lines Listed = 146	
Dipole Moments/Debye: $\mu_a = 3.599$ $\mu_b =$ $\mu_c =$			
Rot. Const./ MHz: A = B = 4529.76 C =			

The set of experimental lines used in the calculation was obtained from the references in W. J. Lafferty and F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 441.

The dipole moment was remeasured by the above authors as 3.724D. An old value, 3.599D, was used in the present calculations, so the intensities should be multiplied by a factor of 1.071.

Species Tag: 52003	Page 1 of 1	Version: 1	Date: Dec, 1979
Species Name: Cyanoacetylene Carbon - 13 isotope on atom 3 H ¹³ CCCN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} cm ² MHz	
Q (Spin-Rot., T = 300K) = 4254.75		J Max. = 99	Lines Listed = 144
Dipole Moments/Debye: $\mu_a = 3.599$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 4408.44 C =			

The set of experimental lines used in the calculation was obtained from the references in W. J. Lafferty and F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 441.

The dipole moment was remeasured by the above authors as 3.724D. An old value, 3.599D, was used in the present calculations, so the intensities should be multiplied by a factor of 1.071.

Species Tag: 52004	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanoacetylene Nitrogen - 15 isotope, HCCC ¹⁵ N		Author: R. L. Poynter	
		Min. Int. = 4×10^{-6} mm ² MHz	
Q (Spin-Rot., T = 300K) = 1415.581	J Max. = 99	Lines listed = 99	
Dipole Moments/Debye: $\mu_a = 3.599$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 4416.75 C =			

The set of experimental lines used in the calculation was obtained from the references in W. J. Lafferty and F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 441.

The dipole moment was remeasured by the above authors as 3.724D. An old value, 3.599D, was used in the present calculations, so the intensities should be multiplied by a factor of 1.071.

Species Tag: 52005	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanoacetylene Deuterium Isotope DCCCN		Author: R. L. Poynter	
		Min. Int. = 10^{-7} cm^2MHz	
Q (Spin-Rot., T = 300K) = 4443.03		J Max. = 99	Lines Listed = 156
Dipole Moments/Debye: $\mu_a = 3.724$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 4221.58 C =			

The set of experimental lines used in the calculation was obtained from the references in W. J. Lafferty and F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 441.

Species Tag: 53001	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Acrylonitrile $C_2 H_3 CN$		Author: R. L. Pcynter	
		Min. Int. = 10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 26198.1	J Max. = 40	Lines Listed = 3697	
Dipole Moments/Debye: $\mu_a = 3.68$ $\mu_b = 1.25$ $\mu_c = 0.$			
Rot. Const./MHz: A = 49850.712 B = 4971.0849 C = 4513.8005			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

- C. C. Costain and B. P. Stoicheff, 1959, J. Chem. Phys. 30, 777.
M. C. L. Gerry and G. Winnewisser, 1973, J. Mol. Spect. 48, 1.
M. C. L. Gerry, K. Yamada, and G. Winnewisser, 1979, J. Phys. Chem. Ref. Data 8, 107.

The dipole moment was measured by W. W. Wilcox, J. H. Goldstein, and J. W. Simmons, 1954, J. Chem. Phys. 22, 516.

Species Tag: 53002	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Chlorine Monoxide $^2\Pi$ states Chlorine - 37 isotope ^{37}ClO $v = 0$		Author: H. M. Pickett	
		Min. Int. = 10^{-10}	cm^2MHz
Q (Spin-Rot., T = 300K) = 3403.2		J Max. = 25	Lines Listed = 926
Dipole Moments/Debye: $\mu_a = 1.239$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = $\beta = 1828.70$ C =			

The experimental and calculational method lines are from R. K. Kakar, E. A. Cohen, M. Geller, 1978, J. Mol. Spec. 70, 243. The partition function was scaled from ^{35}ClO by the ratio of the B values.

Species Tag: 54001	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Acrylonitrile		Author: R. L. Poynter	
Carbon - 13 isotope on atom 1		Min. Int. = 4×10^{-9} nm ² MHz	
CH ₂ CH ¹³ CN			
Q (Spin-Rot., T = 300K) = 26333.	J Max. = 10	Lines Listed = 118	
Dipole Moments/Debye: $\mu_a = 3.68$ $\mu_b = 1.25$ $\mu_c = 0$			
Rot. Const./MHz: A = 49781. B = 4948.153 C = 4494.485			

The methods used on this species were the same as for the parent species, given in tag #53001. The measurements were taken from C. C. Costain and B. P. Stoicheff, 1959, J. Chem. Phys. 30, 77.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 54002	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Acrylonitrile carbon - 13 isotope on atom 2 $\text{CH}_2^{13}\text{CHCN}$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-9} nm^2MHz	
Q (Spin-Rot., T = 300K) = 26665.		J Max. = 10	Lines Listed = 117
Dipole Moments/Debye: $\mu_a = 3.68$ $\mu_b = 1.25$ $\mu_c = 0$			
Rot. Const./MHz: A = 48645. B = 4948.700 C = 4485.145			

The methods used on this species were the same as for the parent species, given in tag #53001. The measurements were taken from C. C. Costain and B. P. Stoicheff, 1959, J. Chem. Phys. 30, 77.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 54003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Acrylonitrile Carbon - 13 isotope on atom 3 $^{13}\text{CH}_2\text{CHCN}$		Author: R. L. Poynter	
		Min. Int. = 3×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 27087.5		J Max. = 10	Lines Listed =
Dipole Moments/Debye: $\mu_a = 3.68$ $\mu_b = 1.25$ $\mu_c = 0$			
Rot. Const./MHz: A = 49180. B = 4837.34 C = 4398.07			

The methods used on this species were the same as for the parent species, given in tag #53001. The measurements were taken from C. C. Costain and B. P. Stoicheff, 1959, J. Chem. Phys. 30, 77.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 54004	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Acrylonitrile Deuterium isotope on atom 2 CH ₂ DCN		Author: R. L. Poynter	
		Min. Int. = 9×10^{-10} cm ² MHz	
Q (Spin-Rot., T = 300K) = 29697.8	J Max. = 10	Lines Listed = 322	
Dipole Moments/Debye: $\mu_a = 3.68$ $\mu_b = 1.25$ $\mu_c = 0$			
Rot. Const./MHz: A = 40198.6 B = 4934.35 C = 4388.41			

The methods used on this species were the same as for the parent species, given in tag #53001. The measurements were taken from C. C. Costain and B. P. Stoicheff, 1959, J. Chem. Phys. 30, 77.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 55001	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide C_2H_5CN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-12} nm ² MHz	
Q (Spin-Rot., T = 300K) = 111848.	J Max. = 20	Lines Listed = 2799	
Dipole Moments/Debye: $\mu_a = 3.850$ $\mu_b = 1.23$ $\mu_c = 0.$			
Rot. Const./MHz: A = 27663.66 B = 4714.144 C = 4235.041			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

J. Burie, J. Demaison, A. Dubrille, and D. Boucher, 1978, J. Mol. Spect. 72, 275.

D. R. Johnson, F. J. Lovas, C. A. Gottlieb, E. W. Gottlieb, M. M. Litvak, M. Guelin, and P. Thaddeus, 1977, Ap. J. 218, 370.

H. Mader, H. M. Heise, and H. Dreizler, 1973, Z. Naturforsch. 29a, 164.

The dipole moment was taken from H. M. Heise, H. Lutz, and H. Dreizler, 1974, Z. Naturforsch. 29a, 1345.

Species Tag: 56001	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide carbon - 13 isotope on atom 1 $\text{CH}_3\text{CH}_2^{13}\text{CN}$		Author: R. L. Poynter	
		Min. Int. = 2×10^{-12} nm ² MHz	
Q (Spin-Rot., T = 300K) = 112466.		J Max. = 10	Lines Listed = 1189
Dipole Moments/Debye: $\mu_a = 3.840$ $\mu_b = 1.370$ $\mu_c = 0$			
Rot. Const./MHz: A = 27634.94 B = 4689.805 C = 4214.746			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

H. M. Heise, H. Lutz, and H. Driezler, 1973, Z. Naturforsch. 29a, 1345.

The dipole moment was assumed the same as in the parent species, tag #55001.

Species Tag: 56002	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide Carbon - 13 isotope on atom 2 $\text{CH}_3^{13}\text{CH}_2\text{CN}$		Author: R. L. Poynter	
		Min. Int. = 2×10^{-12} cm^2MHz	
Q (Spin-Rot., T = 300K) = 113691.		J Max. = 10	Lines Listed = 1197
Dipole Moments/Debye: $\mu_a = 3.840$ $\mu_b = 1.370$ $\mu_c = 0.$			
Rot. Const./MHz: A = 27045.40 B = 4697.868 C = 4207.046			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

H. M. Heise, H. Lutz, and H. Driezler, 1973, Z. Naturforsch. 29a, 1345.

The dipole moment was assumed the same as in the parent species, tag #55001.

Species Tag: 56003	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide Carbon - 13 isotope on atom 3 $^{13}\text{CH}_3\text{CH}_2\text{CN}$		Author: R. L. Poynter	
		Min. Int. = 2×10^{-12} nm ² MHz	
Q (Spin-Rot., T = 300K) = 115302.		J Max. = 10	Lines Listed = 1189
Dipole Moments/Debye: $\mu_a = 3.840$ $\mu_b = 1.370$ $\mu_c =$			
Rot. Const./MHz: A = 27342.174 B = 4597.939 C = 4133.707			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

H. M. Heise, H. Lutz, and H. Driezler, 1973, Z. Naturforsch. 29a, 1345.

The dipole moment was assumed the same as in the parent species, tag #55001.

Species Tag: 56004	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide Nitrogen - 15 - isotope $\text{CH}_3\text{CH}_2\text{C}^{15}\text{N}$		Author: R. L. Poynter	
		Min. Int. = 2×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 38458.	J Max. = 20	Lines Listed = 1621	
Dipole Moments/Debye: $\mu_a = 3.840$ $\mu_b = 1.370$ $\mu_c = 0.$			
Rot. Const./MHz: A = 27541.953 B = 4574.771 C = 4119.4300			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

H. M. Heise, H. Mader, and H. Dreizler, 1976, Z. Naturforsch. 31a, 1228.

The dipole moment was assumed the same as in the parent species, tag #55001.

Species Tag: 56005	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide Deuterium isotope on methyl carbon atom trans to CN group CH ₂ D CH ₂ CN - S		Author: R. L. Poynter	
		Min. Int. = 2×10^{-12} nm ² MHz	
Q (Spin-Rot., T = 300K) = 118802.	J Max. = 10	Lines Listed = 1178	
Dipole Moments/Debye: $\mu_a = 3.840$ $\mu_b = 1.370$ $\mu_c = 0$			
Rot. Const./MHz: A = 27650.795 B = 4425.061 C = 4000.763			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

H. Mader, H. M. Heise, and H. Dreizler, 1973, Z. Naturforsch. 29a, 164.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 56006	Page 1 of 1	Version: 1	Date: Jan. 1980
Species Name: Ethyl Cyanide Deuterium Isotope on methyl carbon gauche to CN group $\text{CH}_2\text{DCH}_2\text{CN}-a$	Author: . L. Poynter		
	Min. Int. = 4×10^{-13} nm^2MHz		
Q (Spin-Rot., T - 300K) - 121064.	J Max. = 10	Lines Listed = 1214	
Dipole Moments/Debye: $\mu_a = 3.840$	$\mu_b = 1.370$	$\mu_c = 0$	
Rot. Const./MHz: A = 25022.568	B = 4583.422	C = 4110.245	

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

H. Mader, H. M. Heise, and H. Dreizler, 1973, Z. Naturforsch. 29a, 164.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 60001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbonyl Sulfide OCS		Author: R. L. Poynter	
		Min. Int. = 4×10^{-7} cm^2MHz	
Q (Spin-Rot., T = 300K) = 1028.17		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 0.715$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 6081.4924 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment has been remeasured by J. M. L. J. Reinartz and A. Dymanus, 1974, Chem. Phys. Lett. 24, 346.

Species Tag: 60002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monosulfide $^{28}\text{Si} \ ^{32}\text{S}$		Author: R. L. Poynter	
		Min. Int. = 9×10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 688.940		J Max. = 97	Lines Listed = 97
Dipole Moments/Debye: $\mu_a = 1.730$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 9077.45 C =			

The experimental measurements are summarized in E. Tiemann, 1976, J. Phys. Chem. Ref. Data, 5, 1147.

The dipole moment was measured by J. Hoelt, F. J. Lovas, E. Tiemann, and T. Törring, 1969, Z. Naturforsch. 24a, 1422.

Species Tag: 61001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbonyl Sulfide Carbon - 13 isotope $O^{13}CS$		Author: R. L. Poynter	
		Min. Int. = 4×10^{-7} cm^2/MHz	
Q (Spin-Rot., T = 300K) = 1031.492		J Max. = (99)	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = .715$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 6061.916 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 61002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monosulfide Silicon - 29 isotope $^{29}\text{Si} \ ^{32}\text{S}$		Author: R. L. Poynter	
		Min. Int. = 9×10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 701.830		J Max. = 98	Lines Listed = 98
Dipole Moments/Debye: $\mu_a = 1.730$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 8910.66 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 62001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbonyl Sulfide Sulfur - 34 isotope OC ³⁴ S		Author: R. L. Poynter	
		Min. Int. = 4 x 10 ⁻⁷ nm ² MHz	
Q (Spin-Rot., T = 300K) = 1053.926		J Max. = (99)	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 0.715$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 5932.8411 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 62002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Carbonyl Sulfide		Author: R. L. Poynter	
Oxygen - 18 isotope ^{18}OCS		Min. Int. = 4×10^{-17} nm ² MHz	
Q (Spin-Rot., T = 300K) = 1096.031	J Max. = (99)	Lines Listed = 99	
Dipole Moments/Debye: $\mu_a = 0.715$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 5704.8608 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 62003	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monosulfide Silicon - 30 isotope $^{30}_{\text{Si}} \ ^{32}_{\text{S}}$		Author: R. L. Poynter	
		Min. Int. = 8×10^{-6} $\text{cm}^2 \text{MHz}$	
Q (Spin-Rot., T = 300K) = 714.275		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 1.730$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 8755.33 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 62004	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Silicon Monosulfide Sulfur - 34 - isotope $^{28}_{\text{Si}} \ ^{34}_{\text{S}}$		Author: R. L. Poynter	
		Min. Int. = 9×10^{-6} nm ² MHz	
Q (Spin-Rot., T = 300K) = 708.329	J Max. = 99	Lines Listed = 99	
Dipole Moments/Debye: $\mu_a = 1.730$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 8828.86 C =			

The measurements were taken from the data set given in F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 64001	Page 1 of 1	Version: 1	Date: Oct. 1979
Species Name: Diatomic Sulfur $v = 0, {}^3\Sigma_g^-$ ground state		Author: H. M. Pickett	
		Min. Int. = 4.0×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 965.37		J Max. = 35	Lines Listed = 34
Dipole Moments/Debye: $\mu_a =$ $\mu_b =$ magnetic $\mu_c =$			
Rot. Const./MHz: A = B = 8831.2 C =			

The experimentally measured lines are given in H. M. Pickett and T. L. Boyd, 1979, J. Mol. Spectry. 75, 53. The calculational method used is from T. Amano and E. Hirota, 1974, J. Mol. Spectry. 53, 346. For S_2 , the parameter $\mu(1)$ is assumed to be zero. Only transitions between states with $J = N$ and $J = N - 1$ are listed because the energies of the $J = N + 1$ states are uncertain by more than 1 GHz. All the neglected transitions from $J = N + 1$ states lie above 450 GHz for $J < 35$.

The intensities of the magnetic dipole allowed transitions were calculated using the g values given in Pickett and Boyd. Hund's case (b) nomenclature is used even though the states are closer to Hund's case (a) for low J. The $J = N + 1$ states correlate with $\Sigma = 0$ and the $J = N - 1$ and $J = N$ states correlate with $\Sigma = \pm 1$ states of $p = \pm (-1)^J$ inversion parity, respectively.

The value of Q is determined from a sum over states to $J = 50$.

Species Tag: 64002	Page 1 of 1	Version: 1	Date: Jan, 1980
Species Name: Sulfur Dioxide SO_2		Author: R. L. Poynter	
		Min. Int. = 5×10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 5898.91	J Max. = 40	Lines Listed = 1902	
Dipole Moments/Debye: $\mu_a = 0.$ $\mu_b = 1.633$ $\mu_c = 0.$			
Rot. Const./MHz: A = 60778.558 B = 10317.913 C = 8799.652			

The experimental measurements were analyzed using the methods described in W. H. Kirchhoff, 1972, J. Mol. Spect. 41, 333. The measurements were taken from the following papers:

F. J. Lovas, 1978, J. Phys. Chem. Ref. Data, 7, 1445. (A complete summary and references to all available data).

The dipole moment was taken from D. Patel, D. Margolese, and T. R. Dyke, 1979, J. Chem. Phys. 70, 2740.

Species Tag: 75001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene		Author: R. L. Poynter	
HC ₅ N		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4695.48		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 4.330$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1331.33 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was measured by the same authors.

Species Tag: 76001	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene Carbon - 13 isotope on atom 1 HCCCC ¹³ CN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4742.326	J Max. = 99	Lines Listed = 99	
Dipole Moments/Debye: $\mu_a = 4.330$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1318.18 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 76002	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene		Author: R. L. Poynter	
Carbon - 13 isotope on atom 2 HCCC ¹³ CCN		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4699.78		J Max. = 99	Lines Listed = 99
Dipole Moments/ Debye: $\mu_a = 4.329$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1330.11 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 76003	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene Carbon - 13 isotope on atom 3 HCC ¹³ CCCN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4700.345		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 4.329$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1329.95 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 76004	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene Carbon - 13 isotope on atom 4 CH ¹³ CCCCN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4744.09		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 4.329$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1317.689 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 76005	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene Carbon - 13 isotope on atom 5 H ¹³ CCCCN		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4820.96		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: μ_a 4.329 μ_b μ_c			
Rot. Const./MHz: A = B = 1296.68 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 76006	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene Nitrogen - 15 isotope HC_5^{15}N		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} cm ² /MHz	
Q (Spin-Rot., T = 300K) = 4813.68		J Max. = 99	Lines Listed = 99
Dipole Moments/Debye: $\mu_a = 4.329$ μ_b $\mu_c =$			
Rot. Const./MHz: A = B = 1298.639 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was measured by the same authors.

Species Tag: 76007	Page 1 of 1	Version: 1	Date: Dec. 1979
Species Name: Cyanodiacetylene Deuterium isotope DC ₅ N		Author: R. L. Poynter	
		Min. Int. = 2×10^{-7} nm ² MHz	
Q (Spin-Rot., T = 300K) = 4918.13	J Max. = 99	Lines Listed = 99	
Dipole Moments/Debye: $\mu_a = 4.329$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 1271.056 C =			

The experimental measurements were taken from A. J. Alexander, H. W. Kroto, and D. R. M. Walton, 1976, J. Mol. Spect. 62, 175.

The dipole moment was assumed to be the same as the parent species.

Species Tag: 80001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Hydrogen Bromide Bromine - 79 isotope H ⁷⁹ Br		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 7×10^{-3} nm ² MHz	
Q (Spin-Rot., T = 300K) = 101.214		J Max. =	Lines Listed = 19
Dipole Moments/Debye: $\mu_a = 0.828$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 250357.6 C =			

The experimental data were taken from F. A. Van Dijk and A. Dymanus, 1969, Chem. Phys. Lett. 4, 170.

The dipole moment was also reported by these same authors in a later publication, and updated in F. A. Van Dijk and A. Dymanus, 1971, 26th Symposium on Molecular Structure and Spectroscopy, Columbus, Ohio, R12, p. 80.

Species Tag: 82001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Hydrogen Bromide Bromine - 81 isotope H^{81}Br		Author: R. L. Poynter & H. M. Pickett	
		Min. Int. = 7×10^{-3} nm ² MHz	
Q (Spin-Rot., T = 300K) = 101.257	J Max. =	Lines Listed = 19	
Dipole Moments/Debye: $\mu_a = 0.828$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 250250.2 C =			

The experimental data were taken from F. A. Van Dijk and A. Dymanus, 1969, Chem. Phys. Lett. 4, 170.

The dipole moment was also reported by these same authors in a later publication, and updated in F. A. Van Dijk and A. Dymanus, 1971, 26th Symposium on Molecular Structure and Spectroscopy, Columbus, Ohio, R12, p. 80.

Species Tag: 95001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Bromine Monoxide Bromine - 79 isotope $2\pi_{3/2}$ state 79 BrO		Author: H. M. Pickett	
Q (Spin-Rot., T = 300K) = 3897.60		J Max. = 49	Lines Listed = 850
Dipole Moments/Debye: $\mu_a = 1.765$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 12830.4 C =			

The 63 GHz spectrum of BrO is referenced in F. J. Lovas and E. Tiemann, 1974, J. Phys. Chem. Ref. Data 3, 653. The other lines were measured at JPL by E. A. Cohen and H. M. Pickett, and a new fit was made which included these lines.

Species Tag: 97001	Page 1 of 1	Version: 1	Date: Feb. 1980
Species Name: Bromine Monoxide Bromine - 81 isotope $^2\Pi_{3/2}$ state 81 BrO		Author: H. M. Pickett	
		Min. Int. = 10^{-9} nm ² MHz	
Q (Spin-Rot., T = 300K) = 3913.9	J Max. = 49	Lines Listed = 850	
Dipole Moments/Debye: $\mu_a = 1.794$ $\mu_b =$ $\mu_c =$			
Rot. Const./MHz: A = B = 12777.08 C =			

The 63 GHz spectrum of BrO is referenced in F. J. Lovas and E. Tiemann, 1974, J. Phys. Chem. Ref. Data 3, 653. The other lines were measured at JPL by E. A. Cohen and H. M. Pickett, and a new fit was made which included these lines.