

## N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM  
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT  
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED  
IN THE INTEREST OF MAKING AVAILABLE AS MUCH  
INFORMATION AS POSSIBLE

THE ROTATION OF URANUS

Richard M. Goody


Harvard University, Cambridge, MA 02138

(NASA-CR-163987) THE ROTATION OF URANUS  
(Harvard Univ.) 11 p HC A02/MF A01 CSCL 03C

N81-19987

Unclas

G3/91 41732



HISTORICAL REVIEW

From the start of this century until the mid-1970's the rotation rate of Uranus was reported to be 10.8 h in a retrograde sense (see, for example, Allen, 1955), but a cursory examination of the origin of this datum reveals that little confidence should be placed in it.

Three independent techniques for measuring the rotation rate are available, each very difficult and not including the most direct method of observing the motion of features across the disc. Visual observers have reported features from time to time (see Alexander, 1965, for a full historical account), but the mean diameter of Uranus is only 3.6 arcsec and large high contrast features are rare in the visible spectrum, if they exist at all.

The three available methods are: use of theoretical interior models together with observations of the oblateness,  $f$ , and the gravitational moment,  $J_2$ ; periodic fluctuations in the brightness; spectrographic measurements of Doppler shifts (line tilts). The first report of a rotation rate close to 11 h, based on a theoretical analysis of the planet's figure, was by Berstrand (1909; some of the early work is not easy to find and where it is only of historical importance I have relied upon Alexander's reports). At that time, there were no measurements of  $J_2$  and no reliable data on the oblateness. The crucial early work was the spectrographic determination of  $10.8 \pm 0.3$  h by Lowell and Slipher (1912). While these observers worked with exemplary care, a reanalysis of their data by Hayes and Belton (1977) shows no significant line tilts if all the data are taken together.

PIERCE HALL  
HARVARD UNIVERSITY  
29 OXFORD STREET Goody  
CAMBRIDGE, MASSACHUSETTS 02138

### THE FIGURE OF URANUS

For an equilibrium rotating body the gravitational moment,  $J_2$ , and the oblateness,  $f$ , are, to first order, related by (Cook, 1973):

$$J_2 = 2f/3 - m/3,$$

$$f = 5m/2[1 + (5/2 - 15\beta/4)^2],$$


$$m = 4\pi^2 r^3 / T^2 GM,$$

$$\beta = C/Mr^2,$$

where  $T$  is the rotational period,  $r$  the radius of the planet,  $M$  the mass,  $C$  the principal moment of inertia and  $G$  the universal gravitational constant.

These equations provide two relationships between four quantities. Given any two, therefore, the others can be calculated.  $J_2$  can be determined from the precession of satellite orbits, but the precession is very small and the measurement was extremely difficult prior to the discovery of the rings. Oblateness must be measured directly. It is about 2%, and its magnitude was in doubt until recent years.  $\beta$  can be calculated given a model of the interior of the planet, and it lies between 0.15 and 0.30 for plausible models


Figure 1. The appearance of Uranus and its satellites on 9 May 1981 (from the *Astronomical Almanac*, p. F62).



(Brown and Goody, 1980; Podolak, 1976; Podolak and Cameron, 1974). Houzeau, in 1856, used speculations about the interior structure of Uranus to obtain  $\beta$  and hence the first estimate of the period, between 7.25 h and 12.5 h.

Despite a great deal of effort over the next 100 years, Houzeau's figure was not easily bettered. Figure 2 indicates the state of affairs in 1976. The two best measurements of  $J_2$  from satellite orbit analysis are shown. The oblateness determination of Dollfuss (1970) is a summary of all ground-based work to date. That of Danielson, *et al.* (1972) is based upon measurements made by Stratoscope II. The broken lines are lines of constant  $\beta$  from a range of interior models of Podolak (1976). On the basis of these data Podolak favored a period  $\sim 18$  h.

Figure 2. The relationship between  $J_2$ ,  $f$ ,  $T$ , and  $\beta$  together with measurements available in 1976. The broken lines are of constant  $\beta$  for a range of interior models. The equations indicate that the lines should be curved but the curvature has been neglected (after Podolak, 1976).



Occultation measurements on the rings of Uranus now allow very precise determinations of  $J_2$ . Nicholson, *et al.* (1978) find  $J_2 = 3.43 \pm 0.02 \times 10^{-3}$ , while more extensive work by Elliot, *et al.* (1980) gives  $J_2 = 3.354 \pm 0.005 \times 10^{-3}$ . Simultaneously, reliable data for the oblateness have become available. Franklin, *et al.* (1980) reanalyzed the Stratoscope II data and found  $f = 0.022 \pm 0.001$ . This result depends upon an understanding of the difference between polar and equatorial limb darkening, but the same criticism cannot be made of the work of Elliot, *et al.* (1980), which is based upon stellar occultations. Elliot, *et al.* find  $f = 0.024 \pm 0.003$ .

If we adopt Franklin, *et al.*'s value of  $f$  and Elliot, *et al.*'s value of  $J_2$  we have

$$T = 16.7 \pm 0.5 \text{ h} ,$$

$$\beta = 0.216 \pm 0.004 .$$

#### BRIGHTNESS VARIATIONS

The foregoing work on the figure of Uranus seems conclusive in its indications, but it is an indirect method and confirmation from a more direct approach is desirable. Brightness fluctuations offer one possibility. A periodic change in the brightness of a planet or satellite with a period on the order of tens of hours is most likely to be associated with the appearance and disappearance of bright to dark features at the limb. Our knowledge of the rotational period of Neptune is mainly based upon such measurements although for Neptune the variations are very large, up to 2 mag in the J-K color index, according to Belton, Wallace and Howard (1981).

The history of brightness measurements on Uranus is summarized in Table 1. Most of the references can be found in Alexander (1965); the editorial comments are my own. The table speaks for itself and shows that no convincing case for brightness variations of Uranus has been established in the published literature.

The most accurate data are those of Lockwood and Thompson (1978), working at Lowell Observatory. They made measurements with filters at and adjacent to methane bands at 6190 and 7261 Å, because other measurements suggest that variability is at a maximum in strong methane bands. It is generally believed that clouds may

occasionally form above the level of methane absorption and give rise to local brightening. Lockwood and Thompson conclude that there are no periodic variations in excess of  $0.003 \text{ mag}$  except perhaps in the range of 23 to 25 h where it is hard to reach a conclusion because of the difficulty of working with irregularly spaced data taken at 24-hour intervals.

Despite this difficulty, a period close to 24 h is, in fact, claimed by a group working at the University of Texas. This work is reported in the abstract only (Smith and Slavsky, 1979): the methods are the same as those used by Slavsky and Smith (1978) for Neptune. I am indebted to David Slavsky for the following details.

Observations were made with filters in and adjacent to the  $6190 \text{ \AA}$  methane band in Texas, Chile and South Africa. Periodic variations of the brightness of Uranus of  $0.006 \text{ mag}$  were recorded in contrast to  $0.002 \text{ mag}$  for a comparison star. The phase for simultaneous measurements in Chile and South Africa agreed in sidereal time but not in local time, and the entire data set is consistent with a period of 23.87 h.

Table 1. The period of Uranus from brightness variations.

<u>Date</u>	<u>Observer(s)</u>	<u>Period</u>	<u>Comments</u>
1884-85	Muller	---	No variations
1915	Waterfield	21 d	Probably insignificant
1916-17	Campbell	10.82 h	0.15 mag variations
1918	Campbell	---	No variations
1921-26	Wirtz	---	No variations
1926	Perenago	10.82 h	Poor statistics
1926	Perenago	---	No variations
1927	Slavenas	10.82 h	Poor statistics
1928	Stebbins/Jacobsen	---	No variations
1928	Gussow	---	No variations
1934-35	Sterne/Calder	10.82 h	Marginal statistics
1976	Belton	21.48 h	Reanalysis of Campbell's data
1977	Lockwood/Thompson	---	Less than $0.003 \text{ mag}$ variations except for 23-25 h period
1980	Smith/Slavsky	23.87 h	Unpublished; $0.006 \text{ mag}$ variations

### SPECTROGRAPHIC METHODS

These methods are also more direct than those based upon the planetary figure. In their simplest form the spectrograph slit is placed along the equator and a measurement made of the tilt of a spectral line.

There are many pitfalls. Line tilts are typically a few degrees. For the observations of Moore and Menzel the diameter of Uranus at the photographic plate was  $\sim 0.2$  mm and the exposure time was 1-2 hours. The lateral distance between the two ends of the line averaged  $5 \mu\text{m}$  or only 0.5 grains, depending upon the characteristics of the film used.

Modern instruments give great improvements. In 1976 and 1977, Robert Brown and I (Brown and Goody, 1977; 1980) worked with the KPNO 4-meter telescope with a Cassegrain echelle to obtain both high spectral dispersion and a large image. The detector was a Kron camera having extremely fine grain combined with linearity over a wide dynamic range. Most importantly, we took steps to allow for the effect of seeing on the recorded line tilts.

The correction function is shown in Fig. 3. The recorded line tilt decreases as the seeing disc increases in size. For  $s$  arcsec the recorded line tilt must be increased in the ratio  $G(0)/G(s)$  before interpretation as a Doppler shift. For a seeing disc of 2 arcsec this ratio is about 1.6. For the long exposures of Moore and Menzel the correction factor (which they did not apply) must be at least this great, although this only serves to increase the discrepancy between their value and all modern determinations.

Seeing can be measured from scans across the spectrum in continuum regions, but use of this result requires that seeing and guiding errors be isotropic. To eliminate the typical bias between errors in RA and Dec we employed automatic guidance on all occasions.

In 1976 we measured the tilts of 23 lines and obtained a rotational period of  $15.57 \pm 0.80$  h, while in 1977 we measured more than 600 lines for a period of  $16.26 \pm 0.34$  h. Both data sets are consistent with

$$T = 16.16 \pm 0.33 \text{ h,}$$


and using measured values of  $J_2$  we find

$$\beta = 0.212 \pm 0.004 .$$

We were also able to confirm that the rotation axis corresponded with the pole of the satellite orbits.

The work was only one of a number of simultaneous attempts to improve spectrographic determinations of Uranus' period. Trauger, Roessler and Münch (1978) used a whole-disc approach which avoids seeing corrections. It requires a knowledge of incoherent scattering processes, however, about which we cannot be confident. Trafton (1977) employed an ingenious variant of the normal spectrographic technique which is, however, subject to large errors. I shall therefore restrict my discussion to the work of Münch and Hippelein (1980) and Hayes and Belton (1977).

Figure 3. Seeing correction to line tilt based on measurements of synthetic spectra. For perfect seeing  $G(0)$  is  $7.65 \times 10^{-2}$  (after Brown and Goody, 1977).





The former obtain a period of  $15.0^{+4.0}_{-2.6}$  h, consistent with our work, and they provided a quasi-analytical basis for the seeing correction. Hayes and Belton obtained  $24 \pm 3$  h, consistent with the work of Smith and Slavsky, but irreconcilable with our determination.

Hayes and Belton used essentially the same equipment as we did and employed a similar method of analysis. It is difficult to avoid the conclusion that one piece of work or the other contains numerical errors. With elaborate numerical algorithms such errors are regrettably easy to make and hard to detect. In a subsequent paper, Belton, Wallace, Hayes and Price (1980) mention "two serious sources of error" in the earlier work but state that their results for Uranus were unaffected.

#### THE ROTATION OF URANUS

The weighted means of our data and that obtained from the planetary figure are

$$T = 10.31 \pm 0.27 \text{ h,}$$

$$\beta = 0.214 \pm 0.003 .$$

The values of  $J_2$  and  $f$  are now firmly established. For the rotational period to be wrong the equilibrium theory of the figure must be inapplicable. It works very well for all other planets for which data exist.

The work of Slavsky and Smith cannot be evaluated until it is published and the only established discrepancy with the above result is, therefore, the work of Hayes and Belton. Unfortunately, it will be a long time before the spectrographic work can be repeated successfully, even assuming that investigators will exist with the desire to do so. The aspect of the planet is becoming increasingly unfavorable. In 1985 the North Pole will point directly toward the Sun and seven or eight more years must elapse before there is a substantial component of the rotation vector orthogonal to the line of sight.

The possibility of detecting the motion of features across the disc remains. Nisenson, *et al.* (1981) have reported on the use of speckle imaging techniques to obtain images of Titan with a resolu-

tion of  $0.12 \text{ arcsec}$ . With this equipment Uranus can be imaged in the  $6190 \text{ \AA}$  methane band and with more suitable image intensifiers, also in the  $7261 \text{ \AA}$  band. If large cloud systems appear above the level of methane absorption it may be possible to detect them.

Finally, the Voyager II flyby in January 1985 may show features on the disc, in which case the controversy over the Uranus rotation period should be finally resolved.

With each value of the period goes a value for  $\beta$  and for the principal moment of inertia of the planet. This datum is crucial for models of the interior structure of Uranus, as discussed at this colloquium by Dr. Hubbard. In terms of the models of Podolak (1976) and Podolak and Cameron (1974) the value  $\beta = 0.214$  is compatible only with envelopes containing the largest anticipated concentrations of volatiles.

#### ACKNOWLEDGEMENTS

I am indebted to Robert Brown for past collaborations. My work is supported by the National Aeronautics and Space Administration under Grant NGL-22-007-228.

#### REFERENCES

- Alexander, A. F. O'D (1965). *The Planet Uranus: A History of Observation, Theory and Discovery*. Faber and Faber, London.
- Allen, C. M. (1955). *Astrophysical Quantities*. Athlone Press, London, p. 157.
- Belton, M. J. S., Wallace, L. and Howard, S. (1981). The periods of Neptune: Evidence for atmospheric motions. Submitted to *Icarus*.
- Belton, M. J. S., Wallace, L., Hayes, S. H. and Price, M. J. (1980). Neptune's rotation period: A correction and a speculation on the difference between photometric and spectrographic results. *Icarus* 42, 71-88.
- Brown, R. A. and Goody, R. M. (1977). The rotation of Uranus. *Ap. J.* 217, 680-687.
- Brown, R. A. and Goody, R. M. (1980). The rotation of Uranus II. *Ap. J.* 235, 1066-1070.
- Campbell, L. (1936). The rotation of Uranus. *Harvard C. O. Bull.* 904, 32-35.
- Cook, A. H. (1973). *Physics of the Earth and Planets*. Wiley, New York, pp. 35, 132.
- Danielson, R. E., Tomasko, M. G. and Savage, B. D. (1972). High resolution imagery of Uranus obtained by Stratoscope II. *Ap. J.* 178, 887-900.

- Dollfus, A. (1970). Diamètres des planètes et satellites. In *Surfaces and Interiors of Planets and Satellites* (A. Dollfus, Ed.). Academic Press, New York, pp. 46-139.
- Elliot, J. L., French, R. G., Foregl, J. A., Elias, J. H., Mink, D. and Liller, W. (1981). Orbits of nine Uranian rings. Submitted to *Ap. J.*
- Franklin, F. A., Avis, C. C., Colombo, G. and Shapiro, I. I. (1980). The geometric oblateness of Uranus. *Ap. J.* 236, 1031-1034.
- Greenberg, R. (1975). The dynamics of Uranus' satellites. *Icarus* 24, 325-332.
- Hayes, S. H. and Belton, M. J. S. (1977). The rotational periods of Uranus and Neptune. *Icarus* 32, 383-401.
- Lockwood, G. W. and Thompson, D. T. (1978). A photometric test of rotational periods for Uranus and time variations of methane-band strengths. *Ap. J.* 221, 689-693.
- Lowell, P. and Slipher, V. M. (1912). Spectroscopic discovery of the rotation period of Uranus. *Lowell Obs. Bull.* 2, No. 3.
- Moore, J. H. and Menzel, D. H. (1930). The rotation of Uranus. *Pub. Astron. Soc. Pac.* 42, 330-335.
- Münch, G. and Hippelein, H. (1980). The effects of seeing on the reflected spectrum of Uranus and Neptune. *Astron. Astrophys.* 81, 189-197.
- Nicholson, P. D., Pensson, S. E., Matthews, K., Goldreich, P. and Neugebauer, G. (1978). The rings of Uranus: Results of the 1978 April 10 occultation. *Astron. J.* 83, 1240-1248.
- Nisenson, P., Apt. J., Goody, R. and Horowitz, P. (1981). Radius and limb darkening of Titan from speckle imaging. Submitted to *Astron. J.*
- Podolak, M. (1976). Methane rich models of Uranus. *Icarus* 27, 473-476.
- Podolak, M. and Cameron, A. G. W. (1974). Models of the giant planets. *Icarus* 22, 123-148.
- Slivsky, D. and Smith, H. J. (1978). The rotation period of Neptune. *Ap. J.* 226, L49-52.
- Smith, H. J. and Slavsky, D. B. (1979). Rotation period of Uranus. *B.A.A.S.* 11, 568.
- Trafton, L. (1977). Uranus' rotational period. *Icarus* 32, 402-412.
- Trauger, J. T., Roesler, F. L. and Münch, G. (1978). A redetermination of the Uranus rotation period. *Ap. J.* 219, 1079-1083.