

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NGR-05-020-559

STRUCTURE OF THE SOLAR OSCILLATION WITH PERIOD NEAR 160 MINUTES

(NASA-CR-168911) STRUCTURE OF THE SOLAR
OSCILLATION WITH PERIOD NEAR 160 MINUTES
(Stanford Univ.) 17 p HC A02/MF A01

N82-25074

CSCI 03B

Unclass

G3/92 20282

Philip H. Scherrer and John M. Wilcox
SUIPR Report No. 865

February 1982



INSTITUTE FOR PLASMA RESEARCH
STANFORD UNIVERSITY, STANFORD, CALIFORNIA

**Structure of the Solar Oscillation With Period
Near 160 Minutes**

Philip H. Scherrer and John M. Wilcox

**Institute for Plasma Research
Stanford University
Stanford, California 94305**

**Stanford University, Institute for Plasma Research
SUIPR Report # 865
February 1982**

Structure of the Solar Oscillation With Period
Near 160 Minutes

Philip H. Scherrer and John M. Wilcox

Institute for Plasma Research
Stanford University
Stanford, California 94305

ABSTRACT

The solar oscillation with period near 160 minutes is found to be unique in a spectrum computed over the range of periods from about 71 to 278 minutes. Our best estimate of the period is 160.0095 ± 0.001 minutes, which is different from 160 minutes ($1/9$ of a day) by a highly significant amount. The width of the peak is approximately equal to the limiting resolution that can be obtained from an observation lasting 6 years, which suggests that the damping time of the oscillations is considerably longer than 6 years. A suggestion that this peak might be the result of a beating phenomenon between the five minute data averages and a solar oscillation with period near five minutes is shown to be incorrect by recomputing a portion of the spectrum using 15 second data averages.

Oscillations of the sun with a period near 160 minutes were discovered by Severny et al.(1976) using a technique in which the Doppler shift of the central portion of the solar disk was compared with that of an outer annulus, using a Babcock solar magnetograph specially modified for this observation. This discovery was further described by Kotov et al.(1978). Similar observations using resonant scattering to measure the mean velocity of the sun as a whole were reported by Brookes et al. (1976).

These reports were met with some skepticism for three reasons: (1) The observed amplitude of less than 1 m/s is very small and near the limits of observing capability, (2) a period of 160 minutes is exactly one ninth of a day, and could therefore appear in a power spectrum as a harmonic of the power at 24 hours that is present as an artifact in data obtained from observatories at mid-latitudes, i.e. when the observations must be interrupted every night, and (3) the source of the oscillation is not understood. It could be a g-mode, but if so one would expect to observe a number of adjacent g-modes, just as there are many p-modes with periods near 5 minutes.

Observations at the Stanford Solar Observatory (Scherrer et al., 1979, 1980) have helped to obviate the

first two problems. The third has not yet been solved.

This report is of a more detailed analysis of the observations. We start the analysis by combining the observations obtained at the Stanford Solar Observatory (for the observational arrangement see Scherrer et al. 1982) during the summers of 1977 through 1980 with those obtained at the Crimean Astrophysical Observatory by V.A. Kotov, A.B. Severny and T.T. Tsap during the summers 1974 through 1979. For this computation the Stanford observations were in the form of 5 minute averages and the Crimean observations were 10 minute averages. The observations on each day were fit with a parabola to remove a daily drift that at Stanford is of the order of one m/s per hour. The residuals were then normalized to have a standard deviation equal to one, so as to give equal weight to each day's observations from each observatory. While this procedure has the unpleasant effect of adding noise to the window function defined by the times of observation, it does allow us to combine data from both observatories in one analysis. The Stanford and Crimean observations were combined as a single data set to compute harmonic amplitude spectra using a simple least-squares method to find the Fourier coefficients. For an observation duration of 6 years, the spectral resolution is about 5 nanoHz so we can compute the spectrum in 2 nanoHz steps to

identify all peaks corresponding to oscillations with lifetimes of more than several years. To examine a large range of possible long period oscillations, we computed the spectrum from 60 to 235 microHz. This corresponds to periods of about 71 to 278 minutes.

Figure 1 shows a small part of the resulting spectrum in the range from 159.9 to 160.1 minutes (104.10 to 104.22 microHz.) The peak near 160 minutes is near the center of this figure. In order to more clearly define this peak, part of the spectrum was recomputed with a resolution of 0.1 nanoHz. This high-resolution spectrum is shown in figure 2. The centroid of the central peak in Figures 1 and 2 is at 104.1605 microHz (160.0095 minutes). Thus 160.0095 ± 0.001 minutes is our best estimate of the period of this solar oscillation, which is different from 160.000 minutes by a highly significant amount. This analysis is consistent with the agreement in the phase of the oscillation at the two observatories reported by Scherrer et al. (1979, 1980) and the "impressively good" agreement observed at the South Pole by Fossat et al. (1981; see also Grec et al., 1980). The first two objections mentioned above are thereby apparently resolved.

The central peak shown in Figures 1 and 2 is surrounded

on each side by a peak displaced by 32 nanoHz, which is the splitting associated with an interval of one year. This interval appears in the spectrum because the observations are concentrated during the summer months.

The full width at half maximum of the central peak in Figure 2 is 5.3 nanoHz, which is approximately the limiting resolution to be obtained from an observation lasting 6 years. This suggests that the damping time of the oscillation is considerably longer than 6 years.

Figure 3 shows the full spectrum in the range from 71 to 278 minutes (60 to 235 microHz), computed in steps of 2 nanoHz. The central peak shown in Figures 1 and 2 now appears in Figure 3 as an isolated line whose amplitude is several times larger than that of any other peak within this frequency range (with the exception of the two yearly satellite peaks explained above). The singular nature of the oscillation at 160.0095 minutes is apparent in Figure 3. To our knowledge an accepted theoretical explanation of this situation does not yet exist. For a concise review of the theoretical situation see de Jager (1981).

Childress and Spiegel (1981) have proposed that this oscillation may be described as a strange attractor. They suggest that such a signature would include long "periods"

(compared to that of the radial fundamental mode), erratic behavior, and intermittency. The physical mechanism may involve instabilities driven in a thin layer.

The amplitude of the oscillation can not be examined in the spectrum described above since the data was normalized. Figure 4 shows the spectrum computed in the same range as Figures 1 and 2 using only five minute averages of the Stanford data. For this computation the individual day's observations were not normalized. This spectrum is similar to the spectra computed from the combined data of Stanford and the Crimea shown in Figures 1 and 2. The amplitude of the central peak in Figure 4 corresponds to a velocity of 17 cm/sec, which is consistent with earlier analyses and with the South Pole observations (Grec et al., 1980, Fossat et al., 1981).

Philippe Delache (1981) has suggested that the solar oscillation with a period near 160 minutes might be understood as a beating between the five minute intervals in which the data were averaged for the above computations and one of the solar oscillation modes with period near five minutes discovered by Claverie et al. (1979). Although some objections to this suggestion could be proposed, the basic test is to compute the spectrum using other than five minute

data averages. The spectrum in Figure 5 was computed using the same data as in Figure 4, but using 15 second data averages. These two spectra are essentially identical. It therefore appears that the suggestion of Delache can be excluded.

Acknowledgements

We thank A.B. Severny, V.A. Kotov and T.T. Tsap for sharing their data with us, and for many interesting discussions at the Crimean Astrophysical Observatory and at the Stanford Solar Observatory.

This work was supported in part by the Office of Naval Research under Contract N00014-76-C-0207, by the National Aeronautics and Space Administration under Grant NGR05-020-559 and Contract NAS5-24420, by the Atmospheric Sciences Section of the National Science Foundation under Grant ATM77-20580 and by the Max C. Fleischmann Foundation.

References


- Brookes, J.R., Isaak, G.R., and Van der Raay, H.B.: 1976,
Nature 259, 92.
- Claverie, A., Isaak, G.R., McLeod, C.P., van der Raay H.B. and
Roca Cortes, T.: 1979, Nature, 282, 591.
- Childress, S., and Spiegel, E.A.: 1981, "A Prospectus for
a Theory of Variable Variability", preprint #A14 of Department
of Astronomy and Astrophysics, Columbia University.
- de Jager, C: 1981, Solar Physics, 74, 11.
- Delache, P.: 1981, "Sur l' observation des oscillations du Soleil
de periodes voisines de 160 mn.", preprint, Departement
D'Astrophysique, Universite De Nice.
- Fossat, E., Grec, G., and Pomerantz, M.: 1981, Solar Physics ,74,
59.
- Grec, G., Fossat, E. and Pomerantz, M.: 1980, Nature, 288, 541.
- Scherrer, P., Wilcox, J.M., Kotov, V.A., Severny A.B. and
Tsap, T.T.: 1979, Nature, 277, 635.
- Scherrer, P., J.M. Wilcox, A.B. Severny, V.A. Kotov and
T.T. Tsap: 1980, Astrophys. J., 237, L97.
- Scherrer, P., Wilcox, J.M., Christensen-Dalsgaard, J. and
Gough, D.O.: 1982, Solar Physics, (these proceedings).


Figure Captions

- Figure 1 Harmonic amplitude spectrum of the combined solar velocity observations from Stanford and the Crimea in the range from about 159.9 to 160.1 minutes. The vertical scale is arbitrary because the daily residuals at each observatory were normalized to have a standard deviation equal to one. The central peak has a period of 160.0095 minutes and is surrounded on each side by a peak displaced by 32 nanoHz, which is the splitting associated with an interval of one year (see text). This spectrum was computed with a resolution of 2 nanoHz.
- Figure 2 Same spectrum as Figure 1, but computed with a resolution of 0.1 nanoHz. The vertical dashed line is drawn at a period of 160 minutes, and it is clear that the central peak is significantly removed from this period.
- Figure 3 Spectrum similar to that computed in Figure 1 but over an extended range from about 71 to 278 minutes (60 to 235 microHz), computed in steps of 2 nanoHz. The central peak shown in Figure 1 at a period of 160.0095 minutes is now seen to be unique within the period range shown in Figure 3.
- Figure 4 Spectrum computed in the range of periods from about 159.9 to 160.1 minutes using only Stanford observations that were

not normalized on each day. The maximum of the central peak corresponds to a velocity of 17 cm/sec. This spectrum was computed using 5 minute averages.


Figure 5 The same as Figure 4, except that in response to a suggestion by Delache (1981) the spectrum was computed using 15 second averages. The central peaks in Figures 4 and 5 are very similar.






ORIGINAL PAGE IS
OF POOR QUALITY


(VELOCITY)²
(arbitrary)




(VELOCITY)²
(arbitrary)




(VELOCITY)²
(arbitrary)



(VELOCITY)²
(arbitrary)



(VELOCITY)²
(arbitrary)



PERIOD
(minutes)

