

nasa CR-166,113

**NASA Contractor Report 166113**

# ICASE

NASA-CR-166113  
19830017160

A COMPARATIVE STUDY OF FINITE ELEMENT AND  
FINITE DIFFERENCE METHODS FOR CAUCHY-RIEMANN  
TYPE EQUATIONS

George J. Fix  
and  
Milton E. Rose

Contract No. NAS1-17070 and NAS1-17130  
March 1983

INSTITUTE FOR COMPUTER APPLICATIONS IN SCIENCE AND ENGINEERING  
NASA Langley Research Center, Hampton, Virginia 23665

Operated by the Universitite Space Research Association

**NASA**  
National Aeronautics and  
Space Administration  
**Langley Research Center**  
Hampton, Virginia 23665

**LIBRARY COPY**

MAY 26 1983

LANGLEY RESEARCH CENTER  
LIBRARY, NASA  
HAMPTON, VIRGINIA



NF02447

A COMPARATIVE STUDY OF FINITE ELEMENT AND FINITE DIFFERENCE METHODS  
FOR CAUCHY-RIEMANN TYPE EQUATIONS

George J. Fix  
Carnegie-Mellon University

Milton E. Rose  
Institute for Computer Applications in Science and Engineering

ABSTRACT

A least squares formulation of the system  $\text{div} \underline{u} = \rho$ ,  $\text{curl} \underline{u} = \underline{\zeta}$  is surveyed from the viewpoint of both finite element and finite difference methods. Closely related arguments are shown to establish convergence estimates.

---

The research reported in this paper was supported by the National Aeronautics and Space Administration under NASA Contract No. NAS1-17070 for the first author and by NASA Contract Nos. NAS1-17130 and NAS1-17070 for the second author while they were in residence at ICASE, NASA Langley Research Center, Hampton, VA 23665.

## Introduction

This paper concerns the three dimensional Cauchy-Riemann type equations

$$\operatorname{div} \underline{u} = \rho, \operatorname{curl} \underline{u} = \underline{\zeta} \quad \text{in } D$$

(1)

$$\underline{u} \cdot \underline{n} = 0 \quad \text{on } \Gamma;$$

$D$  is a bounded domain in  $\mathbb{R}^3$  with boundary  $\Gamma$  on which  $\underline{n}$  is the outward normal. The functions  $\rho, \underline{\zeta}$  are prescribed and satisfy the compatibility conditions

$$\int_D \rho d\pi = 0, \operatorname{div} \underline{\zeta} = 0 \quad \text{in } D;$$

(2)

these express necessary conditions that the overdetermined first-order system (1) has a solution.

The numerical solution of these equations will be studied from both the finite element and finite difference points of view. Indeed, the major goal of this paper is to show how both approaches rest on very similar foundations. In so doing we hope our study may provide a point of contact between those familiar with the, largely separate, literature about each method.

In the case of the finite element method convergence estimates will be shown to result quite directly from proof techniques already common to the finite element literature. In contrast, many of these same techniques have played little or no role in the analysis of finite difference schemes and one of our principal objectives lies in clarifying their relevance to finite difference methods.

The Cauchy-Riemann type equations (1) were chosen for study because they are representative of a type of first-order systems that arise in problems from electromagnetics and fluid dynamics. For such systems, arbitrary finite difference and finite element approximations to (1) are generally unsuitable. This is certainly true of all but a few finite element schemes based on Galerkin formulations [1], while simple finite difference approximations can present, among other difficulties, special problems in incorporating boundary conditions accurately.

In Part A notations used in this paper are introduced and the basic intergral identity

$$(3) \quad \int_D [ |\text{curl}\underline{u}|^2 + |\text{div}\underline{u}|^2 ] = \int_D |\text{grad}\underline{u}|^2$$

is derived. This identity has found use in many applications (see, for example, [3] where it is used in a discussion of the Navier-Stokes equations). The derivation given in Section A.1 differs from (3) in appearance in order to highlight the structural properties of (1) when viewed as a first-order system.

The fact that (1) is well posed is an immediate consequence of (3) and the Lax-Milgram Theorem. This is described in Section A.2 both for completeness and because it shows the fundamental role the least squares ideas can play in discussing overdetermined systems like (1).

The derivation of a stable and optimally convergent finite element scheme is almost immediate from the material developed in Part A. The arguments, while standard, are included in Part B for completeness. It is interesting to note that, unlike other least squares approximations to first-order systems, these do not impose any restrictions on the finite element spaces [2].

In Part C an analogous development is given for the Keller box-scheme based, instead, on a least squares summation formulation. Here a summation by parts formula is used to obtain a summation identity analogous to (3) and results in a proof that the scheme is second order accurate. Of special interest is the fact that this development, also, is not restricted to uniform grids nor to grids obtained by images of uniform grids under a global mapping function. Thus, like the finite element schemes, it can also be used on irregular grids subject only to standard geometric constraints. Finally, we remark that the difference scheme employed here (c. f. [10] [12]) involves the variables on the same, rather than on staggered, grids in contrast to [5], [9].

## Part A

### A Least Squares Formulation

#### A1. An Integral Identity

In the following,  $\underline{x} = (x_1, x_2, x_3)$  is a point in  $\mathbb{R}^3$ ,  $\partial_i \equiv \frac{\partial}{\partial x_i}$ , and  $(ijk)$  indicates that the indices  $i, j, k$  are restricted to an even permutation of  $(123)$ . Thus

$$\operatorname{div} \underline{u} = \sum_{i=1}^3 \partial_i u_i,$$

$$\operatorname{curl} \underline{u} = (\partial_j u_k - \partial_k u_j), \quad (ijk),$$

$$\operatorname{grad} \underline{u} = (\partial_i u_j).$$

With

$$(\underline{u}, \underline{v}) \equiv [\text{grad} \underline{u} : \text{grad} \underline{v}] \equiv \sum_{i=1}^3 \text{grad} u_i \text{grad} v_i,$$

we define

$$\|\underline{u}\|^2 \equiv \int_D (\underline{u}, \underline{u}) d\pi,$$

$$\|\underline{u}\|_0^2 \equiv \int_D |\underline{u}|^2 d\pi$$

and

$$\|\underline{u}\|_1^2 = \|\underline{u}\|^2 + \|\underline{u}\|_0^2.$$

The system (1) may be written in matrix form as

$$(A.1.1) \quad L\underline{u} \equiv \sum_{i=1}^3 A_i \partial_i \underline{u} = \underline{f}$$

where  $\underline{f} = (\rho, \underline{\zeta})^T$  and

$$A_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Set

$$(A.1.2) \quad B_i = A_k^T A_j, \quad (ijk).$$

It is easy to verify that

$$(A.1.3) \quad \begin{aligned} A_i^T A_i &= I, \\ i &= 1, 2, 3, \\ B_i^T &= -B_i \end{aligned}$$

where  $I$  is the  $3 \times 3$  identity matrix.

Recalling the definition  $(\underline{u}, \underline{v}) \equiv [\text{grad}\underline{u}; \text{grad}\underline{v}]$ , integration by parts leads to the identity

$$(A.1.4) \quad (\underline{L}\underline{u})^T \underline{L}\underline{v} = (\underline{u}, \underline{v}) + \text{div}\underline{q} - \Omega(\underline{u})\underline{v}$$

where, if  $\underline{q} = (q_1, q_2, q_3)$ ,

$$(A.1.5) \quad q_i \equiv (B_k^T \partial_j \underline{u} + B_j \partial_k \underline{u})^T \cdot \underline{v}, \quad (ijk),$$

and  $\Omega(\underline{u})$  is a vector with components  $\Omega_i$  given by

$$(A.1.6) \quad \Omega_i(\underline{u}) = (B_k^T \partial_j \partial_i \underline{u} + B_k \partial_i \partial_j \underline{u}^T), \quad (ijk).$$

Suppose  $\underline{u}$  is smooth; since  $B_k^T = -B_k$ , then  $\Omega(\underline{u}) = 0$ . Also, in terms of the components of  $\underline{u}$  and  $\underline{v}$  the component  $q_i$  of  $\underline{q}$  is easily verified to have the form

$$q_i = v_i (\partial_j u_j + \partial_k u_k) - (v_j \partial_j + v_k \partial_k) u_i, \quad (ijk)$$

Let

$$(A.1.7) \quad \hat{q}_i = v_i (\partial_j u_j + \partial_k u_k) + u_i (\partial_j v_j + \partial_k v_k)$$

so that

$$q_i = \hat{q}_i - (\partial_j (v_j u_i) + \partial_k (v_k u_i)).$$

Since the expression in brackets is a surface divergence then

$$\int_{\Gamma} \underline{q} \cdot \underline{n} d\sigma = \int_{\Gamma} \hat{\underline{q}} \cdot \underline{n} d\sigma.$$

Integrating (A.1.4) and employing Gauss' theorem

$$\int_D (\underline{Lu})^T \underline{Ly} d\pi = \int_D (\underline{u}, \underline{y}) d\pi + \int_{\Gamma} \hat{\underline{q}} \cdot \underline{n} d\sigma$$

Suppose  $\underline{u} \cdot \underline{n} = \underline{v} \cdot \underline{n} = 0$  on  $\Gamma$ ; the preceding remark implies that  $\hat{\underline{q}} \cdot \underline{n} = 0$  on  $\Gamma$  so that

$$(A.1.9) \quad \int_D (\underline{Lu})^T \underline{Ly} d\pi = \int_D (\underline{u}, \underline{v}) d\pi.$$

## A2. Norm Estimates and Uniqueness

The goal here is to use the basic identity (A.1.9) and the Lax-Milgram Theorem to show existence and uniqueness for the system

$$(A.2.1) \quad \underline{Lu} = \underline{f} \quad \text{in } D, \quad \underline{u} \cdot \underline{n} = 0 \quad \text{on } \Gamma$$

where  $L$  is defined in (A.1.1). To do this we must first formulate (A.2.1) in terms of a bilinear form  $a(\cdot, \cdot)$  on an appropriate space  $V$ . In particular we put

$$(A.2.2) \quad a(\underline{v}, \underline{w}) = \int_D \underline{Lv}^T \underline{Lw} d\pi = \int_D \{ \text{div} \underline{v} \cdot \text{div} \underline{w} + \text{curl} \underline{v} \cdot \text{curl} \underline{w} \} d\pi$$

Moreover, we let

$$(A.2.3) \quad V = \{ \underline{v} \in \vec{L}^2(D) : \underline{v} \cdot \underline{n} = 0 \quad \text{on } \Gamma, \quad \text{grad} \underline{v} \in \vec{L}^2(D) \}$$

with

$$(A.2.4) \quad \|\underline{v}\| = \left\{ \int_D (\underline{v}, \underline{v}) d\pi \right\}^{1/2} = \left\{ \int_D [\text{grad} \underline{v} : \text{grad} \underline{v}] d\pi \right\}^{1/2}$$



This is a Hilbert space with the associated inner product

$$(A.2.5) \quad \langle \underline{u}, \underline{v} \rangle = \int_D (\underline{u}, \underline{v}) d\pi = \int_D (\text{grad} \underline{u} : \text{grad} \underline{v}) d\pi.$$

That the bilinear form  $a(\cdot, \cdot)$  satisfies the conditions of the Lax-Milgram Theorem [13] follows immediately from the integral identity (A.1.9), which can be written

$$(A.2.6) \quad a(\underline{u}, \underline{v}) = \int_D (\underline{u}, \underline{v}) d\pi.$$

Indeed, putting  $\underline{v} = \underline{u}$  we obtain

$$(A.2.7) \quad a(\underline{u}, \underline{u}) = \|\underline{u}\|^2,$$

while

$$(A.2.8) \quad |a(\underline{u}, \underline{v})| \leq \|\underline{u}\| \|\underline{v}\|$$

is also clear.

It follows that given any bounded linear functional  $G(\cdot)$  on  $V$  there is a unique  $\underline{u} \in V$  for which

$$(A.2.9) \quad a(\underline{u}, \underline{v}) = G(\underline{v}) \quad \text{all } \underline{v} \in V;$$

moreover,

$$(A.2.10) \quad \|\underline{u}\| \leq \|G\|.$$

Our final task is to choose  $G(\cdot)$  so that (A.2.9) is equivalent to (A.2.1) -

(A.2.2). Indeed, given  $\rho \in L^2(D)$ ,  $\underline{\zeta} \in \vec{L}^2(D)$  we put

$$(A.2.11) \quad G(\underline{v}) = \int_D \underline{f}^T \text{Lud}\pi = \int_D \{\rho \text{div} \underline{v} + \underline{\zeta} \cdot \text{curl} \underline{v}\} d\pi.$$

Observe that

$$(A.2.12) \quad |G(\underline{v})| \leq \|\underline{f}\|_0 \|\underline{L}\underline{v}\|_0 = \|\underline{f}\|_0 \|\underline{v}\|$$

so that

$$(A.2.13) \quad \|G\| \leq \|\underline{f}\|_0 = \{\|\rho\|_0^2 + \|\underline{\zeta}\|_0^2\}^{1/2}.$$

Moreover, (A.2.9) is equivalent to

$$(A.2.14) \quad \int_D |\underline{L}\underline{u} - \underline{f}|^2 d\pi = \int_D \{|\operatorname{div}\underline{u} - \rho|^2 + |\operatorname{curl}\underline{u} - \underline{\zeta}|^2\} d\pi$$

be minimized over  $\underline{v} \in V$ . Thus, if the data  $\rho, \underline{\zeta}$  satisfy the compatibility conditions

$$(A.2.15) \quad \int_D \rho d\pi = 0, \quad \operatorname{div}\underline{\zeta} = 0,$$

then the min in (A.2.14) will be zero and the minimizing function  $\underline{u} \in V$  will satisfy (A.2.1).

In conclusion, it follows that if

$$(A.2.16) \quad \rho \in L_0^2(D) \equiv \{\psi \in L^2(D) : \int_D \psi d\pi = 0\}$$

$$(A.2.17) \quad \underline{\zeta} \in \vec{L}_{\operatorname{div}}^2(D) \equiv \{\underline{v} \in \vec{L}^2(D) : \operatorname{div}\underline{v} = 0 \in D\}$$

are given then there is a unique  $\underline{u} \in V$  such that (A.2.1) holds. Moreover,

$$(A.2.18) \quad \|\underline{u}\| \leq (\|\rho\|_0^2 + \|\underline{\zeta}\|_0^2)^{1/2}.$$

## Part B

A Finite Element TreatmentB1. Least Squares Formulation

Since the infinite dimensional problem (A.2.1) - (A.2.2) has a natural characterization (via the Lax-Milgram Theorem) in terms of a least squares formulation, it is reasonable to consider approximations based on these ideas. Indeed, Let

$$(B.1.1) \quad V_h \subseteq V$$

be a finite dimensional subspace parameterized by  $h > 0$ . We seek a  $\underline{u}_h \in V_h$  which minimizes

$$(B.1.2) \quad \int_D |\underline{L}\underline{v}^h - \underline{f}|^2 = \int_D \{ |\operatorname{div}\underline{v}^h - \rho|^2 + |\operatorname{curl}\underline{v}^h - \underline{\zeta}|^2 \} d\pi$$

as  $\underline{v}^h$  varies over  $V_h$ . Observe that if  $a(\cdot, \cdot)$  and  $G(\cdot)$  are defined as in Section A.2, then  $\underline{u}_h$  is a minimizing function if and only if

$$(B.1.3) \quad a(\underline{u}_h, \underline{v}^h) = G(\underline{v}^h) \quad \text{all } \underline{v}^h \in V_h.$$

Moreover, application of the Lax-Milgram Theorem to  $V_h$  shows  $\underline{u}_h \in V_h$  is uniquely determined by (B.1.3). Once a basis is chosen for  $V_h$ , (B.1.3) reduces to a set of symmetric positive definite algebraic equations.

## B2. Error Estimates

Combining (A.2.9) and (B.1.3) we see that

$$(B.2.1) \quad a(\underline{u} - \underline{u}^h, \underline{v}^h) = 0 \quad \text{all } \underline{v}^h \in V_h.$$

This orthogonality condition is central to all error estimates. Indeed, first note that if  $\tilde{\underline{u}} \in V_h$  is given, then (B.2.1) gives

$$(B.2.2) \quad a(\underline{u} - \tilde{\underline{u}}, \underline{u} - \tilde{\underline{u}}) = a(\underline{u} - \underline{u}_h, \underline{u} - \underline{u}_h) + a(\underline{u}_h - \tilde{\underline{u}}, \underline{u}_h - \tilde{\underline{u}}).$$

Thus

$$\|\underline{u} - \tilde{\underline{u}}\|^2 = \|\underline{u} - \underline{u}_h\|^2 + \|\underline{u}_h - \tilde{\underline{u}}\|^2.$$

It follows that  $\underline{u}_h$  is a best approximation in the sense that

$$(B.2.3) \quad \|\underline{u} - \underline{u}_h\| = \inf \|\underline{u} - \tilde{\underline{u}}\| \leq \inf \|\underline{u} - \tilde{\underline{u}}\|_1,$$

where the inf is taken over all  $\tilde{\underline{u}}$  in  $V_h$ .

In particular, if  $V_h$  consist of piecewise linear elements, then (B.2.3) gives

$$(B.2.4) \quad \|\underline{u} - \underline{u}_h\| \leq Ch \|\underline{u}\|_2,$$

where  $h$  is a generic mesh spacing. Here  $\|\cdot\|_2$  is the Sobolev norm containing all derivatives up to second order.

To establish  $L_2$  estimates we use the standard "duality argument." The starting point is the following adjoint problem for  $\underline{w} \in V$ , where the error  $\underline{u} - \underline{u}_h$  is the data:

$$(B.2.5) \quad a(\underline{v}, \underline{w}) = \int_D \underline{v}^T [\underline{u} - \underline{u}_h] d\pi \quad \text{all } v \in V.$$

Suppose for the moment that (B.2.5) can be solved for  $\underline{w}$ , and

$$(B.2.6) \quad \|\underline{w}\|_2 \leq C \|\underline{u} - \underline{u}_h\|_0.$$

In this case we put  $\underline{v} = \underline{u} - \underline{u}_h$  in (B.2.5) to get

$$(B.2.7) \quad a(\underline{u} - \underline{u}_h, \underline{w}) = \int_D |\underline{u} - \underline{u}_h|^2 = \|\underline{u} - \underline{u}_h\|_0^2;$$

using orthogonality (i.e. (B.2.1)) we get

$$(B.2.8) \quad a(\underline{u} - \underline{u}_h, \underline{w} - \underline{w}^h) = \|\underline{u} - \underline{u}_h\|_0^2$$

for any  $\underline{w}^h$  in  $V_h$ . Thus

$$(B.2.9) \quad \|\underline{u} - \underline{u}_h\| \|\underline{w} - \underline{w}^h\| \geq \|\underline{u} - \underline{u}_h\|_0^2.$$

We select  $\underline{w}_0^h \in V_h$  so that

$$(B.2.10) \quad \|\underline{w} - \underline{w}_0^h\| \leq Ch \|\underline{w}\|_2.$$

Thus, with (B.2.6) we get

$$(B.2.11) \quad \|\underline{u} - \underline{u}_h\|_0 \leq Ch \|\underline{u} - \underline{u}_h\|$$

Therefore, if linear elements are used

$$\|\underline{u} - \underline{u}_h\|_0 \leq Ch^2 \|\underline{u}\|_2.$$

The final task is to check (B.2.5) for solvability as well as the a priori bound (B.2.6). Rewriting (B.2.5) we get

$$(B.2.12) \quad \int_D \underline{L} \underline{v}^T \underline{L} \underline{w} d\pi = \int_D \underline{v}^T (\underline{u} - \underline{u}_h) d\pi \quad \text{all } \underline{v} \in V.$$

Suppose  $\underline{v} \in V$  and  $\underline{v} = 0$  on  $\Gamma$ . Then integration by parts gives

$$(B.2.13) \quad \int_D \underline{v}^T \underline{L}^* \underline{L} \underline{w} d\pi = \int_D \underline{v}^T (\underline{u} - \underline{u}_h) d\pi \quad \text{all } \underline{v} \in V,$$

where

$$\underline{L}^* \underline{L} = \text{curl curl-grad div} = \underline{\Delta}.$$

Thus defining  $\underline{w}$  by

$$\underline{L}^* \underline{L} \underline{w} = \underline{u} - \underline{u}_h \quad \text{in } D$$

$$\underline{w} = 0 \quad \text{on } \Gamma$$

it follows that  $\underline{w}$  satisfies (B.2.5). Moreover, the a priori bound (B.2.6) follows from the theory of second order elliptic equations [8].

## Part C

A Finite Difference ApproachC1. Notational Preliminaries;Box Variables.

The notations about to be introduced are most naturally interpreted when  $D$  can be subdivided into cells  $\{\pi\}$  each of which is a rectangular box. In a later section we shall indicate how more general subdivisions may be treated explicitly.

Following Keller [7] we call  $v$  a box variable if it is defined at the vertices of cells. For our purpose the importance in employing box variables lies in the fact that any average value of a function taken over either a cell volume, a face, or an edge of a cell can be approximated in terms of box variables by means of the trapezoidal rule. Certain other properties to be described then provide a finite difference calculus by means of which summation by parts leads to results similar to those established in Section A.

We employ the notation:  $v^1$  indicates the centered average and  $v_{,1}$  the centered divided difference with respect to  $x_1$ , i.e.,

$$v^1 = (v(x_1 + \Delta x_1 / 2) + v(x_1 - \Delta x_1 / 2)) / 2$$

(C.1.1)

$$v_{,1} = (v(x_1 + \Delta x_1 / 2) - v(x_1 - \Delta x_1 / 2)) / \Delta x_1 .$$

For smooth  $v$ , therefore,

$$v^1 = \left( \int_{x_1 - \Delta x_1 / 2}^{x_1 + \Delta x_1 / 2} v(x) dx \right) \div \Delta x_1 + O(\Delta x_1^2);$$

(C.1.2)

in particular,

$$(C.1.3) \quad v_{,k}^{ij} = \left( \int_{\pi} \partial_k v dx_i dx_j dx_k \right) / \Delta\pi + O(h^2)$$

where  $\Delta\pi = \Delta x_1 \Delta x_2 \Delta x_3$  and  $h = \min\{\Delta x_i\}$ .

The algebraic identity

$$(C.1.4) \quad (vw)_{,i} = v_{,i}^i w + w_{,i}^i v$$

then yields a summation by parts formula while the definition

$$v_{,ij} = \left[ v(x_i + \Delta x_i / 2, x_j + \Delta x_j / 2) + v(x_i - \Delta x_i / 2, x_j - \Delta x_j / 2) - v(x_i - \Delta x_i / 2, x_j + \Delta x_j / 2) - v(x_i + \Delta x_i / 2, x_j - \Delta x_j / 2) \right] \div (\Delta x_i \Delta x_j)$$

shows that

$$(C.1.5) \quad v_{,ij} = v_{,ji}$$

The definitions

$$(C.1.6) \quad \text{div}_{\underline{h}} \underline{u} \equiv \sum_{i=1}^3 u_{,i}^{jk}$$

$$\text{curl}_{\underline{h}} \underline{u} \equiv \left( u_{,k}^{ij} \quad -u_{,j}^{ik} \right) \quad (ijk)$$

provide finite volume approximations to  $\text{div} \underline{u}$  and  $\text{curl} \underline{u}$  respectively, i.e.,

$$(C.1.7) \quad \text{div}_{\underline{h}} \underline{u} = \left( \int_{\underline{p}} \text{div} \underline{u} d\pi \right) \div \Delta\pi + O(h^2)$$

$$\text{curl}_{\underline{h}} \underline{u} = \left( \int_{\pi} \text{curl} \underline{u} \delta\pi \right) \div \Delta\pi + O(h^2).$$



We propose to examine the finite difference approximations to (1) given, in terms of box variables in a cell, by

$$\begin{aligned} \text{div}_{\underline{h}} \underline{u} &= \rho^{ijk} \\ \text{curl}_{\underline{h}} \underline{u} &= \underline{\zeta}^{ijk} \end{aligned} \quad (\text{ijk})$$

(C.1.8a)

The boundary conditions  $\underline{u} \cdot \underline{n} = 0$  are imposed by

$$\underline{u}^{ij} \cdot \underline{n}_k = 0 \quad (\text{ijk})$$

(C.1.8b)

where  $\underline{u}^{ij}$  is the trapezoidal approximation to the average value of  $u$  on a face  $\sigma_k$  whose normal is  $\underline{n}_k$ . When box variables are understood we shall often write  $\underline{u} \cdot \underline{n} = 0$  to mean the condition expressed by (C.1.8b).

## C2. A Summation Identity

Define, using box variables,

$$L_{\underline{h}} \underline{u} \equiv A_1 \underline{u}_{,1}^{23} + A_2 \underline{u}_{,2}^{13} + A_3 \underline{u}_{,3}^{12},$$

(C.2.1)

$$\underline{f}_{\underline{h}} \equiv (\rho^{123}, \underline{\zeta}^{123})$$

where the coefficient matrices are the same as in the definition of  $L\underline{u}$  in (A.1.1). The box-variables scheme (C.1.8) expressing  $\text{div} \underline{u} = \rho$ ,  $\text{curl} \underline{u} = \underline{\zeta}$  may then be written as

$$(C.2.2) \quad L_{\underline{h}} \underline{u} = \underline{f}_{\underline{h}}.$$

Next, define

$$\text{grad}_{\underline{h}} \underline{u} \equiv (\underline{u}_{,1}^{23}, \underline{u}_{,2}^{13}, \underline{u}_{,3}^{12})$$

and

$$(C.2.3) \quad (\underline{u}, \underline{v})^{\underline{h}} \equiv (\text{grad}_{\underline{h}} \underline{u} : \text{grad}_{\underline{h}} \underline{v}).$$

The summation-by-parts formula (C.1.4) then leads to

$$(C.2.4) \quad (L_{\underline{h}} \underline{u})^T (L_{\underline{h}} \underline{v}) = (\underline{u}, \underline{v})^{\underline{h}} + \text{div}_{\underline{h}} \underline{q}^{\underline{h}} - \Omega_{\underline{h}}(\underline{u}) \underline{v}^{\underline{h}}$$

where  $\underline{q}^{\underline{h}}$  is the vector with components

$$(C.2.5) \quad q_i^{\underline{h}} = (B_{k-,j}^T u^k + B_{j-,k} u^j) v^{ik}, \quad (ijk)$$

and

$$\Omega_{\underline{h}} \underline{v}^{\underline{h}} = \sum_{i=1}^3 (B_{k-,ij} u^k + B_{k-,ji}^T u^k) v^{ijk}, \quad (ijk).$$

Since  $\underline{u}$  is a box-variable  $\underline{u}_{,ij}^k = \underline{u}_{,ji}^k$  ((C.1.5)) and, since  $B_k^T = -B_k$ , then  $\Omega_{\underline{h}} \underline{v}^{\underline{h}} = 0$ .

Next, multiply (C.2.4) by  $\Delta\pi$  and sum over  $D$  using the summation analogue of Gauss' theorem to obtain

$$(C.2.6) \quad \sum_D (L_{\underline{h}} \underline{u})^T L_{\underline{h}} \underline{v} \Delta\pi = \sum_D (\underline{u}, \underline{v})^{\underline{h}} \Delta\pi + \sum_{\Gamma} \underline{q}^{\underline{h}} \cdot \underline{n} \Delta\pi.$$

By expressing  $\underline{q}^{\underline{h}}$  as given by (C.2.5) in terms of the components of  $\underline{u}$  and  $\underline{v}$  (as in (A.1.7)) the reader may verify that the boundary contribution vanishes when  $\underline{u} \cdot \underline{n} = \underline{v} \cdot \underline{n} = 0$  on  $\Gamma$ .

Hence, defining

$$(C.2.7) \quad a_h(\underline{u}, \underline{v}) \equiv \sum_{D_h} (\underline{u}, \underline{v})^h \Delta\pi$$

we may state: for any box-variables  $\underline{u}$  and  $\underline{v}$  satisfying (C.1.8b), i.e.,  $\underline{u} \cdot \underline{n} = \underline{v} \cdot \underline{n} = 0$  on  $\Gamma$ ,

$$(C.2.8) \quad a_h(\underline{u}, \underline{v}) = \sum_D (L_h \underline{u})^T L_h \underline{v} \Delta\pi.$$

### C3. A Convergence Estimate

The box-scheme (C.2.2) is an overdetermined system of algebraic equations under the boundary conditions  $\underline{u} \cdot \underline{n} = 0$  on  $\Gamma$ . Consider a solution  $\underline{u}^h$  as determined by the least squares problem

$$(C.3.1) \quad \min_{\underline{u}} \sum_D (L_h \underline{u} - \underline{f}_h)^T (L_h \underline{u} - \underline{f}_h) \Delta\pi$$

with  $\underline{u} \cdot \underline{n} = 0$  on  $\Gamma$ .

Using (C.2.8) the Euler equations arising from this problem leads to:  $\underline{u}^h$  provides a least-square solution of  $L_h \underline{u} = \underline{f}_h$  if

$$(C.3.2) \quad a_h(\underline{u}^h, \underline{v}) = \sum_D \underline{f}_h^T L_h \underline{v} \Delta\pi$$

for any box-variable  $\underline{v}$  satisfying  $\underline{v} \cdot \underline{n} = 0$ . Write

$$(C.3.3) \quad \|\underline{u}^h\|^h \equiv (a_h(\underline{u}^h, \underline{u}^h))^{1/2}.$$

Since only central averages and differences are involved in the definition of  $L_h \underline{u}$  it follows that if  $\underline{u}$  is a solution of  $L_h \underline{u} = \underline{f}$  which has

continuous and bounded mixed third derivatives then  $L_h(\underline{u}-\underline{u}^h) = O(h^2)$ , i.e.

$$(C.3.4) \quad \|\underline{u}-\underline{u}^h\|^h = O(h^2).$$

Define

$$(C.3.5) \quad \|\underline{u}\|_0^h = \left[ \sum_{\pi \in D} \sum_{\sigma \in \pi} \underline{u}^T(\sigma) \underline{u}(\sigma) \right]^{1/2}$$

where  $\underline{u}(\sigma)$  indicates the box-variable approximation to the average value of  $\underline{u}$  over a face  $\sigma$  of a cell  $\pi$ . For the continuous problem Friedrichs' inequality, when  $\underline{u} \cdot \underline{n} = 0$ , yields

$$\|\underline{u}\|_0 \leq \gamma \|\underline{u}\|$$

for some constant  $\gamma$ . The same argument which establishes this inequality may also be followed using summation and difference operators in place of integration and differential operators. The result is

$$(C.3.6) \quad \|\underline{u}\|_0^h \leq \gamma \|\underline{u}\|^h.$$

Using (C.3.4) we thus obtain

$$\|\underline{u}-\underline{u}^h\|_0^h = O(h^2),$$

i.e. the box-scheme is second order accurate in the  $\|\cdot\|_0^h$  norm.

#### C4. A Remark About More General Cells

As indicated earlier the definitions of  $\text{div}_h \underline{u}$ ,  $\text{curl}_h \underline{u}$  can be made independent of any assumption about the shape of a cell  $\pi$ . Properly interpreted, so also can the summation by parts formula (C.1.4) as well as (C.1.5). A little reflection will convince the reader that the convergence proof just given applies as well for irregularly shaped subdomains.

The following describes explicit representations for the box-scheme on irregular cells.

Let  $\sigma_\nu$  denote an oriented face of  $\pi$ ,  $\nu = 1, 2, \dots, 6$ . Applying (C.1.7) to a smooth function  $\underline{u}$ , and employing Gauss' theorem,

$$(C.4.1) \quad \Delta\pi \cdot \text{div}_h \underline{u} \approx \int_\pi \text{div} \underline{u} d\pi = \sum_\nu \int_{\sigma_\nu} \underline{u} \cdot d\sigma \approx \hat{\underline{u}}(\underline{\sigma}_\nu) \cdot \Delta\sigma_\nu$$

where  $\hat{\underline{u}}(\underline{\sigma}_\nu)$  indicates the value of  $\underline{u}$  at the centroid of  $\sigma_\nu$ . By approximating  $\hat{\underline{u}}(\underline{\sigma}_\nu)$  by the average of its values at the vertex points of  $\sigma_\nu$  an expression for  $\text{div}_h \underline{u}$  results when  $\underline{u}$  is a box-variable.

Similarly,

$$\text{curl}_h \underline{u} \cdot \underline{n}_1 \approx \int_\pi (\text{curl} \underline{u} \cdot \underline{n}_1) d\pi / \Delta\pi.$$

Apply Stoke's theorem on a face  $d\underline{\sigma}_\nu$  and let  $d\underline{s}_\nu$  indicate an element of arc length in the direction  $\underline{t}_{\nu 1} = d\underline{\sigma}_\nu \times \underline{n}_1$ ; the result is

$$(C.4.2) \quad \text{curl}_h \underline{u} \cdot \underline{n}_1 \approx \sum_\nu \int_{\sigma_\nu} (\underline{u} \cdot \underline{t}_{\nu 1}) ds_\nu dx_1 / \Delta\pi \approx \sum_\nu (\underline{u} \cdot \underline{t}_{\nu 1}) \Delta\sigma_\nu / \Delta\pi$$

where  $\underline{u} \cdot \underline{t}_{\nu 1}$  is evaluated at the centroid of the face  $\sigma_\nu$  having the area  $\Delta\sigma_\nu$ . By evaluating  $\underline{u} \cdot \underline{t}_{\nu 1}$  as the average of its values at the vertices of  $\sigma_\nu$  (C.4.2) provides an interpretation of  $\text{curl}_h \underline{u}$  in terms of box-variables.

Concluding Remarks

A comparison of the convergence proofs employed in Parts B and C is of interest. The finite-element approach allowed the integral identity (A.1.9) to be used directly but utilized estimates arising from the adjoint problem. The finite difference approach, on the other hand, required the development of a summation identity corresponding to (A.1.9); Friedrichs' inequality then provided in the required convergence estimate.

Both proofs are independent of any assumptions about the type of cells  $\pi$  upon which approximations are based. However, the numerical implementation of the finite-element approach in such cases may be simpler to employ than the box-scheme because of extensive and readily available software for finite-element methods. The above remarks suggest that this software could be adapted to the finite difference method as well.

Both methods lead to sparse matrices which may be solved by direct algebraic techniques. An iterative scheme for least squares problems due to Kaczmarz [6] and Tanabe [11] provides an alternative approach and has been employed in [4] to treat the finite difference problem in two dimensions. Progress in developing fast iterative methods has been reported to us in personal communications by our colleagues (Grosch and Phillips) and will be described elsewhere.

## REFERENCES

- [1] G. J. FIX, M. D. GUNXBURGER, R. A. NICOLAIDES. On mixed finite element methods for first order elliptic systems, *Numer. Math.*, 37 (1981), pp. 29-48.
- [2] G. J. FIX, M. D. GUNXBURGER, R. A. NICOLAIDES. On finite element methods of the least squares type, *Comp. Math. Appl.*, 5 (1979), pp. 87-98.
- [3] P. A. GARIART and P. A. RAVIART. *Finite Element Approximations to the Navier-Stokes Equations*, Springer-Verlag, New York, 1980.
- [4] T. B. GATSKI, C. E. GROSCHE and M. E. ROSE. A numerical study of the two-dimensional Navier-Stokes equations in vorticity-velocity variables, *J. Comput. Phys.*, Vol. 48, No. 1, (1982), pp. 1-22.
- [5] M. GHIL and R. BALGOVIND. A fast Cauchy-Riemann solver, *Math. Comp.*, Vol. 33, No. 146, April 1979, pp. 585-635.
- [6] S. KACZMARZ. *Bull. Acad. Pol. Sci. Lett.*, A, (1937), pp. 355.
- [7] H. B. KELLER. A new difference scheme for parabolic equations, *Numerical Solutions of Partial Differential Equations II*, Academic Press, Inc., New York, 1971.

- [8] J. L. LIONS and E. MAGNES. Nonhomogeneous Boundary Value Problems, Springer-Verlag, 1973.
  
- [9] H. LOMAX and E. D. MARTIN. Fast direct numerical solutions of the nonhomogeneous Cauchy-Riemann equations, J. Comp. Phys., 15 (1974).
  
- [10] M. E. ROSE. A 'unified' treatment of the wave equation and the Cauchy-Riemann equations, SIAM J. Numer. Anal., Vol. 19, No. 4, August 1982.
  
- [11] K. TANABE. Numer. Math., Vol. 17, (1971), pp. 203.
  
- [12] B. WENDROFF. The structures of certain finite difference schemes, SIAM Rev., 3 (1961), pp. 237-242.
  
- [13] K. YOSIDA. Functional Analysis, Springer-Verlag, New York, 1965.



1 Report No NASA CR-166113		2 Government Accession No		3 Recipient's Catalog No	
4 Title and Subtitle A COMPARATIVE STUDY OF FINITE ELEMENT AND FINITE DIFFERENCE METHODS FOR CAUCHY-RIEMANN TYPE EQUATIONS				5 Report Date March 1983	
7 Author(s) George J. Fix and Milton E. Rose				6 Performing Organization Code	
9 Performing Organization Name and Address Institute for Computer Applications in Science & Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23665				8 Performing Organization Report No 83-4	
12 Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, DC 20546				10 Work Unit No	
				9 Contract or Grant No NAS1-17070 & NAS1-17130	
				13 Type of Report and Period Covered Contractor Report	
				14 Sponsoring Agency Code	
15 Supplementary Notes Technical Monitor: Robert H. Tolson Final Report					
16 Abstract A least squares formulation of the system $\text{div} \underline{u} = \rho$ , $\text{curl} \underline{u} = \underline{\zeta}$ is surveyed from the viewpoint of both finite element and finite difference methods. Closely related arguments are shown to establish convergence estimates.					
17 Key Words (Suggested by Author(s)) finite element, finite difference, Cauchy-Riemann equations, least squares				18 Distribution Statement  Unclassified - Unlimited  Subject Category 64	
19 Security Classif (of this report) Unclassified		20 Security Classif (of this page) Unclassified		21 No of Pages 24	22 Price A02

**End of Document**