

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NASA CR-170601

WASHINGTON UNIVERSITY
DEPARTMENT OF EARTH AND PLANETARY SCIENCES

(NASA-CR-170601) COPERNICUS OBSERVATIONAL
SEARCHES FOR OH AND H₂O IN DIFFUSE CLOUDS
Final Technical Report (Washington Univ.)
40 p HC A03/MF A01

N84-12004

CSCI 03A

G3/89 Unclass
44511

NASA Grant NSG 5187
Final Technical Report



Submitted By:

Wm Hayden Smith
Principal Investigator

Final Technical Report for NASA Grant NSG 5187

The above named grant resulted in a paper on the abundances of OH and H₂O in the interstellar toward the chosen stars. The Copernicus observations were successfully carried out, the data were reduced, and analysed. We found that the lower limits for the OH/H₂O ratio are, at present, consistent with the theoretical models. The effort we made here has led to a better understanding of the validity of the molecule producing mechanisms towards diffuse clouds, and has uncovered through continued laboratory study, even more favorable transitions of H₂O with which to establish the interstellar abundance of H₂O in diffuse clouds. The enclosed paper has now been submitted for publication.

I would like to thank NASA for their support in this program.

Invention Report

No inventions were made during the course of this research.

COPERNICUS OBSERVATIONAL SEARCHES FOR OH AND H₂O
IN DIFFUSE CLOUDS

by

Wm. Hayden Smith*
McDonnell Center for Space Sciences
Department of Chemistry
Washington University
St. Louis, MO 63130

and

Theodore P. Snow, Jr.
Laboratory for Atmospheric and Space Physics
University of Colorado
Boulder, CO 80309

submitted to Ap.J. (July, 1978)

ABSTRACT

An intensive search for OH and H₂O in the directions of σ Sco, α Cam, and \circ Per was undertaken with the Copernicus satellite. Multiple scans were carried out over the wavelength region for the expected absorption features due to the OH D-X and H₂O \tilde{C} - \tilde{X} transitions. The feature due to OH was detected marginally towards σ Sco, and only an upper limit can be given towards α Cam. H₂O was not detected in any of the stars at the signal level accumulated. The OH abundance towards σ Sco and the respective lower limits for the OH/H₂O ratios are discussed with regard to the extant models for the steady state abundances of OH and H₂O, and shown not to be inconsistent with ion-molecule schemes.

*Guest Investigator with the Princeton University telescope on the Copernicus satellite, which is sponsored and operated by the National Aeronautics and Space Administration.

I. INTRODUCTION

The effectiveness of molecule-producing processes in diffuse clouds can hardly be questioned. The molecules observed to date include H_2 , CO, OH, CH, CH^+ , CN and C_2 . Reviews of possible formation mechanisms have been given by Watson (1975), Herbst and Klemperer (1973), and Dalgarno and Black (1976), among others. In spite of the amount of information already in hand and the formation models calculated, with the exception of molecular hydrogen (Hollenbach and Salpeter, 1971), there is not yet a fully understood formation mechanism for any given molecule. While schemes can be devised which produce large amounts of any of the above molecules, the lack of adequate laboratory data on relevant processes usually requires the use of poorly-supported estimates for various rate constants. A possible exception to this situation may be found in the ion-molecule scheme for the formation of OH and H_2O , where the abundances and ratios of OH/ H_2O are determined by processes which would appear to be relatively well understood. Both molecules have transitions from the ground state in the ultraviolet. Observation of OH and H_2O in various lines-of-sight then offers a possibly significant test for that particular ion-molecule formation scheme.

Since a few lines-of-sight have already been investigated (Crutcher and Watson, 1976; Chaffee and Lutz, 1977;


ORIGINAL LITERATURE
OF POOR QUALITY

Now, 1976a, 1976b, 1977), it is known that OH frequently exists at detectable levels and the H_2O is less abundant. We have therefore obtained very extensive new data on σ Sco and α Cam for both molecules and also have carried out further scans of the H_2O transition in the spectrum of α Per. After a brief discussion of the ion-molecule scheme for OH and H_2O production, we will detail our observations and the conclusions derived from them.



II. ION-MOLECULE REACTION FORMATION OF OH AND H₂O

Using the terminology of Dalgarno and Black (1976), we may write the following reaction scheme for OH and H₂O.

First, the charge exchange



(Field and Steigman, 1971) initiates the reaction scheme. The O⁺ thus formed is removed by reaction with H₂ in several successive reactions which are interrupted by recombination with ambient electrons, and hence are very sensitive to the depletion of elements producing the electrons. Two reactions ultimately produce H₂O and OH:



The branching ratio for this latter reaction has recently been given as approximately 10 in favor of OH by Herbst (1978), rather than the value of 1 assumed by many earlier studies (Watson, 1975; Herbst and Klemperer, 1973; Smith and Zweibel, 1976; and Glassgold and Langer, 1975). Destruction processes for OH and H₂O have been studied by Smith and

Zweibel and by Smith and Stella (1975). For a typical cloud of 100 H atoms cm^{-3} with a depletion factor f of 0.2, and an obscuration of 2 magnitudes at 950 \AA , the ratio of OH to H_2O is given by

$$n(\text{OH})/n(\text{H}_2\text{O}) = \left(\frac{\beta_{\text{H}_2\text{O}}}{\beta_{\text{OH}} + \lambda_{\text{OH}}} \right) \frac{1}{(1-\alpha)(1-f)}$$

which has been simplified from Equation 10 of Smith and Zweibel (1976). In this equation $\beta_{\text{H}_2\text{O}}$ and β_{OH} , the photodissociation rates, are given by Smith and Zweibel in their Table 1, and λ_{OH} is given by Herbst and Klemperer (1973) as $1.6 \times 10^{-9} \text{ cm}^3/\text{sec}$ times n_{ct} . According to Herbst (1978), the branching ratio α can be as high as 10, while the commonly adopted value 1 is actually a lower limit. The higher ratio in favor of OH increases $n(\text{OH})/n(\text{H}_2\text{O})$ by a factor of 5, or in the above simplified equation, gives $n(\text{OH})/n(\text{H}_2\text{O})$ as about 25, probably accurate to within a factor of 2 considering the approximations involved.

IV. RESULTS AND CONCLUSIONS

In Table 1 we summarize the extant ultraviolet observations of OH and searches for H_2O in various directions, including some earlier results and cloud parameters. Although the transitions of both molecules are well characterized as to wavelength, the strengths of the features have been the subject of considerable discussion in the literature since we initiated these observations. The OH D-X transitions' line strengths have been calculated and subsequently measured indirectly. Ray and Kelly (1975) computed an f -value of 0.0015, while Chaffee and Lutz (1977) derived an f -value for the 1220 Å line from their observation of the near ultraviolet transitions of OH near 3078 Å and the equivalent width for the OH D-X lines measured by Snow towards ζ Per. The recommended f -value for the transition is 0.0052 from the observational results. This value is much smaller than expected for an allowed Rydberg transition. The situation with regard to the f -value for the H_2O Rydberg transition near 1239 Å is also rife with uncertainty. Smith and Zweibel (1976) estimated from literature data an f -value which was believed to be correct to within a factor of 2. The f -value was 0.018. Very recently, Smith and Parkinson (1978) found an upper limit to the line f -value of 0.023, consistent with the earlier estimate. For the present, $f = 0.018$ is adopted.

The OH/H₂O limits given in the last column of Table 1 are consistent with the models, but only if Herbst's (1978) branching ratio strongly favoring the production of OH is correct. Hence, the present data appear to substantiate a high value for this branching ratio. These limits, however, refer only to the ground rotational state. For $T_{\text{kin}} \leq 16^\circ\text{K}$ (Snow, 1978), most of the molecules will be in this state, so this should not introduce much error. In any case, there is insufficient data available on collisional excitation cross-sections, radiative lifetimes, and the exact kinetic temperature to make any evaluation of rotational excitation very reliable.

Another factor which comes into play was recently discussed by Sandell (1977); viz. the shielding of molecules by the grains may be much less efficient than has previously been supposed. Grains as described by Lillie and Witt (1976) will have strongly forward scattering phase functions allowing the star light to penetrate much further into obscured clouds than would be assumed for isotropically scattering grains. In this case, the fact that longer wavelengths of light are much more effective in photodissociating H₂O than OH will result in larger OH/H₂O ratios. In order to treat this problem correctly, the transfer of radiation must be handled as indicated by Sandell. Overall, the theoretical situation has evolved to the point that predictions are extremely model-dependent. Our upper limits

then are of considerably reduced value although at least indicative. A ray of hope exists that detection of H_2O may yet be possible through the features accessible to the Copernicus satellite. A stronger Rydberg series near 1115 \AA may contain features sharp and strong enough for such observations (Johns, 1978). A careful search there could increase the lower limit of the ratio to allow a significant test of model predictions..

Work done at Princeton was supported by the National Aeronautics and Space Administration Contract NAS5-23576.

ORIGINAL PAGE IS
OF POOR QUALITY

Table 1. OH and H₂O Observations

HD	E(B-V)	log N(H)* cm ⁻²	N(H ₂)/N(H)*	n _H ⁺ cm ⁻³	OH Wλ(mA)	No. Scans	H ₂ O Wλ(mA)	No. Scans	N(OH)/N(H ₂ O)**
23180	0.32	21.21	0.50	10 ³ : 10 ³	Q: 5.8±2.0 P: < 2.0	32	< 0.90	87	> 23.0
24398	0.33	21.20	0.59	10 ³ : 10 ³	Q: 3.6±1.7	31	--	--	--
30614	0.32	21.09	0.35	--	Q: < 1.8	93	< 1.7	60	--
147165	0.40	21.37	0.05	740	Q: 0.71±.6	120	< .37	120	> 6.9
149757	0.32	21.15	0.63	120	Q: 3.5±.42 P: 2.3±.42	30	< .76	30	> 16.4

*hydrogen column densities are from Savage et al. (1977).

**density of hydrogen nuclei are from CO rotational equilibrium analyses by Snow (1976b, 1977); Smith, Stecher, and Krishnamoorthy (1978); and Snow and Jenkins (1978).

The ratio N(OH)/N(H₂O) given here refers only to the ground rotational state.

ORIGINAL PAGE IS
OF POOR QUALITY

REFERENCES

ORIGINAL PAGE IS
OF POOR QUALITY

- Chaffee, F.H., Jr., and Lutz, B.L., 1977. Ap. J., 213, 394.
- Crutcher, R.M., and Watson, W.D., 1976. Ap. J. (Letters), 203, L123.
- Dalgarno, A., and Black, J.H., 1976. Rep. Prog. Phys., 39, 573.
- Field, G.B., and Steigman, G., 1971. Ap. J., 166, 59.
- Glassgold, A.E., and Langer, W.D., 1975. Ap. J., 197, 347.
- Herbst, E., 1978. Preprint.
- Herbst, E., and Klemperer, W., 1973. Ap. J., 185, 505.
- Hollenbach, D.J., and Salpeter, E.E., 1971. Ap. J., 163, 155.
- Johns, J.W.C., 1978. Private communication.
- Lillie, C.F., and Witt, A.N., 1976. Ap. J., 208, 64.
- Ray, S., and Kelly, H.P., 1975. Ap. J. (Letters), 202, L57.
- Rogerson, J.B., Spitzer, L., Drake, J.F., Dressler, K., Jenkins, E.B., Morton, D.C., and York, D.G., 1973. Ap. J. (Letters), 181, L97.
- Sandell, G., 1977. Preprint.
- Savage, B.D., Bohlin, R.C., Drake, J.F., and Budich, W., 1977. Ap. J., 216, 291.
- Smith, A.M., Stecher, T.P., and Krishna Swamy, K.S., 1975. Ap. J., 220, 138.
- Smith, P.L., and Parkinson, W.H., 1978. Private communication.
- Smith, W.H., and Stella, G., 1975. J. Chem. Phys., 63, 2395.
- Smith, W.H., and Zweibel, E.G., 1976. Ap. J., 207, 758.

Snow, T.P., 1976a. Ap. J., 204, L127.

ORIGINAL PAGE IS
OF POOR QUALITY

Snow, T.P., 1976b. Ap. J., 204, 759.

Snow, T.P., 1977. Ap. J., 216, 724.

Snow, T.P., 1978: Ap. J. (Letters), 220, L93.

Snow, T.P., and Jenkins, E.B., 1978. In preparation.

Watson, W.D., 1975. Atomic and Molecular Physics and the
Interstellar Matter (Amsterdam, North-Holland).

York, D.G., and Miller, A.E., 1974. Memo to Users of Copernicus
Data, Dec. 9.

Authors' Addresses:

W. M. Smith
McDonnell Center for Space Sciences
Department of Chemistry
Washington University
St. Louis, MO 63130

T. P. Snow, Jr.
Laboratory for Atmospheric and Space Physics
University of Colorado
Boulder, CO 80309

ORIGINAL PAGE IS
OF POOR QUALITY

FIGURE CAPTIONS

Figure 1 - Average of 87 scans of the H_2O feature in the spectrum of α Per. The error envelope represents the 2σ uncertainty due to photon statistics. The expected position of the line is indicated.


Figure 2 - The averaged OH data on α Cam.

Figure 3 - The averaged H_2O data on α Cam.


Figure 4 - The averaged OH data on σ Sco. The feature is marginally present.

Figure 5 - The averaged H_2O data on σ Sco.


ORIGINAL PAGE IS
OF POOR QUALITY




ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 10
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 10
OF POOR QUALITY

