

---

# **Numerical Modeling of D-Mapping with Applications to Chemical Kinetics**

---

S.K. Dey

---

March 1984

---

# **Numerical Modeling of D-Mapping with Applications to Chemical Kinetics**

---

S. K. Dey, Ames Research Center, Moffett Field, California



National Aeronautics and  
Space Administration

**Ames Research Center**  
Moffett Field California 94035

# NUMERICAL MODELING OF D-MAPPINGS WITH APPLICATIONS TO CHEMICAL KINETICS

S. K. Dey

Computational Fluid Dynamics Branch, NASA Ames Research Center,  
Moffett Field, California 94035

**Abstract.** Numerical modeling of D-mappings has been studied and applied to solving nonlinear stiff systems. These mappings have been locally linearized for convergence analysis, and some applications have been made to chemical kinetics.

**Keywords.** Nonlinear equations; numerical methods; biomedical.

## INTRODUCTION

A numerical method was developed by Dey (1977) for solving nonlinear systems, some applications of which were later made to stiff systems (Dey, 1982). Convergence analysis was done using nonlinear D-mappings (Dey, 1981).

It is extremely difficult to represent this analysis computationally. Local linearization for such an analysis, which rendered computational modeling of D-mappings feasible, was suggested by Lomax (1983). In this article we discuss linearized modeling of D-mappings and some applications of the method.

## D-MATRICES AND D-MAPPINGS

If a sequence of square matrices of the same order satisfy the following condition,

$$\lim_{k \rightarrow \infty} A_k A_{k-1} \dots A_1 = 0 \quad (1)$$

each  $A_k$  is called a D-matrix. A D-matrix is not necessarily a convergent matrix, and conversely.

**Theorem 1.** A sufficient condition that  $A_k$  is a D-matrix is that

$$\|A_k\|_q \leq \alpha < 1 \quad (2)$$

$\forall k > K$  and that  $q$  is the same  $\forall k$ .

Theorem 1 is easily proved. Let

$$u^k = (u_1^k, u_2^k, \dots, u_j^k)^T \in D_k,$$

$k = 1, 2, \dots$ , ( $u_j^k$  = value of  $u_j$  at some  $k$ th iteration).

Let us consider a chained linear spaces  $D_k \subseteq D_{k-1} \subseteq \dots \subseteq D \subset \mathbb{R}^n$ .  $\mathbb{R}^n$  =  $n$ -dimensional real space. Let  $u^k, u^* \in D_k \forall k$  and

$$G_k: D_{k+1} \times D_k \rightarrow D_{k+1}. \text{ If}$$

$$\begin{aligned} G_k(u^{k+1}, u^k) &= G_k(u^*, u^*) \\ &= A_k(u^{k+1} - u^*) + B_k(u^k - u^*) \end{aligned} \quad (3)$$

and  $\forall k > K$ , and if  $(I - A_k)^{-1}B_k$  is a D-matrix,  $G_k$  is called a D-mapping (Dey, 1981).

If we now consider a nonstationary iterative scheme of the form

$$u^{k+1} = G_k(u^{k+1}, u^k) \quad (4)$$

and if  $G_k: D_{k+1} \times D_k \rightarrow D_{k+1}$  is a D-mapping, then

$$\lim_{k \rightarrow \infty} u^k = u^* \quad (5)$$

where  $u^* = G_k(u^*, u^*) \forall k$  (Dey, 1981).

## LOCALLY LINEARIZED D-MAPPING

Let us linearize (4) on  $D_k \times D_k$ , using first-order approximation of  $G_k(u^{k+1}, u^k)$  near  $(u^k, u^k)$ . Then,

$$u^{k+1} = G_k(u^k, u^k) + G'_k(u^{k+1} - u^k) \quad (6)$$

where  $G'_k$  is the Fréchet derivative of  $G_k$  on  $D_k \times D_k$ . Equation (6) may be expressed as

$$u^{k+1} = A_k u^k + b_k \quad (7)$$

where  $A_k = -(I - G'_k)^{-1}G'_k$ ,

$b_k = (I - G'_k)^{-1}G_k(u^k, u^k)$ . We have assumed that  $(I - G'_k)$  is invertible. Now we may prove a second theorem.

**Theorem 2.** If (i)  $|A_j - A^*| < E$ , where  $E$  is a matrix consisting of elements that are positive and arbitrarily small and (ii)  $|b_j - b^*| < \epsilon$ ,  $\epsilon$  is a vector consisting of elements that are positive and arbitrarily small, then (5) is true (convergence) if  $A_k$  is a D-matrix (Dey, 1983a).

**Theorem 3.** If  $G_k^i$  is a D-matrix, so is  $A_k$  (Dey, 1983a).

This principle may now be applied computationally.

#### PERTURBED FUNCTIONAL ITERATION

Let a nonlinear system be expressed as

$$\begin{aligned} u &= G_0(u) \\ u &\in D, \quad G_0: D \rightarrow D \end{aligned} \quad (8)$$

A Gauss-Seidel-type iteration for the solution may be expressed as

$$\begin{aligned} u^{k+1} &= G(u^{k+1}, u^k) \\ G: D \times D &\rightarrow D, \quad u^k \in D \quad \forall k \end{aligned} \quad (9)$$

A perturbed iterative scheme (Dey, 1977) may be expressed as (in the element form)

$$u_j^{k+1} = \omega_j^{k+1} + G_j(u^{k+1}, u^k) \quad (10)$$

where

$$\begin{aligned} \omega_j^{k+1} &= [G_j(G_j^{k+1,k}) - G_j^{k+1,k}] \\ &\times [1 - \partial_j G_j^{k+1,k}]^{-1} \end{aligned} \quad (11a)$$

$$G_j^{k+1,k} = G_j(u_1^{k+1}, \dots, u_{j-1}^{k+1}, u_j^k, \dots, u_J^k) \quad (11b)$$

$$\begin{aligned} G_j(G_j^{k+1,k}) &= G_j(u_1^{k+1}, \dots, u_{j-1}^{k+1}, \\ &\quad G_j^{k+1,k}, u_{j+1}^k, \dots, u_J^k) \end{aligned} \quad (11c)$$

$$\begin{aligned} \partial_j(G_j^{k+1,k}) &= [\partial G_j / \partial u_j] \\ &\quad u_1^{k+1}, u_2^{k+1}, \dots, \\ &\quad u_{j-1}^{k+1}, G_j^{k+1,k}, \\ &\quad u_{j+1}^k, \dots, u_J^k \end{aligned} \quad (11d)$$

The  $\omega_j$  term is a perturbation parameter which accelerates the rate of convergence of (9) and stabilizes the numerical algorithm.

It has been proved (Dey, 1981) that if  $G$  is a D-mapping on  $D_{k+1} \times D_k$ , a necessary and sufficient condition for convergence is

$$\lim_{k \rightarrow \infty} |\omega_j^k| = 0 \quad \forall j \quad (12)$$

Following Theorems 2 and 3 we may prove that if  $G_k^i$  is a D-matrix, then (12) is true (linearized sense). Recent results (Dey, 1983b) using local linearization indicates that if

$$\max_{j,m} |G_{jm}| \leq \beta/J \quad (13)$$

where  $G_{jm} = \partial G_j / \partial u_m$  and  $0 \leq \beta < 1$ ,  $G$  in (9) is a D-mapping. In order that (13) may be correct, certain input parameters for the system (e.g., mesh size and time-step) have to be chosen in special ways. If this cannot be found a convex-type operation may be defined as follows:

$$\hat{G}_j(u, u) = (1 - \alpha_j)u_j + \alpha_j G_{j1}(u, u) \quad (14)$$

Assuming  $G_{jj}(u, u) \neq 1$ , it has been found that  $\hat{G}$  is a D-mapping (locally linearized) for the following:

$$\begin{aligned} 1. \quad \alpha_j &= (-1)^p (1 - G_{jj})^{-1} \quad \text{if } G_{jm} = 0, \\ m \neq j \quad \text{and } p &= 0 \quad \text{if } G_{jj} < 1, \quad p = 1 \quad \text{if } G_{jj} > 1. \end{aligned}$$

$$2. \quad \alpha_j = \min_{\substack{1 \leq m \leq J \\ m \neq j}} \left( \frac{\beta/J}{|G_{jm}|}, (-1)^p \frac{(1 + \beta/J)}{1 - G_{jj}} \right)$$

$$\begin{aligned} \text{if } G_{jm} \neq 0, m \neq j \quad \text{and } p &= 0 \quad \text{if } G_{jj} < 1, \\ p &= 1 \quad \text{if } G_{jj} > 1. \end{aligned} \quad (15a)$$

where  $\beta$  is such that

$$1 > \beta \geq J \left( 1 + \frac{|1 - G_{jj}|}{|G_{jj}|} \right)^{-1} \quad (15b)$$

$$3. \quad \alpha_j = 1 \quad \text{if (13) is satisfied.}$$

The algorithm of perturbed functional iteration including a linearized convergence analysis may be briefly expressed as follows. At each iteration level, compute  $G_{jm}$ ,  $m = 1, 2, \dots, J$ . If (13) is satisfied, set  $\alpha_j = 1$ ; otherwise, compute  $\alpha_j$  using (15). If  $\alpha_j \neq 1$ , replace  $G_j$  by  $\hat{G}_j$ , as given by (14). Compute  $\omega_j$  using (11a)-(11d) and compute  $u_j$  at the new iteration level by (10). If (12) is satisfied at some iteration level, convergence is found; if  $G_{jj} = 1$ , the method fails.

In general, for a  $J \times J$  system the method requires (i)  $J^2 + J$  functionals to be computed for convergence analysis, (ii) partial linearization along the diagonal, and (iii) no Jacobians.

It has been proved analytically (Dey, 1977) that in the vicinity of the root, the method should display a superlinear rate of convergence.

## A DEGENERATE IMPLICIT CODE

Let a nonlinear model be represented by

$$du/dt = f(u), \quad u = (u_1 u_2 \dots u_J)^T \quad (16)$$

$u(0) = u_0$  (initial condition). Approximating (16) by a two-point backward-difference scheme, we get:

$$u_j^{n+1} = u_j^n + \Delta t f_j(u^{n+1}), \quad \Delta t = \text{time-step} \quad (17)$$

This nonlinear system may now be solved by the above method which forms a degenerate implicit code (since the one-step, matrix-inversion principle is not used for solution). If a convex-type operation of the form (14) is used, (16) becomes

$$du/dt = (I - \alpha)du/dt + \alpha f(u) \quad (18)$$

where  $\alpha = \text{diag}(\alpha_1, \alpha_2, \dots, \alpha_J)$  and  $I$  = the identity matrix. If (16) is a stiff system, (condition number of  $f'(u) \gg 1$ ), it generally requires  $\Delta t$  to be very small if a functional iteration of the form (9) is applied for solution. In (18),  $\alpha$  scales the elements in the Jacobian matrix  $f'(u)$ , and using (15) for  $\alpha_j$ 's mean (for a given  $\Delta t$ ), D-mapping is found so that perturbed functional iteration converges.

## APPLICATIONS

### Application 1.

(Bui, 1979)  $\dot{u}_1 = -10,004 u_1 + 10,000 u_2^4$ ;  $\dot{u}_2 = u_1 - u_2 - u_2^4$ ;  $u_1(0) = u_2(0) = 1$ . An approximate solution is  $u_2(t) = \{10,004 \exp(-3t)/[10,008 - 4 \exp(-3t)]\}^{1/3}$ .  $u_1(t) = (10,000/10,004)u_2^4$ . As  $t \rightarrow \infty$ ,  $u_1(t) \rightarrow 0$ ,  $u_2(t) \rightarrow 0$ . Linearizing this system near  $u_1(0), u_2(0)$ , the condition number is  $10^4$ .

Using linearized convergence analysis for the degenerate implicit code we got  $\Delta t \approx 10^{-5}$ , a sufficient condition for convergence if  $\alpha_j = 1$ . Introducing (18) and computing  $\alpha_j$ 's in a subroutine using (15) we used  $\Delta t = 10^{-3}$  to  $10^8$ ; correct results were found. No program interruption was caused. Details may be found in Dey (1983b).

### Application 2. Irradiation of Neutral Water

The model developed by Chatterjee and Magee and the analysis of its numerical solution are given in Chatterjee and others (1983). The equations and the rate constants are given in Table 1. For our present analysis we linearized the system and computed  $\alpha_j$ 's. Stiffness was measured by Strate (1983) at  $t = 0, 0.1, 1$ , and  $10$ . Condition numbers are, respectively,  $10^{19}$ ,  $10^{12}$ ,  $10^{12}$ ,  $10^{10}$  (approximately). This may be seen to be true in Fig. 1. This pattern of solution was ana-

lyzed by Chatterjee and Magee and was found to be valid. Here, difference equations were formed by approximating the derivatives by using the two-point trapezoidal rule. D-mappings were introduced, and time-accurate solutions were computed with  $\Delta t = 10^{-8}$ ,  $10^{-6}$ .

## CONCLUSION

Numerical solutions of stiff systems are generally obtained by using multistep implicit codes (Miranker, 1981) which require inversion of matrices obtained by computing Jacobians. This has been avoided in the technique explained here. However, the code is dependent on the Jacobians for its convergence analysis. Such a linearized analysis seems to be quite effective, and, in contrast with its nonlinear counterpart, the complete analysis can be done computationally. More applications are under consideration.

## ACKNOWLEDGMENT

I thank my 11-year-old son Charlie Dey for writing codes for the first problem in TRS-80 color computer.

## REFERENCES

- Bui, T. D. (1979). Some A-stable and L-stable methods for the numerical integration of stiff ordinary differential equations. *J. ACM*, 20, 3.
- Chatterjee, A., J. L. Magee, and S. K. Dey (1983). The role of homogeneous reactions in the radiolysis of water. *Radiat. Res.* (to appear).
- Dey, S. K. (1977). Perturbed iterative solution of nonlinear equations with applications to fluid dynamics. *J. Comp. Appl. Math.*, 3, 1, 15-30.
- Dey, S. K. (1981). Convergence analysis of a perturbed iterative scheme for solution of nonlinear systems. *Int. J. Math. Sci.*, 4, 2, 383-391.
- Dey, S. K. (1982). Applications of perturbed functional iterations to nonlinear stiff initial value chemical kinetic problems. *Proceedings of the International Conference on Stiff Computations*, Park City, Utah, April 12-14.
- Dey, S. K. (1983a). Numerical modeling of linearized D-mappings with applications to stiff systems. NASA TM (to be published).
- Dey, S. K. (1983b). Simulations of D-mappings in flow models. *Proceedings of Modeling and Simulation on Microcomputers*. 83 Society of Computer Simulation, San Diego, Calif., Jan. 27-30.
- Lomax, H. (1983). NASA Ames Research Center, private communication.
- Miranker, W. L. (1981). *Numerical Methods for Stiff Equations*. D. Reidel Publ. Co., Boston.
- Strate, G. (1983). NASA Ames Research Center, private communication.

Table 1 Differential Equations for Transient Species and  
Radiation Products in Irradiated Neutral Water

$\frac{d(H)}{dt}$	$= -2k_1(H)^2 - k_2(e^-_{Aq})(H) - k_5(H)(OH) + k_7(H_3O^+)(e^-_{Aq}) - k_9(H)(H_2O_2)$ $+ k_{12}(OH)(H_2) - k_{13}(HO_2)(H) - k_{17}(H)(O_2)$	+ 0.55 I
$\frac{d}{dt}(e^-_{Aq})$	$= -k_2(e^-_{Aq})(H) - 2k_3(e^-_{Aq})^2 - k_4(e^-_{Aq})(OH) - k_7(H_3O^+)(e^-_{Aq})$ $- k_{10}(e^-_{Aq})(H_2O_2) - k_{14}(e^-_{Aq})(O_2)$	+ 2.65 I
$\frac{d}{dt}(OH)$	$= -k_4(e^-_{Aq})(OH) - k_5(H)(OH) - 2k_6(OH)^2 + k_9(H)(H_2O_2) + k_{10}(e^-_{Aq})(H_2O_2)$ $- k_{11}(OH)(H_2O_2) - k_{12}(OH)(H_2) - k_{15}(HO_2)(OH)$	+ 2.70 I
$\frac{d}{dt}(H_3O^+)$	$= -k_7(H_3O^+)(e^-_{Aq}) - k_8(H_3O^+)(OH^-) - k_{18}(H_3O^+)(O_2^-) + k_{19}(H_2O) + k_{20}(HO_2)$	+ 2.65 I
$\frac{d}{dt}(H_2O)$	$= k_5(H)(OH) + k_8(H_3O^+)(OH^-) + k_9(H)(H_2O_2) + k_{11}(OH)(H_2O_2) + k_{12}(OH)(H_2)$ $+ k_{15}(HO_2)(OH) + k_{18}(H_3O^+)(O_2^-) - k_{19}(H_2O)$	- 4.10 I
$\frac{d}{dt}(H_2)$	$= k_1(H)^2 + k_2(e^-_{Aq})(H) + k_3(e^-_{Aq})^2 - k_{12}(OH)(H_2)$	+ 0.45 I
$\frac{d}{dt}(H_2O_2)$	$= k_6(OH)^2 - k_9(H)(H_2O_2) - k_{10}(e^-_{Aq})(H_2O_2) - k_{11}(OH)(H_2O_2) + k_{13}(HO_2)(H)$ $+ k_{16}(HO_2)^2$	+ 0.70 I
$\frac{d}{dt}(OH^-)$	$= k_2(e^-_{Aq})(H) + 2k_3(e^-_{Aq})^2 + k_4(e^-_{Aq})(OH) - k_8(H_3O^+)(OH^-)$ $+ k_{10}(e^-_{Aq})(H_2O_2) + k_{19}(H_2O)$	
$\frac{d}{dt}(HO_2)$	$= k_{11}(OH)(H_2O_2) - k_{13}(HO_2)(H) - k_{15}(HO_2)(OH) - 2k_{16}(HO_2)^2 + k_{17}(H)(O_2)$ $+ k_{18}(H_3O^+)(O_2^-) - k_{20}(HO_2)$	
$\frac{d}{dt}(O_2)$	$= -k_{14}(e^-_{Aq})(O_2) + k_{15}(HO_2)(OH) + k_{16}(HO_2)^2 - k_{17}(H)(O_2)$	
$\frac{d}{dt}(O_2^-)$	$= k_{14}(e^-_{Aq})(O_2) - k_{18}(H_3O^+)(O_2^-) + k_{20}(HO_2)$	

where

$k_1 = 10^{10}$	$k_2 = 2.5 \times 10^{10}$	$k_3 = 6 \times 10^9$	$k_4 = 3 \times 10^{10}$
$k_5 = 2.4 \times 10^{10}$	$k_6 = 4 \times 10^9$	$k_7 = 2.3 \times 10^{10}$	$k_9 = 3 \times 10^{10}$
$k_9 = 10^8$	$k_{10} = 1.2 \times 10^{10}$	$k_{11} = 5 \times 10^7$	$k_{12} = 6 \times 10^7$
$k_{13} = 10^{10}$	$k_{14} = 1.9 \times 10^{10}$	$k_{15} = 10^{10}$	$k_{16} = 2 \times 10^6$
$k_{17} = 10^{10}$	$k_{18} = 3 \times 10^{10}$	$k_{19} = 5.5 \times 10^{-6}$	$k_{20} = 10^6$




Fig. 1 Concentrations of species (Ex 2) vs. time for  $I = 6.667 \times 10^{-7}$  in the logarithmic scale up to  $t = 30$  sec. (Here steady state is reached for all the species.)

1 Report No NASA TM-84332	2 Government Accession No	3 Recipient's Catalog No	
4 Title and Subtitle NUMERICAL MODELING OF D-MAPPINGS WITH APPLICATIONS TO CHEMICAL KINETICS		5 Report Date March 1984	
		6 Performing Organization Code	
7 Author(s) S. K. Dey		8 Performing Organization Report No A-9256	
		10 Work Unit No T-6469	
9 Performing Organization Name and Address  Ames Research Center Moffett Field, CA 94035		11 Contract or Grant No	
		13 Type of Report and Period Covered Technical Memorandum	
12 Sponsoring Agency Name and Address  National Aeronautics and Space Administration Washington, DC 20546		14 Sponsoring Agency Code	
15 Supplementary Notes  Point of contact: H. Lomax, MS 202A-1, Ames Research Center, Moffett Field, CA 94035, (415)965-6607 or FTS 448-6607			
16 Abstract  Numerical modeling of D-mappings has been studied and applied to solving nonlinear stiff systems. These mappings have been locally linearized for convergence analysis, and some applications have been made to chemical kinetics.			
17 Key Words (Suggested by Author(s))  Nonlinear equations Numerical methods Biomedical		18 Distribution Statement  Unlimited  Subject category - 64	
19 Security Classif (of this report) Unclassified	20 Security Classif (of this page) Unclassified	21 No of Pages 8	22 Price* A02