

NASA-TM-85967 19840018455

FOR REFERENCE

NOT TO BE TAKEN FROM THIS ROOM

Detection of Shocked Atomic Gas in the Kleinmann-Low Nebula

M.W. Werner, M.K. Crawford,
R. Genzel, D.J. Hollenbach,
C.H. Townes, and
Dan M. Watson

LIBRARY COPY

JUN 12 1984

LANGLEY RESEARCH CENTER
LIBRARY, NASA
HAMPTON, VIRGINIA

May 1984

National Aeronautics and
Space Administration

NF00826

Detection of Shocked Atomic Gas in the Kleinmann-Low Nebula

M. W. Werner

D. J. Hollenbach, NASA Ames Research Center, Moffett Field, California

M. K. Crawford

R. Genzel, University of California, Berkeley, California

Dan M. Watson, California Institute of Technology, Pasadena, California



National Aeronautics and
Space Administration

Ames Research Center

Moffett Field, California 94035

N84-26523[#]

Detection of Shocked Atomic Gas in the Kleinmann-Low Nebula

M.W. Werner¹, M.K. Crawford², R. Genzel², D.J. Hollenbach¹,

C.H. Townes², and Dan M. Watson³

ABSTRACT

We have mapped the $63\mu\text{m } ^3P_1 \rightarrow ^3P_2$ fine structure line emission of neutral atomic oxygen at the center of the Orion nebula with a resolution of $30''$. There are three main emission peaks. One is associated with the region of strongest thermal radio continuum radiation close to the Trapezium cluster, and probably arises at the interface between the HII region and the dense Orion molecular cloud. The other two [OI] line emission peaks, associated with the Kleinmann-Low nebula, are similar in both distribution and velocity to those of the $2\mu\text{m } S(1)$ line of molecular hydrogen and of the high-velocity wings of rotational CO emission. The [OI] line profiles in the KL region are resolved at a resolution of 45 km s^{-1} and have an intrinsic width of about 50 km s^{-1} , while the line profile in the vicinity of the Trapezium is unresolved. We show that the [OI] emission from the KL nebula can be produced in the shocked gas associated with the mass outflows in this region and is an important coolant of the shocked gas.

Subject Headings: infrared: sources - infrared: spectra - nebulae: Orion nebula.

I. Introduction

The Kleinmann-Low (KL) nebula in Orion is the site of a violent mass outflow associated with the formation of massive stars (c.f., the review by Genzel and Downes 1982). Infrared emission lines from highly excited states of H_2 , OH, and CO have been detected from this region (cf., Gautier et al. 1976; Storey et al. 1981; Beck et al. 1982); this emission is probably excited in shocks occurring where dense shells driven by the mass outflow strike the ambient molecular cloud material. In this paper, we

¹ NASA Ames Research Center, Moffett Field, CA. 94035

² UC Berkeley, Dept. of Physics, Berkeley, CA. 94720

³ California Institute of Technology, Pasadena, CA. 91125

report mapping of the central 2.5' of the Orion-KL region with 30" resolution and show spectra taken at 45 km s⁻¹ resolution which together reveal emission in the atomic oxygen $^3P_1 \rightarrow ^3P_2$ ground state fine structure transition associated with the KL nebula. This transition is expected to be the dominant cooling transition for dense ($10^8 \geq n \geq 10^3$) atomic interstellar gas for temperatures between 100 and 5000 K.

II. Observations and Results

1) Observations

The observations were carried out with the 91cm telescope of the Kuiper Airborne Observatory in 1983 February and 1983 October, using the UC Berkeley tandem Fabry-Perot spectrometer described by Storey, Watson and Townes (1980). The beam had a FWHM of 30", with an effective area of a 38" diameter disk. In February 1983, the instrumental profile was Lorentzian with a FWHM of 170 km s⁻¹. A chopper spacing of 4' in azimuth was used (approximately east-west). Spectral scans were obtained at approximately forty points (typical integration time 1 minute per point) on a grid spaced by 20" in both right ascension and declination. Each spectral scan included the line and the adjacent continuum, so that maps of line and continuum intensity were produced simultaneously with good relative positioning. Relative positions across the map are accurate to 5". Absolute positions were obtained by assuming that the continuum peak is at R.A. = 5^h32^m46^s.8, Dec. = -05°24'25" (1950, see Wynn-Williams et al. 1984). Repeated measurements of the line intensity at a point near the map center agreed to $\pm 10\%$, and it is this reproducibility, rather than the signal-to-noise ratio, which is the limiting uncertainty in the quality of the map presented in Figure 1. Observations were made during two flights in October, 1983, at velocity resolutions of 60 and 45 km s⁻¹ towards several points on the map made in February; the 45 km s⁻¹ spectra are shown in Figure 2. The beam size was again 30", and the chopper spacing was 5' in azimuth, which was again approximately east-west. Radial velocities were calibrated by measuring the relative wavelength separation between the [OI] lines in Orion (rest wavelength 63.18372 μ m) and an H₂O line at 63.3236 μ m. The line fluxes in Figures 1 and 2 were calibrated by measurements of the line to continuum ratio toward the continuum peak. We assumed a flux density

of 7×10^4 Jy for the KL continuum in a $30''$ beam (scaling by 0.6 the flux densities given by Werner et al. 1976). The overall calibration uncertainty is estimated to be $\pm 25\%$.

2) Spatial Distribution

Figure 1 shows the spatial distribution of $63\mu\text{m}$ line and continuum flux over the region mapped, compared with other relevant data. The [OI] line emission shows a peak $10''$ SW of the Trapezium cluster and a double-peaked ridge near the KL nebula. The peak near the Trapezium is very similar in position and shape to the 5 GHz radio continuum emission. [OI] line emission in this region probably comes from dense ($n_{\text{H}} \sim 10^{4-5} \text{ cm}^{-3}$), warm ($T \sim$ a few hundred K) gas at the interface between the HII region and the molecular cloud. This interface region will be discussed in a forthcoming paper (Ellis and Werner, 1984; see also Jaffe and Pankonin, 1978). The [OI] peaks in the KL region are close to but somewhat interior to the emission lobes of hot molecular hydrogen (Figure 1b) and are also similar to the distribution of high-velocity rotational CO emission (cf. Erickson et al. 1982). The lower resolution data of Naylor et al. (1982) and Ellis and Werner (1984) also clearly show an extension of the $63\mu\text{m}$ emission from the Trapezium toward the BN-KL region but lack sufficient resolution to identify a separate peak.

3) Kinematics

Figure 2 shows [OI] spectra taken with a resolution of 45 km s^{-1} toward the radio peak west of the Trapezium, H_2 peaks 1 and 2 and the KL peak (the center of the $63\mu\text{m}$ continuum emission). The line toward the radio peak is unresolved; the line shape is identical with the instrumental profile. The LSR velocity of the line is $8.5 \pm 3 \text{ km s}^{-1}$ which is in good agreement with the velocity centroid of the 6300\AA line of [OI] at this position (Münch and Taylor, 1974). The line profiles near the KL nebula are resolved and the lines are significantly blueshifted. The line width at H_2 peak 1 is $67 \pm 5 \text{ km s}^{-1}$ FWHM, and the peak of the line is at $v_{\text{LSR}} = 0 \pm 3 \text{ km s}^{-1}$. The line widths at H_2 peak 2 and the KL peak are $61 \pm 5 \text{ km s}^{-1}$ and they are centered at $v_{\text{LSR}} = +3 \pm 3 \text{ km s}^{-1}$. Most of the difference between these line profiles and the unresolved line toward the radio peak is probably due to the influence of high velocity [OI] emission, blueshifted by about 20 to 50 km s^{-1} . The presence of strong blueshifted wings is also characteristic of the H_2 $2\mu\text{m}$ lines toward peaks 1 and 2 (Nadeau et al. 1982). The

intrinsic width of the [OI] line towards the KL region may be estimated, by comparing the line width observed at 45 km s^{-1} with that at 60 km s^{-1} resolution, to be about FWHM 50 km s^{-1} at H_2 peak 1 and 40 km s^{-1} toward the KL peak. These are lower limits to the width of the emission associated with the KL nebula itself since the profile probably includes a narrow component of line emission from the interface [OI] gas in front of KL.

III. Discussion

1) Properties of the [OI]- Emitting Gas in the KL-nebula

The [OI]-emitting gas in the KL nebula has a large velocity dispersion and its emission has a spatial distribution similar to that of the $2\mu\text{m}$ emission from hot hydrogen. It is, therefore, likely that the $63\mu\text{m}$ [OI] emission is produced in shocks due to the mass outflow from the center of the KL nebula. The [OI] line is of substantial importance to the gas cooling in the shocked region since its luminosity is of the same order of magnitude as the luminosities of the H_2 and CO lines. The peak surface brightness of the line near H_2 peak 1 is $3.3 \pm 1 \times 10^{-2} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1}$, after subtraction of a background of about $2.5 \times 10^{-2} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1}$ which may originate in the interface to the HII region. The total $63\mu\text{m}$ line luminosity from the shocked region is $50 \pm 25 L_\odot$, which is only about 3-4 times smaller than the total luminosity of all H_2 infrared lines ($\sim 200 L_\odot$, c.f., Scoville et al. 1982, Beckwith et al. 1983), or the luminosity of the far-IR CO lines ($150 L_\odot$). The minimum column density of [OI] gas required to produce the observed surface brightness is $4.5 \times 10^{17} \text{ cm}^{-2}$, assuming statistical equilibrium at high density ($n \geq 5 \times 10^5 \text{ cm}^{-3}$) and $T \gg 230 \text{ K}$. This is probably a good assumption for the parameters of the [OI] emitting gas in the KL nebula. The ratio of $63\mu\text{m}$ intensity to that of the second [OI] line at $145\mu\text{m}$ toward the center of the KL nebula is between 20 and 40 (Ellis and Werner, 1984). This range is consistent with optically thin emission from dense gas ($n_{\text{H}} \sim 10^{5 \pm 1} \text{ cm}^{-3}$) at high temperature ($T \sim 300$ to 1500 K) and is also characteristic of shocked gas. With a width of 50 km s^{-1} , the line has negligible opacity ($\tau_{63} \sim 10^{-2}$). Assuming all oxygen atoms are in the form of OI and a cosmic abundance of oxygen ($\text{O}/\text{H} = 6.6 \times 10^{-4}$), the lower limit to the column density of hydrogen nuclei in the [OI] emitting region is $7 \times 10^{20} \text{ cm}^{-2}$, which corresponds to a mass of $\sim 0.15 M_\odot$. This is

significantly larger than the amount of very hot hydrogen gas behind the molecular shock ($T \geq 2000$ K; $N_{\text{H}} \sim 5 \times 10^{19} \text{ cm}^{-2}$, c.f., Beckwith et al. 1983) but less than the hydrogen column density estimated for the medium temperature molecular gas in this region ($T \sim 500\text{--}1000$ K; $N_{\text{H}} \sim 2 \text{ to } 6 \times 10^{21} \text{ cm}^{-2}$, cf. Watson et al. 1984).

2) [OI] Emission from the Cloud Shock

The mass outflow in the KL nebula drives shocked, neutral gas into the ambient interstellar medium. As pointed out by Castor, McCray, and Weaver (1975), Hollenbach (1982), and Chernoff, Hollenbach, and McKee (1982, hereafter CHM), two separate shock fronts will exist under these circumstances. The "wind shock" occurs at the point where the stellar wind collides with the inner edge of the shocked, expanding material. A second shock, (the "cloud shock") occurs where the outer edge of the expanding material strikes the ambient molecular cloud. Draine and Roberge (1982) and CHM have modeled the cloud shock as a C-type shock in which the magnetic field effectively "cushions" the shock front. These models have successfully reproduced the intensities of the high excitation molecular lines of H_2 , OH and CO observed in the KL region. In such a shock, the [OI] $63\mu\text{m}$ line is predicted to be two or more orders of magnitude fainter than the integrated H_2 emission, because any preshock atomic oxygen is rapidly converted to H_2O in the hot post-shock region. By contrast, the present observations show that the $63\mu\text{m}$ line is within an order of magnitude as intense as the total H_2 emission. The column density of medium temperature ($T > 500$ K) C-shocked gas, however, is sufficient to produce the [OI] emission if $\sim 10\%$ of the oxygen could be maintained as [OI]. Thus, if our present understanding of the chemistry in this region is inadequate, it is possible that the [OI] emission could in fact be produced in the molecular shock.

3) The Wind Shock

Alternatively, the [OI] emission may come from the wind shock (Hollenbach, 1984; Draine and Roberge, 1982). If the outflow has a characteristic speed in excess of 100 km s^{-1} , it strikes the inner surface of the compressed shell at a speed in excess of 50 km s^{-1} . This is sufficient to dissociate and significantly ionize the wind material, which leads to formation of a J-type wind shock front with sharp density and temperature discontinuities and high post-shock temperatures ($T \gg 10^4$ K). As the gas

behind the shock front cools and recombines, it passes through a range of temperatures from approximately 5000 to 100 K in which [OI] $63\mu\text{m}$ emission is the dominant coolant for the largely atomic gas. Because the $63\mu\text{m}$ line is the dominant coolant in this temperature range, about 5000 k of energy ($k = 1.38 \times 10^{-16} \text{ erg K}^{-1}$) is radiated in the line for each nucleon which crosses the shock front. Thus the flux of material into the shocks, $n_{\text{H}} v \text{ (cm}^{-2} \text{ sec}^{-1}\text{)}$, is related to the surface brightness, I , of the $63\mu\text{m}$ line by:

$$I = 5000 \frac{k}{4\pi} n_{\text{H}} v \text{ erg s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1}, \quad (1).$$

where $n_{\text{H}} \text{ [cm}^{-3}\text{]}$ is the preshock hydrogen density in the wind, and v is the relative velocity between wind and shell. Detailed calculations over the range $40 \text{ km s}^{-1} < v < 200 \text{ km s}^{-1}$ and $10 \text{ cm}^{-3} < n_{\text{H}} < 10^{+5} \text{ cm}^{-3}$ show that the [OI] $63\mu\text{m}$ intensity lies within a factor of two of this linear relationship for $n_{\text{H}} v \leq 10^{12} \text{ cm}^{-2} \text{ s}^{-1}$ (Hollenbach, 1984). Assuming that the wind shock in the Orion-KL region falls within this parameter range, the observed [OI] $63\mu\text{m}$ intensity corresponds to a value of $n_{\text{H}} v = (6 \pm 2) \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1}$. In this limit, the total luminosity of the $63\mu\text{m}$ line gives a direct estimate of the mass loss rate \dot{M} from the central object and is about $7 L_{\odot}$ if $\dot{M} = 10^{-3} M_{\odot} \text{ y}^{-1}$. For the observed total [OI] luminosity of $50 \pm 25 L_{\odot}$, the corresponding mass loss rate is $(7 \pm 3.5) \times 10^{-3} M_{\odot} \text{ y}^{-1}$, which is comparable with or somewhat higher than other estimates of the mass loss rate in the KL region. If the [OI] emission arises predominantly in the wind shock, while the H_2 and CO infrared emission lines arise in the molecular shock, differences in the details of the line profiles and spatial distribution are expected. The [OI] data presented above are consistent with these expectations but do not exclude the possibility that the [OI] emission comes from the cloud shock. The [OI] $63\mu\text{m}$ emission line may be one of the few direct observational probes of J-type wind shocks in embedded infrared sources. If [OI] emission from the wind shock can be unambiguously identified, its intensity may be an important measure for determining mass loss rates in outflow sources.

Acknowledgements:

We thank the staff of the Kuiper Airborne Observatory for their excellent support. We are grateful to J.B. Lugten and H.B. Ellis for help with the observations and data reduction. This work was supported by NASA grant NAG2-208 (UCB) and by the Astrophysics Division of NASA Ames.

References

- Beck, S.C., Bloemhof, E.E., Serabyn, E., Townes, C.H., Tokunaga, A.T., Lacy, J.H., and Smith, H.A., 1982, *Ap. J. (Letters)*, **253**, L83.
- Beckwith, S., Evans, N.J., Gatley, I., Gull, G., and Russell, R.W., 1983, *Ap. J.*, **264**, 152.
- Castor, J.M., McCray, R., and Weaver, R., 1975, *Ap. J. (Letters)*, **200**, L207.
- Chernoff, D.F., Hollenbach, D.J., and McKee, C.F., 1982, *Ap. J. (Letters)*, **259**, L97.
- Draine, B.T., and Roberge, W.G., 1982, *Ap. J. (Letters)*, **259**, L91.
- Ellis, H.B., and Werner, M.W., 1984, in preparation.
- Erickson, N.R., Goldsmith, P.F., Snell, R.L., Berson, R.L., Huguenin, G.R., Ulich, B.L., and Lada, C.J., 1982, *Ap. J. (Letters)*, **261**, L103.
- Gautier, T.N., Fink, U., Treffers, R.R., and Larson, H.P., 1976, *Ap. J. (Letters)*, **207**, L129.
- Gautier, T.N., 1979, Ph.D. Thesis, University of Arizona.
- Genzel, R., and Downes, D., 1982, *Regions of Recent Star Formation*, ed. R.S. Roger and P.E. Dewdney (Dordrecht: Reidel), p.251.
- Hollenbach, D.J., 1982, *Ann. N.Y. Acad. Sci.*, **395**, 242.
- Hollenbach, D.J., 1984, in *Proceedings of the Protostars and Planets Symposium*, to be published in *Icarus*.
- Jaffe, D.T., and Pankonin, V., 1978, *Ap. J.*, **226**, 869.
- Johnston, K.J., Palmer, P., Wilson, T.L., and Bieging, J.H., 1983, *Ap. J. (Letters)*, **271**, L89.

Münch, G., and Taylor, K., 1974, *Ap. J. (Letters)* 192, L93.

Nadeau, D., Geballe, T.R., and Neugebauer, G., 1982, *Ap. J.*, **253**, 154.

Naylor, D.A., Emery, R., Fitton, B., Furniss, I., Jennings, R.E., and King, K.J., 1982, in *Regions of Recent Star Formation*, ed. R.S. Roger and P.E. Dewdney, (Dordrecht: Reidel), 73.

Scoville, N.Z., Hall, D.N.B., Kleinmann, S.G., and Ridgway, S.T., 1982, *Ap. J.*, **253**, 136.

Storey, J.W.V., Watson, D.M., and Townes, C.H., 1980, *Int. J. IR and MM Waves*, **1**, 15.

Storey, J.W.V., Watson, D.M., Townes, C.H., Haller, E.E., and Hansen, W.L., 1981, *Ap. J. (Letters)*, **248**, L109.

Werner, M.W., Gatley, I., Harper, D.A., Becklin, E.E., Loewenstein, R.F., Telesco, C.M., and Thronson, H.A., 1976, *Ap. J.*, **204**, 420.

Wynn-Williams, C.G., Genzel, R., Becklin, E.E., and Downes, D., 1984, *Ap. J.*, in press.

Figure Captions

Figure 1a (left): Contours of integrated $63\mu\text{m}$ [OI] line emission toward the Orion-KL region with a resolution of $30''$ (heavy lines). Contour units are $2.5 \times 10^{-17} \text{ W cm}^{-2}$ ($9.4 \times 10^{-3} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1}$). The dashed lines are contours of 5 GHz radio continuum emission at a resolution of $28''$ (from Johnston et al. 1983, contour unit 55 mJy per beam). The crosses mark the positions of the $63\mu\text{m}$ continuum peak (the KL-nebula, upper) and of $\Theta^1\text{C Ori}$ (lower), the brightest star in the Trapezium cluster.

Figure 1b (middle): Contours of integrated [OI] emission (heavy lines), superposed on contours of $\nu = 1 \rightarrow 0$ (S(1) emission from H_2 at $2\mu\text{m}$ (Gautier 1979), $30''$ beam, contour unit $6 \times 10^{-4} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1}$). The northwestern, stronger peak is H_2 peak 1, the southeastern peak is H_2 peak 2.

Figure 1c (right): Contours of $63\mu\text{m}$ continuum emission (heavy contours 1, 2, 4, 6, 8, 10, 12, 14, 16, 18 times $3.8 \times 10^3 \text{ Jy}$ or $1.6 \times 10^{-17} \text{ W cm}^{-2}$), superposed on the 5 GHz radio continuum map.

Figure 2: Spectra of $63\mu\text{m}$ [OI] emission toward several positions in the Orion-Nebula region: $10''$ west of Trapezium (bottom), H_2 peak 2 (second from bottom), $63\mu\text{m}$ continuum peak (third from bottom) and H_2 peak 1 (top). Beam size is $30''$ and the instrumental profile is a Lorentzian of FWHM 45 km s^{-1} (indicated by thin lines in the upper three spectra). The bottom line is unresolved. The curvature in the continuum on the left and right of the [OI] line is due to telluric H_2O absorption. The vertical scale is given separately on the y-axis for each spectrum. Typical integration times are 6 minutes per spectrum.

Author's Addresses

M.W. Werner and D.J. Hollenbach

NASA Ames Research Center

Moffett Field, CA. 94035

M.K. Crawford, R. Genzel, and C.H. Townes

University of California, Dept. of Physics,
Berkeley, CA. 94720

Dan M. Watson

Downs Laboratory of Physics

California Institute of Technology

Pasadena, CA. 91125




Fig. 1




Fig. 2

1. Report No. NASA TM-85967		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle DETECTION OF SHOCKED ATOMIC GAS IN THE KLEINMANN-LOW NEBULA				5. Report Date June 1984	
				6. Performing Organization Code ATP	
7. Author(s) M. W. Werner,* M. K. Crawford, [†] R. Genzel, [†] D. J. Hollenback,* C. H. Townes, [†] and F. M. Watson [‡]				8. Performing Organization Report No. A-9774	
				10. Work Unit No. T5726	
9. Performing Organization Name and Address *Ames Research Center, Moffett Field, CA 94035 [†] University of California, Department of Physics, Berkeley, CA 94720, [‡] California Institute of Technology, Pasadena, CA 91125				11. Contract or Grant No.	
				13. Type of Report and Period Covered Technical Memorandum	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, DC 20546				14. Sponsoring Agency Code 352-02-03	
15. Supplementary Notes Preprint Series #16. Supported by NASA grants. Point of Contact: L. C. Haughney, Moffett Field, CA 94035, M/S 211-12, (415) 965-5339, FTS 448-5339					
16. Abstract We have mapped the $63 \mu\text{m } ^3\text{P}_1 - ^3\text{P}_2$ fine structure line emission of neutral atomic oxygen at the center of the Orion nebula with a resolution of 30". There are three main emission peaks. One is associated with the region of strongest thermal radio continuum radiation close to the Trapezium cluster, and probably arises at the interface between the HII region and the dense Orion molecular cloud. The other two [OI] line emission peaks, associated with the Kleinmann-Low nebula, are similar in both distribution and velocity to those of the $2 \mu\text{m S}(1)$ line of molecular hydrogen and of the high-velocity wings of rotational CO emission. The [OI] line profiles in the KL region are resolved at a resolution of 45 km s^{-1} and have an intrinsic width of about 50 km s^{-1} , while the line profile in the vicinity of the Trapezium is unresolved. We show that the [OI] emission from the KL nebula can be produced in the shocked gas associated with the mass outflows in this region and is an important coolant of the shocked gas.					
17. Key Words (Suggested by Author(s)) Astronomy: Infrared: Sources Infrared: Spectra Nebulae: Orion nebula				18. Distribution Statement Unlimited Subject category 89	
19. Security Classif. (of this report) Unclassified		20. Security Classif. (of this page) Unclassified		21. No. of Pages 14	
				22. Price* A03	

End of Document