

ON A POSSIBILITY OF INELASTICITY PARTIAL COEFFICIENT  $K_{\gamma}$   
 DETERMINATION IN  $\pi C$  AND  $\pi Pb$  INTERACTIONS AT  $10^{14}$  EV  
 (EXPERIMENT "PAMIR" I)

Pamir Collaboration:

Borisov A.S., Cherdyntseva K.V., Guseva Z.M., Denisova V.G.,  
 Dunaevsky A.M., Kanevskaya E.A., Maximenko V.M., Nam R.A.,  
 Pashkov S.V., Puchkov V.S., Slavatinsky S.A., Zimin M.V.,  
 Smirnova M.D., Smorodin Yu.A., Zelevinskaya N.G. and Zhdanov  
 -P.N. Lebedev Physical Institute, Academy of Sciences of  
 USSR, Moscow

Mukhammadshin R.A., Nedel'ko O.E., Nikolaeva L.P.,  
 -Institute for Nuclear Research, Academy of Sciences of  
 USSR, Moscow

Amineva T.P., Baradzei L.T., Ivanenko I.P., Iljina N.P.,  
 Lazareva T., Managadze A.K., Mikhailova I.A., Murzina E.A.,  
 Pomelova E.I., Popova E.G., Rakobolskaya I.V., Roganova T.M.,  
 Ryabova N.G., Sveshnikova L.G.

-Institute of Nuclear Physics, Moscow University, Moscow  
 Cananov S.D., Chadranyan E.Kh., Khisanishvili L.A., Ladarina N  
 Leptukh G.G., Roinishvili N.N., Svanidze M.S.

-Institute of Physics, Georgian Academy of Sciences,  
 Tbilisi

Asimov Z.A., Bobodjanov I.B., Gubar' N.E., Gulov Yu.A.,  
 Normuratov F., Shoboronov Kh.

-Institute of Physics and Engineering, Tajik Academy of  
 Sciences, Dushanbe

Azimov S.A., Khalilov D.A., Mulladjanov E.G., Nuritdinov Kh.,  
 Talipov D.A., Yuldashbaev T.S.

-Institute of physics and Engineering, Uzbek. Academy of  
 Sciences, Tashkent

Bakhtigireev S.E., Dobrotin N.A., Emeljanov Yu.A., Zaitseva E  
 -Institute of High Energy Physics, Kazakh. Academy of  
 Sciences, Alma-Ata

Bielawska H., Linke M., Malinowski J., Novicka A., Podgorzak,  
 Tomaszewski A., Wlodarczyk Z., Wrotniak A.

-Institute of Physics, University of Lodz, Poland

and Sharvadze Z.SH.\*, Institute of Physics, Georgian Academy  
 of Sciences, Tbilisi

The investigation of hadron-nuclear interactions in  
 "Pamir" experiment is carried out by means of X-ray emulsion  
 chambers of two types: carbon (C) and lead (Pb) ones [1].

While comparing the results from the chambers of both  
 types [2] it was found a discrepancy in  $\langle n_h \rangle$  and  $\langle E_h^{(0)} \rangle$   
 values.

---

\* This article only

Here  $\langle n_h \rangle$  is an average hadron multiplicity normalized to one  $\gamma$ -h-family fitted the following selection criterion A:  $\sum E_Y \geq 100$  Tev,  $n_h \geq 0$ ,  $E_k^{(v)} \geq 4$  Tev;  $\langle E_k^{(v)} R \rangle$  is a space-energetic characteristics of hadrons in families fitted the criterion B:  $\sum E_Y \geq 100$  Tev,  $n_h > 3$ ,  $E_k^{(v)} \geq 4$  Tev;  $E_k^{(v)}$  is the energy registered in hadron block.  $n_h$  values are corrected on interaction probability in each chamber.

Tables I and II represent our experimental results, where  $\nu$  and  $\chi$  parametres defined as

$$\nu = \frac{\langle n_h \rangle_{Pb}}{\langle n_h \rangle_c} \quad (1) \quad ; \quad \chi = \frac{\langle E_k^{(v)} R \rangle_{Pb}}{\langle E_k^{(v)} R \rangle_c} \quad (2)$$

Table I

	$N_{fam}$	$\langle n_h \rangle$	$\beta - 1$
C	169	$3.0 \pm 0.2$	$1.0 \pm 0.1$
Pb	41	$4.2 \pm 0.3$	$1.1 \pm 0.1$
$\nu$	-	$1.4 \pm 0.2$	

Table II

	$N_{fam}$	$E_k^{(v)} R_{num}$
C	33	$350 \pm 30$
Pb	22	$455 \pm 70$
$\chi$	-	$1,3 \pm 0,2$

$N_{fam}$  is a number of families in C and Pb chambers fitted the above mentioned criteria,  $\beta - 1$  is a slope of integral energy spectrum of hadrons from families selected by A-criterion

In the present paper we connect the observed discrepancy between  $\langle n_h \rangle$  and  $\langle E_k^{(v)} R \rangle$  in C and Pb chambers with the difference in values of effective coefficients of energy transfer to the soft component  $K_{eff}$  for C and Pb chambers. The following considerations can be a ground of this suggestion:

1) It is known that a probability of hadron registration (in case of power like spectrum) is equal to  $\langle K_Y^{\beta-1} \rangle$ . Thus, the ratio of multiplicities

$$\nu = \frac{\langle n_h \rangle_{Pb}}{\langle n_h \rangle_c} = \frac{\langle K_Y^{\beta-1} \rangle_{Pb}}{\langle K_Y^{\beta-1} \rangle_c}$$

As  $\beta - 1 \approx 1$  (see Table I) we have:

$$\nu = \langle K_Y \rangle_{Pb} / \langle K_Y \rangle_c \quad (3)$$

2)  $\langle E_k^{(v)} R \rangle$  value can be written down as  $\langle K_Y \rangle \langle E_k R \rangle$  (in case of  $K_Y$  independent of  $E_k$ ). Hence:  $\chi = \langle K_Y \rangle_{Pb} / \langle K_Y \rangle_c$  (4).

To test this considerations some simulations on nuclear electromagnetic cascades in the atmosphere were made on the basis of fireball scaling model (S-model) [3]. Hadrons passage through the chamber was imitated in the simulations. For each hadron the value of  $K_Y$  coefficient was found by means of  $f(K_Y)$  distribution function taken in the form of  $f(K_Y) \sim K_Y^\alpha \cdot \exp(-K_Y/\beta)$  [4]. The parametres  $\alpha$  and  $\beta$  define the momenta of this distribution. Two series of simulation with various  $\langle K_Y \rangle$  values were carried out. One of them with  $\langle K_Y \rangle = 0.17$  imitated the situation in C-chamber, the other with  $\langle K_Y \rangle = 0.30$  in Pb-chamber. The latter value is greater than

the real one in Pb-chamber, was chosen especially to emphasize the effects connected with  $K_\gamma$  variations. The ratio of simulation values of  $\langle K_\gamma \rangle$  is equal to

$$\langle K_\gamma \rangle_{\text{Pb}} / \langle K_\gamma \rangle_{\text{C}} = 1.76 \quad (5)$$

A hadron was considered registered one in case of  $E_h^\gamma = K_\gamma E_h \geq 4$  Tev.

For  $\langle E_h^\gamma R \rangle$  determination we selected the families with number of registered hadrons  $n_h > 3$ . Table III represents the obtained results.

Table III

	$N_{\text{fam}}$	$E_h K$ Tev mm
C	126	285 $\pm$ 20
Pb	275	510 $\pm$ 15
$\chi$		1,80 $\pm$ 0.1

Table IV

	$N_{\text{fam}}$	$\langle n \rangle$	$\beta - 1$
C	505	2.3 $\pm$ 0.1	0.9 $\pm$ 0.05
Pb	505	3.9 $\pm$ 0.1	1.0 $\pm$ 0.05
$\gamma$		1.7 $\pm$ 0.1	

As it is seen from Table III

$$\chi = \langle E_h^{(\gamma)} R \rangle_{\text{Pb}} / \langle E_h^{(\gamma)} R \rangle_{\text{C}} = 1.80 \pm 0.10 \quad (6)$$

and that is in good agreement with value (5).

The incoming to the installation hadron families with  $n_h > 3$  and  $\sum E_h \geq 250$  Tev were selected to calculate  $\gamma$  parametre. The last condition provides registration of the same events in both sets of simulation (in "both chambers").

Table IV represents the obtained values of mean multiplicities and energy spectra slope.  $\gamma$  value which is equal to  $1.70 \pm 0.10$  is also in agreement with the given ratio of  $\langle K_\gamma \rangle$  (5). Thus, the experimental estimation of  $\chi$  and  $\gamma$  values in the two types of chambers gives an opportunity to get a ratio of  $\langle K_\gamma \rangle$  coefficients for lead and carbon nuclei.

Returning to preliminary experimental evaluations of  $\chi$  and  $\gamma$  parametres (Table 1) one can see that the coefficients ratio turned to be roughly equal to  $1.2 \pm 1.5$ . This ratio corresponds to  $K_\gamma$  coefficients for pion-nuclear interactions as the majority of hadrons in families are  $\pi$ -mesons.

Experiments carried out in the low energy range yield that  $K_{\pi^0}$  partial inelastic coefficient dependence on atomic number of target nucleus is described as  $K_{\pi^0} \sim A^\alpha$ . Using this dependence and  $\langle K_\gamma \rangle_{\text{Pb}} / \langle K_\gamma \rangle_{\text{C}}$  value we can estimate  $\alpha$ -index at energy  $\sim 10^4$  ev. Here we should bear in mind, that  $K_\gamma^{\text{Pb}} \approx K_{\pi^0}^{\text{Pb}}$ , while  $K_\gamma^{\text{C}}$  can considerably differ from  $K_{\pi^0}^{\text{C}}$ . However, it was shown in simulations [5]  $K_\gamma$  value in carbon chamber in  $E_h^{(\gamma)}$  energy range from 3 up to 30 Tev is similar to  $K_{\pi^0}$  value in pion-nucleon interactions. In such case

$$\frac{\langle K_\gamma \rangle_{\text{Pb}}}{\langle K_\gamma \rangle_{\text{C}}} \approx A^{\alpha} \text{Pb} \quad \text{and} \quad \alpha \approx 0.06 \pm 0.02. \quad \text{This value is}$$

not in contradiction to  $\alpha$ -measureings at lower energies

and confirms a possibility of the suggested method.

To determine  $\alpha$ -value with 20% accuracy it is necessary to have approximately 100 families with  $\sum E_{\gamma} \geq 100$  Tev in each chamber as well as to make precise simulations of connection between partial  $K_{\pi^0}$  and effective  $K_{\gamma}$  inelasticity coefficients in chambers of both types.

#### References

1. Trudy FIAN, v.154, 1984, (in Russian)
2. Pamir Collaboration. Proc. of International Symposium of Cosmic Rays and Particle Physics, Tokyo, P292, 1984
3. Wrotniak I.A. Zess.Nauk.Univers.Lodz, vol.60, p.73, 1977
4. Malinowski A. et.al. Pamir Coll. Workshop, p.49, UL, Lodz, 1980
5. Wlodarczyk Z. et.al. Proc. of International Symp. of Cosmic Rays and Particle Physics, Tokyo, p 451, 1984