

510-46
N89-25808⁴⁸
187409⁷¹

MEASUREMENTS OF H₂O IN THE TERRESTRIAL MESOSPHERE AND IMPLICATIONS FOR EXTRA-TERRESTRIAL SOURCES; J.J. Olivero, Department of Meteorology, The Pennsylvania State University, University Park, PA 16802

Water vapor is an important minor constituent in the terrestrial mesosphere (approx. 50 km to 85 km or 10⁰ mbar to 10⁻² mbar) and nearly coincident D-region ionosphere (~60 km to 90 km). Its photolysis products control ozone above the ozone mixing ratio peak; they are radiatively active, especially in the infrared; water sublimates to cause polar mesospheric clouds and possibly other suggested aerosol phenomena; the molecules cluster to form massive ions which dominate the chemistry of the lower ionosphere; and, finally, as the principal hydrogen compound below the mesosphere, it controls the global hydrogen budget and H escape rate.

During the last decade several measurement techniques have been employed to observe H₂O at these atmospheric levels; these are presented in Figure 1 with symbols given in Table 1.

Five of these data sets were produced by ground-based microwave radiometry (B1, SC, B2, OL, TS). These results were all contributed by a collaboration between The Pennsylvania State University and the Naval Research Laboratory, joined recently by the Jet Propulsion Laboratory. The observations cover all seasons but are limited to a small range of latitudes - 34N to 24N - over North America.

The second group (AK, GR, OE, RD, SN, WA) involve various aircraft and rocket techniques and were observed over a much wider range of latitudes.

Ground-based microwave measurements at mid-latitudes strongly support a very dry mesopause-lower thermosphere. Other measurements, [6] and [7], can be interpreted as being consistent with a high altitude or external source of H₂O. The latter process was not seriously considered until very recently-- [12], [13] and [14]--however, these new hypotheses have profound implications for the upper atmospheres of all the planets [15].

The remainder of the talk will discuss the apparent qualitative limits that H₂O measurements to date place upon a downward H₂O flux above the mesosphere.




Figure 1. A collection of measurements of mesospheric water vapor from the past decade; symbols referenced in Table 1.

Table 1.

KEY TO SYMBOLS IN FIGURE 1

<u>SYMBOL</u>	<u>REFERENCE</u>	<u>COMMENT</u>
B1	Bevilacqua, et al. (1983)..... [1]	GROUND-BASED
SC	Schwartz, et al. (1983)..... [2]	MICROWAVE
B2	Bevilacqua, et al. (1985)..... [3]	RADIOMETRY
OL	Olivero, et al. (1986)..... [4]	MID-LATITUDES
TS	Tsou (1986)..... [5]	
AK	Arnold and Krankowsky (1979).. [6]	MISCELLANEOUS
GR	Grossmann, et al. (1982)..... [7]	TECHNIQUES
OE	O'Brien and Evans (1981)..... [8]	-
RO	Rogers, et al. (1977)..... [9]	RANGE OF
SN	Swider and Narcisi (1975)..... [10]	NON-POLAR
WA	Waters, et al. (1980)..... [11]	LATITUDES

REFERENCES

1. Bevilacqua, R. M., Olivero, J. J., Schwartz, P. R., Gibbins, C. J., Bologna, J. M. and Thacker, D. L. (1983) J. Geophys. Res., 88, 8523-8534.
2. Schwartz, P. R., Croskey, C. L., Bevilacqua, R. M. and Olivero, J. J. (1983) Nature, 305, 294-205.
3. Bevilacqua, R. M., Wilson, W. J., Ricketts, W. B., Schwartz, P. R. and Howard, R. J. (1985) Geophys. Res. Lett., 12, 397-400.
4. Olivero, J. J., Tsou, J. J., Croskey, C. L., Hale, L. C. and Joiner, R. G. (1986) Geophys. Res. Lett., 13, 197-200.
5. Tsou, J. J. (1986) Microwave Radiometric Measurements of Mesospheric Water Vapor: Ground-Based Observations in Both Solar Absorption and Atmospheric Emission Modes, Ph.D. dissertation in Meteorology, The Pennsylvania State University, University Park, PA, 185 pp.
6. Arnold, F. and Krankowsky, D. (1977) Nature, 268, 218-219.
7. Grossman, K. U., Frings, W., Offerman, D. O., Joos, W., Andre, L. and Kopp, E. (1982) Concentrations of some minor neutral constituents in the mesosphere and lower thermosphere above northern Europe, XXIV COSPAR Asembly, Ottawa.
8. O'Brien, R. S. and Evans, W. F. J. (1981) J. Geophys. Res., 86, 12101-12107.
9. Rogers, J. W., Stair, A. J., Jr., Degges, J. C., Wyatt, C. L. and Baker, D. J. (1977) Geophys. Res. Lett., 4, 366-368.
10. Swider, W. and Narcisi, R. S. (1975) J. Geophys. Res., 80, 655-664.
11. Waters, J. W., Gustincic, J. J., Swanson, P. N. and Kerr, A. R. (1980) Measurements of upper atmospheric H₂O emission at 183 GHz, in Atmospheric Water Vapor, edited by A. Deepak, T. O. Wilkerson, and L. H. Ruhnke, Academic Press, New York, 229-240.
12. Frank, L. A., Sigwarth, J. B. and Craven J. D. (1986) Geophys. Res. Lett., 13, 307-310.

13. Frank, L. A., Sigwarth, J. B. and Craven, J. D. (1986) Geophys. Res. Lett., 13, 307-310.
14. Dubin, M. (1986) Noctilucent clouds-the problems of origin, American Geophysical Union - Spring Meeting, Baltimore.
15. Donahue, T. M. (1986) Geophys. Res. Lett., 13, 555-558.