N91-71533

Workshop on Early Detection of Stratospheric Changes. March 5 - 7, 1986

SUMMARY OF THE RESULTS FROM ATMOS C.B. Farmer

The ATMOS experiment was flown for the first time in April, 1985, as part of the Spacelab 3 payload. The instrument is a Michelson interferometer operating in the absorption (solar occultation) mode, covering the 2 to 16 microns wavelength range (600 to 4500 cm⁻¹) at a resolution of 0.01 cm⁻¹. During the Spacelab 3 mission ATMOS recorded 12 sunset and 4 sunrise occultations, on April 30th, and May 1st and 2nd. The sunsets occurred at latitudes between 25 and 33° N and the sunrises between 47 and 49° S.

The reduction and analysis of the data has been carried out with the ATMOS computer system, a dedicated facility at JPL with remote terminals at several coinvestigator sites in the US and in Europe. The analysis is currently proceeding towards the final determination of vertical profiles of:

- 1. Density, temperature and pressure;
 - 2. Concentration of:

ì

- (i) minor gases
- (ii) nitrogen trace species
- (iii) halogen source gases
- (iv) halogen products
- (v) other trace gases;
- 3. Winds in the mesosphere and lower thermosphere;
- 4. Isotopic abundances.

In addition to these, useful upper limits (i.e. at or below the concentration levels predicted by current photochemical models) are being determined for other key species such as ClO, HOCl, HO₂, and H_2O_2 .

An atlas of high signal-to-noise coadded spectra of the sun and of the upper atmosphere, at tangent altitude intervals of approximately two scale heights, is being prepared for publication later this year.

The Table summarises the altitude ranges over which the profiles of concentration can be retrieved for each of the molecular constituents identified in the ATMOS spectra thus far. In general, the precision of the retrieved values of mixing ratio is about 10%, and can be better than 5% for constituents which show a large number of spectral features (for example, 0₃, CO_2 , N₂O, HCl, HNO₃ etc.). The systematic uncertainties (excluding molecular spectral parameters) are estimated to be between 10 and 20%, depending on the spectral region involved. It is expected that these uncertainties will be reduced considerably as the refinement of the analyses proceeds. Nevertheless, the precision achieved at the present time is sufficient to reveal longitudinal variability (e.g. H₂O, O₃), hemispheric (or diurnal) differences, and vertical structure. The analysis of the first data set has necessitated a careful verification and redefinition of the microwindows assigned to the species retrievals. In turn the ATMOS spectra are proving to be of great value in clarifying deficiencies in the basic molecular spectral parameters.

Constituent	Altitude Range (km)	Comments/Isotopes
N_2 CO_2 CO H_2O O_3 N_2O CH_4	< $18 - 35$ 5 - 140 5 - 120 5 - 85 < $10 - 90$ 5 - 65 5 - 80	Pressure at the tangent altitude T,P,p; Cl2, Cl3, Ol6, Ol7, Ol8 Cl2, Cl3, Ol6, Ol7, Ol8 HDO between 5 and 35 km O ₃ (16,16,18) < 20 to 42 km Cl2, Cl3
NO NO2 N2O5 HNO3 HNO4	< 18 - 140 < 18 - 50 20 - 38 18 - 50 18 - 40	Max. 1.6 x 10 ⁻⁹ at 32 km (sunrise) Max. 3 x 10 ⁻¹⁰ at 28 km (sunset)
CF ₄ CC1 ₄ F11, 12, 22 CH ₃ C1 CH ₃ CC1 ₃ HC1 HF COF ₂ C10N0 ₂	5 - 40 5 - 20 5 - 35 18 - 35 12 - 60 < 15 - 55 18 - 40 18 - 35	Several bands observed Several Q branches 3×10^{-9} above 45 km 6×10^{-10} above 45 km 2 bands; max. ~ 1.3 x 10^{-10} near 32 km 5 bands; max. ~ 1.5 x 10^{-9} near 30 km
осs нсn с ₂ н ₂ с ₂ н ₆	< 18 - 20 < 18 - 30 5 - 14 5 - 10	2 bands observed

TABLE 1 MOLECULAR SPECIES IDENTIFIED IN THE ATMOS SPECTRA

February 1986

<u>م</u>

-2-

· ...

• •