

NASA Conference Publication 3146

Nutritional Requirements for Space Station *Freedom* Crews

Helen W. Lane
*NASA Lyndon B. Johnson Space Center
Houston, Texas*

Barbara L. Rice
and Christine F. Wogan
*KRUG Life Sciences, Inc.
Houston, Texas*

Report of a panel discussion sponsored by
the Biomedical Operations and Research Branch,
Medical Sciences Division, Lyndon B. Johnson
Space Center, Houston, Texas, and held at
the Lunar and Planetary Institute
Houston, Texas
February 4–5, 1991

NASA

National Aeronautics and
Space Administration
Office of Management
Scientific and Technical
Information Program

1992



Table of Contents

Conference Participants v

Introduction 1

Nutritional Recommendations 1

Assessment of Nutritional Status: Crewmembers 3

Assessment of Nutritional Status: Food Analysis 4

Suggestions for Future Research 4

Summary 6

References 6

Appendix
 Conference Agenda

NUTRITIONAL REQUIREMENTS FOR SPACE STATION *FREEDOM* CREWS

A Panel Discussion held February 4-5, 1991
at the Lunar and Planetary Institute
Sponsored by the Biomedical Operations and Research Branch
NASA - Johnson Space Center, Houston, TX

PANELISTS

Chairs

Donald McCormick, Ph.D.
Department of Biochemistry
Emory School of Medicine

Sherwood L. Gorbach, M.D.
Department of Community Health
Tufts University School of Medicine

Lindsay H. Allen, Ph.D.
Department of Nutritional Sciences
University of Connecticut

Alfred Merrill, Ph.D.
Department of Biochemistry
Emory School of Medicine

Lynn Bailey, Ph.D.
Food Science and Human Nutrition
University of Florida at Gainesville

James Olson, Ph.D.
Dept of Biochemistry and Biophysics
Iowa State University

Janet Greger, Ph.D.
Department of Nutritional Sciences
University of Wisconsin

Dale Schoeller, Ph.D.
Clinical Nutrition Research Unit
University of Chicago

Scott Grundy, M.D.
Center of Human Nutrition
U.T. SW Medical Center

John Suttie, Ph.D.
Department of Biochemistry
University of Wisconsin

Robert Jacob, Ph.D.
Department of Biochemistry
USDA Western Nutrition Research Ctr

Connie Weaver, Ph.D.
Department of Foods and Nutrition
Purdue University

Janet King, Ph.D.
Department of Nutritional Sciences
University of California

Robert Wolfe, Ph.D.
Department of Metabolism
U. T. Medical Branch

Roy Martin, Ph.D.
Department of Foods and Nutrition
University of Georgia

Conference Organizers

Helen W, Lane, Ph.D.
Biomedical Operations and Research Branch
NASA - Johnson Space Center

Barbara L. Rice, R.D., L.D.
KRUG Life Sciences, Inc.
Johnson Space Center

Technical Editor
Christine F. Wogan
KRUG Life Sciences, Inc.
Johnson Space Center

OTHER PARTICIPANTS

Clarence Alfrey, M.D.
Methodist Hospital
Houston, TX

Charles Bourland, Ph.D.
Man-Systems Division
NASA-Johnson Space Center

Nitza M. Cintron, Ph.D.
Biomedical Operations and Research Branch
NASA-Johnson Space Center

Theda Driscoll
Methodist Hospital
Houston TX

G. William Fortner, Ph.D.
KRUG Life Sciences
Houston, TX

Mary Ann Frey, Ph.D.
NASA Headquarters
Washington, DC

Kimberly Glaus-Late
Lockheed
Houston, TX

John Greenleaf, Ph.D.
Life Science Division
NASA-Ames Research Center

Susan Hill
Lockheed
Houston, TX

Carolyn Leach Huntoon, Ph.D.
Space and Life Sciences Directorate
NASA-Johnson Space Center

Vickie Kloeris
Shuttle Food Systems
NASA-Johnson Space Center

Jane McAvin
Lockheed
Houston, TX

Jeannie Nillen
KRUG Life Sciences
Houston, TX

Robert W. Phillips, D.V.M., Ph.D.
NASA-Johnson Space Center

Victor Schneider, M.D.
University of Texas
Health Sciences Center
Houston, TX

M. Rhea Seddon, M.D.
Astronaut Office
NASA-Johnson Space Center

Robert D. Taylor, Ph.D.
KRUG Life Sciences
Houston, TX

Timothy T. White
Space and Life Sciences Directorate
NASA-Johnson Space Center

Troy Whitehurst
Boeing Aerospace
Houston, TX

T. C. Yang, Ph.D.
Biomedical Operations and Research Branch
NASA-Johnson Space Center

Gloria Zerkus
Lockheed
Houston, TX

Introduction

Current evidence strongly supports the role of nutrition in maintaining the health and optimal performance of astronauts during space flight and return to Earth. Significant research, with provision of valid data, is essential to improve understanding of how space flight affects human nutritional requirements. This workshop was convened to review the present state of knowledge of space nutrition, and to solicit recommendations from experts in the field of nutritional science. The panel was presented with an overview of physiological issues associated with space flight, and reviewed the food systems for the Space Shuttle and Space Station *Freedom* (see appended agenda). The panel then offered recommendations to the nutritional support plan for 90- to 180-day *Freedom* missions. These recommendations are presented here in four parts: nutritional recommendations; assessment of nutritional status (crewmembers) and nutrient content (diet); suggestions for future research; and summary.

Nutritional Recommendations

Dietary composition is important for the Astronaut Corps both with respect to their performance during space flight and after landing. The dietary guidelines reviewed by the Surgeon General [1] and the National Research Council [2] should be used as the starting point; these guidelines can be modified on the basis of data from previous space flights and from bed rest studies. The chosen menus for each 14-day cycle should reflect these nutritional requirements. Means of monitoring diet and gustatory changes should be included in the space station hardware. Specific requirements are as follows:

Fluid. Water intake should be adequate to reduce the incidence of kidney stones; a tentative guideline could be 1 mL per kcal of consumed energy.

Calories. Calorie consumption should be sufficient to maintain weight and body composition, with continuous monitoring during space flight. A 70 kg man exercising 1 to 2 hours per day is expected to require about 3,000 kcal/day.

Protein. It is difficult to recommend a protein-intake level because of the calcium and muscle loss observed during space flight. A high protein intake may slow muscle loss, but increases calcium loss. The panel therefore recommended adherence to the RDA of 12 to 15% protein, with further study of this nutrient. The type of protein should be balanced between animal and plant sources in a ratio of approximately 60:40 to facilitate meeting the requirements for other nutrients.

Fat. This nutrient should constitute between 30 and 35% of total calories, with a distribution among polyunsaturated/monounsaturated/saturated fats of 1:1.5-2:1. This relatively high percentage is recommended based on consideration of (1) the usual dietary pattern of astronauts, (2) palatability and satiety, (3) caloric density, and (4) concern that higher carbohydrate and protein levels could influence calcium loss.

Carbohydrate. By difference, the carbohydrates should provide approximately 50% of the total calories, and should be derived from foods that emphasize complex carbohydrates. Simple sugars should constitute no more than 10% of the total carbohydrate intake. The current consumption of 10 to 15 g of dietary fiber/day may be adequate; because of practical considerations, however, this point warrants further study.

Calcium. Bone and calcium losses are problematic, but it is not clear that calcium consumption of >800 to 1,000 mg/day will be beneficial. The source of calcium can be natural foods or calcium citrate. Those individuals who do not voluntarily consume >800 mg/d should take supplements. To augment bone retention, food supplies should be fortified to provide RDA levels of vitamin D and fluoride.

Vitamins. Crewmembers who will fly missions of 90 to 180 days should be provided with RDA levels of vitamins. Because stress levels are expected to be high before and during flight, and vitamin C turnover correspondingly high, the higher of the two RDA levels for vitamin C (100 mg/day) should be used. Vitamins A and B12, in contrast, are stored by the liver in sufficient quantities to maintain necessary levels, although B12 levels should be monitored in vegetarian astronauts. Consumption and maintenance of adequate levels of thiamine and vitamin K are important because of thiamine's role in high calorie intake and increased exercise levels, and vitamin K's association with calcium and bone metabolism.

Trace elements. As to mineral content, food supplied should also provide the RDA levels of zinc, selenium, and iodine, as well as safe and adequate levels of copper, manganese, fluoride, and iron. Because space flight has been associated with decreased erythropoiesis and increased serum ferritin concentrations, the use of iron supplements should be prohibited, and maximum intake held at the male RDA (10 mg of Fe per day).

Although many micronutrients can be provided by vitamin-mineral supplements, the panel recommended that food be the primary source of nutrients, with supplements to be used only when absolutely necessary. Natural foods contain other essential nonnutritive substances such as fiber and carotenoids. Consumption of natural foods also provides a sense of psychological well-being that will be important during long space missions. In light of potential skeletal changes during long missions, the intake of low-fat dairy products is to be encouraged whenever possible. Dairy products include multiple micronutrients such as vitamins A and D, riboflavin, zinc, and calcium. Because body vitamin D stores will become depleted in the absence of sunlight, dietary vitamin D levels may require an increase, or perhaps a means of ultraviolet irradiation of crewmembers should be considered. Provision and consumption of fresh fruits and vegetables should also be encouraged.

Assessment of Nutritional Status: Crewmembers

Establishing a baseline nutritional status before long-term space flights will be necessary in order to quantify the effects of microgravity, particularly in light of limited opportunities for in-flight assessment. Preflight dietary habits may well affect both adaptation to space flight and recovery upon return. The panel suggested offering nutritional consultations to astronauts in an effort to encourage compliance through education. Such interviews might include discussion of operational health issues, such as nutrition's role in in-flight stresses such as bone loss and exposure to radiation.

Nutritional status should absolutely be assessed before and after flight, and if at all possible during flight. A means of measuring body mass in flight must be available. In addition to interviews and other educational opportunities, tests can be performed on blood, urine, and fecal samples, perhaps in conjunction with Medical Operations collections. Such tests would include:

From blood samples:

- RBC and plasma folate concentrations
- ascorbate concentration
- Vit E, retinoids and carotenoids (performed together on HPLC)
- whole blood superoxide dismutase
- thrombotic potential (for Vit K)*
- erythrocyte glutathione reductase activity (for riboflavin)
- erythrocyte transketolase activity (for thiamine)
- plasma pyridoxal phosphate concentration (for Vit B6)
- 25-hydroxyvitamin D (Calcifediol)
- osteocalcin*
- iron status assays

From urine samples:

- iodine
- gamma-carboxyglutamate (GLA)*
- ascorbate
- flavins
- 4-pyridoxic acid (B6)

From fecal samples:

- occult blood

*Given the possible influence of vitamin K status on calcium metabolism and/or osteoporosis, osteocalcin (B6P) and urinary GLA should be monitored before (during if possible) and after flight.

It is especially important to exercise better control in recording food intake in order to ensure adequate consumption of micronutrients. Methods of monitoring actual intake easily and unobtrusively need to be developed.

Assessment of Nutritional Status: Food Analysis

The amount and type of some vitamins in foods can be estimated from existing food composition tables; however, direct analysis of Space Station *Freedom* diets should be performed for levels of folate, ascorbic acid (vitamin C), thiamine (B1), pyridoxine (B6), riboflavin (B2), and vitamins E and K. In general, compositional table data for these compounds is incomplete; furthermore, levels of these vitamins will vary with food-processing techniques and storage conditions. Some compounds (folate, pyridoxine, and riboflavin) are light sensitive; ascorbate and vitamin E are susceptible to oxidation. Direct analyses should also include the concentrations of thiamine and vitamin K in foods, and their stability under the proposed storage conditions.

Direct analysis of zinc, copper, iodine, and selenium may also be required, as the food-composition table data for these minerals are inadequate for the foods to be provided in space.

In view of observations that serum ferritin levels increase with age, the use of iron supplements should be discouraged, and iron intake should be monitored carefully.

The planned use of iodine as an antimicrobial agent in the water supply mandates consideration of iodine intake from water as well as food, with the concern that increased iodine levels may affect thyroid function. The use of recycled water for consumption should be considered from a nutritional standpoint as well, because of the probable presence of calcium and other trace elements in the potable water supply.

Suggestions for Future Research

Maintaining lean body mass in space. Quantifying lean body mass in space, and understanding its role in maintaining strength, endurance, and optimal performance, is complicated by the lack of "hard" data on protein intake and exercise. As a first step, protein intake should be quantified, and then these values associated with total food intake, type and duration of exercise, net protein synthesis, and urinary calcium excretion. Accurate recording of exercise regimens (resistance vs. aerobic exercise, physical workloads, and length and frequency of exercise periods) must be instituted, as well as recording when meals are consumed relative to the exercise period. Other useful information can be provided by measuring mediators of protein catabolic response. Although this study will require data from human subjects, these data could be supplemented by studying rats that consume varying levels of protein before and after space flight. In addition, a more precise measure is needed to determine the number of calories required to maintain lean body mass and avoid deposition of fat.

Maintaining the skeleton in space. In order to achieve a better understanding of the mechanisms of skeletal loss in space, ideal intake (and form) of calcium, phosphorus, vitamin D, fluoride and zinc should be determined, as well as the ideal type and duration of exercise in terms of its effects on total body calcium, bone density, calcium kinetics, and hormonal response.

Preflight nutritional status. The role of preflight nutrition, including fluid intake, high protein intake, and glycogen loading, should be evaluated, in particular in relation to the amelioration of motion sickness symptoms.

Intake during flight. Food and fluid intake, and duration and intensity of exercise, must be quantified in flight. The described shift in food preferences in space should be explored.

Postflight recovery. This period should be extended to 3 to 6 months after landing, and more detail collected on recovery of skeletal mass in relation to all possible mediating factors, and recovery of muscle mass and function.

Micronutrients. The effect of microgravity on micronutrient utilization and turnover must be determined. In particular, micronutrients known to be involved in erythropoiesis (folate, iron, vitamin B12, and copper), and those known or presumed to be involved in bone stability (vitamins D, K, and A, zinc, fluoride, and manganese) should be assessed.

Although the panel recognized that some of these important nutritional issues can be addressed only by a mission specifically devoted to nutritional research, they emphasized that some baseline information can and should be collected during Space Shuttle flights in order to better define nutritional requirements for longer space missions.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June 1992	3. REPORT TYPE AND DATES COVERED Conference Publication		
4. TITLE AND SUBTITLE Nutritional Requirements for Space Station <i>Freedom</i> Crews			5. FUNDING NUMBERS	
6. AUTHOR(S) Helen W. Lane, Barbara L. Rice, and Christine F. Wogan				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biomedical Operations and Research Branch National Aeronautics and Space Administration Lyndon B. Johnson Space Center Houston, TX 77058			8. PERFORMING ORGANIZATION REPORT NUMBER S-672	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, D.C. 20546-001			10. SPONSORING / MONITORING AGENCY REPORT NUMBER NASA CP-3146	
11. SUPPLEMENTARY NOTES Helen W. Lane: Lyndon B. Johnson Space Center, Houston, TX; Barbara L. Rice and Christine F. Wogan: KRUG Life Sciences, Inc., Houston, TX.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Unclassified - Unlimited Subject Category 54			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This CP reports the results of a meeting convened in Houston, Texas on February 4-5, 1991. The purpose of this panel, sponsored by the Biomedical Operations and Research Branch, was to set preliminary nutritional requirements for crewmembers flying 90- to 180-day missions on Space Station <i>Freedom</i> . NASA was strongly encouraged to provide nutritional education to the Astronaut Corps, with emphasis on how nutritional status affects performance during and after space flight. Specific recommendations included providing crewmembers with in-flight feedback on nutritional intake, weight, and strength, and incorporating issues of energy intake, body weight, body composition, strength, and protein intake in the flight medicine program. Exercise must be considered an integral part of any plan to maintain nutritional status, especially those modes that stress the skeleton and maintain body weight. Nutrient intake, amount of exercise, and drugs ingested must be recorded daily; high priority should be given to development of fully automated record systems that minimize astronauts' effort. A system of nutritional supplements should be developed to provide a method of reducing intake deficits that become apparent. Finally, postflight monitoring should include bone density, muscle mass and function, and iron status at 3 and 6 months after landing.				
14. SUBJECT TERMS Nutrition, Energy Requirements, Food Analysis, Micronutrients			15. NUMBER OF PAGES 12	
			16. PRICE CODE A03	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT	