



Effect of Carbon Dioxide Enrichment on Radish Production Using Nutrient Film Technique (NFT)

C. L. Mackowiak, L. M. Ruffe, N. C. Yorio and R. M. Wheeler

(NASA-TM-107198) EFFECT OF CARBON
DIOXIDE ENRICHMENT ON RADISH
PRODUCTION USING NUTRIENT FILM
TECHNIQUE (NFT) (NASA) 17 p

N94-27411

Unclas

⁶³
~~51~~/51 0000297



Effect of Carbon Dioxide Enrichment on Radish Production Using Nutrient Film Technique (NFT)

*C. L. Mackowiak, L. M. Ruffe, N. C. Yorio, The Bionetics Corporation
R. M. Wheeler, Biomedical Operations and Research Office
John F. Kennedy Space Center, NASA*



TABLE OF CONTENTS

SECTION	PAGE
TABLE OF CONTENTS.....	i
ABSTRACT	ii
LIST OF TABLES AND FIGURES	iii
ACKNOWLEDGEMENTS	iv
PRODUCT DISCLAIMER	v
INTRODUCTION	1
MATERIALS AND METHODS	1
Environmental conditions	2
Culture	2
RESULTS AND DISCUSSION	3
Biomass.....	3
Water uptake and pH control	4
CONCLUSIONS	5
TABLES	6
FIGURES	7
REFERENCES	9

ABSTRACT

Radish plants (*Raphanus sativus* L. cvs. Cherry Belle, Giant White Globe, and Early Scarlet Globe) were grown in four different CO₂ enriched environments, 0.04, 0.10, 0.50, and 1.00 kPa (400, 1000, 5000, 10000 ppm). Cultivar responses to CO₂ treatments varied, where cv. Cherry Belle showed no significant response to CO₂ enrichment, cv. Giant White Globe was moderately affected, and Early Scarlet Globe was strongly affected. Enrichment at 0.10 kPa led to greater root dry matter (DM) than 1.00 kPa for cv. Giant White Globe, whereas 0.10 kPa produced greater storage root, shoot, and root DM than 1.00 kPa for cv. Early Scarlet Globe. The data suggest that 1.00 kPa CO₂ may be detrimental to the growth of certain radish cultivars. Root:shoot ratios tended to increase with increasing CO₂ concentration. Water use efficiency (g biomass/kg H₂O) increased with increasing CO₂ enrichment, up to 0.5 kPa but then declined at the 1.00 kPa treatment. The total nitric acid used to maintain nutrient solution pH was lowest at the 1.00 kPa treatment as well, suggesting a decreased demand of nutrients by the plants at the highest CO₂ level.

LIST OF TABLES AND FIGURES

TABLE		PAGE
1	Inorganic composition of the nutrient solutions	6
2	Yield parameters of radish as influenced by CO ₂	6
FIGURE		
1	Effect of CO ₂ on radish plant dry mass	7
2	Effect of CO ₂ on water and nitric acid requirements	8

ACKNOWLEDGEMENTS

We would like to thank Elise Blaese and the Student Life Science Training Program (SLSTP) for screening several radish cultivars, from which three of the best yielding were selected for this study.

PRODUCT DISCLAIMER

This report, in whole or part, may not be used to state or imply the endorsement by NASA employees of a commercial product, process or service, or used in any other manner that might mislead.



INTRODUCTION

Interest has been generated by NASA in developing Controlled Ecological Life Support Systems (CELSS) for long-duration space missions and lunar habitations. On a smaller scale, salad crops have been given attention as a means to provide fresh food aboard Space Station Freedom and also to aid in the understanding of plant production in space. The project "Salad Machine" includes lettuce as one of the crops that will be considered for production on the Space Station. However, information pertaining to its culture in hydroponic systems is limited. Yield response surfaces for temperature, radiation (intensity and quality), and photoperiod have been investigated (Cracker et al., 1983; Inada et al., 1991), but CO₂ responses have not been thoroughly defined.

The atmosphere of human inhabited space craft tend to have CO₂ concentrations far above normal Earth ambient (0.035 kPa) levels. It is not unusual to report levels as great as 0.60 kPa in Space Shuttle and Mir quarters (D. Weigrefe, Bionetics Corp., 1993, personal communication; V. Polyakov, Cosmonaut, 1991, personal communication). Most CO₂ enrichment studies involving plants have focused on concentrations that are twice ambient levels. Carbon dioxide enrichment on radish production has shown that 0.06 - 0.07 kPa CO₂ resulted in improved yields over ambient concentrations (Sionit et al., 1982; Idso et al., 1988). However, effects of much higher CO₂ levels, that might occur in space habitats, have not been studied. It has been suggested that CO₂ enrichment much beyond 0.10 kPa (1000 ppm) may cause growth inhibition, resulting in yield reductions in some plants, e.g. cucumber (Peet, 1986). However, CO₂ levels at 0.50 kPa have revealed only minor effects in two soybean cultivars, with growth at 0.50 kPa being ~ 10% less than that at 0.10 kPa (Wheeler et al., 1993). Thus CO₂ effects may be species and cultivar dependent.

The objectives of this study were to 1) grow three cultivars of radish, with four levels of CO₂ enrichment, and 2) determine CO₂ effects on radish production and water use efficiencies (WUE) under moderate light conditions.

MATERIALS AND METHODS

A series of four 16-day radish (*Raphanus sativus* L.) studies was conducted in a 1.8 m x 2.4 m walk-in growth chamber (EGC Inc., Chagrin Falls, OH). Each study was set up as a particular CO₂ treatment, and the CO₂ treatments were not repeated. Within each treatment cvs. Cherry Belle (CB), Giant White Globe (GWG), and Early Scarlet Globe (ESG) were grown at a density of 24 plants per square meter. Analysis of variance (ANOVA) was used to analyze

the cultivars individually. Mean separation was performed using Scheffe's test (SAS version 6.07, SAS Institute Inc. Cary, NC).

Environmental Conditions:

The four CO₂ treatments were 0.04, 0.10, 0.50, and 1.00 kPa (400, 1000, 5000, 10000 ppm). The 0.04 kPa treatment was chosen over normal ambient (0.035 kPa) to avoid periodic CO₂ increases resulting from human activity around the growth chamber. Levels of CO₂ in the chamber were monitored with an infrared gas analyzer (ANARAD Inc., Santa Barbara, CA) and controlled within 5% of setpoint with a dedicated computer system. Levels of CO₂ were 5% (coefficients of variation or CV) of the set point.

Irradiance in the chamber was provided with thirty 96-inch VHO Vita-Lite fluorescent lamps (Duro-Test Inc., North Bergen, NJ). Photosynthetic photon flux (PPF) levels for all tests averaged $230 \mu\text{mol m}^{-2} \text{s}^{-1} \pm 3\% \text{ CV}$ using a 20 h light/4 h dark photoperiod. Temperatures for all tests averaged $23.9 \pm 0.3 \text{ C}$ light and $23.0 \pm 0.4 \text{ C}$ dark. Relative humidities were constant during the light/dark cycle and averaged $69 \pm 1\%$.

Culture:

Plants were grown in eight trapezoidal-shaped, plastic culture trays (Mackowiak et al., 1989). Two kinds of plant support (tray) inserts were tested. The first insert was the same type used for soybean and wheat production studies (Mackowiak, et al., 1989) and consisted of rows of juxtapositioned black on white plastic in which was positioned a double piece of Nyltex™ plastic fabric for wicking nutrient solution at the bottom of the tray to the seed. The second insert was similar but had holes containing the plastic strips rather than rows. This design had been used in lettuce production studies (Prince and Knott, 1989). Since insert style did not affect plant performance, it was not factored into the results and will not be discussed further.

The nutrient solution was composed of a modified half-strength Hoagland mix, and was monitored and replenished daily with a nutrient concentrate to maintain the solution electrical conductivity at $0.12 \pm 0.02 \text{ S m}^{-1}$ (Table 1). Elemental concentrations were monitored weekly using ICP spectroscopy. Nutrient solution pH was controlled automatically to 5.8 units with additions of 0.39 M nitric acid. Water was manually added to the reservoir each day to maintain a constant volume (80 L).

Twelve dry seeds were planted directly onto inserts for each tray. White acrylic germination covers were used to maintain high humidity around the seedlings for the first four days. At 7 days after planting (DAP), extra plants were removed, resulting in six plants per

tray. Both GWG and CB had three replicates (3 trays) and due to chamber space, ESG had two replicates.

Plants were harvested at 16 DAP, when leaf and storage root fresh mass (FM), and leaf area were recorded. Leaf, storage root, and fibrous root tissue were dried at 70 C for 48 h prior to measuring dry mass (DM).

RESULTS AND DISCUSSION

Biomass:

Results showed that CO₂ effects on yield were cultivar dependent for radish, just as had been reported for soybean (Wheeler et al., 1993). Cultivar CB showed no significant differences in growth parameters under the tested CO₂ levels (Table 2). Similar findings have been seen with cv. Sativas when testing CO₂ concentrations up to 1.0 kPa, but there were negative effects on growth at 3.0 kPa (Pfeufer and Krug, 1984).

There appeared to be a trend of greater leaf area at 0.10 kPa CO₂ enrichment for GWG and ESG (Fig 1a), although the differences were not significant (Table 2). Moderate enrichment has been shown to increase leaf area in many other crops; however, this was mainly associated with more extensive branching and less from individual leaf size (Lawlor and Mitchell, 1991).

There was some effect of CO₂ on root dry mass for both GWG and ESG, where CO₂ at 0.10 kPa produced significantly more root dry mass than at 1.00 kPa (Table 2). Soybean root biomass increased by 143% when CO₂ was raised from 0.035 to 0.67 kPa (Rogers et al., 1992). There also appeared to be a CO₂ effect with cv. ESG relative to storage root fresh mass and dry mass, where the 0.10 kPa treatment was significantly greater than the 1.00 kPa CO₂ treatment (Table 2). GWG and CB showed no significant differences for those variables. Enrichment to 0.07 kPa CO₂ has been reported to increase root:shoot ratios for radish, along with other root crops (Idso et al., 1988). We found similar results at 0.1 kPa CO₂; however, root:shoot ratios declined at the 1.00 kPa treatment (Fig 1b). The most productive treatments for storage root yields, namely, GWG at 0.50 kPa and ESG at 0.10 kPa CO₂, also had relatively high storage root standard errors. In fact, the coefficient of variation for GWG and ESG was about 38% greater than it was for CB, across CO₂ treatments. The greater variation may suggest that there was a higher degree of heterozygosity in ESG and GWG seed, which was accentuated by CO₂ enrichments.

Partitioning into the storage root per se, was not affected by CO₂ concentration; however, there were definite cultivar differences, where CB had consistently greater harvest index values than either GWG or ESG (Fig 1c). Although GWG had the lowest harvest index, it usually

had the largest storage roots and total biomass. Subsequently, it also had the greatest crop growth rate ($\text{g m}^{-2} \text{d}^{-1}$) and conversion efficiency per plant. Since CB had lower leaf area values, it may be a candidate for denser plantings. It also seems to be less affected by varying CO_2 concentrations (Table 2), leading to very predictable yields under the range of CO_2 and growing conditions used in these studies. It would not be surprising to learn that the variation in cultivar responses to CO_2 enrichment may also occur with other environmental parameters, so cultivar selection would depend on the the culture environment.

In a CELSS, great importance would be placed on efficient use of mass, volume, and energy. Sionit et al. (1982) have found that a combination of high PPF ($1200 \mu\text{mol m}^{-2} \text{s}^{-1}$) with CO_2 enrichment (0.07 kPa) improved radish yields, when using a 14 h light/10 h dark photoperiod. However, others have shown that the best light conversion efficiency for radish was with a 20 h light/04 h dark photoperiod combined with a thermoperiod that varied 5 C between light and dark cycles (Inada et al., 1991). In other words, plants that can produce highest yields in high PPF environments may not be the most energy (photon) efficient.

Water and pH Control:

Since all cultivars shared the same nutrient delivery system, water use (total evapotranspiration) represented all the cultivars combined, for any single CO_2 treatment. Water uptake was normalized to the following units: g total biomass/kg H_2O (i.e. water use efficiency or WUE). Increasing CO_2 above 0.04 kPa increased WUE, but WUE decreased at levels above 0.10 kPa CO_2 (Fig 2a). These results were similar to those reported for soybean (Wheeler et al., 1993); however, in this study, the relationship between total plant dry mass and WUE were highly correlated ($r^2 = 0.97$).

When NO_3^- is taken up by the plant, the release of HCO_3^- causes an increase in solution pH (Schon, 1992). Measurement of the nitric acid used for pH control in this study, resulted in an indirect method for determining NO_3^- uptake. Nitric acid use was normalized to the following units: g total biomass/mmmole of acid (i.e. acid use efficiency or AUE). AUE was greatest at 0.10 kPa CO_2 , which corresponded with the greatest total biomass (Fig 2b). Unlike WUE, AUE did not correlate well with total biomass ($r^2 = 0.55$), but it correlated well with leaf area ($r^2 = 0.82$). Nitrate has been found to accumulate in leaf vacuoles of leafy vegetables, such as lettuce, for subsequent use as a storage pool (Blom-Zandstra, 1989). Our results suggest that there may have been luxury consumption of nitrate, which was stored in leaf vacuoles, leading to lower AUEs in treatments having smaller leaf areas.

CONCLUSIONS

Effects of CO₂ on radish production in a hydroponic system were cultivar dependent, where CO₂ enrichment to 0.10 - 0.50 kPa improved yields and tended to increase leaf area for GWG and ESG. Carbon dioxide enrichment above 0.04 kPa had no effect on yields or on any of the other harvest parameters for CB. CB had a harvest index twice the value of GWG, but GWG produced up to 70% more storage root than CB, over all CO₂ treatments. Early Scarlet Globe ranked between the other two cultivars for a majority of harvest parameters. Total chamber WUE was similar to results found in previous CO₂ enrichment studies using soybean, where efficiency was greatest at moderately elevated CO₂ levels and was poorest at the lowest and highest CO₂ levels (i.e. 0.04 and 1.00 kPa). Acid Use Efficiency was greatest at 0.10 kPa CO₂ and lowest at 1.00 kPa, which may be related to inefficient use of nitrate at the highest CO₂ concentration. Based on yields, WUE, and AUE, it appears that CO₂ enrichment to 1.0 kPa was detrimental to GWG and ESG. The available literature suggests that other environmental parameters, i.e., photoperiod, irradiance, and thermoperiod, might also be adjusted, along with CO₂, for optimal radish production.

Table 1. Inorganic composition of the nutrient solution and replenishment solutions per liter.

SALT	NUTRIENT SOLUTION CONCENTRATION	REPLENISHMENT CONCENTRATION
	(mM)	(mM)
Ca(NO ₃) ₂	2.5	9.0
KNO ₃	2.5	17.5
KH ₂ PO ₄	0.5	5.0
MgSO ₄	1.0	8.0
	(μM)	(μM)
Fe-EDTA	50.00	90.00
H ₃ BO ₃	4.75	24.00
MnCl ₂	3.70	18.50
ZnSO ₄	0.52	3.20
CuSO ₄	0.64	2.60
(NH ₄) ₆ Mo ₇ O ₂₄	0.02	0.05

Table 2. Effect of CO₂ concentration on the yield parameters of three radish cultivars.

Cultivar*	CO ₂	SR**	SR	TOP	ROOT	LA**
	(kPa)	(g FM/plant)	(g DM/plant)	(g DM/plant)	(g DM/plant)	(cm)
CB	0.04	8.39 (0.53)	0.52 (0.04)	0.43 (0.04)	0.05 (0.01)	85 (7)
CB	0.10	8.51 (0.88)	0.56 (0.06)	0.47 (0.04)	0.05 (0.01)	84 (8)
CB	0.50	8.32 (0.50)	0.52 (0.03)	0.43 (0.04)	0.05 (0.01)	84 (6)
CB	1.00	8.20 (0.77)	0.51 (0.04)	0.44 (0.04)	0.03 (0.00)	76 (6)
<i>Significance</i>		<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
GWG	0.04	10.8 (1.37)	0.61 (0.07)	1.66 (0.15)	0.22 (0.02)	366 (33)
GWG	0.10	11.9 (1.40)	0.72 (0.08)	2.04 (0.12)	0.28 (0.03)	400 (24)
GWG	0.50	14.2 (2.02)	0.80 (0.10)	1.94 (0.14)	0.25 (0.03)	373 (21)
GWG	1.00	10.2 (1.64)	0.57 (0.09)	1.49 (0.11)	0.18 (0.02)	301 (20)
<i>Significance</i>		<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>p < 0.05</i>	<i>ns</i>
ESG	0.04	9.5 (0.65)	0.59 (0.04)	0.76 (0.06)	0.09 (0.01)	160 (14)
ESG	0.10	12.7 (1.72)	0.87 (0.11)	1.12 (0.12)	0.12 (0.01)	195 (22)
ESG	0.50	11.5 (1.11)	0.77 (0.07)	0.84 (0.07)	0.07 (0.01)	174 (13)
ESG	1.00	7.5 (0.96)	0.47 (0.06)	0.67 (0.07)	0.05 (0.01)	144 (11)
<i>Significance</i>		<i>p < 0.05</i>	<i>p < 0.05</i>	<i>p < 0.05</i>	<i>p < 0.05</i>	<i>ns</i>

* CB = Cherry Belle, GWG = Giant White Globe, and ESG = Early Scarlet Globe.

** SR = storage root; LA = Leaf area. Standard errors are in parentheses.




Fig 1a. Effect of CO₂ on radish leaf area. CB = Cherry Belle; GWG = Giant White Globe; ESG = Early Scarlet Globe.




Fig 1b. Effect of CO₂ on radish root:shoot ratio. Cultivars same as in 1a.




Fig 1c. Effect of CO₂ on radish harvest index. Cultivars same as in 1a.




Fig. 2a. Effect of CO₂ on water use efficiencies (WUE). Since all cultivars shared the same nutrient tank, the values represent the combined evapotranspiration for all cultivars.




Fig. 2b. Effect of CO₂ on acid use efficiencies (AUE) from pH control. Since all cultivars shared the same nutrient tank, the values represent the combined use by all cultivars.

REFERENCES

- Blom-Zandstra, M. 1989. Nitrate accumulation in vegetables and its relationship to quality. *Ann. Appl. Biol.* 115:553-561.
- Cracker, L.E. and M. Seibert. 1982. Light energy requirement for controlled environment growth of lettuce and radish. *ASAE* 25(1):214-216.
- Idso, S.B., B.A. Kimball, and J.R. Mauney. 1988. Effects of atmospheric CO₂ enrichment on root:shoot ratios of carrot, radish, cotton, and soybean. *Agric. Eco. Env.* 21:293-299.
- Inada, K. and Y. Yabumoto. 1991. Effects of light quality, daylength, and periodic temperature variation on the growth of lettuce and radish plants. *Jap. J. Crop Sci.* 58:689-694.
- Lawlor, D.W. and R.A.C. Mitchell. 1991. The effects of increasing CO₂ on crop photosynthesis and productivity: a review of field studies. *Plant Cell and Env.* 14:807-818.
- Mackowiak, C.L., L.P. Owens, C.R. Hinkle, and R.P. Prince. 1989. Continuous hydroponic wheat production using a recirculating system. *NASA Tech. Mem.* 102784.
- Pfeufer, B. and H. Krug. 1984. Effects of high CO₂ concentrations on vegetables. *Acta Hort.* 162:37-44.
- Prince, R.P., W.M Knott. 1989. CELSS breadboard project at the Kennedy Space Center. In: D.W. Ming and D.L. Henninger (eds.) *Lunar Base Agriculture: Soils for Plant Growth*. *Agron. Soc. Amer.*, pp. 155-163.
- Peet, M.M. 1986. Acclimation to high CO₂ in monoecious cucumbers. I. Vegetative and reproductive growth. *Plant Physiol.* 80:59-62.
- Rogers, H.H. C. M. Peterson, J.N. McCrimmon, and J.D. Cure. Response of plant roots to elevated atmospheric carbon dioxide. *Plant Cell and Env.* 15:749-752.
- Schon, M. 1992. Tailoring nutrient solutions to meet the demands of your plants. In *Proceed. Ann. Conf. Hydroponics*. HSA, Orlando FL. pp 1-7.
- Sionit, N., H. Hellmers, and B.R. Strain. 1982. Interaction of atmospheric CO₂ enrichment and irradiance on plant growth. *Agron. J.* 74:721-725.
- Wheeler, R.M. C.L. Mackowiak, L.M. Siegrist, and J.C. Sager. 1993. Supraoptimal carbon dioxide effects on growth of soybean [*Glycine max* (L.) Merr.]. *J. Plant Physiol.* (In press).



REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE March 1994	3. REPORT TYPE AND DATES COVERED Technical Memorandum	
4. TITLE AND SUBTITLE Effect of Carbon Dioxide Enrichment on Radish Production Using Nutrient Film Technique (NFT)		5. FUNDING NUMBERS	
6. AUTHOR(S) C.L. Mackowiak, L.M. Ruffe, N.C. Yorio and R.M. Wheeler		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Bionetics Corporation, Mail Code: BIO-3 (CLM, LMR, NCY) and NASA Biological Research and Life Support Office, Mail Code: MD-RES (RMW) J.F. Kennedy Space Center, FL 32899		10. SPONSORING / MONITORING AGENCY REPORT NUMBER TM-109198	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Biomedical Operations and Research Mail Code: MD Kennedy Space Center, FL 32899		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Unclassified - Unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Radish plants (<u>Raphanus sativus</u> L. cvs. Cherry Belle, Giant White Globe, and Early Scarlet Globe) were grown in four different CO ₂ enriched environments, 0.04, 0.10, 0.50, and 1.00 kPa (400, 1000, 5000, 10000 ppm). Cultivar responses to CO ₂ treatments varied, where cv. Cherry Belle showed no significant response to CO ₂ enrichment, cv. Giant White Globe was moderately affected and Early Scarlet Globe was strongly affected. Enrichment at 0.10 kPa led to greater root dry matter (DM) than 1.00 kPa for cv. Giant White Glove, whereas 0.10 kPa produced greater storage root, shoot, and root DM than 1.00 kPa for cv. Early Scarlet Globe. The data suggest that 1.00 kPa CO ₂ may be detrimental to the growth of certain radish cultivars. Root:shoot ratios tended to increase with increasing CO ₂ concentration. Water use efficiency (g biomass/kg H ₂ O) increased with increasing CO ₂ enrichment, up to 0.5 kPa but then declined at the 1.00 kPa treatment. The total nitric acid used to maintain nutrient solution pH was lowest at the 1.00 kPa treatment as well, suggesting a decreased demand of nutrients by the plants at the highest CO ₂ level.			
14. SUBJECT TERMS radish, carbon dioxide, cultivar, supraoptimal, CELSS, water use efficiency			15. NUMBER OF PAGES 15
17. SECURITY CLASSIFICATION OF REPORT Unclassified			16. PRICE CODE
18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	