

LUMINOUS RADIO-QUIET SOURCES IN THE W3(MAIN) CLOUD CORE

C. G. WYNN-WILLIAMS, E. F. LADD, J. R. DEANE, AND D. B. SANDERS

Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

ABSTRACT We have resolved $450\mu\text{m}$ and $800\mu\text{m}$ emission from the W3(Main) star forming region into three major peaks, using $8'' - 14''$ beams with the James Clerk Maxwell Telescope on the summit of Mauna Kea. One of the submillimeter sources is identified with W3 - IRS5, a well-known candidate protostar. However, to our surprise, we find that none of the submillimeter peaks coincides with any of the prominent compact HII regions in the area. We estimate that the three submillimeter sources together contribute 30 - 50% of the total bolometric luminosity of the region, and speculate that the contribution of luminous radio-quiet sources to the total luminosity of HII region/molecular cloud complexes may be larger than is often assumed.

INTRODUCTION

The spectral energy distributions of HII region/molecular cloud complexes indicate that the bulk of their luminosity is emitted in the far infrared and that this far infrared and submillimeter emission is generated by cool ($T = 30-60$ K) dust (see e.g. Wynn-Williams & Becklin 1974; Chini, Krugel, & Wargau 1987). Because of the large beam sizes typically used for far-infrared and submillimeter continuum observations, it has been difficult to determine whether the OB stars ionizing the HII regions are the sources of the luminosity, or whether this luminosity is generated independently within the nearby molecular material. We present new submillimeter continuum observations of the W3(Main) region with high spatial resolution (< 0.1 pc). These new observations show that the distribution of submillimeter flux is not similar to the distribution of radio continuum emission, and suggest that the total luminosity generated by HII region/molecular cloud complexes is not dominated by the O and B stars which generate the observed HII regions.

OBSERVATIONS

Submillimeter observations of the W3(Main) region were made with the 15 m JCMT in 1992 November. The continuum maps were obtained using the facility UKT-14 bolometer system with passbands centered at approximately $450\mu\text{m}$ and $800\mu\text{m}$. Our results are shown in Figure 1, along with maps of $20\mu\text{m}$

(Wynn-Williams et al. 1972) and a 5 GHz radio continuum emission (Harris & Wynn-Williams 1976). The submillimeter continuum emission breaks up into three main emission centers—one in the east, and two in the west. We designate the sources SMS 1, 2, and 3 from E to W in order of decreasing right ascension.

DISCUSSION

SMS 1 is resolved and nearly circular at half power, with low flux level extensions to the east in the direction of IRS 3/W3B and north in the direction of IRS 1/W3A (see Figure 2a). The 450 μm and 800 μm centroid positions are consistent with the 20 μm position of IRS 5 (Wynn-Williams et al. 1972), several H₂O maser groupings (Genzel et al. 1987), and the radio continuum source W3(M) (Colley 1980).

SMS 2 lies close to the 20 μm source IRS 4 (Wynn-Williams et al. 1972), and near to the compact HII region W3(C) (Wynn-Williams 1971; see Figure 2b). We find that the position of SMS 2 lies 7.5'' from the center of W3(C). The size of the positional discrepancy is sufficiently large that we are confident that SMS 2 is not associated with W3(C). IRS 4 lies 4'' from the centers of both W3(C) and SMS 2, nearly on a line between these two sources. Therefore we conclude that there are at least two distinct major sources of emission in this area (W3(C) and SMS 2), and quite likely an additional unrelated infrared source (IRS 4).

SMS 3 is more extended than the other two submillimeter sources, with FWHM size of 30'' \times 16'' ($\alpha \times \delta$). No 20 μm emission was detected in this region to a point source detection threshold of 150 Jy (Wynn-Williams et al. 1972), nor was radio continuum emission detected greater than 6 mJy/2'' beam (Harris & Wynn-Williams 1976).

Combining our data with infrared results (Wynn-Williams et al. 1972; Werner et al. 1980), we have estimated the 20–800 μm luminosities for these three sources, as well as for IRS 1 (which was not detected as a distinct source in our submillimeter maps) and the entire region. The total luminosity for the region is estimated to be $5.2 \times 10^5 L_{\odot}$. IRS 1 and IRS 5 each account for about 30% of the total. SMS 1 and SMS 2 account for an additional 6% each. However, it should be noted that the luminosity of SMS 2 may contain some additional contribution from either IRS 4 or the source associated with W3(C), and therefore this luminosity should be regarded as an upper limit to the luminosity of SMS 2.

IRS 5 has long been recognized as a candidate high-mass protostar, based on its high luminosity, infrared energy distribution, and relatively weak radio continuum emission (see e.g. Wynn-Williams et al. 1972; Hackwell et al. 1978; Werner et al. 1980; Wynn-Williams 1982). We have found two more sources in the W3(Main) cloud that exhibit behavior similar to that of IRS 5. While both have luminosities about a factor of 5 lower than that of IRS 5, they are not associated with detected radio continuum emission. These radio-quiet sources account for at least 35% (SMS 1/IRS 5 + SMS 3) and up to 50% (SMS 1/IRS 5 + SMS 2 + SMS 3) of the total luminosity generated in the W3(Main) core. With the addition of extended emission probably generated by lower-luminosity, non-ionizing sources, the luminosity from W3(Main) could be roughly equally




Fig. 1 Submillimeter images of the W3 (Main) region at 450 μm (**top left**) and 800 μm (**bottom left**) compared with the 20 μm map from Wynn-Williams et al. (1972; **top right**) and the 5 GHz contours from Harris & Wynn-Williams (1976; **bottom right**). The submillimeter beam sizes are shown in the lower left of each relevant panel. Contours for the 450 μm map begin at 15 Jy / 8" beam and increment by 15 Jy / 8" beam. Contours for the 800 μm map begin at 1 Jy / 14" beam and increment by 1 Jy / 14" beam.




Fig. 2 a) Contours of the 450 μm emission near IRS 5. The 450 μm contours begin at 5 Jy / 8" beam and increment by 5 Jy / 8" beam. The positions of the 20 μm source, the radio continuum source W3 (M) and the H₂O maser centers are indicated by a circle, box and cross respectively. b) Comparison of our 450 μm contours with the 15 GHz emission from Colley (1980) toward the SMS 2 / IRS 4 / W3 (C) region. The 450 μm contours begin at 5 Jy / 8" beam and increment by 5 Jy / 8" beam. The position of the 20 μm and 5 μm sources are indicated by a circle and a triangle respectively.

divided between sources associated with HII regions and sources which have little or no ionized environs.

ACKNOWLEDGMENTS

We thank the operators of the JCMT for their assistance in obtaining the sub-millimeter continuum data. C. G. W.-W. acknowledges support from NSF grant AST89-19563. E. F. L. was supported by the James Clerk Maxwell Telescope Fellowship.

REFERENCES

- Chini, R., Krugel, E., & Wargau, W. 1987, *A&A*, 181,378
- Colley, D. 1980, *MNRAS*, 193,495
- Genzel, R., Downes, D., Moran, J. M., Johnston, K. J., Spencer, J. H., Walker, R. C., Haschick, A., Matveyenko, L. I., Kogan, L. R., Kostenko, V. I., Rönnäng, B., Rydbeck, O. E. H., & Moiseev, I. G. 1978, *A&A*, 66,13
- Hackwell, J. A., Gehrz, R. D., Smith, J. R., & Briotta, D. A. 1978, *ApJ*, 221,797
- Harris, S., & Wynn-Williams, C. G. 1976, *MNRAS*, 174,649
- Werner, M. W., Becklin, E. E., Gatley, I., Neugebauer, G., Sellgren, K., Thronson, H. A., Harper, D. A., Loewenstein, R., & Moseley, S. H. 1980, *ApJ*, 242,601
- Wynn-Williams, C. G. 1982, *ARA&A*, 20,587
- Wynn-Williams, C. G. 1971, *MNRAS*, 151,397
- Wynn-Williams, C. G., & Becklin, E. E. 1974, *PASP*, 86,5
- Wynn-Williams, C. G., Becklin, E. E., & Neugebauer, G. 1972, *MNRAS*, 160,1

DMT
12
E 127

Science with High Spatial Resolution Far-Infrared Data

LIST OF AUTHORS

C.H. Anderson	139
H.H. Aumann	1
Tj.R. Bontekoe	23, 133
Y. Cao	55
B. Das	17
J.R. Deane	173
N.A. Devereux	79
P.E. Dewdney	161
Rh. Evans	117
J.W. Fowler	1
S.K. Ghosh	17
R.A. Gonsalves	47
D.A. Harper	97
B. Heikkila	91
C.E. Heiles	161
G. Helou	85, 111
J.P. Kennealy	47
D.J.M. Kester	23, 133
G.R. Knapp	97
R.M. Korte	47
E.F. Ladd	173
W.D. Langer	139
J.F. Lees	97
D. Lester	33
C. Loup	133
K.A. Marsh	111
A.P. Marston	11, 103
P.G. Martin	161
J.M. Mazarella	69
R.K. Piña	61
S.D. Price	47
T.A. Prince	55
R.C. Puetter	61
T.N. Rengarajan	17
W.T. Reach	161
C. Rogers	161
M.P. Rupen	97
R.S. Sample	47
D.B. Sanders	173
M.F. Schlapfer	169
J.A. Surace	69, 123
S. Terebey	123
R.P. Verma	17
R.G. Walker	169
Z. Wang	153

L.B.F.M. Waters	133
W.R. Webber	91
R.W. Wilson	139
C.G. Wynn-Williams	173
C. Xu	85

Science with High Spatial Resolution Far-Infrared Data

LIST OF ATTENDEES

George Aumann
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Jean-Phillipe Bernard
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

T. Romke Bontekoe
 European Space Agency
 Astrophysics Division
 Postbus 299
 2200 AG Noordwijk
 THE NETHERLANDS

Yu Cao
 California Institute of Technology
 MS 103-33
 Pasadena, CA 91125

Thomas Chester
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

James Deane
 Inst. for Astronomy
 2680 Woodlawn Drive
 Honolulu, HI 96822

N. Devereux
 New Mexico State University
 Dept. of Astronomy
 Box 30001 - Dept 4500
 Las Cruces, NM 88003

Rhodri Evans
 University of Toledo
 Dept. of Physics and Astronomy
 Toledo, OH 43606

John Fowler
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

W. Gong
 Queen Mary College
 Dept. of Physics
 Miles End Road
 London E1 4NS
 UNITED KINGDOM

John Hackwell
 The Aerospace Corp.
 P.O. Box 92957
 M2-266
 Los Angeles, CA 90009

Bryant Heikkila
 New Mexico State University
 Dept. of Astronomy
 Las Cruces, NM 88003

George Helou
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Do Kester
 University of Groningen
 Space Research Dept
 WSN Gebouw
 P.O. Box 800
 9700 Av Groningen
 THE NETHERLANDS

Ned Ladd
 University of Hawaii
 Institute for Astronomy
 2680 Woodlawn Drive
 Honolulu, HI 96822

William Langer
 Jet Propulsion Laboratory
 MS 169-506
 4900 Oak Grove Drive
 Pasadena, CA 91109

Joanna F. Lees
 University of Chicago
 Dept. of Astronomy
 5640S. Ellis Ave
 Chicago, IL 60637

Daniel Lester
 University of Texas
 Dept. of Astronomy
 RLM Hall
 Austin, TX 78712

Debbie Levine
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Steve Lord
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Kenneth Marsh
 Jet Propulsion Laboratory
 MS 163-327
 4900 Oak Grove Drive
 Pasadena, CA 91109

A.P. Marston
 Drake University
 Dept. of Physics and Astronomy
 Des Moines, IA 50311

Peter Martin
 University of Toronto
 Canadian Inst. Theor. Astrophysics
 Toronto, ON, M5S 1A7 CANADA

Joseph M. Mazarella
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Mehrdad Moshir
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Stephan D. Price
 Phillips Laboratory
 Optical Physics Division
 Hanscom AFB, MA 01731-5000

Thomas Prince
 California Institute of Technology
 MS 220-47
 Pasadena, CA 91125

Richard Puetter
 University of California, San Diego
 CASS
 Code C-011
 LaJolla, CA 92093

Michael P. Rupen
 National Radio Astron.Obs.
 P.O. Box O
 Socorro, NM 87801

Martin Schlapfer
 Jamieson Science and Engineering
 Suite 204
 5321 Scotts Valley Drive
 Scotts Valley, CA 95066

James Schombert
 California Institute of Technology
 Infrared Processing and Analysis Center
 MS 100-22
 Pasadena, CA 91125

Jason Surace
 Institute for Astronomy
 University of Hawaii
 2680 Woodlawn Drive
 Honolulu, HI 96822

Susan Terebey
California Institute of Technology
Infrared Processing and Analysis Center
MS 100-22
Pasadena, CA 91125

Ann Wehrle
California Institute of Technology
Infrared Processing and Analysis Center
MS 100-22
Pasadena, CA 91125

R.P Verma
Tata Institute of Fundamental Research
Homi Bhabha Road
Bombay 400005
INDIA

Z. Wang
Center for Astrophysics
MS 66
60 Garden Street
Cambridge, MA 02138

L.B.F.M. Waters
Laboratory for Space Research
P.O. Box 800
9700 Av Groningen
THE NETHERLANDS

Michael Werner
Jet Propulsion Laboratory
MS 233-303
4900 Oak Grove Drive
Pasadena, CA 91109

William Webber
New Mexico State University
Dept. of Astronomy
Las Cruces, NM 88003

Gareth Wynn-Williams
University of Hawaii
Institute for Astronomy
2680 Woodlawn Drive
Honolulu, HI 96822

Cong Xu
Max-Plank-Inst fur Kernphysik
Astrophysics Group
Postfach 103980
D-6900 Heidelberg 1
GERMANY

Erick Young
University of Arizona
Steward Observatory
Tucson, AZ 85721

1. Report No. 94-5	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Infrared Processing and Analysis Center (IPAC) Science With High Spatial Resolution Far-Infrared Data Proceedings of a Workshop Held June 14-16, 1993		5. Report Date April 1994	
		6. Performing Organization Code	
7. Author(s) S. Terebey and J. Mazzarella		8. Performing Organization Report No. 94-5	
9. Performing Organization Name and Address JET PROPULSION LABORATORY California Institute of Technology 4800 Oak Grove Drive Pasadena, California 91109		10. Work Unit No.	
		11. Contract or Grant No. NAS7-918	
		13. Type of Report and Period Covered Proceedings	
12. Sponsoring Agency Name and Address NATIONAL AERONAUTICS AND SPACE ADMINISTRATION Washington, D.C. 20546		14. Sponsoring Agency Code	
		15. Supplementary Notes	
16. Abstract <p>Data at far-infrared wavelengths are difficult to obtain and usually have much lower spatial resolution than data which are routinely obtained at other wavelengths. However, the many astrophysical problems that can only be well addressed in the far-infrared have inspired efforts to extract high spatial resolution information from instruments including the Infrared Astronomical Satellite (IRAS) and the Kuiper Airborne Observatory (KAO). The IRAS data also provide a fundamental archive since IRAS produced a nearly complete survey of the infrared sky. Although the IRAS survey strategy was not designed with the specific purpose of constructing images of the sky, the ingenuity of software developers has recently resulted in innovative techniques which are allowing astronomers to push the limits of these and other far-infrared data.</p> <p>The goal of this workshop was to discuss new science and techniques relevant to high spatial resolution processing of far-infrared data, with particular focus on high resolution processing of IRAS data. Users of the maximum correlation method, maximum entropy, and other resolution enhancement algorithms applicable to far-infrared data gathered at the Infrared Processing and Analysis Center (IPAC) on the Caltech campus in Pasadena for two days in June 1993 to compare techniques and discuss new results. During a special session on the third day, interested astronomers were introduced to IRAS HIRES processing, which is IPAC's implementation of the maximum correlation method to the IRAS data. Individuals who desire further information about IRAS or HIRES processing are encouraged to contact IPAC.</p>			
17. Key Words (Selected by Author(s)) Far-infrared data; high spatial resolution data processing; resolution enhancement; maximum entropy.		18. Distribution Statement ORIGINAL PAGE IS OF POOR QUALITY	
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 187	22. Price

