GUIDELINES FOR LIGHTING OF PLANTS IN CONTROLLED ENVIRONMENTS

23

240-74

N96-18163

Gerald Dietzer, Department of Horticulture, University of Maryland, College Park, MD, Robert Langhans, Department of Floriculture, Cornell University, Ithaca, NY, John Sager, Code MD-RES-L, Kennedy Space Center, FL, L. Art Spomer, Department of Horticulture, University of Illinois, Urbana, IL, Ted Tibbitts, Department of Horticulture, University of Wisconsin, Madison, WI

The organizing committee outlined draft guidelines for plants to provide a focus for the discussions at the workshop. These were distributed to the participants before the meeting. It was recognized that there was insufficient data to support many of the particular quantities presented. The guidelines served as a basis for discussion amongst the workshop attendees and led to a number of recommendations that were recorded by the session chairpersons. The organizing committee indicated they would incorporate the recommendations and suggestions into a revised set of guidelines for additional discussion. Interested participants were then asked to indicate their willingness to review this revised set of guidelines to lead toward the future development of guidelines for lighting in controlled environments. It was understood that these guidelines will not be standards and will require upgrading and modifications as lamps and equipment become available and as new insights are obtained on plants response to light.

Revised draft guidelines are included as Tables 1 and 2 that have been developed by the organizing committee following the suggestions obtained at the workshop. Table 1 are the guidelines for growth chambers and Table 2 for greenhouses. These have been distributed to the participants that indicated a willingness to review proposals that were developed. It is hoped that these proposals will lead to the development of guidelines that will have general acceptance by plant scientists.

TABLE 1. GUIDELINES FOR LIGHTING IN GROWTH CHAMBERS

The purpose of these guidelines is to help writers of specifications, engineers, and architects, who have to make recommendations for the installation of lighting in growth chambers. It is not the intent of these guidelines to mandate the lighting a researcher may need for specific projects, but rather guidelines that indicate reasonable lighting that can grow acceptable crops any time of the year.

PHOTOSYNTHETICALLY ACTIVE RADIATION (PAR)

A daily average irradiance of 26 mol m⁻² day⁻¹ will effectively grow most species of higher plants. This equates to an instantaneous irradiance of 300 μ mol m⁻² s⁻¹ for 24 hours or 600 μ mol m⁻² s⁻¹ for 12 hours. For comparison in the continental United States, the average annual daily irradiance is about 26 mol m⁻² for Madison, WI and Washington, DC. In the summer the maximum daily irradiance is 62 mol m⁻² at Phoenix, AZ and in the winter the minimum irradiance is 8 mol m⁻² at Madison, WI (see Table 1). The maximum solar irradiance around midday of 2000 μ mol m⁻² s⁻¹ is transient and is not necessary for normal plant growth since the plants response is based on the average daily irradiance.

UNIFORMITY

Less than $\pm 10\%$ variation on a horizontal plane over the growing area at the plant canopy height. The variation should be based on measurements taken in the center of each square meter of the plant growing area.

SPECTRAL

280-320 nm (Ultraviolet-B)	Unspecified, but in general the effects of UV-B are deleterious to plant growth and development. However, some plants, such as members of the Solonaceae, may require a small quantity ($\sim 3 \text{ W m}^{-2}$) to avoid abnormal development.
320-400 nm (Ultraviolet-A)	Unspecified, but may have an additive effect with the requirement for blue.
400-500 nm (Blue)	An absolute quantity for elongation control is required for most higher plants (\geq 30 µmol m ⁻² s ⁻¹).
500-600 nm (Green)	Not necessary for photosynthesis, but contributes to photosynthesis and is a significant component of most radiation sources.
600-700 nm (Red)	Optimize output for maximal photosynthesis. Monochromatic red will cause abnormal development in some species.
700-750 nm (Far-red)	Enhancement of flowering, stem elongation, etc. of certain species (as a function of the red/far-red ratio) with the quantity centered around 725 nm equal tc or greater than the quantity centered around 660 nm.

TOTAL IRRADIANCE (Over the range of 280-50,000 nm)

A ratio of total irradiance to PAR of 0.50 or less W m⁻² per μ mol m⁻² s⁻¹ (2.3 W m⁻² per W m⁻² PAR) is desirable to reduce thermal heating of plants and soil. The solar radiation ratio is less than 0.50 W m^{-s} per μ mol m⁻² s⁻¹ PAR. A ratio below 0.50 cannot be obtained with most lamps without a barrier and adequate ventilation or a luminaire specifically designed to dissipate infra-red radiation. For example, the ratios for metal halide lamps without a barrier, with an acrylic barrier, and with an acrylic barrier with 5 cm of water was shown to be 0.60, 0.53, and 0.24 W m⁻² per μ mol m⁻² s⁻¹ PAR, respectively. Low temperature sources such as low pressure sodium lamps and light emitting diodes (LEDs) without barriers have been shown to have ratios of 0.41 and 0.28, respectively.

TABLE 2. GUIDELINES FOR INSTALLATION OF SUPPLEMENTAL LIGHTING IN GREENHOUSES

The purpose of these guidelines is to help writers of specifications, engineers, and architects, who have to make recommendations for the installation of supplemental lighting in greenhouses. It is not the intent of these guidelines to mandate the lighting a researcher may need for specific projects, but rather guidelines that indicate reasonable lighting that can grow acceptable crops any time of the year.

PHOTOSYNTHETICALLY ACTIVE RADIATION (PAR)

A daily average irradiance of 26 mol m⁻² day⁻¹ from sunlight plus added supplementation will effectively grow most species of higher plants. Lamp lighting should be used to supplement sunlight and provide 26 mol m⁻² day⁻¹. Installations of lighting providing greater than 200 μ mol m⁻² s⁻¹ generally add too much heat to the greenhouse environment and the extra luminaries add too much shade. Lighting can be provided during the sunlight hours or during the night period depending upon the plant's photoperiod requirements, and/or depending upon the most cost-effective time to activate the lamps. Shading systems should be utilized under high sunlight conditions to reduce the average irradiance to 26 moles day⁻¹.

UNIFORMITY

Less than $\pm 15\%$ variation on a horizontal plane over the growing area at the plant growing canopy height. The variation should be based on measurements taken in the center of each square meter of the total lighted area. Installation of a uniform lighting system in the greenhouse is difficult. It is seldom possible to obtain this uniformity on the outside edges of the growing area, particularly against the walls of the greenhouse.

SPECTRAL

There are no special spectral requirements for the supplemental lighting for photosynthesis in greenhouses. Sunlight should supply the balance of wavelengths required by plants. Most glazings remove some portion of the ultraviolet radiation from sunlight and thus certain plant species, including most *Solonaceous* species, may have some abnormal development (oedema) as a consequence. However daylength extensions should use lamps high in red and far-red.

TOTAL IRRADIANCE (Over the range of 280-50,000 nm)

Recommend that supplemental lighting produce no more than 0.6 Wm^{-2} of total irradiation for each μ mol m⁻² s⁻¹ of PPF.

REPORT DOC	Form Approved OMB No. 0704-0188				
gathering and maintaining the data needed, and com collection of information, including suggestions for re Davis Highway, Suite 1204, Arlington, VA 22202-4302	pleting and reviewing the collection of i educing this burden, to Washington Hea t, and to the Office of Management and	nformation. Send comments regain dquarters Services, Directorate for Budget, Paperwork Reduction Proje	viewing instructions, searching existing data sources, ding this burden estimate or any other aspect of this information Operations and Reports, 1215 Jefferson ect (0704-0188), Washington, DC 20503.		
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE March 1994	3. REPORT TYPE AND Conference Publication Ma	rch 27-30, 1994		
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS		
. (N/			NASW 4527 (NASA Code UL)		
6 . AUTHOR(S) Ted W. Tibbitts (Ed)					
7. PERFORMING ORGANIZATION NAME	8. PERFORMING ORGANIZATION REPORT NUMBER				
University of Wisconsin Department of Horticult Madison, WI 53706			None		
9. SPONSORING / MONITORING AGENC	AND ADDRESSIES	1	10. SPONSORING / MONITORING		
U.S. Department of Agri	culture	,	AGENCY REPORT NUMBER		
Agricultural Experiment National Aeronautics an National Science Founda Department of Energy	ion	CP-3309			
Department of Energy Wisconsin Center for Dep 11. SUPPLEMENTARY NOTES	mand Side Research				
12a. DISTRIBUTION / AVAILABILITY STA	TEMENT		12b. DISTRIBUTION CODE		
Unclassified-Unlimited					
Subject Category: 54					
13. ABSTRACT (Maximum 200 words)					
plant research and is g carefully for animal an reevaluate the technolog	aining recognition d human research. gy that is availabl effective use of 1 r plants but also t	as a significant Thus this worksh e today and to w ighting in contr o initiate inter	op was convened to ork toward developing olled environments with		
14. SUBJECT TERMS Photosynthesis, Plant r Ultraviolet, Animal req distribution filter, he	uirements, Lighting	, lamps, lightin	15. NUMBER OF FAGES 405 8 16. PRICE CODE		
	SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFIC	CATION 20. LIMITATION OF ABSTRACT		
Unclassified	S. THIS FACE	U. ADJINACI	UL		
NSN 7540-01-280-5500		• • • • • • • • • • • • • • • • • • • •	Standard Form 298 (Rev. 2-89)		

Jungara	0			-	
Prescribed by 298-102	ANSI	Std	Z39-18		
290-102					