

# There aren't Non-standard Solutions for the Braid Group Representations of the QYBE Associated with 10-D Representations of SU(4).

Huang Yijun, Yu Guochen and Sun Hong\*

Department of Foundation the First Aeronautical College of Air Force

Xinyang Hennan 464000 P. R. China .

## Abstract

In this paper by employing the weight conservation and the diagrammatic techniques we show that the solutions associated with the 10-D representations of SU (4) are standard alone .

## 1 Introduction

It is well known that the quantum Yang-Baxter equations ( QYBE) play an important role in various theoretical and mathematical physics, such as completely integrable system in (1+1)-dimensions, exactly solvable models in statistical mechanics, the quantum inverse scattering method and the conformal field theories in 2-dimensions . Recently, much remarkable progress has been made in constructing the solutions of the QYBE associated with the representations of lie algebras . It is shown that for some cases except the standard solutions, there also exist new solutions, but the others have not non-standard solutions . In reference 11, we derived the braid group representations associated with the 10-dimensional representation of SU (4) and corresponding trigonometric and rational solutions . In this paper, the classical limit of the braid group representations is checked . Then it is shown that the solutions associated with the 10-dimensional representations are standard alone .

## 2 Classical Limit

It is well known that in the classical limits as  $q \rightarrow 1$  the standard solution of QYBE require that<sup>[10]</sup>

$$S \Big|_{q \rightarrow 1} = P [ I + (q-1) r ] + o[ (q-1)^2 ] \quad (2.1)$$

and

$$\Phi_v^T r \Phi_v = C_v = 2C_R - C_{E_v} \quad (2.2)$$

where P is the permutation operator and  $\Phi_v$  stands for the normalized classical eigenvectors, r is the classical r-matrix,  $C_R$  and  $C_{E_v}$  are the Casimirs . The eigenvalues are given by

$$\lambda_v = (\pm) q^c \quad (2.3)$$

In Ref . (11), We have derived the braid group representations associated with the 10-D

---

\* Address: Jinan 250023

representations of SU (4) . The Casimir eigenvalues of S -matrix were given by

$$\lambda_1 = q^{-3}, \lambda_2 = -q, \lambda_3 = q^3 \quad (2.4)$$

From the result of Ref. (11) we know that there are some fundamental submatrices,  $A_1, A_2^{(1)}, A_3^{(1)}, A_4^{(2)}, A_4^{(3)}$  and  $A_6^{(2)}$ , and others can be expressed by direct sum of the fundamental submatrices. So we discuss only the classical limits of this submatrices.

For example, we discuss only  $A_3^{(1)}$ :

$$A_3^{(1)} = \begin{bmatrix} 0 & 0 & q \\ 0 & q^{-1} & qw \\ q & qw & (1-q^2)w \end{bmatrix} \quad (2.5)$$

$$A_3^{(1)} \Big|_{q \rightarrow 1} = \begin{bmatrix} 0 & 0 & q \\ 0 & 2-q & 4(1-q) \\ q & 4(1-q) & 0 \end{bmatrix} \quad (2.6)$$

$$r_3^{(1)} = \begin{bmatrix} 1 & -4 & 0 \\ 0 & -1 & 4 \\ 0 & 0 & 1 \end{bmatrix} \quad (2.7)$$

$$\Phi_1^T = \frac{1}{\sqrt{6}} (1 \ 2 \ 1), \quad \Phi_2^T = \frac{1}{\sqrt{2}} (1 \ 0 \ -1), \quad \Phi_3^T = \frac{1}{\sqrt{3}} (1 \ -1 \ 1) \quad (2.8)$$

and

$$\Phi_v^T r_3^{(1)} \Phi_v = C_v = \begin{cases} 3 & v=1 \\ 1 & v=2 \\ 3 & v=3 \end{cases} \quad (2.9)$$

Therefore the solutions of QYBE are standard.

### 3 About absence of the nonstandard solution

From Ref. (11) we have known that so long as  $u_6 = u_2 = u_{-4} = u_{-6}$ , there exists the alone solution. In Ref. (11), we have

$$\begin{cases} u_4 q_8^{2(2.4)} + w_{10}^{(4.6)} p_8^{2(2.6)} = w_{10}^{(4.6)} p_6^{2(2.4)} \\ u_6 q_8^{2(2.4)} + w_{10}^{(4.6)} u_4^2 = u_4 w_{10}^{2(2.6)} + w_{10}^{(4.6)} p_{10}^{2(4.6)} \\ u_4^2 = P_{10}^{(4.6)} p_6^{2(4.6)} \end{cases} \quad (3.1)$$

$$\left\{ \begin{array}{l} u_6 p_8^{(2,6)} = p_{10}^{2(4,6)} \\ u_2 p_8^{(2,6)} = p_6^{2(2,4)} \end{array} \right. \quad (3.2)$$

$$\left\{ \begin{array}{l} u_{-5} q_{-10}^{2(-6,-5)} + w_{-9}^{(-5,-4)} p_{-10}^{2(-6,-4)} = w_{-9}^{(-5,-4)} p_{-11}^{2(-6,-5)} \\ u_{-4} q_{-10}^{2(-6,-5)} + w_{-9}^{(-5,-4)} u_{-5}^2 = w_{-9}^{2(-5,-4)} u_{-5} + w_{-9}^{(-5,-4)} p_{-9}^{2(-5,-4)} \\ u_{-5}^2 = p_{-11}^{(-6,-5)} p_{-9}^{(-5,-4)} \end{array} \right. \quad (3.3)$$

$$\left\{ \begin{array}{l} u_{-6} p_{-10}^{(-6,-4)} = p_{-11}^{2(-6,-5)} \\ u_{-4} p_{-10}^{(-6,-4)} = p_{-9}^{2(-5,-4)} \end{array} \right. \quad (3.4)$$

$$\left\{ \begin{array}{l} u_0 q_{10}^{2(-6,0)} + w_{-6}^{(-6,0)} p_0^{2(-6,6)} = w_{-6}^{(-6,0)} p_6^{2(0,6)} \\ u_5 q_{10}^{2(-6,0)} + u_0^2 w_6^{(0,6)} = u_0 w_6^{2(0,6)} + w_6^{(0,6)} p_6^{2(0,6)} \\ u_0^2 = p_6^{(0,6)} p_{-6}^{(-6,0)} \end{array} \right. \quad (3.5)$$

$$\left\{ \begin{array}{l} u_{-6} p_0^{(-6,6)} = p_{-6}^{2(-6,0)} \\ u_6 p_0^{(-6,6)} = p_6^{2(0,6)} \end{array} \right. \quad (3.6)$$

From eq. (3.1), we have

$$u_4 = p_{10}^{(4,6)} ; p_6^{(2,4)} = p_{10} \quad (3.7)$$

From eq. (3.2) + eq. (3.6), we have

$$u_2 = u_6 \quad (3.8)$$

$$u_{-5} = p_{-9}^{(-5, -4)}, \quad p_{-9}^{(-5, -4)} = p_{-11}^{(-6, -5)} \quad (3.9)$$

$$u_{-4} = u_{-6} \quad (3.10)$$

$$u_0 = p_6^{(0, 6)}, \quad p_6^{(0, 6)} = p_{-6}^{(-6, 0)} \quad (3.11)$$

$$u_{-6} = u_6 \quad (3.12)$$

From eq. (3.8), (3.10) and (3.12) we have

$$u_6 = u_2 = u_{-4} = u_{-6} \quad (3.13)$$

Therefore the solutions of the QYBE are standard alone.

## References

- [1] C. N. Yang, Phys. Rev. Lett. 19 (1967) 1312.
- [2] R. J. Baxter, Ann. Phys. 70 (1972) 1.
- [3] A. B. Zamolodchikov and Al. B. Zamolochikov, Ann. Phys. 120 (1979) 253.
- [4] L. D. Faddeev, Sov. Sci. Rev. Math. Phys. cl (1980) 107.
- [5] Molin Ge, Yongshi wu and Kang Xue, ITP-SB-90-02.
- [6] Molin Ge, and Kang Xue, ITP-SB-90-20.
- [7] Y. Q. Li, Weight Conservation and Braid Group Representations. Ph. D. thesis, Lanzhou University (1989).
- [8] Molin Ge, Changpu Sun, Luyu Wang and Kang Xue, Weight conservation and Quantum Group Construction of the Braid Group Representation, J. Phys. A.
- [9] Molin Ge, Y. Q. Wang and K. Xue, Int. J. Mod. Phys. (1990).
- [10] N. Yu, Reshetikhin, LOMI E -4-87, E-14-87 (1988).
- [11] Guochen Yu, Xifu Wang and Kang Xue, Commun. Theor. Phys. 18 (1992).