

Reprinted from

U.S. NAGW-4358

11-52-012

(C) 1995

65043

BRAIN RESEARCH

Brain Research 695 (1995) 275-278

Short communication

Role of lateral parabrachial nucleus in the inhibition of water intake produced
by right atrial stretch

Lynne E. Ohman, Alan Kim Johnson *

Departments of Psychology and Pharmacology and the Cardiovascular Center, University of Iowa, Iowa City, IA 52242-1407, USA



ELSEVIER

BRAIN RESEARCH

SCOPE AND PURPOSE

BRAIN RESEARCH provides a medium for prompt publication of articles in the fields of neuroanatomy, neurochemistry, neurophysiology, neuroendocrinology, neuropharmacology, neurotoxicology, neurocommunications, behavioural sciences, molecular neurology and biocybernetics. Clinical studies that are of fundamental importance and have a direct bearing on the knowledge of the structure and function of the brain, the spinal cord, and the peripheral nerves will also be published.

TYPE OF PAPERS

1. **Research Reports** reporting results of original fundamental research in any branch in the brain sciences. It is expected that these papers will be published about six months after acceptance.
2. **Short Communications** reporting on research which has progressed to the stage when it is considered that the results should be made known quickly to other workers in the field. The maximum length allowed will be 1500 words or equivalent space in tables and illustrations. It is expected that short communications will be published about three months after acceptance.

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted in *quadruplicate* and be complete in all respects (including four copies of all illustrations), and deal with original material not previously published, or being considered for publication elsewhere. Manuscripts submitted under multiple authorship are reviewed on the assumption that all listed authors concur with the submission and that a copy of the final manuscript has been approved by all authors and tacitly or explicitly by the responsible authorities in the laboratories where the work was carried out. If accepted, the manuscript shall not be published elsewhere in the same form, in either the same or another language, without the consent of the Editors and Publisher. Submission of a paper to **Brain Research** is understood to imply that it has not previously been published (except in abstract form) and that it is not being considered for publication elsewhere.

The Publisher and Editor-in-Chief regret that they are unable to return copies of submitted manuscripts except in the case of rejected articles, where only the original manuscript and figures are returned to the author.

Research reports and short communications should be sent to:

Professor Dr. D.P. Purpura
Brain Research Office of the Dean
Albert Einstein College of Medicine
Jack and Pearl Resnick Campus
1300 Morris Park Avenue
Bronx, NY 10461, USA
Tel.: (1) (718) 430-2387
Fax: (1) (718) 597-4072
E-mail: brain@aecom.yu.edu

Correspondence relating to proofs, publication and reprints should be sent to:

Brain Research
Elsevier Science B.V.
P.O. Box 1527
1000 BM Amsterdam
The Netherlands
Tel.: (31) (20) 485 3462
Fax: (31) (20) 485 3239
E-mail: r.wattimena@elsevier.nl

Information about Neuroscience Protocols: Dr. F.G. Wouterlood, Department of Anatomy, Faculty of Medicine, Van der Boechorststraat 7, 1081 BT Amsterdam, The Netherlands. Fax: (31) (20) 444 8054; E-mail: fg.wouterlood.anat@med.vu.nl

EDITORIAL BOARD

Editor-in-Chief: Dominick P. Purpura (Bronx, NY, USA)

G.K. Aghajanian (New Haven, CT, USA)

B.W. Agranoff (Ann Arbor, MI, USA)

A.J. Aguayo (Montreal, Que., Canada)

P. Andersen (Oslo, Norway)

E.C. Azmitia (New York, NY, USA)

M.V.L. Bennett (Bronx, NY, USA)

L.I. Benowitz (Boston, MA, USA)

J.M. Besson (Paris, France)

A. Björklund (Lund, Sweden)

F.E. Bloom (La Jolla, CA, USA)

D. Choi (St. Louis, MO, USA)

R.E. Coggeshall (Galveston, TX, USA)

H. Collewijn (Rotterdam, The Netherlands)

W.M. Cowan (Bethesda, MD, USA)

A.C. Cuello (Montreal, Que., Canada)

M.S. Cynader (Vancouver, BC, Canada)

J.E. Dowling (Cambridge, MA, USA)

J.J. Dreifuss (Geneva, Switzerland)

B. Droz (Lausanne, Switzerland)

R. Dubner (Bethesda, MD, USA)

S.B. Dunnett (Cambridge, UK)

P.C. Emson (Cambridge, UK)

S.J. Enna (Kansas City, KS, USA)

J. Feldman (Los Angeles, CA, USA)

H.L. Fields (San Francisco, CA, USA)

F.H. Gage (La Jolla, CA, USA)

J. Glowinski (Paris, France)

P.S. Goldman-Rakic (New Haven, CT, USA)

S. Grillner (Stockholm, Sweden)

B. Gustafsson (Göteborg, Sweden)

A. Hamberger (Göteborg, Sweden)

U. Heinemann (Köln, FRG)

K.-P. Hoffmann (Bochum, FRG)

T.G.M. Hökfelt (Stockholm, Sweden)

R.L. Isaacson (Binghamton, NY, USA)

M. Ito (Saitama, Japan)

B.L. Jacobs (Princeton, NJ, USA)

E.G. Jones (Irvine, CA, USA)

P. Kalivas (Pullman, WA, USA)

K. Kogure (Saitama-ken, Japan)

G.F. Koob (La Jolla, CA, USA)

L. Kruger (Los Angeles, CA, USA)

J. LaVail (San Francisco, CA, USA)

C.L. Masters (Parkville, Vic., Australia)

M. Mattson (Lexington, KY, USA)

B.S. McEwen (New York, NY, USA)

E.G. McGeer (Vancouver, BC, Canada)

R.Y. Moore (Pittsburgh, PA, USA)

P. Morell (Chapel Hill, NC, USA)

W.T. Norton (New York, NY, USA)

J.M. Palacios (Barcelona, Spain)

M. Palkovits (Budapest, Hungary)

A.J. Patel (London, UK)

R. Quirion (Verdun, Que., Canada)

C.S. Raine (Bronx, NY, USA)

G. Raisman (London, UK)

P. Rakic (New Haven, CT, USA)

H.J. Ralston III (San Francisco, CA, USA)

S.I. Rapoport (Bethesda, MD, USA)

C.E. Ribak (Irvine, CA, USA)

P. Rudomín (Mexico, DF, Mexico)

M. Schachner (Zürich, Switzerland)

B.K. Siesjö (Lund, Sweden)

E.J. Simon (New York, NY, USA)

S.H. Snyder (Baltimore, MD, USA)

C. Sotelo (Paris, France)

J. Stone (Sydney, NSW, Australia)

D.F. Swaab (Amsterdam, The Netherlands)

L. Swanson (Los Angeles, CA, USA)

L. Terenius (Stockholm, Sweden)

H. Thoenen (Planegg, FRG)

M. Tohyama (Osaka, Japan)

S. Tucek (Prague, Czech Republic)

K. Unsicker (Heidelberg, FRG)

S.G. Waxman (New Haven, CT, USA)

F.G. Wouterlood (Amsterdam, The Netherlands)

R.J. Wurtman (Cambridge, MA, USA)

I.S. Zagon (Hershey, PA, USA)

W. Zieglgänsberger (Munich, FRG)

R.S. Zukin (Bronx, NY, USA)

GUIDELINES FOR THE SUBMISSION OF MANUSCRIPTS

These can be found in the "front matter" of every last issue of each volume of **Brain Research**.

The preferred medium of submission is on disk with accompanying manuscript (see "Electronic manuscripts").

Short communication

Role of lateral parabrachial nucleus in the inhibition of water intake produced by right atrial stretch

Lynne E. Ohman, Alan Kim Johnson *

Departments of Psychology and Pharmacology and the Cardiovascular Center, University of Iowa, Iowa City, IA 52242-1407, USA

Received 18 January 1994; accepted 11 July 1995

Abstract

Rats with either bilateral electrolytic or sham lesions of the ventrolateral portion of the lateral parabrachial nucleus (VLLPBN) were implanted with latex balloons that lay at the right superior vena cava/atrial junction (RSVC/AJ). Water intake in response to isoproterenol was measured both with and without inflation of the balloon. Water intake of the sham-lesioned rats was significantly depressed by balloon inflation during the first hour of the experiment. In contrast, water intake in the VLLPBN-lesioned rats was unaffected by balloon inflation. These results suggest that the VLLPBN is involved in the processing of afferent input from stretch-activated RSVC/AJ receptors.

Keywords: Atrial volume receptor; Water intake; Lateral parabrachial nucleus; Isoproterenol; Rat

The role of vascular volume receptors located on the low pressure side of the systemic circulation (i.e., located in the atria and great veins; a.k.a. cardiopulmonary receptors) in the control of water intake is unclear. There is some evidence, albeit largely circumstantial, that input from these receptors may facilitate water intake in states of extracellular volume depletion [7,23]. In contrast, however, evidence suggesting that the volume receptors contribute to satiety (i.e., the termination of drinking) is more complete. Activation of mechanoreceptors in the rat by inflation of a small balloon in the right superior vena cava/atrial junction (RSVC/AJ) has been reported to attenuate spontaneous night-time water intake, as well as drinking in response to 24 h water deprivation, treatment with the hyperoncotic colloid polyethylene glycol and isoproterenol [10]. This manipulation had no effect on drinking induced by the infusion of hypertonic saline. Similar findings have been reported in the dog, with the exception that volume receptor activation in this species also attenuates water intake stimulated by cellular dehydration [19].

It is likely that volume information is integrated into the 'visceral' neural network that is involved in the control of

drinking behavior. Anatomical evidence indicates that the lateral parabrachial nucleus (LPBN) is one site where peripherally derived inputs might converge. The LPBN, particularly the ventrolateral portion of the LPBN (VL-LPBN), receives the most prominent ascending projection of both the area postrema and nucleus of the solitary tract [16,17,26] and is involved through a rich communication network with hypothalamic nuclei known to be important in the maintenance of body fluid balance [6,17]. In addition, functional data exist linking the LPBN to the control of water intake [4,20,21].

The purpose of this experiment was to test the hypothesis that the VLLPBN mediates, at least in part, the reduction of water intake following activation of RSVC/AJ receptors. Isoproterenol, which to a degree mimics extracellular volume depletion (e.g., see Ref. [23] for review and references), was chosen as the experimental dipsogen.

This experiment was carried out on male Sprague-Dawley rats weighing between 300–420 g at the time of testing. The animals received either a bilateral electrolytic or sham lesion of the VLLPBN. Lesions were made by passing a 1.6 mA direct current through a stainless steel electrode for 4 s. Following a 2-week recovery period, atrial balloons were implanted in both groups of rats following the procedure described by Kaufman [10]. Briefly, the balloon was constructed of latex securely tied over the end of PE 50 tubing. The latex and PE tubing

* Corresponding author. Department of Psychology, University of Iowa, 11 Seashore Hall E., Iowa City, IA 52242-1407, USA. Fax: (1) (319) 335-0191.

were covered with silastic tubing so that only the tip of the balloon was allowed to expand. The balloon was introduced through the right external jugular vein, and was advanced until the tip lay near the RSVC/AJ. The tubing was anchored to the right clavicle, as well as the muscle near the external jugular vein, and was subcutaneously tunneled to the nape of the neck and secured.

After a 1-week recovery period, drinking tests were begun. The atrial balloon was inflated in half of the rats in each group. This was accomplished by manual infusion of 0.05 ml water immediately prior to isoproterenol (30 $\mu\text{g}/\text{kg}$ s.c.) administration. The balloons remained expanded throughout the 120-min test session. Rats in the balloon deflation condition simply received the injection of isoproterenol. Cumulative water intake was recorded at 15, 30, 60 and 120 min after dipsogen delivery. The balloons were deflated at the end of the test period. The conditions were reversed on a subsequent test day so that water intake in response to isoproterenol was measured in all rats under both balloon inflation and deflation conditions. Forty-eight hours intervened between consecutive tests.

Upon completion of testing, the lesioned and sham-lesioned animals were transcardially perfused with physiological saline followed by 10% buffered formalin. The brains were sectioned and stained with Cresyl violet in order to histologically verify the lesion placement. Histological criteria for a good lesion have been described previously [20]. The chest cavity of each animal was opened to reveal the right superior vena cava and atrium. The balloon was inflated and placement was visually verified. Rats that did not drink at least 1 ml during the balloon deflation condition were eliminated from the study. A repeated measure ANOVA was used to compare water intake during balloon inflation and deflation conditions for each group at each time point. Differences between group water intake were analyzed using one-way ANOVAs. Sta-




Fig. 2. Drinking in response to isoproterenol is shown for the VLLPBN (LESION) and sham (SHAM) lesioned rats at 15, 30, 60, and 120 min after dipsogen delivery. Water intake during the balloon inflation condition is depicted by hatched bars, solid bars represent water intake while the balloon was deflated. * Indicates a significant difference between inflation and deflation conditions. Significance is $P < 0.05$.

tistical significance was defined as $P < 0.05$. All values are reported as means \pm S.E.M.

Six rats were determined to have both good VLLPBN lesions and accurate balloon placements. The lesions are comparable to those presented in previous publications [20,21] from our laboratory and a representative VLLPBN lesion from the present study is presented in Fig. 1. The brains of sham-lesioned animals were examined to ensure there was no damage to the VLLPBN, and balloon placement was also verified. Eleven rats met both criteria. Water intakes are presented in Fig. 2. Very little drinking occurred within the first 15 min following isoproterenol administration and balloon inflation did not significantly alter water intake in either the lesioned (inflation: 1.4 ± 0.9 ml; deflation: 0.4 ± 0.4 ml) or sham-lesioned (inflation: 0.1 ± 0.1 ml; deflation: 0.4 ± 0.3 ml) group. However, 30 min after drug administration, water intake was significantly decreased in sham-lesioned rats during balloon inflation (inflation: 1.1 ± 0.4 ml; deflation 3.0 ± 0.7 ml).



Fig. 1. Typical bilateral lesions (arrows) of the ventrolateral portion of the lateral parabrachial nuclei (VLLPBN).

Balloon expansion had no effect on drinking behavior of the VLLPBN-lesioned rats at this time point (inflation: 5.1 ± 2.2 ml; deflation: 5.3 ± 1.6 ml). A similar trend was seen at 60 min post-isoproterenol delivery. Drinking in the sham-lesioned rats was significantly suppressed by RSVC/AJ stimulation (3.3 ± 0.9 ml) as compared to the control condition (6.7 ± 1.8 ml), whereas water intake in the lesion group was unaltered by balloon expansion (inflation: 10.3 ± 2.3 ml; deflation: 10.2 ± 1.9 ml). By the end of the experiment, 2 h after isoproterenol administration, water intake was no longer significantly decreased by RSVC/AJ expansion in the sham-lesioned rats (inflation: 5.5 ± 1.8 ml; deflation: 7.2 ± 1.7 ml). Water intake in the VLLPBN-lesioned rats remained unaffected by this stimulus (inflation: 11.0 ± 2.5 ml; deflation: 10.8 ± 2.4 ml). Although there was a tendency for the VLLPBN-lesioned rats to drink more than the sham-lesioned animals, this trend did not reach statistical significance during the balloon deflation condition.

These results confirm previous work indicating that isoproterenol-induced water intake is attenuated by stretch of the RSVC/AJ [10,19] and suggests that the VLLPBN is critically involved in the mediation of the attenuation of drinking to this manipulation. It is unlikely that the reduced drinking exhibited by the sham-lesioned rats was the result of a debilitating effect of balloon expansion, as water intake was unaltered by this manipulation in the VLLPBN-lesioned rats. These data support the hypothesis that activation of cardiopulmonary volume receptors by stretch of the RSVC/AJ signals contribute to the termination of drinking, but it appears that this inhibitory signal adapts. That is, by the end of the experiment, the cumulative water intake of the sham-lesioned rats was not different between experiment conditions. Adaptation and resetting of vascular stretch receptors on both the high and low pressure sides of the circulation are well-described phenomena (see Refs. [2] and [27] for reviews).

It is reasonable to hypothesize that the VLLPBN is involved in volume receptor-induced inhibition of water intake. This region is heavily innervated by both the area postrema and nucleus of the solitary tract [16,26,28], nuclei known to receive input from cardiac and aortic arch baroreceptors via the vagus and glossopharyngeal nerves [3,9]. In addition, the lateral subdivisions of the parabrachial nucleus share reciprocal projections with the median preoptic nucleus, paraventricular nucleus, central nucleus of the amygdala, and other forebrain nuclei involved in the control of body fluid balance [6,17]. Therefore, the LPBN appears to be a component of a neural network capable of integrating information related to cardiovascular status and body fluid balance. Furthermore, the VLLPBN has been previously implicated in the attenuation of drinking. Bilateral electrolytic or chemical lesions of this nucleus augment drinking in response to centrally and peripherally administered dipsogens which simulate an extracellular fluid deficit [4,20,21]. It will be necessary to repeat the

present balloon inflation studies in rats where chemical lesions (e.g., as in Ref. [4]) are used in order to discern whether the present effects are due to destruction of neurons with soma located in the LPBN or disruption of fibers of passage.

While the data presented here suggest that the VLLPBN plays a role in the volume receptor-induced attenuation of water intake in response to isoproterenol, it is not clear whether this is a neurally or humorally mediated event. There is evidence to suggest that the observed decrease in drinking is the result of direct neural stimulation of central nervous system (CNS) centers involved in drinking. Moore-Gillon and Fitzsimons [19] reported that inflation of a balloon implanted at the left pulmonary vein and atrial junction in the dog reduced water intake in response to isoproterenol and that blockade of the left vagus nerve prohibited this inhibitory action. Further, unilateral cervical vagotomy significantly increased water intake in the rat following the administration of isoproterenol [18]. Together these findings suggest that decreased drinking following volume receptor stimulation is the result of impulses traveling to the CNS via the vagus nerve.

Alternatively, it is possible that this information is relayed to the brain via a humoral rather than neural mechanism. Isoproterenol-induced water intake is believed to be largely mediated by the renin-angiotensin system [5,8,23]. Left atrial stretch has been demonstrated to inhibit renin release in the dog [15,25], suggesting that the observed attenuation of water intake might be due to reduced plasma renin levels. This is unlikely, however, since Kaufman found no significant change in plasma renin activity during right atrial expansion by balloon in the rat [11]. However, atrial natriuretic peptide (ANP) is also released by atrial distention, and may underlie this phenomenon [14]. Administration of ANP has been demonstrated to attenuate drinking in response to a number of dipsogenic challenges [1,12,13], and water intake itself has been reported to elevate circulating levels of the peptide [24]. A high density of ANP binding sites has been found in the area postrema, a circumventricular organ [22], and circulating ANP acting through this nucleus may activate CNS pathways involving the VLLPBN to inhibit drinking. It should be noted, however, that RSVC/AJ distention does not diminish water intake in response to all dipsogenic stimuli [10], which would be expected if ANP release via atrial stretch was solely responsible for the termination of drinking.

In summary, activation of RSVC/AJ receptors has been shown to attenuate isoproterenol-induced water intake in intact rats and this inhibition is abolished by ablation of the VLLPBN. These findings support the hypothesis that the VLLPBN is involved in the inhibition of drinking mediated by volume receptor stimulation. These results add to the growing body of evidence suggesting that the VLLPBN is involved in the termination of thirst stimulated by a variety of dipsogenic challenges.

Acknowledgements

This research was supported by National Heart, Lung and Blood Institute, NIH Grants HL 14338 and HL 44546.

References

- [1] Antunes-Rodrigues, J., McCann, S.M., Rogers, L.C. and Samson, W.K., Atrial natriuretic factor inhibits dehydration- and angiotensin II-induced water intake in the conscious, unrestrained rat, *Proc. Natl. Acad. Sci. USA*, 82 (1985) 8720–8723.
- [2] Chapeau, M.W. and Abboud, F.M., Mechanisms of adaptation and resetting of baroreceptor reflex. In R. Hainsworth and A.L. Mark (Eds.), *Cardiovascular Reflex Control in Health and Disease*, W.B. Saunders, Philadelphia, 1993, pp. 165–193.
- [3] Ciriello, J. and Calaresu, F., Projections from buffer nerves to the nucleus of the solitary tract: an anatomical and electrophysiological study in the cat, *J. Auton. Nerv. Syst.*, 3 (1981) 299–310.
- [4] Edwards, G.L. and Johnson, A.K., Enhanced drinking after excitotoxin lesions of the parabrachial nucleus in the rat, *Am. J. Physiol.*, 261 (1991) R1039–R1044.
- [5] Evered, M.D. and Robinson, M.M., Increased or decreased thirst caused by inhibition of angiotensin-converting enzyme in the rat, *J. Physiol.*, 348 (1984) 573–588.
- [6] Fulwiler, C.E. and Saper, C.B., Subnuclear organization of the efferent connections of the parabrachial nucleus in the rat, *Brain Res. Rev.*, 7 (1984) 229–259.
- [7] Hosutt, J.A., Rowland, N. and Stricker, E.M., Hypotension and thirst in rats after isoproterenol treatment, *Physiol. Behav.*, 21 (1978) 593–598.
- [8] Houpt, K.A. and Epstein, A.N., The complete dependence of beta-adrenergic drinking on the renal dipsogen, *Physiol. Behav.*, 7 (1971) 897–902.
- [9] Kalia, M. and Sullivan, J.M., Brainstem projections of sensory and motor components of the vagus nerve in the rat, *J. Comp. Neurol.*, 211 (1982) 248–264.
- [10] Kaufman, S., Role of right atrial receptors in the control of drinking in the rat, *J. Physiol.*, 349 (1984) 389–396.
- [11] Kaufman, S., Influence of right atrial stretch on plasma renin activity in the conscious rat, *Can. J. Physiol. Pharmacol.*, 65 (1987) 257–259.
- [12] Kaufman, S. and Monckton, E.A., Effect of peripherally administered atriopeptin III on water intake in rats, *J. Physiol.*, 396 (1988) 379–387.
- [13] Lappe, R.W., Dinish, J.L., Bex, F., Michalak, K. and Wendt, R.L., Effects of atrial natriuretic factor on drinking responses to central angiotensin II, *Pharmacol. Biochem. Behav.*, 24 (1986) 1573–1576.
- [14] Ledsome, J.R., Wilson, N., Courneya, C.A. and Rankin, A.J., Release of atrial natriuretic peptide by atrial distention, *Can. J. Physiol. Pharmacol.*, 63 (1985) 739–742.
- [15] Lee, M.E., Thrasher, T.N. and Ramsay, D.J., Elevated cardiac pressure inhibits renin release after atrial hypotension in conscious dogs, *Am. J. Physiol.*, 247 (1984) R953–R959.
- [16] Loewy, A.D. and Burton, H., Nuclei of the solitary tract: efferent connections to the lower brain stem and spinal cord of the cat, *J. Comp. Neurol.*, 181 (1978) 421–450.
- [17] Loewy, A.D. and McKellar, S., The neuroanatomical basis of central cardiovascular control, *Fed. Proc.*, 39 (1981) 2495–2503.
- [18] Moore-Gillon, M.J., Effects of vagotomy on drinking in the rat, *J. Physiol.*, 308 (1980) 417–426.
- [19] Moore-Gillon, M.J. and Fitzsimons, J.T., Pulmonary vein atrial junction stretch receptors and the inhibition of drinking, *Am. J. Physiol.*, 242 (1982) R452–R457.
- [20] Ohman, L.E. and Johnson, A.K., Lesions in lateral parabrachial nucleus enhance drinking to angiotensin II and isoproterenol, *Am. J. Physiol. (Regulatory Integrative Comp. Physiol. 20)*, 251 (1986) R504–R509.
- [21] Ohman, L.E. and Johnson, A.K., Brain stem mechanisms and the inhibition of angiotensin-induced drinking, *Am. J. Physiol. (Regulatory Integrative Comp. Physiol. 25)*, 256 (1989) R264–R269.
- [22] Quirion, R., Dalpé, M., De Lean, A., Gutkowska, J., Cantin, M. and Genest, J., Atrial natriuretic factor (ANF) binding sites in brain and related structures, *Peptides*, 5 (1984) 1167–1172.
- [23] Rettig, R., Ganten, D. and Johnson, A.K., Isoproterenol-induced thirst: renal and extra-renal mechanisms, *Am. J. Physiol. (Regulatory Integrative Comp. Physiol. 10)*, 241 (1981) R152–R157.
- [24] Salazar, F.J., Granger, J.P., Joyce, M.L.M., Burnett, Jr., J.C., Bove, A.A. and Romero, J.C., Effects of hypertonic saline infusion and water drinking on atrial peptide, *Am. J. Physiol. (Regulatory Integrative Comp. Physiol. 20)*, 251 (1986) R1091–R1094.
- [25] Schultz, H.D., Fater, D.C., Sundet, W.D., Geer, P.G. and Goetz, K.L., Reflexes excited by acute stretch of atrial vs pulmonary receptors in conscious dogs, *Am. J. Physiol. (Heart Circ. Physiol. 11)*, 242 (1982) H1065–H1076.
- [26] Shapiro, R.E. and Miselis, R.R., The central neural connections of the area postrema of the rat, *J. Comp. Neurol.*, 234 (1985) 344–364.
- [27] Thoren, P., Resetting of cardiogenic reflexes in hypertension. In R. Hainsworth and A.L. Mark (Eds.), *Cardiovascular Reflex Control in Health and Disease*, W.B. Saunders, Philadelphia, 1993, pp. 195–216.
- [28] Vander Kooy, D. and Koda L.Y., Organization of the projections of a circumventricular organ: the area postrema in the rat, *J. Comp. Neurol.*, 219 (1983) 328–338.

Elsevier Science B.V.

Fax: (31) (20) 485 3239

Phone: (31) (20) 485 3462

Postal Address:

Brain Research

Elsevier Science B.V.
P.O. Box 1527, 1000 BM Amsterdam
The Netherlands

Courier Service Address:

Brain Research

Elsevier Science B.V.
Molenwerf 1, 1014 AG Amsterdam
The Netherlands

* * *

If you need information about your accepted manuscript, proof, etc. then phone or FAX us at the above numbers, stating the journal name and article code number. We can also FAX this journal's Instructions to Authors to you.

NEW AND FORTHCOMING TITLES IN ELSEVIER'S NEUROSCIENCE PROGRAMME

Progress in Brain Research Series

Neuroscience: From the Molecular to the Cognitive — Vol. 100, edited by F. Bloom

Biological Function of Gangliosides — Vol. 101, edited by L. Svennerholm

The Self-Organizing Brain — Vol. 102, edited by M.A. Corner and J. van Pelt

Neural Regeneration — Vol. 103, edited by F.J. Seil

Neuroscience Protocols — Module 3, edited by F. Wouterlood

Neuroscience Protocols — Module 4, edited by F. Wouterlood

OPAL Module 3: Immune Cells, by H.-P. Hartung and X. Zhang

OPAL Module 4: Endocrine Cells, by B. Cooke

For more information or a free OPAL Preview Disk, please contact Joyce Hobbelink, Elsevier Science, PO Box 1527, 1000 BM Amsterdam, The Netherlands, Fax: (31) (20) 485 3342, e-mail: j.hobbelink@elsevier.nl

SUBSCRIPTION AND PUBLICATION DATA 1995

Brain Research (including **Molecular Brain Research**, **Developmental Brain Research**, **Cognitive Brain Research** and **Brain Research Reviews**) will appear weekly and be contained in 53 volumes (108 issues): **Brain Research**, Volumes 669–705 (37 volumes in 74 issues), **Molecular Brain Research**, Volumes 26–32 (7 volumes in 14 issues), **Developmental Brain Research**, Volumes 84–90 (7 volumes in 14 issues), **Cognitive Brain Research**, Volume 3 (1 volume in 4 issues) and **Brain Research Reviews**, Volumes 20–21 (2 volumes in 6 issues). Please note that Volumes 26 and 27 of **Molecular Brain Research** were published ahead of schedule in 1994, in order to reduce publication time. The volume remains part of the 1995 subscription year. **Discussions in Neuroscience**, Volume 11 (1 volume in 4 issues) will be mailed as an integral part to all full-set subscribers to **Brain Research** in 1995. **Separate subscriptions: Molecular Brain Research**, Vols. 26–32, **Developmental Brain Research**, Vols. 84–90, **Cognitive Brain Research**, Vol. 3, and **Brain Research Reviews**, Vols. 20 and 21, may also be ordered separately. Prices are available from the Publisher upon request. Subscriptions are accepted on a prepaid basis only, unless different terms have been previously agreed upon.

Subscription orders can be entered only by calendar year (Jan.–Dec.) and should be sent to Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands, Tel.: (31) (20) 485 3642, Fax: (31) (20) 485 3598, or to your usual subscription agent.

Postage and handling charges include surface delivery except to the following countries where air delivery via SAL (Surface Air Lift) mail is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, P.R. China, Singapore, South Africa, South Korea, Taiwan, Thailand and USA. For all other countries airmail rates are available upon request.

Claims for missing issues must be made within six months of our publication (mailing) date, otherwise such claims cannot be honoured free of charge. In the United States and Canada: for further information concerning this or any other Elsevier Science journal, contact Elsevier Science Inc. Attn: Journal Information Center, 655 Avenue of the Americas, New York, NY 10010, U.S.A. Tel.: (1) (212) 633-3750; Telefax: (1) (212) 633-3990; Telex: 420-643 AEP UI.

Information for Advertisers. Advertising orders and enquiries can be sent to Elsevier Science B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands, Tel.: (31) (20) 485 3796; Fax: (31) (20) 485 3810. Office address for courier services: Molenwerf 1, 1014 AG Amsterdam, The Netherlands. For the U.K.: T.G. Scott and Son Ltd., Vanessa Bird, Portland House, 21 Narborough Road, Cosby, Leicestershire LE9 5TA, U.K., Tel.: (44) (116) 2750 520/2753 333; Fax: (44) (116) 2750 522. For the U.S.A. and Canada: Weston Media Associates, Daniel Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, U.S.A., Tel.: (1) (203) 261-2500; Fax: (1) (203) 261-0101.

ADONIS Identifier. This Journal is in the ADONIS Service, whereby copies of individual articles can be printed out from CD-ROM on request. An explanatory leaflet can be obtained by writing to ADONIS B.V., P.O. Box 839, 1000 AV Amsterdam, The Netherlands.

