

Chiron and the Centaurs: escapees from the Kuiper belt

Alan Stern & Humberto Campins

10-3-92
10-3-92
013373

The Centaurs—a group of objects orbiting chaotically among the giant planets of our Solar System—appear to be a population transitional in size between typical short-period comets and the large Kuiper-belt objects that orbit beyond Neptune. They promise to reveal much about the origin of and interrelationships between the icy bodies of the outer Solar System.

THE outer Solar System has long appeared to be a largely empty place, inhabited only by the four giant planets, Pluto and a transient population of comets. In 1977 however, a faint and enigmatic object—2060 Chiron—was discovered¹ moving on a moderately inclined, strongly chaotic 51-year orbit which takes it from just inside Saturn's orbit out almost as far as that of Uranus. It was not initially clear from where Chiron originated.

Following Chiron's discovery, almost 15 years elapsed before other similar objects were discovered; five more have now been identified¹. Based on the detection statistics implied by these discoveries, it has become clear that these objects belong to a significant population of several hundred (or possibly several thousand) large icy bodies moving on relatively short-lived orbits between the giant planets. This new class of objects, known collectively as the Centaurs, are intermediate in diameter between typical comets (1–20 km) and small icy planets such as Pluto (~2,300 km) and Triton (~2,700 km). Although the Centaurs are interesting in their own right, they have taken on added significance following the recognition that they most probably originated in the ancient reservoir of comets and larger objects located beyond the orbit of Neptune known as the Kuiper belt.

Origin of the Centaurs

The first clue to the origin of the Centaurs came about as a result of dynamical studies of Chiron's orbit. At first discovered in the late 1970s³, and forcefully reiterated in more modern calculations⁴, Chiron's orbit is highly unstable to perturbations by the giant planets. As a result, Chiron's orbital lifetime among the giant planets is short, leading to the conclusion that its origin was in a more stable reservoir, either in the asteroid belt, or beyond the giant planets. The discovery of a coma around Chiron^{5–7}, in the late 1980s, indicated the presence of surface volatiles which could not have survived the age of the Solar System in the comparatively warm asteroid belt⁸; such volatiles therefore strongly indicate that Chiron originated in a distant reservoir, beyond the giant planets.

A second line of evidence relating to Chiron's origin came about from simulations of cometary dynamics. These studies^{9–12} demonstrated that the dominant dynamical class of short-period comets, called the Jupiter-family comets (JFC) cannot be derived from the classical, Oort-cloud cometary reservoir. The reason for this is that their characteristically low orbital inclinations cannot be efficiently produced by the action of planetary perturbations on orbits initially in the inclination-randomized (that is, nearly spherical) Oort cloud.

Instead, the JFC seem to derive from a dynamically stable reservoir concentrated near the plane of the planetary system. Any such reservoir for the JFC must satisfy the criterion that the loss rate of objects from it be low enough that the reservoir can persist for the age of the Solar System. Because the dynamical clearing time for orbits between the planets is characteristically one to two orders of magnitude shorter than the age of the Solar System^{12–14}, there are few regions of space that provide stable,

candidate source locations for the JFC. One such region is the zone beyond Neptune's orbit (at 30 AU, astronomical units) where nonlinear perturbations by Neptune and the other giant planets can excite orbital eccentricities on timescales comparable to the age of the Solar System. Once orbital eccentricities are excited sufficiently to cause objects to cross Neptune's orbit, a fraction of these objects become temporarily trapped on Centaur-like orbits among the giant planets. Other such objects are dynamically transported by subsequent encounters with the other giant planets onto orbits that pass within 1–2 AU of the Sun¹⁵, where they generate comae and become easily detectable. A second possible source region for the JFC is the slowly dynamically evaporating jovian Trojan clouds, whose dynamics are controlled by the stability of a narrow phase space surrounding the leading and trailing lagrangian points of Jupiter. However, recent dynamical simulations¹⁶ show that the jovian Trojan clouds are not as effective as the so-called trans-neptunian zone in populating the JFC and Centaur populations. Following historical suggestions dating back as far as the 1940s that a disk-shaped reservoir of planetesimals and other small objects might reside beyond the Neptune^{17,18}, the trans-neptunian region has been dubbed the Kuiper belt, or in analogy to debris belts around the other stars, the Kuiper disk.

The pivotal breakthrough concerning the reality of the hypothesized Kuiper belt came in 1992, with the discovery of a faint (R-band magnitude near 23), 180-km diameter object¹⁹ designated 1992QB₁. 1992QB₁ orbits the Sun in a stable, nearly circular orbit some 14 AU beyond Neptune. In the four years since 1992QB₁ was found, over three dozen similar objects have been discovered in the trans-neptunian region². Estimates^{2,20} indicate that some of these objects have diameters approaching 400 km. Based on the efficiency with which such objects are being detected and their surface density on the sky, it has been estimated that around 7×10^4 objects with diameters greater than 100 km orbit between 30 and 50 AU (ref. 2). Here we refer to these larger objects populating the Kuiper Belt as QB_s.

Following the discoveries of numerous QB_s in the Kuiper belt, the Hubble Space Telescope (HST) was used to conduct a search for the much smaller, and much fainter, cometary nuclei which must be present in this region if it is indeed a source of the JFCs. Last year, Cochran *et al.*²¹ reported exciting evidence, near the limit of HST's capabilities, for numerous objects with V-band magnitudes of ~28.6, corresponding to comet-like diameters of a few kilometres to ~10 km. This evidence corresponds to a population of several hundred million comet-sized objects. If this result is coupled with models¹² that predict the ratio of the population detected in the region searched by Cochran to the entire trans-neptunian zone, a total population is calculated of several billion comets in the 30–50 AU region. As such, it appears that the Kuiper belt is indeed the source region of most JFC, as dynamical simulations predicted^{9,14}.

Taken together, the discovery of both QB_s and comet-sized

objects in the Kuiper-belt region indicates that the Kuiper belt supplies a wide size range of objects onto orbits in the giant-planet region. Based both on expectations resulting from the planetesimal accretion codes, and the observational evidence for many more comets than QB_s, it appears that a power-law-like source population histogram exists in the Kuiper belt, with many more small bodies than QB_s. Because the dynamical transport process that brings objects from the belt to planet-crossing orbits is essentially independent of the mass of the object being transported¹⁵, it is expected that the population of objects ranging in size from Centaurs down to JFC orbiting among the giant planets is representative of the population of objects in the 30–50 AU zone from which they are derived.

Physical attributes of the Centaurs

It is now established that the slow leakage of objects from the Kuiper belt due to planetary perturbations creates a population of objects on comparatively short-lived, planet-crossing orbits in the giant-planet region between 5 and 30 AU from the Sun. Studies of the dynamical evolution of orbits dislodged from the Kuiper belt¹⁵ predict a characteristic equilibrium population of objects on planet-crossing orbits that is $\sim 10^{-4}$ of the population of the Kuiper-belt reservoir from which they are derived. These studies also predict that the median lifetime of such orbits is of the order of 5×10^7 years. Such findings imply that a population of $\sim 5 \times 10^5$ to perhaps 10^6 comets, and ~ 30 – 300 Centaur objects of diameter 100 km or larger, are orbiting between the giant planets.

Chiron and its recently discovered cohort of Centaurs are thus now seen to be objects derived from the Kuiper belt. Table 1 summarizes the orbital attributes of the six known Centaurs; Fig. 1 depicts the orbits of these objects and their dynamical context in the outer Solar System.

As escapees from the Kuiper belt, the Centaurs are an important population for study. Indeed, owing to their greater proximity to the Sun, the brightest centaurs are some 5–7 astronomical magnitudes (factors of ~ 100 to ~ 600) brighter than the brightest

TABLE 1 Orbital characteristics of the known Centaurs

Object	Semi-major axis	Perihelion distance	Eccentricity	Inclination	Present opposition V magnitude
2060 Chiron	13.70 AU	8.46 AU	0.38	25°	15.5
5145 Pholus	20.30 AU	8.68 AU	0.57	7°	17.9
1993HA ₂	24.73 AU	11.84 AU	0.52	16°	21.0
1994TA	16.82 AU	10.69 AU	0.31	5°	23.8
1995DW2	25.03 AU	18.84 AU	0.25	4°	21.9
1995GO	18.14 AU	6.79 AU	0.62	18°	20.3

These characteristics are taken from ref. 53. Although the basic orbital properties of the third to sixth objects are known, they have not yet been named because, by IAU convention, objects must first be observed for long enough to produce astrometrically reliable orbits.

Kuiper-belt objects, which enables more detailed studies of the Centaurs than are possible with the QB_s and Kuiper-belt comets. Additionally, being closer to the Sun, the Centaurs experience greater heating, which generates characteristic perihelion surface temperatures in the range ~ 120 to 150 K (ref. 22); by contrast, Kuiper-belt objects probably never experience surface temperatures in excess of 60 – 70 K. Therefore, because vapour pressure depends exponentially on the temperature of the ice, Centaurs are much more likely than Kuiper-belt objects to show sublimation-generated activity. Although such heating causes the surfaces of the Centaurs to evolve chemically and physically over long time-scales²³, it also causes the surface ices to sublime, and thus reveal valuable insights into the nature of these objects.

Unfortunately, although the Centaurs are brighter than Kuiper-belt objects, they are still faint in absolute terms, so considerable dedication is required to obtain physical information on them. As a result, comparatively little work has been done to reveal their compositions, colours, shapes, rotational properties and other attributes (Table 2). Despite the great deficits in our knowledge about the physical and chemical characteristics of this unique population, several important pieces of information are emerging.

First, with regard to the derived sizes of the Centaurs discovered to date, roughly half appear to be near 60 km in diameter, but 2060 Chiron^{22,24} and 5145 Pholus²⁵ are much larger, with ~ 180 -km diameters that are comparable to typical QB_s being discovered in the trans-neptunian zone. Second, infrared spectroscopy and colour photometry have given the first clues about the surface compositions of these objects. The first clearly detected spectral




FIG. 1 The orbits of the giant planets (black lines), the six known Centaurs (red lines) and those Kuiper-belt objects with well established orbits (green lines). The dot on each orbit depicts the current location of the object. For scale, Jupiter's orbit is approximately 10 AU across. Abbreviations on the figure as follows: GO, 1995GO; TA, 1994TA; DW2, 1995DW2; Chiron, 2060 Chiron; HA2, 1993HA₂; Pholus, 5145 Pholus. (Figure courtesy of H. F. Levison)

TABLE 2 Physical characteristics of the known Centaurs

Object	Diameter* (km)	Geometric albedo (in V)	Rotational period	Rotational amplitude	V - J colour†	Detected activity
2060 Chiron	182 ± 10	0.11–0.20	5.92 h	9%	1.13 ± 0.04	Yes
5145 Pholus	185 ± 22	0.04 ± 0.02	9.98 h	20%	2.53 ± 0.06	No
1993HA ₂	62 km†	0.05			2.07 ± 0.40	No
1994TA	28 km†	0.05				No
1995DW2	68 km†	0.05				No
1995GO	60 km†	0.05				No

* The sizes of Chiron and Pholus were obtained^{27,24,25} from thermal fluxes, with computed albedos based on their sizes and V magnitudes. Chiron's albedo is strictly an upper limit owing to a possible, small, residual coma contribution.

† These sizes were computed by assuming a V-band geometric albedo of 5%, a value which is commonly found for cometary surfaces.

‡ Colour data are discussed in the text^{27,31,32}. Note a V - J colour of 1.116 would be identical to the Sun⁴⁷, thereby indicating a neutrally coloured surface. Higher V - J colours indicate red surfaces.

absorption feature among the Centaurs was a deep 2.25- μ m absorption on Pholus²⁶. This feature has been associated with light organic solids mixed with ices^{27,28}. Importantly a 2.04- μ m absorption features has also now been detected, both in Pholus and Chiron²⁹, which Cruikshank *et al.* identify³⁰ in Chiron as an absorption band of water-ice. Third, although none of the three Centaurs that have been explored in the infrared have displayed any statistically significant colour variation with rotational phase³¹, they do show striking colour differences between one another (refs 27, 32 and Fig. 2). Indeed, whereas Chiron's intrinsic colour is grey (that is, neutral) throughout the 0.3–2.5 μ m band, Pholus, which lies in a similar orbit and is similar in size, is extremely red. 1993HA₂, though smaller and in a more distant orbit, is also very red compared to Chiron. How much of these differences between various Centaurs is due to evolutionary mechanisms (as opposed to intrinsic attributes) is not yet clear, but it is well established that long-term exposure to cosmic rays and solar ultraviolet radiation darkens (and initially reddens) surfaces containing light-weight organics, in turn creating a more complex chemical mélange.

Additionally, the determination of well constrained albedos for 2060 Chiron, and more particularly 5145 Pholus (because it is not active), provides useful information for predicting the sizes of QB₁s from their observed magnitudes.

The particular value of Chiron

Chiron is uniquely valuable among the Centaurs because of the long history of its sporadic outbursts. Why has such activity only been detected in Chiron? Possibly Chiron is dynamically younger, and therefore more active than the other objects. Alternatively the other objects may have thicker surface mantles, may be fundamentally different in their composition or simply may have not been observed long enough to expect to detect activity. Chiron's

uniqueness in showing activity is perhaps the most intriguing observable obtained on the Centaurs so far.

Chiron's activity was first recognized when it suffered an outburst that increased its brightness by a factor (in 1989) of just over two^{33,34}. In 1989–91 Chiron was also observed to show a highly variable particulate coma and tail extending as far as 2×10^6 km (refs 35, 36), and a cloud of CN gas³⁷ presumably derived from the photodissociation of some heavier, parent molecule. When these various observations were made, Chiron was still more than 10 AU from the Sun, where the solar radiation field is too weak to sublime water-ice, the common volatile that powers the cometary activity close to the Sun. Although other mechanisms remain plausible, the sublimation of highly volatile ices like CO, N₂ or CH₄ (buried a short distance below the surface) were therefore flavoured as the source of Chiron's activity. Further evidence for the sublimation of such volatiles was obtained through the discovery of even more extreme activity on archival, pre-discovery images of Chiron obtained when it was near its aphelion at 19 AU, and therefore far too cold to sublime anything but highly volatile ices like those mentioned above³⁸. The final confirmation of this hypothesis came in 1995, though the discovery of CO gas itself in Chiron's coma^{39,40}.

The fact that Chiron's activity was greater at its aphelion than it has been at any time since provides compelling evidence that its level of activity is not a simple function of heliocentric distance alone. Instead, Chiron's activity probably involves a complex interaction between the level of insolation reaching its surface, the obliquity of its spin axis, the location of its near-surface volatiles and extensive surface mantling by substances (possibly including silicates, water-ice and carbonaceous materials) which do not strongly sublime that far from the Sun.

Chiron's strong variability and the low gas-production rate inferred from CN and CO gas detections in its coma provide




FIG. 2 Colour-albedo diagram showing the visible-band albedos and visible-infrared (that is, V - J) colours for several Centaurs (Chiron, Pholus and 1993HA₂), several cometary nuclei (Arend-Rigaux, Halley and Neujmin 1), Pluto and the one QB₁ (1993FW) for which applicable data are available. A neutrally coloured surface would have the V - J colour of the Sun, 1.116 (ref. 54). Note that the Centaur Pholus is quite red; in fact, it is far redder than any other object in the Solar System. The V - J colour for 1993FW is an upper limit. V - J for Pluto was obtained from D. P. Cruikshank (personal communication; the V-albedo error bars for Pluto represent its intrinsic rotational lightcurve variation).

strong evidence that Chiron's activity is generated by localized sources covering < 1% of the surface. The case supporting highly localized vents or jets on the surface is further supported by short-term brightness fluctuations in Chiron's coma^{41,42} that occur on timescales consistent with clouds of material being ejected onto suborbital trajectories, and by the detection of complex opacity structures in Chiron's coma during two recently observed stellar occultations by Chiron^{24,43}. It has been pointed out⁴⁴ that these vents or jets may resemble the geysers detected on the surface of Neptune's large, captured satellite, Triton.

Chiron's low gravity creates a situation in which its escape speed ($\sim 10^2 \text{ m s}^{-1}$) is comparable to both the thermal velocity of sublimating gas ($\sim 2 \times 10^2 \text{ m s}^{-1}$), and the estimated muzzle velocity of Triton-like geysers⁴⁵ (of the order of 40–300 m s^{-1}). As a result, some of the gas and entrained particulates ejected from Chiron's surface would be deposited into high suborbital trajectories; much would also escape. Modellers have only begun to explore the range of interesting physical phenomena likely to result in this intermediate regime between freely escaping cometary comae and strongly bound planetary atmospheres⁴⁶. Among these is the distinct possibility that Chiron's neutral colour and comparatively high albedo are the direct result of its activity, which probably causes a thin veneer of icy particulates to rain back onto the surface from suborbital trajectories.

An emerging view

We are witnessing a revolution in our understanding of the content and architecture of the outer Solar System. Whereas a decade ago, the outer Solar System seemed to consist only of the outer planets, the Oort-cloud comets and the then-roguish object Chiron, we now see revealed both the teeming Kuiper belt and its progeny, the comets and Centaur-sized objects orbiting among the outer planets. As a result, we have come to recognize that the

outer Solar System is littered with icy objects intermediate in size between comets and the giant planets.

We have also learned that the early stages of planetary formation, with widespread growth from planetesimals to objects with diameters of several hundreds of kilometres, provide concrete evidence for an ancient era of planet-building in the Kuiper-belt region^{47,48}. For some reason (probably involving the role of Neptune that excited orbital eccentricities that were not conducive to further growth), the era of accretion in the Kuiper belt was prematurely truncated at a stage where intermediate-sized objects had formed⁴⁹. The strong circumstantial evidence for the early formation of numerous objects in the 1,000-km class, of which Pluto and Triton are apparently the sole extant remnants within observational reach^{50,51}, further supports the case for initially strong but eventually arrested planetary accretion in the Kuiper-belt region^{48,52}. As such, the Kuiper belt has become one of the most important regions in the Solar System for studies of planetary origins. The Centaurs and QB_s therefore represent a valuable, relic population of icy objects whose growth was arrested at a fascinating, intermediate stage between comets and small planets.

The Centaurs also serve as bright proxies for distant comets, as laboratories for studying surface processes occurring on comets, Triton and perhaps Pluto, and as nearby proxies for the QB_s and other intermediate-scale bodies that bridge the size gap between comets, Pluto and Triton. As such, they hold special promise for understanding the origin and interrelationships among the icy bodies of the outer Solar System. □

A. Stern is at the Space Science Department, Southwest Research Institute, Boulder, Colorado 80302, USA. H. Campins is at the Department of Astronomy, University of Florida, Gainesville, Florida 32611, USA.

1. Kowal, C. T. *Icarus* **77**, 118–123 (1989).
2. Weissman, P. R. & Levison, H. F. in *Pluto & Charon* (eds Tholen, D. J. & Stern, S. A.) (Univ. Arizona Press, Tucson, in the press).
3. Okawa, S. & Everhart, E. *Astron. J.* **84**, 134–139 (1979).
4. Dones, L., Levison, H. F. & Duncan, M. D. in *The Content of the Solar System* (ed. Rettig, T.) (ASP Conf. Ser., in the press).
5. Tholen, D. J. et al. *IAU Circ. No.* 4554 (1988).
6. Hartmann, W. K., Cruikshank, D. P. & Tholen, D. J. *Icarus* **83**, 1–15 (1989).
7. Meech, K. J. & Belton, M. J. S. *Astron. J.* **100**, 1323–1338 (1990).
8. Stern, S. A. *Publ. Astron. Soc. Pacif.* **101**, 126–132 (1989).
9. Duncan, M. J., Quinn, T. & Tremaine, S. **328**, L69–L73 (1988).
10. Quinn, T. R., Tremaine, S. & Duncan, M. J. *Astrophys. J.* **355**, 667–679 (1990).
11. Levison, H. F. & Duncan, M. J. *Astrophys. J.* **406**, L35–L38 (1993).
12. Duncan, M. J., Levison, H. F. & Budd, S. M. *Astron. J.* **110**, 3073–3081 (1995).
13. Gladman, B. & Duncan, M. J. *Astron. J.* **100**, 1680–1693 (1990).
14. Wisdom, J. & Holman, M. *Astron. J.* **102**, 1528–1538 (1991).
15. Levison, H. F. & Duncan, M. J. *Icarus* (submitted).
16. Levison, H. F. & Shoemaker, E. *Icarus* (submitted).
17. Kuiper, G. P. in *Astrophysics: A Topical Symposium* (ed. Hynek, J. A.) 357–406 (McGraw Hill, New York, 1951).
18. Edgeworth, K. E. *Mon. Not. R. Astron. Soc.* **109**, 600–609 (1949).
19. Jewitt, D. C. & Luu, J. X. *Nature* **362**, 730–732 (1993).
20. Jewitt, D. C. & Luu, J. X. *Astron. J.* **109**, 1867–1873 (1995).
21. Cochran, A. L. et al. *Astrophys. J.* **455**, 342–346 (1995).
22. Campins, H. et al. *Astron. J.* **108**, 2318–2322 (1994).
23. Murma, M. J., Weissman, P. R. & Stern, S. A. in *Protostars and Planets III* (eds Levy, E. H. & Lunine, J. I.) 1177–1252 (Univ. Arizona Press, Tucson, 1993).
24. Bus, S. J. et al. *Icarus* (in the press).
25. Davies, J. K. et al. *IAU Circ. No.* 5698 (1996).
26. Davies, J. K. et al. *Icarus* **102**, 166–169 (1993).
27. Cruikshank, D. P. in *International School of Space Chemistry* (ed. Greenberg, M.) (1994).
28. Wilson, P. D., Sagan, C. & Thompson, W. R. *Icarus* **107**, 288–303 (1994).
29. Luu, J. X., Jewitt, D. C. & Cloutis, E. *Icarus* **108**, 133–144 (1994).
30. Cruikshank, D. P. et al. *Icarus* (submitted).
31. Buie, M. W. & Bus, S. J. *Icarus* **100**, 288–294 (1992).
32. Davies, J. K. & Tholen, D. J. in *The Content of the Solar System* (ed. Rettig, T.) (ASP Conf. Ser., in the press).
33. Tholen, D. J. et al. *IAU Circ. No.* 4554 (1988).
34. Hartmann, W. K., Cruikshank, D. P. & Tholen, D. J. *Icarus* **83**, 1–15 (1989).
35. Meech, K. J. & Belton, M. J. S. *Astron. J.* **100**, 1323–1338 (1990).
36. Larson, S. & Marcialis, R. L. *IAU Circ. No.* 5669 (1992).
37. Bus, S. J. *Science* **251**, 774–776 (1991).
38. Bus, S. J. et al. in *Proc. Distant Comet Activity Workshop* (eds Keller, H. U. & Huebner, W.) 41–43 (Southwest Res. Inst., San Antonio, Texas, 1993).
39. Womack, M. & Stern, S. A. *IAU Circ. No.* 6193 (1995).
40. Womack, M. & Stern, S. A. *Astrophys. J.* (submitted).
41. Luu, J. X. & Jewitt, D. C. *Astron. J.* **100**, 913–932 (1990).
42. Buratti, B. J. & Dunbar, R. S. *Astrophys. J.* **366**, L47–L49 (1991).
43. Elliot, J. L. et al. *Nature* **373**, 46–48 (1995).
44. Stern, S. A., Jackson, A. A. & Boice, D. C. *Astron. J.* **107**, 765–771 (1994).
45. Kirk, R. L. et al. in *Neptune & Triton* (ed. Cruikshank, D.) 949–989 (Univ. Arizona Press, Tucson, 1995).
46. Boice, D. C. et al. in *Proc. Distant Comet Activity Workshop* (eds Keller, H. U. & Huebner, W.) 134–139 (Southwest Res. Inst., San Antonio, Texas, 1993).
47. Stern, S. A. *Astron. J.* (in the press).
48. Davis, D. R. & Farinella, P. *Icarus* (submitted).
49. Stern, S. A. *Astron. J.* **110**, 856–865 (1995).
50. McKinnon, W. B. *Nature* **311**, 355–358 (1984).
51. Stern, S. A. *Icarus* **90**, 271–281 (1991).
52. Stern, S. A., McKinnon, W. B. & Lunine, J. I. in *Pluto & Charon* (eds Tholen, D. J. & Stern, S. A.) (Univ. Arizona Press, Tucson, in the press).
53. Marsden, B. & Williams, G. W. *Minor Planet Electronic Circular*, 1996-E02, 1996.
54. Campins, H., Rieke, G. H. & Lebofsky, M. J. *Astron. J.* **90**, 896–899 (1985).

ACKNOWLEDGEMENTS. We thank our colleagues D. Cruikshank, D. Jewitt, S. Larson, H. Levison, R. Marcialis, B. Marsden, W. McKinnon, W. Merlini, J. Parker and P. Weissman for comments, and A. Fitzsimmons for a helpful review.

CORRESPONDENCE should be addressed to A.S. (e-mail: astern@swri.edu)

