

Chimeric plant calcium/calmodulin-dependent protein kinase gene with a neural visinin-like calcium-binding domain

SHAMEEKUMAR PATIL, D. TAKEZAWA, AND B. W. POOVAIAH*

Laboratory of Plant Molecular Biology and Physiology, Department of Horticulture, Washington State University, Pullman, WA 99164-6414

Communicated by Edwin G. Krebs, University of Washington, Seattle, WA, February 24, 1995

ABSTRACT Calcium, a universal second messenger, regulates diverse cellular processes in eukaryotes. Ca^{2+} and Ca^{2+} /calmodulin-regulated protein phosphorylation play a pivotal role in amplifying and diversifying the action of Ca^{2+} -mediated signals. A chimeric Ca^{2+} /calmodulin-dependent protein kinase (CCaMK) gene with a visinin-like Ca^{2+} -binding domain was cloned and characterized from lily. The cDNA clone contains an open reading frame coding for a protein of 520 amino acids. The predicted structure of CCaMK contains a catalytic domain followed by two regulatory domains, a calmodulin-binding domain and a visinin-like Ca^{2+} -binding domain. The amino-terminal region of CCaMK contains all 11 conserved subdomains characteristic of serine/threonine protein kinases. The calmodulin-binding region of CCaMK has high homology (79%) to α subunit of mammalian Ca^{2+} /calmodulin-dependent protein kinase. The calmodulin-binding region is fused to a neural visinin-like domain that contains three Ca^{2+} -binding EF-hand motifs and a biotin-binding site. The *Escherichia coli*-expressed protein (≈ 56 kDa) binds calmodulin in a Ca^{2+} -dependent manner. Furthermore, ^{45}Ca -binding assays revealed that CCaMK directly binds Ca^{2+} . The CCaMK gene is preferentially expressed in developing anthers. Southern blot analysis revealed that CCaMK is encoded by a single gene. The structural features of the gene suggest that it has multiple regulatory controls and could play a unique role in Ca^{2+} signaling in plants.

Calcium plays a pivotal role as a second messenger by regulating many aspects of cellular signaling in plants and animals. The signal-induced changes in free Ca^{2+} concentration in a cell have been portrayed as a switch turning on various cellular processes (1–5). Protein phosphorylation is one of the major mechanisms by which eukaryotic cells transduce extracellular signals to intracellular responses (6, 7). Ca^{2+} and Ca^{2+} /calmodulin-dependent protein kinases are involved in amplifying and diversifying the action of Ca^{2+} -mediated signals (5, 6, 8–12). In animals, multifunctional Ca^{2+} /calmodulin-dependent protein kinase (CaMKII) is known to play a pivotal role in cellular regulation because of its ability to phosphorylate various proteins upon binding to Ca^{2+} /calmodulin (9, 13). Although not much is known about Ca^{2+} /calmodulin-dependent protein kinases in plants (5, 14, 15), Ca^{2+} -dependent calmodulin-independent protein kinases have been well documented (10, 11). These kinases are characterized by the presence of a calmodulin-like Ca^{2+} -binding domain. This report describes a chimeric Ca^{2+} /calmodulin-dependent protein kinase (CCaMK) gene[†] in plants with some of the structural features resembling both mammalian Ca^{2+} /calmodulin-dependent protein kinases and plant Ca^{2+} -dependent protein kinases.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

MATERIALS AND METHODS

Plant Material. Lily (*Lilium longiflorum* Thunb cv. Nellie White) plants were grown under greenhouse conditions and various parts were excised and frozen in liquid nitrogen.

PCR and cDNA Library Screening. Three lily cDNA libraries made from developing anthers and mature and germinating pollen were used for PCR. Degenerate oligonucleotides corresponding to two highly conserved regions, DLKPEN and FNARRKL, of mammalian Ca^{2+} /calmodulin-dependent protein kinases were used as primers for PCR (16). The amplification reaction mixture contained 1 \times PCR buffer (Cetus), all four dNTPs (each at 200 μM), 50 pmol of each primer, 1.5 mM MgCl_2 , 2 μl of cDNA library (10^9 plaque-forming units/ml), and 2.5 units of *Taq* DNA polymerase in a 100- μl total reaction volume. The cycling profile was 30 cycles of 94°C for 1 min, 48°C for 1 min, and 72°C for 1 min. The specific PCR product of the expected size (471 bp) was subcloned into pBluescriptII KS(+) (Stratagene) and sequenced. This fragment was used to screen the cDNA library (17) from developing anthers (18) to obtain the cDNA clone.

Sequence Analysis. The sequencing of the cDNA was carried out by using the Sanger dideoxynucleotide chain-termination method (19). A search of the GenBank data base (March 1994) was done by using Genetics Computer Group version 7.0 software (20).

The Expression of the CCaMK Gene. The RNase protection assay (17) was performed by using total RNA (20 μg) from various parts of the lily. Total RNA was isolated from leaf, stem, and various organs from immature flower (21). A 612-bp fragment of the CCaMK coding region (nt 1010–1621) was subcloned into pBluescriptII KS(+) plasmid (Stratagene) and used as a template for making the ^{32}P -labeled RNA probe.

Southern Blot Analysis. Lily genomic DNA (5 μg) was digested with various restriction enzymes and transferred to nylon membrane, and Southern blot analysis was carried out by using standard protocols (17).

Expression of CCaMK in *Escherichia coli*. The CCaMK protein was expressed in *E. coli* from the pET3b vector (22). *E. coli* BL21 (DE3)-pLysS was transformed with the pET3b expression vector containing CCaMK cDNA. Bacteria were grown at 35°C in M9 minimal medium supplemented with Casamino acids (2 g/liter), ampicillin (100 mg/liter), and chloramphenicol (25 mg/liter). The protein was induced by adding 0.5 mM isopropyl β -D-thiogalactoside when the OD₆₀₀ reached 0.5–0.7 unit. Three hours after induction, cells were collected by centrifugation, and the protein was then extracted and purified by using a calmodulin-Sepharose 4B affinity column as described by Hagiwara *et al.* (23). The quality of the purified protein was checked by SDS/PAGE.

Abbreviations: CCaMK, chimeric Ca^{2+} /calmodulin-dependent protein kinase; CaMKII, multifunctional Ca^{2+} /calmodulin-dependent protein kinase.

*To whom reprint requests should be addressed.

[†]The sequence reported in this paper has been deposited in the GenBank data base (accession no. U24188).

FIG. 2. Comparison of the deduced amino acid sequence of the C-terminal region (amino acids 338–520) of CCaMK to neural visinin-like Ca^{2+} -binding proteins. Conserved amino acids are boxed; Ca^{2+} -binding domains (I–III) are indicated by solid lines; putative autophosphorylation site is indicated by an asterisk; and the putative biotin-binding site (B) is indicated by a hatched bar. Rahc1, rat hippocalcin (Gen2:Ratp23K); Rav13, rat neural visinin-like protein (Gen2:Ratnvp3); Bov11, bovine neurocalcin (Gen1:Bovpcaln); Rav11, rat neural visinin-like protein (Gen2:Ratnvp1); Chv11, chicken visinin-like protein (Gen2:Ggvilip); Rav12, rat neural visinin-like protein (Gen2:Ratnvp2); Drfr1, *Drosophila* frequenin (Gen2:Drofreq).




FIG. 3. Schematic representation of CcCaMK showing various structural features.

oligonucleotide primers corresponding to two highly conserved regions of mammalian Ca^{2+} /calmodulin-dependent protein kinases. This fragment was not amplified when the cDNA libraries made from mature and germinating pollens were used. The nucleotide sequence of the PCR-amplified fragment contained conserved sequences corresponding to catalytic subdomains VI–XI and part of the calmodulin-binding domain of mammalian CaMKII (28).

A cDNA clone of CcCaMK (2514 bp) was obtained by screening the cDNA library by using the PCR-amplified fragment as a probe and its nucleotide sequence was determined (Fig. 1). The cDNA codes for a polypeptide of 520 amino acids flanked by a 634-bp untranslated region at the 5' end and a 317-bp untranslated region at the 3' end. This polypeptide contains all 11 major conserved subdomains of the catalytic domain of serine/threonine kinases (28). Sequence comparisons revealed that CcCaMK has high homology to Ca^{2+} /calmodulin-dependent protein kinases, especially in the kinase and the calmodulin-binding domains (amino acids 1–338). This region of CcCaMK has highest homology to kinases from apple (Gen3:Mdstpkn), rat (Gen2:Ratpk2g), human (Gen1:Humccdpkb), and fruit fly (Gen2:Drocdpdk), where data in parentheses are names for corresponding genes in GenBank.

The calmodulin-binding region of CcCaMK (ARRKLRAAA-IASVL, residues 325–338) has 79% similarity to the calmodulin-binding domain (ARRKLKGAILTTML, residues 296–309) of α subunit of mammalian CaMKII, a well-characterized Ca^{2+} /calmodulin-dependent protein kinase (29). However, the calmodulin-binding domain of CcCaMK has 43% and 50% similarity to the calmodulin-binding domains of CaMKII homologs of yeast and *Aspergillus*, respectively (30, 31). The helical wheel projection of the calmodulin-binding domain (amino acids 325–338) of CcCaMK formed a basic amphipathic

α -helix (32), a characteristic feature of calmodulin-binding sites (data not shown).

The sequence downstream of the calmodulin-binding region of CcCaMK (amino acids 339–520) does not have significant homology to known Ca^{2+} /calmodulin-dependent protein kinases. Further analysis of this region revealed the presence of three Ca^{2+} -binding EF-hand motifs that had the highest homology (52–54% similarity; 32–35% identity) to a family of genes belonging to visinin-like Ca^{2+} -binding proteins (Fig. 2), found mainly in neural tissue (33–37). Even though four EF-hand motifs are present in the calmodulin-like domain of Ca^{2+} -dependent calmodulin-independent protein kinases, this domain shared only 25% identity with the visinin-like domain of CcCaMK. Out of the six residues of the EF-hand [positions 1(X), 3(Y), 5(Z), 7(–Y), 9(–X), and 12(–Z), where X, Y, and Z refer to Ca^{2+} ligating residues] involved in Ca^{2+} binding, position 7(–Y) is not conserved in CcCaMK. A similar deviation is also observed in visinin-like proteins, wherein the residue at position 9(–X) of the EF-hand motifs of visinin-like proteins (Fig. 2) is not conserved. These differences between the EF-hands of the visinin-like domain of CcCaMK and other Ca^{2+} -binding proteins may affect Ca^{2+} -binding and protein–protein interactions.

Frequenin, neurocalcin, hippocalcin, and visinin-like neural Ca^{2+} -binding proteins are members of a family of Ca^{2+} -sensitive regulators, each with three Ca^{2+} -binding EF-hand motifs. The presence of such proteins has not been reported in plants. These proteins are activated at nanomolar concentrations of Ca^{2+} . At such low levels, calmodulin-dependent pathways are not activated. Frequenin acts as a Ca^{2+} -sensitive activator of a photoreceptor particulate guanylyl cyclase (37). It has also been suggested that frequenin might be involved in activating protein kinases and phosphatases in response to changes in intracellular Ca^{2+} , similar to the action of calmodulin (37).




FIG. 4. (A) Expression of CcCaMK in *E. coli*. The CcCaMK protein was induced in *E. coli* and the cell extract was subjected to SDS/PAGE. Lanes: 1, isopropyl β -D-thiogalactoside-induced cell extract; 2, uninduced cell extract. The size of the protein is marked (in kDa) on the left. (B) Calmodulin binding to CcCaMK. CcCaMK protein (250 ng) was transferred onto a nitrocellulose filter and incubated with ^{35}S -labeled calmodulin (50 nM) in the buffer containing either 5 mM EGTA or 1 mM CaCl_2 . The histogram shows radioactivity (cpm) on the nitrocellulose filter. Bars: 1, 5 mM EGTA; 2, 1 mM CaCl_2 ; 3, 1 mM CaCl_2 /2.5 μM unlabeled calmodulin. Autoradiogram is shown above each bar. (C) Ca^{2+} binding to CcCaMK. Proteins were transferred to a Zeta-Probe membrane and probed with ^{45}Ca . Bands: 1, bovine serum albumin (2 μg); 2, calmodulin (2 μg); 3, CcCaMK (2 μg). Note that the intensity of calmodulin control is less than the intensity of CcCaMK possibly because of inefficient binding of calmodulin to the membrane (39).




FIG. 5. Expression of CCaMK gene in lily. RNase protection assay was performed by using total RNA (20 μ g) from various parts of lily. Total RNA was used from leaf, stem, and various organs from immature flower. Lanes: 1, leaf; 2, stem; 3, anthers from phase II; 4, sepals and petals from phase II; 5, anthers from phase III; 6, sepals and petals from phase III; 7, yeast tRNA control. Phases II and III correspond to stages of anther development as described by Wang *et al.* (40).

An unusual feature of CCaMK is the presence of a putative biotin-binding site (LKAMKMNSLI) within the visinin-like domain (Fig. 2). Such a biotin-binding site has not been observed in neural visinin-like proteins. Although, biotin is known to play a catalytic role in several essential metabolic carboxylation and decarboxylation reactions (38), its role in the regulation of CCaMK is not known. CCaMK also contains two consensus motifs, RXXT/S (Figs. 1 and 2), analogous to the autophosphorylation site of mammalian CaMKII and its homologs (9).

The structural features of the CCaMK gene indicate that it is a chimeric Ca^{2+} - and Ca^{2+} /calmodulin-dependent protein kinase with two discrete regulatory domains, a calmodulin-binding domain and a visinin-like Ca^{2+} -binding domain (Fig. 3). The presence of these distinct domains suggests dual modes of regulation. Furthermore, the presence of a putative biotin-binding site suggests yet another mode of regulation, adding to the functional diversity of CCaMK. The chimeric feature of the CCaMK gene suggests that it has evolved from a fusion of two genes that are functionally different and phylogenetically diverse in origin.




FIG. 6. Southern blot of lily genomic DNA digested with various restriction enzymes and probed with the CCaMK clone. Lanes: 1, *Dra* I; 2, *Eco*RI; 3, *Eco*RV; 4, *Hind*III; 5, *Pst* I; 6, *Xba* I; 7, *Xho* I. Sizes in kb are shown to the right.

To study the functional role of the predicted structural motifs of CCaMK, the *E. coli*-expressed protein was used for Ca^{2+} - and calmodulin-binding assays. The protein was expressed in *E. coli* (Fig. 4A) and purified by calmodulin affinity chromatography to near homogeneity as judged by SDS/PAGE. The calmodulin-binding assay confirmed that calmodulin binds to CCaMK only in the presence of Ca^{2+} (Fig. 4B). Furthermore, when incubated with excess amounts (50-fold) of unlabeled calmodulin, the binding of ^{35}S -labeled calmodulin to CCaMK was effectively reduced, suggesting that calmodulin binding to CCaMK was specific. To determine the functional role of the EF-hand motifs within the visinin-like domain, ^{45}Ca -binding assays were carried out. The results revealed that Ca^{2+} directly binds to CCaMK (Fig. 4C). Moreover, CCaMK also showed a Ca^{2+} -dependent shift in mobility by SDS/PAGE (data not shown). Although, these results suggest that CCaMK has some of the structural properties of both Ca^{2+} -dependent and Ca^{2+} /calmodulin-dependent protein kinases, an in-depth study is required to conclusively demonstrate a functional link among the three structural domains (Fig. 3).

The CCaMK gene was preferentially expressed during phase III (40) of anther development as revealed by the ribonuclease protection assay (Fig. 5). The expression of CCaMK during phase III suggests that it may be involved in microsporogenesis. Some of the EF-hand proteins like calmodulin (41) are ubiquitous and are active in diverse tissues. However, visinin-like proteins are restricted to specialized tissues such as neurons. Interestingly, CCaMK, which has a visinin-like domain, is also expressed in an organ-specific manner. Genomic Southern blot analysis revealed that CCaMK is encoded by a single gene (Fig. 6). Hybridization at low stringency using the CCaMK probe indicated the presence of a CCaMK homolog in other plants, such as *Arabidopsis*, apple, and tobacco (data not shown). We have also cloned a CCaMK homolog from tobacco with structural components similar to lily, including calmodulin-binding and visinin-like domains. These results suggest that the CCaMK-like gene is present in both monocotyledonous and dicotyledonous plants.

The Ca^{2+} -signaling pathway mediated through Ca^{2+} /calmodulin-dependent protein phosphorylation is well established in animals. This report confirms the presence of a Ca^{2+} /calmodulin-dependent protein kinase in plants. However, the presence of a visinin-like Ca^{2+} -binding domain in CCaMK adds an additional Ca^{2+} -sensing mechanism not previously known in these kinases. This feature distinguishes CCaMK from all known Ca^{2+} /calmodulin-dependent protein kinases. The discovery of the CCaMK gene increases our understanding of Ca^{2+} -mediated signal transduction in plants.

We thank Gynheung An for providing the cDNA libraries and V. Paranjape and Z. Liu for their help during this investigation. We also thank Michael Kahn (Institute of Biological Chemistry, Washington State University) for reviewing the manuscript. The support of the National Science Foundation (Grant DCB 9104586), the National Aeronautics and Space Administration (Grant NAG-10-0061), and the Agricultural Experimental Station (Project 0321) to B.W.P. and the Rotary Foundation Scholarship to S.P. is gratefully acknowledged.

1. Rasmussen, H. (1989) *Sci. Am.* (Oct.), 66.
2. Nairn, A. C. & Picciotto, M. R. (1994) *Cancer Biol.* 5, 295–303.
3. Campbell, A. K. (1983) *Intracellular Calcium: Its Universal Role as Regulator* (Wiley, Chichester, U.K.).
4. Trewavas, A. J. & Gilroy, S. (1991) *Trends Genet.* 7, 356–360.
5. Poovaiah, B. W. & Reddy, A. S. N. (1993) *CRC Crit. Rev. Plant Sci.* 12, 185–211.
6. Edelman, A. M., Blumenthal, D. K. & Krebs, E. G. (1987) *Annu. Rev. Biochem.* 56, 567–613.
7. Cohen, P. (1992) *Trends Biochem. Sci.* 17, 408–413.
8. Veluthambi, K. & Poovaiah, B. W. (1984) *Science* 223, 167–169.
9. Colbran, R. J. & Soderling, T. R. (1990) *Curr. Top. Cell Regul.* 31, 181–221.

10. Harper, J. F., Sussman, M. R., Schaller, G. E., Putnam-Evans, C., Charbonneau, H. & Harmon, A. C. (1991) *Science* **252**, 951-954.
11. Roberts, D. M. & Harmon, A. C. (1992) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **43**, 375-414.
12. Poovaiah, B. W. & Reddy, A. S. N. (1987) *CRC Crit. Rev. Plant Sci.* **6**, 47-103.
13. Hanson, P. I. & Schulman, H. (1992) *Annu. Rev. Biochem.* **61**, 559-601.
14. Poovaiah, B. W., Reddy, A. S. N., An, G., Choi, Y. J. & Wang, Z. Q. (1992) in *Progress in Plant Growth Regulation*, ed. Karssen, C. M., VanLoon, L. C. & Vreugdenhil, D. (Kluwer, Dordrecht, The Netherlands), p. 691-702.
15. Watillon, B., Kettmann, R., Boxus, Ph. & Burny, A. (1993) *Plant Physiol.* **101**, 1381-1384.
16. Hunter, T. (1987) *Cell* **50**, 823-829.
17. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY), 2nd Ed.
18. Kim, S.-R., Kim Y. & An, G. (1993) *Plant Mol. Biol.* **21**, 39-45.
19. Sanger, F., Nicklen, S. & Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467.
20. Devereaux, J., Haeberli, P. & Smithies, O. (1984) *Nucleic Acids Res.* **12**, 387-395.
21. Verwoerd, T. C., Decker, B. M. M. & Hoekema, A. (1989) *Nucleic Acids Res.* **17**, 2362.
22. Studier, F. W., Rosenberg, A. H., Dunn, J. J. & Dubendorff, J. W. (1990) *Methods Enzymol.* **185**, 60-89.
23. Hagiwara, T., Ohsako, S. & Yamauchi, T. (1991) *J. Biol. Chem.* **266**, 16401-16408.
24. Fromm, H. & Chua, N.-H. (1992) *Plant Mol. Biol. Rep.* **10**, 199-206.
25. Jena, P. K., Reddy, A. S. N. & Poovaiah, B. W. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 3644-3648.
26. Sikela, J. M. & Hahn, W. E. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 3038-3042.
27. Maruyama, K., Mikawa, T. & Ebashi, S. (1984) *J. Biochem. (Tokyo)* **95**, 511-519.
28. Hanks, S. K., Quinn, A. M. & Hunter, T. (1988) *Science* **241**, 42-52.
29. Colbran, R. J., Schworer, C. M., Hashimoto, Y., Fong, Y. L., Rich, D. P., Smith, M. K. & Soderling, T. R. (1989) *Biochem. J.* **258**, 313-325.
30. Pausch, M. H., Kaim, D., Kunisawa, R., Admon, A. & Thorner, J. (1991) *EMBO J.* **10**, 1511-1522.
31. Kornstein, L. B., Gaiso, M. L., Hammell, R. L. & Bartelt, D. C. (1992) *Gene* **113**, 75-82.
32. O'Neill, K. T. & DeGrado, W. F. (1990) *Trends Biochem. Sci.* **15**, 59-64.
33. Kuno, T., Kajimoto, Y., Hashimoto, T., Mukai, H., Shirai, Y., Saheki, S. & Tanaka, C. (1992) *Biochem. Biophys. Res. Commun.* **184**, 1219-1225.
34. Kobayashi, M., Takamatsu, K., Saitoh, S., Miura, M. & Noguchi, T. (1992) *Biochem. Biophys. Res. Commun.* **189**, 511-517.
35. Lenz, S. E., Henschel, Y., Zopf, D., Voss, B. & Gundelfinger, E. D. (1992) *Mol. Brain Res.* **15**, 133-140.
36. Okazaki, K., Watanabe, M., Ando, Y., Hagiwara, M., Terasawa, M. & Hidaka, H. (1992) *Biochem. Biophys. Res. Commun.* **185**, 147-153.
37. Pongs, O., Lindemeier, J., Zhu, X. R., Theil, T., Engelkamp, D., Krah-Jentgens, I., Lambrecht, H. G., Koch, K. W., Schwemer, J., Rivoisechi, R., Mallart, A., Galceran, J., Canal, I., Barbas, J. A. & Ferrus, A. (1993) *Neuron* **11**, 15-28.
38. Chandler, C. S. & Ballard, F. J. (1988) *J. Biol. Chem.* **263**, 1013-1016.
39. Van Eldik, L. J. & Wolchok, S. R. (1984) *Biochem. Biophys. Res. Commun.* **124**, 752-759.
40. Wang, C.-S., Walling, L. L., Eckard, K. J. & Lord, E. M. (1992) *Am. J. Bot.* **79**, 118-127.
41. Moncrief, N. D., Krestinger, R. H. & Goodman, M. (1990) *J. Mol. Evol.* **30**, 522-562.

