

Lower hybrid oscillations in multicomponent space plasmas subjected to ion cyclotron waves

G. V. Khazanov,¹ E. N. Krivorutsky,² T. E. Moore,¹ M. W. Liemohn,³
and J. L. Horwitz²

Abstract. It is found that in multicomponent plasmas subjected to Alfvén or fast magnetosonic waves, such as are observed in regions of the outer plasmasphere and ring current-plasmapause overlap, lower hybrid oscillations are generated. The addition of a minor heavy ion component to a proton-electron plasma significantly lowers the low-frequency electric wave amplitude needed for lower hybrid wave excitation. It is found that the lower hybrid wave energy density level is determined by the nonlinear process of induced scattering by ions and electrons; hydrogen ions in the region of resonant velocities are accelerated; and nonresonant particles are weakly heated due to the induced scattering. For a given example, the light resonant ions have an energy gain factor of 20, leading to the development of a high-energy tail in the H^+ distribution function due to low-frequency waves.

1. Introduction

1.1. Low-Frequency Wave-Plasma Interaction

Low-frequency Alfvén, fast magnetosonic, and ion cyclotron (IC) waves are commonly found in space plasmas. One example is IC waves generated by ion temperature anisotropies in the ring current [Cornwall, 1965, 1966; Kennel and Petschek, 1966; Liemohn, 1967]. IC waves have also been observed at geostationary orbit [Young *et al.*, 1981; Mauk, 1982; and references therein], in the outer magnetosphere [Anderson *et al.*, 1992], at midaltitude high latitudes [Erlandson *et al.*, 1990], and at ionospheric altitudes [Iyemori and Hayashi, 1989].

Low-frequency waves (LFWs) in the IC range can effectively interact with different components of the magnetospheric plasma through heating and scattering (e.g., reviews by Gendrin [1981, 1983, 1985] and Chang and Andre [1993]. Such LFWs are believed to be responsible for precipitation of ring current protons [Kennel and Petschek, 1966; Cornwall *et al.*, 1970], formation of SAR-arcs [Cornwall *et al.*, 1971] and acceleration of heavy ions [Omura *et al.*, 1985; Tanaka, 1985]. Noncollisional damping of IC waves may lead to the heating of cold electrons and ions in the outer plasmasphere [Galeev, 1975; Gorbachev *et al.*, 1988; Konikov *et al.*, 1989], deduced from the observed increase of plasma temperatures in this region [Decreau *et al.*, 1982; Gringauz, 1983, 1985;

Comfort *et al.*, 1985]. The transport of the dissipating wave energy from the plasmasphere along the geomagnetic field lines into the ionosphere below could account for the observed subauroral rise in the electron temperature [e.g., Kozyra *et al.*, 1986, and references therein].

By interacting with charged particles, the waves influence the behavior of the plasma as a whole, and therefore wave effects must be included in corresponding models. Ganguli and Palmadesso [1987, 1988], Singh [1988], Singh and Torr [1990], Brown *et al.* [1991, 1995], and Lin *et al.* [1992, 1994] took into account effects from the low-frequency (LF) electrostatic turbulence, and they demonstrated that wave-particle interactions lead to significant effects on the evolution of the core plasma distribution functions.

In developing a mathematical model to describe plasma transport in the magnetosphere-ionosphere system that accounts for the active wave processes which occur there, we must develop a general scheme to include an analysis of the dispersion characteristics of the medium in order to choose suitable wave modes, a wave-particle interaction mechanism, and a system of hydrodynamical equations governing macroscopic plasma parameters which properly accounts for the presence of wave-particle interactions. One basis for this scheme's development has been described elsewhere [Khazanov and Chernov, 1988; Konikov *et al.*, 1989; Chernov *et al.*, 1990; Gamayunov *et al.*, 1991, 1992; Gorbachev *et al.*, 1992].

When such LF waves are propagating in a plasma, a ponderomotive force is produced, which leads to significant effects in the plasma. Boehm *et al.* [1990] suggested that this ponderomotive force can be the cause of significant density perturbations in the auroral zone. Allan *et al.* [1991] and Allan [1992] have investigated mass transport caused by the ponderomotive force of hydrodynamic waves in the middle magnetosphere. It was also suggested that significant plasma energization could occur in these regions [Allan, 1993]. Li and Temerin [1993] and Lee and Parks [1996] have used similar ideas to evaluate ion energization by large-amplitude Alfvén waves ($E > 100$ mV/m; $\omega < \Omega_i$) in the auroral zone.

¹Space Sciences Laboratory, NASA/Marshall Space Flight Center, Huntsville, Alabama.

²Department of Physics and Center for Space Plasma and Aeronomic Research, University of Alabama in Huntsville.

³Space Physics Research Laboratory, University of Michigan, Ann Arbor.

It was recently demonstrated by *Khazanov et al.* [1996] that large-amplitude LFWs can generate lower hybrid waves (LHWs) in the auroral zone and ring current region. The ion energization due to the LHWs may be comparable with that produced by the ponderomotive force of the LFWs. The purpose of this paper is to extend the analysis of *Khazanov et al.* [1996] to multicomponent plasmas and to small LFW electric fields.

1.2. LHW Excitation Due to LFWs and Description of the Problem

It is well known that in the electric field of a LFW, electrons and ions move with different velocities. The arising ion-electron relative velocity can lead to a current instabilities and, in particular, LHW excitation. Depending on the frequency of the LFW, the magnitude of the electric field, plasma composition, temperatures of different plasma species, and the external magnetic field, strong or weak LHW turbulence may develop.

In the case of large-amplitude LFWs, strong lower hybrid (LH) turbulence could be produced [*Gamayunov et al.*, 1992; *Khazanov et al.*, 1996]. Strong LH turbulence results in plasma heating and acceleration [*Shapiro et al.*, 1993], the formation of ion conics [*Chang and Coppi*, 1981; *Retterer et al.*, 1986], and possibly the saturation of Alfvén oscillations in the ring current region [*Gamayunov et al.*, 1992]. Weak LH turbulence can also lead to important changes in the plasma state; for example, *Omelchenko et al.* [1994] found that the LH "fan" instability may accelerate ions.

The level of energy density of LFWs which leads to excitation of LHWs in a hydrogen plasma, was discussed by *Khazanov et al.* [1996]. It can be estimated by taking into account that the ion-electron relative velocity needed for LHW excitation is of the same order of magnitude as the thermal ion velocity. This leads to a requirement for the LFW energy density $|E_0|^2$,

$$\frac{E_0^2}{8\pi nT} \geq \frac{\omega_{Bi}^4}{\omega_0^2 \omega_{pi}^2} \quad (1)$$

where ω_{Bi} and ω_{pi} are the cyclotron and plasma frequencies for ions, respectively; nT is the thermal plasma energy density; and ω_0 is the LFW frequency.

Khazanov et al. [1996] assumed that the plasma consists of only electrons and protons and that the LFW electric field is strong enough not only to make the relative proton velocity greater than its thermal velocity, but also to create strong LHW turbulence. From (1), however, it can be seen that the LFW energy density needed for LHW excitation may be lower in the presence of a heavy ion component. Since the magnetosphere and ionosphere have multicomponent plasmas, it can be supposed that LHWs can be excited in such plasmas by LFWs with amplitudes less than that needed for a proton-electron plasma.

This leads us to the problem of LFW interaction with a multicomponent plasma due to LHW excitation. In accordance with this problem, we will study the following questions:

1. How does a small mixture of heavy ions influence the LFW electric field needed for LHW excitation? What are the growth rates of the LHWs and how do they depend on the heavy ion population, LFW electric field, and core plasma parameters? What are these dependencies in the kinetic and hydrodynamic regimes of LHW excitation? (These are discussed in section 2.)

2. What is the physical mechanism that determines the level of LHW energy density? Does the quasilinear interaction of LHWs with oxygen, helium, and hydrogen ions as well as with electrons lead to saturation of the LHW instability, or is it the nonlinear-induced scattering mechanism that determines the LHW energy density level? What is the level of LHW energy? (These are discussed in sections 3.1 and 3.2.)

3. What is the influence of weak LHW turbulence on different plasma species? (This is discussed in section 3.3.)

In section 4 the suppositions used and results obtained are summarized. Fulfillment of these suppositions and the possibility of LFW interaction with magnetospheric plasma due to LHW excitation are analyzed, and an estimation of particle energization for a magnetospheric plasma is conducted. Section 5 contains a short summary of the results.

2. Linear Analysis

2.1. Ion Drift in a LFW Electric Field

Let us consider a multicomponent plasma containing electrons and oxygen, helium, and hydrogen ions subjected to LFW. In a variable electric field \mathbf{E} (for instance, the electric field of an Alfvén or fast magnetosonic wave, which has a component normal to the external magnetic field \mathbf{B}_0) the electrons and ions will acquire different velocities, \mathbf{w}_e and \mathbf{w}_α , due to this field. If the frequency ω_0 of the field is considerably lower than the electron cyclotron frequency, the electron and ion velocities will be given by [e.g., *Akhiezer*, 1975]

$$\begin{aligned} \mathbf{w}_e &= Re \left\{ c \frac{\mathbf{E} \times \mathbf{B}_0}{B_0^2} - i \frac{e(\mathbf{E} \cdot \mathbf{b})\mathbf{b}}{m_e \omega_0} \right\} \\ \mathbf{w}_\alpha &= Re \left\{ \frac{e}{m_\alpha (\omega_{B\alpha}^2 - \omega_0^2)} \left[\omega_{B\alpha} \mathbf{E} \times \mathbf{b} + i \omega_0 (\mathbf{E} \times \mathbf{b}) \times \mathbf{b} \right] \right\} \quad (2) \\ \mathbf{E} &= \mathbf{E}_0 e^{-i\omega_0 t} \end{aligned}$$

where \mathbf{b} is a unit vector in the direction of the external magnetic field \mathbf{B}_0 and $\omega_{B\alpha}$ is the ion cyclotron frequency of the ion species α . In a LFW the parallel electric field component can be small and the perpendicular drift velocities of the electrons and ions may be comparable to or even larger than the parallel electron velocity. This study will be restricted to the case where the parallel electric field and the parallel electron velocity can be neglected.

In a coordinate system where the magnetic field \mathbf{B}_0 is parallel to the z axis and the LF electric field \mathbf{E} is parallel to the x axis, the relative velocity of the ion species α with respect to the electrons, $\mathbf{u}_\alpha = \mathbf{w}_\alpha - \mathbf{w}_e$, is

$$\mathbf{u}_\alpha = \frac{eE_0\omega_0}{m_\alpha (\omega_0^2 - \omega_{B\alpha}^2)} \left[i \sin \omega_0 t + \mathbf{j} \frac{\omega_0}{\omega_{B\alpha}} \cos \omega_0 t \right] \quad (3)$$

For LFW with left-hand and right-hand circular polarization propagating along the z axis, the relative velocity is given by the expression

$$\mathbf{u}_\alpha = \frac{eE_0\omega_0}{\sqrt{2}m_\alpha (\pm\omega_0 - \omega_{B\alpha})\omega_{B\alpha}} [i \sin \omega_0 t + \mathbf{j} \cos \omega_0 t] \quad (4)$$

If ω_0 is not near the cyclotron frequency of the light or heavy ions, $\omega_B^H < \omega_0 < \omega_B^L$, then the ratio of the velocity of the heavy ions u^H to the velocity of light ions u^L is of the order

$$\frac{u^H}{u^L} \sim \frac{\omega_B^L}{\omega_0}$$

for waves with linear or circular polarization. For $\omega_0 \sim \omega_B^H$, again for both cases, we get

$$\frac{u^H}{u^L} \sim \frac{m^H}{m^L} \left(\frac{\omega_0}{\omega_B^H} - 1 \right)^{-1}$$

It may be concluded from the estimations of (3) and (4) that the relative ion velocity in the electric field of LFW with fixed frequency increases with ion mass. The reason is that the relative velocity of the various ions is caused by the different inertia of ions and electrons in the oscillating electric field. For a LFW frequency near ω_B^H the resonance effect also becomes important. The role of the resonance denominator in (3) and (4) can be significant in some cases. For example, IC waves with a frequency near the helium gyrofrequency, generated in the ring current region and propagating in the direction of stronger magnetic field, may move to a region where ω_0 is closer to the oxygen gyrofrequency. For waves observed by the Active Magnetospheric Particle Tracer Explorers (AMPTE) Ion Release Module (IRM) satellite [LaBelle and Treumann, 1992], the relation ω_0 / ω_{Bo^+} was 1.2 and the relative velocity of oxygen ions was 20 times greater than that of hydrogen. It follows from the above argument that in a multicomponent plasma, LFWs due to inertial drift of the heavy ions form a current across the magnetic field.

2.2. Lower Hybrid Wave Generation

Let us consider the LHW generation in the linear approximation. The dispersion equation in this case has the form [e.g., Akhiezer, 1975]

$$1 + \frac{\omega_{pe}^2}{k^2 v_{Te}^2} \left[1 + I_0(x) e^{-x} z_e Z(z_e) \right] + \sum_{\alpha} \frac{\omega_{p\alpha}^2}{k^2 v_{T\alpha}^2} \left[1 + z_{\alpha} Z(z_{\alpha}) \right] = 0$$

$$x = \frac{k_{\perp}^2 v_{Te}^2}{\omega_{Be}^2}; \quad z_e = \frac{\omega}{\sqrt{2} k_z v_{Te}}; \quad z_{\alpha} = \frac{\omega - \mathbf{k}_{\perp} \mathbf{u}_{\alpha}}{\sqrt{2} k v_{T\alpha}}; \quad (5)$$

$$v_T^2 = \frac{T}{m}; \quad \omega_{pe}^2 = \frac{ne^2}{m_e \epsilon_0}; \quad \omega_{p\alpha}^2 = \frac{n_{\alpha} e^2}{m_{\alpha} \epsilon_0}; \quad \omega_{Be} = \frac{eB}{m_e}$$

where $I_0(x)$ is the modified Bessel function; $Z(z_{\alpha})$ is the plasma dispersion function; and ω and \mathbf{k} are the frequency and the wave vector of the LH oscillations, respectively.

In obtaining (5), it was assumed that $\omega_{Ba}^2 \ll \omega^2 \ll \omega_{Be}^2$ and the motion of all ion species is unmagnetized, i.e., $k_{\perp}^2 v_{T\alpha}^2 / \omega_{Ba}^2 \gg 1$. This inequality is easily satisfied, taking into account that the ions with velocity u mainly generate LHWs with wave vector $k \sim \omega/u$, where $\omega \sim (\omega_{Be} \omega_{Bh^+})^{1/2}$ and $u \geq v_{T\alpha}$. The dispersion equation (5) is valid if during a time of order γ^{-1} (where γ is the growth rate of LHW), the ion motion can be considered regular. This leads to the condition

$$\gamma > \omega_0 \quad (6)$$

We will restrict our analysis of (5) to the case $c_{O^+} \ll c_{He^+} \ll c_{H^+}$, where $c_{\alpha} = n_{\alpha}/n$.

2.2.1. LHW generation in the kinetic limit. Let us examine the kinetic limit of LHW generation. Considering

$z_e, z_{H^+}, z_{He^+} \gg 1$, neglecting the first term and u_{H^+}, u_{He^+} in (5), and taking into account only the oxygen term in the imaginary part of (5), the dispersion law becomes

$$\omega^2 = \omega_{LH}^2 \left(c_{H^+} + \frac{c_{He^+}}{4} + \frac{m_{H^+}}{m_e} \cos^2 \theta + \Gamma(\theta) \frac{k^2 v_{Th^+}^2}{\omega_{LH}^2} \right) \quad (7)$$

where $k_{\perp}^2 v_{Te}^2 / \omega_{Be}^2 < 1$, $\omega_{LH} = \sqrt{\omega_{Be} \omega_{Bh^+}}$, θ is the angle between the wave vector \mathbf{k} and the magnetic field, and $2.5 \leq \Gamma(\theta) \leq 4$; $\Gamma(0) \sim 4$. The function $\Gamma(\theta)$ depends on the different ion species' concentrations c_{α} , and the minimum $\Gamma(\theta)$ is at θ , satisfying the condition $m_i / m_e \cos^2 \theta \sim c_{H^+}$.

The growth rate of LHW is

$$\gamma = -\frac{\sqrt{\pi}}{2R(\theta)} \frac{\omega^3}{k^2 v_{Th^+}^2} \left[\frac{T_i}{T_e} I_0(x) e^{-x} z_e e^{-z_e^2} + \sum_{\alpha} c_{\alpha} z_{\alpha} e^{-z_{\alpha}^2} \right] \quad (8)$$

where $R(\theta)$ is a slowly changing function of k and θ ,

$$R(\theta) = \left(1 + \frac{3}{z_e^2} - x \right) \frac{m_{H^+}}{m_e} \cos^2 \theta + \left(1 + \frac{3}{z_{H^+}^2} \right) c_{H^+}$$

It is clear from (8) that a positive γ can be obtained for weak LFW electric fields due to the oxygen ion drift (which is the largest drift) when in the oxygen term, $\omega - \mathbf{k}_{\perp} \mathbf{u}_{O^+} < 0$ and $\omega - \mathbf{k}_{\perp} \mathbf{u}_{O^+} / k_{\perp} v_{To^+} \leq 1$. For a small oxygen concentration, $\gamma > 0$ in a narrow region of wave velocities, $\omega / k_{\perp} \sim u_{O^+}$, with $u_{O^+} > v_{Th^+}, v_{THe^+}$. In this case, the oxygen term is at a maximum and the number of hydrogen and helium ions in resonance with LHW is small. Also, electron Cherenkov damping limits the generation, so LHW excitation is possible only for

$$\frac{T_e m_i}{T_i m_e} \cos^2 \theta \leq 1$$

Figure 1 illustrates these features of LHW generation, showing $\gamma > 0$ only for large θ and a certain range of ω / k .

For a purely hydrogen plasma, $\gamma > 0$ if $\omega - \mathbf{k}_{\perp} \mathbf{u}_{H^+} < 0$ and $\omega - \mathbf{k}_{\perp} \mathbf{u}_{H^+} / k_{\perp} v_{Th^+} \leq 1$, which means $u_{H^+} \sim v_{Th^+}$. This condition leads to much greater required LFW electric fields than for $u_{O^+} > v_{Th^+}$. From (8) the maximum growth rate (neglecting the damping) is




Figure 1. The normalized growth rate of lower hybrid waves (LHWs) as a function of the wave normal angle $[T_i m_e / (T_e m_i \cos^2 \theta)]$ and wave velocity $[\omega / (\sqrt{2} k v_{Th^+})]$.

$$\gamma_{\max} \approx 0.3 c_{O^+} \frac{u_{O^+}^2}{v_{Th^+}^2} \omega \quad (9)$$

The restriction given in (6) leads to an inequality that is easy satisfied when u_{O^+} equals several times v_{Th^+} and $c_{O^+} \geq 10^{-2}$.

Note that LHW excitation is possible if the oxygen velocity is of the order of the thermal hydrogen velocity. Taking $z_{O^+} \gg 1$, the dispersion equation (5) can be presented in the form

$$1 + \delta\epsilon^e(\mathbf{k}, \omega) + \delta\epsilon^{H^+}(\mathbf{k}, \omega) + \delta\epsilon^{He^+}(\mathbf{k}, \omega) - \frac{\omega_{pO^+}^2}{(\omega - \mathbf{k}_\perp \mathbf{u}_{O^+})^2} = 0 \quad (10)$$

where $\delta\epsilon^e$, $\delta\epsilon^{H^+}$, and $\delta\epsilon^{He^+}$ are the electron, hydrogen, and helium dielectric permittivities, respectively.

Assuming $\omega = \mathbf{k}_\perp \mathbf{u}_{O^+} + \eta$, $\eta \ll \omega$, we can find the growth rate from (10)

$$\eta = \frac{\omega_{pO^+}}{\sqrt{1 + \delta\epsilon^e(\mathbf{k}, \mathbf{k}_\perp \mathbf{u}_{O^+}) + \delta\epsilon^{H^+}(\mathbf{k}, \mathbf{k}_\perp \mathbf{u}_{O^+}) + \delta\epsilon^{He^+}(\mathbf{k}, \mathbf{k}_\perp \mathbf{u}_{O^+})}} \quad (11)$$

If $u_{O^+} \approx v_{Th^+}$, the term $\delta\epsilon^{H^+}$ has an imaginary part caused by the presence of the resonant hydrogen ions, and so one root of (11) leads to an instability. For $\omega_{pH^+}^2 / k^2 v_{Th^+}^2 \gg 1$ the estimation of the growth rate is

$$\gamma \sim \frac{c_{O^+}^{1/2}}{4} \mathbf{k}_\perp \mathbf{u}_{O^+} \quad (12)$$

2.2.2. LHW generation in the hydrodynamic limit. Let us examine the hydrodynamic region of LHW generation, $z_\alpha, z_e \gg 1$, $x \ll 1$. For angles $\cos^2 \theta \sim m_e / m_i$, taking into account only the relative motion of oxygen ions, (5) can be transformed into

$$1 + \frac{\omega_{pe}^2}{\omega_{be}^2} \sin^2 \theta - \frac{\omega_{pH^+}^2}{\omega^2} - \frac{\omega_{pHe^+}^2}{\omega^2} - \frac{\omega_{pO^+}^2}{(\omega - \mathbf{k}_\perp \mathbf{u}_{O^+})^2} = 0 \quad (13)$$

Assuming $\omega = \mathbf{k}_\perp \mathbf{u}_{O^+} + \eta$, where $\omega^* = \mathbf{k}_\perp \mathbf{u}_{O^+}$ is the solution of (13) for $c_{O^+} = 0$, the maximum growth rate of LHW comes out of this expression as

$$\eta = \sqrt[3]{1 - \frac{\mathbf{k}_\perp \mathbf{u}_{O^+}}{4} c_{O^+}^{1/3}} \quad (14)$$

The hydrodynamic consideration is valid if $z_{O^+} \gg 1$, i. e.,

$$\frac{u_{O^+}}{v_{To^+}} \gg \frac{4}{c_{O^+}^{1/3}} \quad (15)$$

and $|\eta| > \omega_0$ (see (6)). Both of these conditions are easily satisfied.

Using the estimations for u_{O^+} and u_{H^+} (see (3), (4), and below), (15) may be rewritten as

$$\frac{u_{H^+}}{v_{Th^+}} \gg \frac{m_{H^+}}{m_{O^+}} c_{O^+}^{1/3} \left(\frac{\omega_0}{\omega_{BO^+}} - 1 \right)$$

For a purely hydrogen plasma the LHW hydrodynamic instability is possible if $u_{H^+} / v_{Th^+} \gg 1$. It follows from the last two inequalities that for a multicomponent plasma the LHW hydrodynamic instability occurs for a smaller drift velocity and LFW electric field than for a purely hydrogen plasma.

Note that the obtained results are valid for plane polarized as well as circularly polarized LFW. In the case of circular polarization the oxygen ions act as a beam of oscillators, with a rotation frequency ω_0 (see (4)). The radius of rotation, u_{O^+} / ω_0 , is much greater than the LH wavelength, because

$$\frac{k_\perp^2 u_{O^+}^2}{\omega_0^2} \geq \frac{k_\perp^2 v_{Th^+}^2}{\omega_{BO^+}^2} \gg 1$$

and for this linear analysis of LHW generation the rotation can be neglected.

Comparing the growth rates of (9), (12), and (14), it can be found that the strongest dependence on the oxygen ion density as well as the LFW electric field takes place in the kinetic regime of LHW generation in (9). Of course, in this regime the LFW electric field needed for instability excitation is smaller than in the hydrodynamic case (15). From the above derivation it follows that in a plasma subjected to LFW the heavy plasma species acts as a beam of particles with a relative drift velocity. This heavy ion beam determines the character of the LHW excitation, growth rate, and restrictions on the excitation. It also follows that in a plasma with some oxygen ions, the LHW instability can be excited by LFW with an electric field amplitude less than in a purely hydrogen plasma.

3. Quasi-linear Approximation

The LHW instability excited by the oxygen ion drift in the LFW electric field leads to the development of LH turbulence. The growth of the LHW can be restricted owing to their quasi-linear interaction with different plasma species. If such interaction does not lead to the instability saturation, the level of LH turbulence is determined by nonlinear processes, in particular, by induced scattering of LHW by ions and electrons. Below, these processes are studied and the quasi-steady-state level of the LHW energy density is found.

Let us summarize the characteristics of LHWs excited in a multicomponent plasma subjected to plane polarized LFW, with $\omega_0 \gg \omega_{BO^+}$ and some other results of the previous analysis. From (3) it follows that the main component of the relative velocity, \mathbf{u}_{O^+} , is directed along the y axis in the same phase as the LFW electric field. We will restrict our consideration to weak LFW electric fields (small E_0). It follows from (8) that in this case the LHW may be generated in a narrow region of angles near \mathbf{u}_{O^+} (the LFW electric field needed for excitation of LHW is at a minimum when $\mathbf{k}_\perp \parallel \mathbf{u}_{O^+}$). It was found above that the hydrogen and helium ion motions can be treated as unmagnetized, so the influence of the magnetic fields on their motion will only lead to phase mixing. Taking into account that the thermal velocity of oxygen ions is much smaller than their relative velocity, we can consider their motion as one-dimensional. The region of wave velocities in which the LHW generation is possible is determined by the condition $\gamma > 0$, where the lower limit is given by (8) and the upper limit is u_{O^+} and is narrow for small LFW electric fields (Figure 1).

3.1. Quasi-linear Equations

Let us now obtain the quasilinear equations describing the behavior of the LHW spectral energy density and distribution functions of different species of plasma particles.

The kinetic equation for oxygen ions in the presence of a LFW electric field is

$$\frac{\partial \tilde{f}_{O^+}}{\partial t} + \mathbf{v} \cdot \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{r}} + i \frac{eE_0}{m_{O^+}} \cos \omega_0 t \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} + \frac{e}{m_{O^+}} \mathbf{E} \cdot \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} + \omega_{BO^+} [\mathbf{v} \cdot \mathbf{b}] \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} = 0 \quad (16)$$

Here \tilde{f}_{O^+} is the oxygen ion distribution function and \mathbf{E} is the electric field of LHW. Introducing the relative velocity $\mathbf{v} = \mathbf{v}' - \mathbf{j}E_0 \cos \omega_0 t / B_0$, (16) becomes

$$\frac{\partial \tilde{f}_{O^+}}{\partial t} + (\mathbf{v} + \mathbf{j} \frac{E_0}{B_0} \cos \omega_0 t) \cdot \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{r}} + \mathbf{j} \omega_0 \frac{E_0}{B_0} \sin \omega_0 t \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} + \omega_{BO^+} [\mathbf{v} \cdot \mathbf{b}] \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} + \frac{e}{m_{O^+}} \mathbf{E} \cdot \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} = 0 \quad (17)$$

By averaging (17) over the characteristic time of the LH oscillation, the quasilinear equation for the oxygen ion distribution function can be found

$$\frac{\partial \tilde{f}_{O^+}}{\partial t} - u \omega_0 \sin \omega_0 t \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} = \frac{\pi e^2}{m_{O^+}^2} \frac{\partial}{\partial \mathbf{v}} \int dk |E_k|^2 \delta(\mathbf{k} \cdot \mathbf{v} - \omega_k) \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} \quad (18)$$

where $u = eE_0/m\omega_0$, $\mathbf{v} = \mathbf{v}_y$, and $\delta(x)$ is a delta function. Equation (18) follows from (17) if the oxygen ions drift along the x axis and the magnetic field terms are neglected. The first assumption is valid with an accuracy of $\omega_{BO^+}/\omega_0 \ll 1$, the second with an accuracy of $v_{T0^+}/u \ll 1$. Supposing the LHW spectrum to be one-dimensional, evaluation of the integral in the right-hand side of (18) yields

$$\frac{\partial \tilde{f}_{O^+}}{\partial t} - u \omega_0 \sin \omega_0 t \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} = \frac{\pi e^2}{m_{O^+}^2} \frac{\partial}{\partial \mathbf{v}} \frac{|E_k|^2}{v} \frac{\partial \tilde{f}_{O^+}}{\partial \mathbf{v}} \quad (19)$$

where $|E_k|^2$ is the LHW spectral energy density. It was assumed in (19) that $|v - \partial \omega / \partial k|$, arising after the integration, is equal to v .

Taking into account the phase mixing and supposing the LHW spectrum is one-dimensional, the quasilinear equations for the resonant part of the hydrogen and helium distribution functions [Shapiro and Shevchenko, 1988] can be described in the form

$$\frac{\partial f_{H^+}(\text{He}^+)}{\partial t} = \frac{\pi e^2}{m_{H^+}^2(\text{He}^+)} \frac{1}{v_{\perp}} \frac{\partial}{\partial v_{\perp}} \int \frac{|E_k|^2 \omega_k^2}{k^2 \sqrt{k^2 v_{\perp}^2 - \omega^2}} \frac{1}{v_{\perp}} \frac{\partial f_{H^+}(\text{He}^+)}{\partial v_{\perp}} dk \quad (20)$$

Equations (20) are integrated over the azimuth angle in the plane perpendicular to the magnetic field assuming axial symmetry of the distribution functions. It follows from (20) that the resonant interaction for unmagnetized particles ($\omega = k_{\perp} v_{\perp}$) is possible for hydrogen and helium ions with $v_{\perp} \geq \omega/k_{\perp}$.

The resonant condition for electrons may be satisfied only for Cherenkov interaction. The quasilinear equation determining the electron distribution function evolution is [Shapiro and Shevchenko, 1988]

$$\frac{\partial f_e}{\partial t} = \frac{\pi e^2}{m_e^2} \int dk \frac{\partial}{\partial v_z} k_z \frac{|E_k|^2}{k^2} \frac{J_0^2(x)}{\left| v_z - \frac{\partial \omega}{\partial k} \right|} k_z \frac{\partial f_e}{\partial v_z} \bigg|_{k_z = \frac{\omega}{v_z}} \quad (21)$$

where $J_0(x)$ is the Bessel function. In (21), which is one-

dimensional, the small component of the wave vector along the magnetic field k_z must be prescribed.

The changes in the LHW energy density spectrum are described by an equation which takes into account the energy generation due to oxygen ion drift and energy damping due to the absorption by all other species [Shapiro and Shevchenko, 1988]:

$$\frac{\partial |E_k|^2}{\partial t} = \frac{\pi e^2}{m_{O^+}^2} \frac{\omega^2}{k_{\perp}^2} \frac{\partial f_{O^+}}{\partial \mathbf{v}} |E_k|^2 + \frac{\pi \omega^2}{k_{\perp}^2 |k_z| n} \int dv_{\perp} J_0^2(x) k_z \frac{\partial f_e}{\partial v_z} |E_k|^2 + \frac{\pi \omega^4}{k_{\perp}^2 n} \sum_{\alpha = H^+, \text{He}^+} \frac{m_{H^+}}{m_{\alpha}} \int dv_{\perp} \frac{\partial f_{\alpha}}{\partial v_{\perp}} \frac{|E_k|^2}{\sqrt{k^2 v_{\perp}^2 - \omega^2}} + NLT \quad (22)$$

The last term, NLT , takes into account nonlinear interaction processes of LHW with particles. In this case, it is induced scattering by ions and electrons [Musher et al., 1978]. The system of equations (19)-(22) describes the evolution of distribution functions and LHW electric field energy density.

3.2 LHW Energy Density

Our purpose here is to determine the LHW energy density and the influence of LHW on oxygen, hydrogen, and helium ions and electrons. We will avoid the beginning stage of LHW instability development and restrict the analysis to a time long enough for significant changes to occur in the initial distribution functions of the particles.

Let us examine (20) describing the distribution functions of hydrogen and helium ions. Owing to the interaction with the resonant LHW, these ions are accelerated, and, for large times, it can be supposed that the resonant ions have velocities such that $v_{\perp} \gg \omega/k_{\perp}$. Then (20), for $\omega/k_{\perp} = 0$, have an asymptotic solution for $t \rightarrow \infty$ that does not depend on the initial distribution function and is self-similar in form [Shapiro and Shevchenko, 1988],

$$f_{H^+}(\text{He}^+) = \frac{5 \tilde{n}_{H^+}(\text{He}^+)}{\Gamma(2/5) \tilde{\tau}^{2/5}} \exp[-v_{\perp}^5 / \tilde{\tau}] \quad (23)$$

$$\tilde{\tau} = \frac{25 e^2}{m_{H^+}^2(\text{He}^+)} \int_0^t dt' \int dk \frac{\omega_k^2}{k^3} |E_k|^2$$

where $\Gamma(2/5)$ is the Gamma function and $\tilde{n}_{H^+}(\text{He}^+)$ is the number of resonant particles.

From (23) it follows that the distribution functions of hydrogen and helium ions, and therefore the energy absorption, only weakly depend on fast energy density oscillations and thus are determined by the slow changes in the average energy density. For $T_{H^+} \sim T_{\text{He}^+}$, the thermal velocity of helium ions is half that of the hydrogen ions, so the number of helium ions in the region of resonant velocities (for Maxwellian initial distribution functions) is $(n_{\text{He}^+}/n_{H^+}) \exp(-3v^2/v_{T_{H^+}}^2)$ times less than the number of hydrogen ions. This means that the absorption by hydrogen ions is much greater than by helium ions.

From (21) it follows that the electron distribution function is one-dimensional. Therefore, for large times, a plateau in the region of resonant velocities along the z axis will eventually form and the energy absorption will cease. Thus, for large times, only the term describing the energy absorption by hydrogen ions needs to be kept in (22).

Note that the maximum level of LHW energy density is limited by the nonlinear term in (22). This energy level can be estimated from the balance of the quasilinear oxygen term and the nonlinear term.

The damping rates due to induced scattering by electrons and ions in the case under consideration are of the same order of magnitude [Musher *et al.*, 1978],

$$\gamma^{NL} \sim \frac{\omega_{Be}^2 \omega_{pH+}^2}{\omega_{LH}^2 \omega_{LH}^2} \frac{\epsilon}{nT} \quad (24)$$

where

$$\epsilon = \frac{1}{8\pi} \int |E_k|^2 dk \sim \frac{|E_k|^2}{8\pi} \Delta k \approx \frac{|E_k|^2}{8\pi} \frac{\omega_{LH}}{u}$$

From the equality of the linear growth rate γ (the maximum value of the quasilinear term) and the nonlinear damping rate γ^{NL} , the maximum value of the LHW energy density can be estimated,

$$\frac{|E_k|^2}{8\pi} \sim \frac{\gamma \omega_{LH}^2 u}{\omega_{Be}^2 \omega_{pH+}^2} nT \quad (25)$$

If we suppose that the system of LHWs and particles progresses to a steady-state condition due to quasilinear interactions, then we can consider the slow evolution of the oxygen distribution function and the LHW spectral energy density.

Let us investigate the solution of (20) and (22) with the form

$$\begin{aligned} f_0 &= F(v, t) + a(v) \cos \omega_0 t + b(v) \sin \omega_0 t \\ |E_k|^2 &= \epsilon_k(v, t) + c(v) \cos \omega_0 t + d(v) \sin \omega_0 t \end{aligned} \quad (26)$$

where F and ϵ_k are slowly varying functions of time. We will assume that the growth rate determined by the slowly changing part of the oxygen ion distribution function F and the damping rate caused by hydrogen ions are less than ω_0 . This damping rate can be estimated from the hydrogen term in (22) using the distribution function (23) and the LHW energy density (25) and is in accordance with this assumption.

In this case, the equation for F may be written in the form

$$\frac{\partial F}{\partial t} + \frac{\pi e^2 u^2}{m_{O+}^2} \frac{\partial^2 \epsilon_k}{\partial v^2} \frac{\partial^2 F}{\partial v^2} = \frac{\pi e^2}{m_{O+}^2} \frac{\partial \epsilon_k}{\partial v} \frac{\partial F}{\partial v} \quad (27)$$

The second term on the left-hand side shows that the asymptotic influence of the LFW electric field on the oxygen ion motion is determined by the "friction" of the ions with the LHW electric field. Equation (22) for the slowly varying part of LHW energy density ϵ_k has the form

$$\frac{\partial \epsilon_k}{\partial t} = \frac{\pi m_{H+}}{m_{O+}} \frac{\omega^3}{k^2 n} \epsilon_k \frac{\partial F}{\partial v} + \frac{\pi \omega^4}{k^3 n} \int \frac{dv_{\perp}}{v_{\perp}} \frac{\partial f_{H+}}{\partial v_{\perp}} \epsilon_k \quad (28)$$

As was discussed above, only the absorption by the hydrogen ions is taken into account in (28).

We will suppose below, as is usual in a quasilinear approximation, that the LHW spectrum has a sharp maximum at the wave vector $k \sim \omega/u$. Equation (27) can then be reduced to the following form:

$$\frac{\partial F}{\partial \tau} + \frac{u^2}{2} \frac{\partial^2}{\partial v^2} \frac{1}{v} \frac{\partial^2 F}{\partial v^2} = \frac{\partial}{\partial v} \frac{1}{v} \frac{\partial F}{\partial v} \quad (29)$$

where

$$\tau = \frac{\pi e^2}{m_{O+}^2} \int_0^t \epsilon_k dt$$

Let us estimate the role of the different terms on the time behavior of the oxygen ion distribution function. If we neglect the quasilinear term, then the asymptotic self-similar solution can be found. It does not depend on the initial conditions and can be calculated by the perturbation method as a series of powers of $\eta = v^5 \tau^1$. The term which determines the LHW evolution for $t \rightarrow \infty$ is $F \sim v \tau^{2/5}$. Conversely, if we neglect the term with u_{O+}^2 , the analogous quasilinear solution is $F \sim v^3 \tau^{4/3}$. The first expression shows that the LFW electric field causes a positive value for the term $\partial F / \partial v$ and leads to LHW generation. From the second expression for F it follows that the quasilinear interaction gives a negative $\partial F / \partial v$ term. Such a result means that by neglecting the term with the LFW electric field, we reduce the problem to a completely different one with the oxygen particles being accelerated by the LHW electric field.

We will suppose for further estimations that, for large times, the behavior of the derivative, $\partial F / \partial v$, is determined by the second term in expression (29). This means that the energy flow to the plasma is directly related to the LFW electric field and leads to

$$\frac{\partial F}{\partial v} \approx \frac{n_{O+}}{u^{4/5}} \tau^{-3/5} \quad (30)$$

Taking into account (23) and (30), the equation for the LHW spectral energy density (28) may be reduced to the form

$$\frac{d^2 D}{dx^2} = \frac{\pi \omega}{\omega_0} \frac{dD}{dx} \left[\frac{m_{H+}}{m_{O+}} \frac{n_{O+}}{n} D^{-3/5} - \frac{\tilde{n}}{n} \left(25 \frac{m_{O+}^2}{m_{H+}^2} D \right)^{-3/5} \right] \quad (31)$$

where $x = \omega_0 t$ and

$$D = \frac{\pi e^2}{m_{O+}^2 \omega_0 u^2} \int_0^x \epsilon_k dt$$

and we have assumed $\omega/k \sim u$.

In (31), the asymptotic dependence of the LHW energy density is different for the oxygen and hydrogen terms (the first and second terms on the right-hand side, respectively). The oxygen term yields $E \sim (t - t_0)^{3/2}$, a growth of energy, while the hydrogen term has the form $E \sim (t_0 - t)^{2/3}$, a damping of energy. Examining the powers shows that the oxygen term will dominate for large times and the LHW energy will grow.

The numerical analysis of (31) gives the same results. In the calculations below a maximum time $\omega_0 t \sim 10^3$ is used, supposing that ion velocities are 10^5 – 10^6 cm/s, the region of interaction with LFW is 10^8 – 10^9 cm, and ω_0 is a few s^{-1} . Figure 2 shows that the solution of (31) weakly depends on the initial conditions. The quantity \tilde{n}/n can be changed in magnitude 10–20 times without significant energy changes, allowing a wide range of concentrations of resonant hydrogen ions (Figure 3). The dependence on oxygen ion concentrations is more significant, however. If n_{O+}/n is changed from 10^{-2} to 3×10^{-2} , the LHW energy growth for $\omega_0 t \sim 10^3$ changes 20 times. Owing to the difference of the numerical coefficients in the terms on the right-hand side of (34), the exact dependence of the LHW energy on τ in the derivative in (31) is not very significant




Figure 2. The normalized LHW energy density ϵ/ϵ_0 from (31) for initial conditions of $\epsilon/\epsilon_0=0.5$ (solid curve), $\epsilon/\epsilon_0=0.05$ (dashed curve), and $\epsilon/\epsilon_0=0.005$ (dotted curve), as a function of normalized time $\omega_0 t$, with $\tilde{n}/n=0.005$ and $D^{-2/5}$.

(for $\omega_0 t \leq 10^3$). Even for a time dependence of $\partial F/\partial v \sim \tau^{-3/2}$, the energy grows for $\omega_0 t \sim 10^3$ (Figure 4) and the quasi-steady-state condition cannot be achieved. It follows from the above analysis that the quasilinear interaction does not lead to the saturation of the LHW instability, so the level of LHW energy is determined by the nonlinear process and is given by (25).

3.3. Plasma Energization

Now we can estimate the plasma energization produced by LFW due to LHW excitation. As it was found before, the quasilinear interaction of LHW with helium ions is small owing to the large difference between the wave phase velocity and the thermal velocity of the particles. Since the electrons are magnetized and the region of LHW phase velocities is small, the quasilinear interaction for electrons also can be neglected. The main quasilinear effect is LHW-hydrogen ion interaction. The LH turbulence also gives energy to the plasma due to induced scattering by ions and electrons. Using the spectral energy density from (25) and the hydrogen ion distribution function from (23), we can estimate the resonant hydrogen ion energy

and the dependence of this energy on the time of interaction with LHW,

$$K(t) = \frac{1}{\tilde{n}} \int \frac{mv^2}{2} f_{H^+} dv_{\perp} \approx \frac{m_{H^+} v_{TH^+}^2}{2} \frac{u^2}{v_{TH^+}^2} \left[3.5 \times 10^{-2} \frac{v_{TH^+}^2}{u^2} \gamma t \right]^{2/5} \quad (32)$$

where \tilde{n} is the number of resonant hydrogen ions.

Let us estimate the energy flow Q in the plasma particles due to induced scattering by ions and electrons. For $\omega \sim 3-4\omega_{LH}$ the result is of the same order of magnitude for both species [Musher *et al.*, 1978]. For the stationary state,

$$\int \gamma W_k dk = Q \quad (33)$$

where W_k is the spectral energy density of LH oscillations. Using (25), it is found that

$$\frac{Q}{nT} \sim \frac{\gamma^2 \omega}{\omega_{Be}^2} \quad (34)$$

4. Discussion and Results for Magnetospheric Plasma

Let us discuss the main results and the assumptions that are used in the analyses of LFW interaction with a multicomponent plasma due to LHW excitation and the possibility of their applications to the magnetospheric plasma.

In obtaining the LHW frequency (8), it was supposed that the LFW frequency is small compared with the LHW frequency and growth rate. If the LFW is an ion cyclotron wave (ICW) with a frequency near the helium cyclotron frequency [Young *et al.*, 1981] or oxygen cyclotron frequency [Fraser *et al.*, 1992], the frequency of LHW is $\omega = 2\omega_{LH}$ (7), γ is determined by (9), and the condition $\gamma > \omega_0$ for $\omega_0 \approx \omega_{BHe^+}$ leads to the inequality $c_{O^+} u_{O^+}^2 / v_{TH^+}^2 \gg 10^{-2}$. For the oxygen drift velocity $u_{O^+} \sim 3v_{TH^+}$ this condition is satisfied even for an oxygen ion concentration $c_{O^+} \sim 1\%$. For $\omega_0 \approx \omega_{BO^+}$ the inequality is 4 times weaker.




Figure 3. The normalized LHW energy density from (31) for the initial condition $\epsilon/\epsilon_0=0.005$ as a function of $\omega_0 t$ with $D^{-2/5}$. The solid curve is for $\tilde{n}/n=0.1$, the dashed curve is $\tilde{n}/n=0.05$, and the dotted curve is $\tilde{n}/n=0.005$.




Figure 4. The normalized LHW energy density ϵ/ϵ_0 from (31) for the initial condition $\epsilon/\epsilon_0=0.005$ as a function of $\omega_0 t$ with $\tilde{n}/n=0.005$. The solid curve is for $D^{-7/5}$, the dashed curve is $D^{-4/5}$, and the dotted curve is $D^{-2/5}$.

Another assumption used in obtaining the LHW growth rate is about the uniformity of the plasma, external magnetic field, and the electric field of LFW. The wavelength of the LHW can be estimated as $\lambda \sim 2\pi u_{O^+} / \omega_{LH}$, and even for $L \sim 7$ it is of order $\lambda \sim 10^5$ cm. Therefore this assumption is usually fulfilled, because the ICW wavelength is greater than 10^7 cm.

Let us discuss the LFW electric fields needed for LHW excitation. LHW excitation in a plasma with an oxygen mixture of 1-3% is possible if the oxygen ion drift velocity u_{O^+} is 3.5-4 times greater than the thermal velocity of hydrogen ions. In this case, the LHW Cherenkov damping on hydrogen ions is small and the development of the instability is possible. Taking into account that the instability develops on a timescale much smaller than the LFW period, the LHW generation is possible even if the oxygen ions have such a drift velocity only during a short part of the LFW period.

As was found in section 2.1 in (3), the drift velocity of a particle in a LFW electric field depends on the mass of the particle and the ratio of the LFW electric field frequency to the particle's cyclotron frequency. Expressions (3) and (4) are correct if $|\omega_0 - \omega_{B\alpha}| \gg k_0 v_{T\alpha}$, where k_0 is the wave vector of LFW. Owing to the large LFW wavelength, the right-hand side of this inequality is small. For ICW with a frequency of the order of the helium ion cyclotron frequency and an oxygen ion temperature 1-2 eV, $k_0 v_{T_{O^+}} \sim 5 \times 10^{-4}$. Therefore (3) and (4) are correct for ICW frequencies close to the particle's gyrofrequency.

Now let us estimate the possibility of LHW generation using observed LFW magnetic and electric fields. The plasma composition and the hydrogen ion temperature are not given in the papers cited below, so it was assumed that oxygen concentration is of order 1% and the hydrogen ion temperature is 1 eV. First, let us present the cases to be discussed.

Young et al. [1981], for an event observed on board GEOS 1 (August 16, 1977; $L \sim 7$; 1200 LT), found that the electric fields were 2.9 mV/m (1120 UT) and 4.3 mV/m (1215 UT); the wave frequency was $f = 0.6$ Hz, the frequency range was $\Delta f \sim 0.1$ Hz, and the helium gyrofrequency $f_{He^+} = 0.45$ Hz.

The event observed by ISEE 1 and 2 (August 22, 1978; $L \sim 7$, 1700 LT) is described by *Fraser et al.* [1992]. For this event the electric field was found to be 1.4 mV/m; the wave frequency $f = 0.12$ Hz, and the oxygen gyrofrequency $f_{O^+} \approx 0.06$ Hz.

In the case described by *LaBelle et al.* [1988] and *LaBelle and Treumann* [1992] (AMPTE/IRM observation, June 6, 1985; $L \sim 4.5$; 1547 UT) the electric field in full is not given. Our estimation based on this data leads to an electric field amplitude of ~ 1.8 mV/m. The wave frequency in this case is close to the oxygen ion gyrofrequency, $f = 0.15$ Hz, $f_{O^+} = 0.125$ Hz; and the frequency region is $\Delta f \sim 0.15f$.

In all of these cases, the oxygen ion drift velocity is 3.5 or more times greater than the hydrogen ion thermal velocity, and so LHW can be excited. In the last two examples the resonance denominator in the expression for the oxygen ion drift velocity (3) is significant. Taking into account that the listed observations were obtained in the region near the equator for $L \sim 5-7$, it can be found that the difference $\omega_0 - \omega_{BO^+}$ does not change significantly in a region of order R_E along the magnetic field lines.

The next supposition used is that the LFW is monochromatic. In two of the cases cited above (*Fraser et al.* [1992] and *LaBelle et al.* [1988] and *LaBelle and Treumann* [1992]) it is

quasi-monochromatic with $\Delta f / f \sim 0.1$. *Young et al.* [1981] reported that the ICW emissions observed on board GEOS 1 and 2 "are often quasi-monochromatic ($\Delta f / f \sim 0.1-0.2$)."

From the previous analysis it can be concluded that LHW excitation due to LFW activity is a possible mechanism of LFW interaction with multicomponent magnetospheric plasma, at least at $L > 5$ near the equatorial region. As discussed by *Khazanov et al.* [1996], the same mechanism of interaction is possible in the auroral zone.

Now we will estimate the plasma energization due to the discussed mechanism for magnetospheric plasma. For this calculation, let us take the magnetospheric plasma in the region of the plasmopause ($L \sim 4-5$) with the densities of the hydrogen, helium, and oxygen ions in the ratio 100:10:1, respectively. The plasma is assumed to be isothermal with a temperature of 1 eV. For the LFW with the frequency 0.15 Hz and oxygen ion gyrofrequency $f_{O^+} = 0.125$ Hz, the needed electric field amplitude of LFW, E_0 (see (8)), is 1.8 mV/m.

The LHW frequency in that region, given by (7), is $\omega \sim (0.5-1) \times 10^{-3} \text{ s}^{-1}$, and the growth rate is proportional to the oxygen ion concentration c_{O^+} (9), $\gamma \sim 5c_{O^+}\omega$. The wavelength of LH oscillation is $\lambda = 2\pi u / \omega \sim 10^4$ cm, and the envelope of generation is $\Delta \lambda \sim \lambda$.

We will suppose that the LFW electric field as well as the external magnetic field and plasma parameters are the same in a region of order $\ell \sim 10^8-10^9$ cm. Then the LHW turbulence in this region can be treated as uniform. The tracing time for hydrogen ions for such a region is of order 10^2-10^3 s. It can be taken that during this time the LFW and LHW activities are quasi-steady. Then, from (32), the energy obtained by the resonate hydrogen ions during the tracing time can be found. It is of order 20-40 eV. The number of resonant hydrogen ions can be estimated supposing that their initial distribution is Maxwellian. It is equal to the number of particles with velocities above the minimum LHW phase velocity. In the case under consideration, the LHW phase velocity is $3.5 v_{Th^+}$ and the number of resonant hydrogen ions is $\sim 0.5\%$. The energy of the core plasma, found from (34), is doubled for the period of interaction owing to induced scattering of LHW.

5. Summary

The preceding analysis demonstrates that LFWs in a multicomponent plasma can generate LH oscillations. The energy density of the LFW electric field needed is significantly less than in the case of a purely hydrogen plasma. The oxygen ions play the role of a beam, and in the linear approximation the concentration dependence of the growth rate and the kinetic and hydrodynamic limits of LHW generation are the same as those of a beam instability. The quasilinear effects do not lead to saturation of the instability, and the energy density level is determined by the nonlinear processes of induced scattering.

In the region of resonant velocities the hydrogen ions are accelerated and the concentration of accelerated ions weakly depends on the oxygen ion concentration. The nonresonant particles are weakly heated owing to induced scattering of LH oscillations.

For a given region of the magnetosphere ($L \sim 4-5$) with LFW electric fields of 2 mV/m, plasma temperatures of 1-2 eV, and an oxygen ion concentration of $\sim 1\%$, the resonant hydrogen ions ($\sim 0.5\%$) obtain energies of 20-40 eV. Therefore the discussed mechanism may lead to the growth of a tail in the

hydrogen distribution function in the presence of low-frequency activity.

Acknowledgments. This work was supported at MSFC by the NASA Office of Space Science under UPN 432-20-00, at the University of Alabama in Huntsville by NSF grant ATM-9523699, and at the University of Michigan under NASA GSRP grant NGT-51335. G. V. Khazanov held a National Research Council-Marshall Space Flight Center Senior Research Associateship while this work was performed.

The Editor thanks M. Schulz and another referee for their assistance in evaluating this paper.

References

- Akhiezer, A. I. (Ed.), *Plasma Electrodynamics*, Pergamon, Tarrytown, NY, 1975.
- Allan, W., Ponderomotive transport in the magnetosphere, *J. Geophys. Res.*, **97**, 8483, 1992.
- Allan, W., Plasma energization by the ponderomotive force of magnetospheric standing Alfvén waves, *J. Geophys. Res.*, **98**, 11,383, 1993.
- Allan, W., J. R. Manuel, and E. M. Poulter, Magnetospheric cavity modes: Some nonlinear effects, *J. Geophys. Res.*, **96**, 11,461, 1991.
- Anderson, B. J., R. E. Erlandson, and L. J. Zanetti, A statistical study of Pc 1-2 magnetic pulsations in the equatorial magnetosphere; 1. Equatorial occurrence distributions, *J. Geophys. Res.*, **97**, 3075, 1992.
- Boehm, M. H., C. W. Carlson, J. P. McFadden, J. H. Clemmons, and F. S. Mozer, High-resolution sounding rocket observation of large-amplitude Alfvén waves, *J. Geophys. Res.*, **95**, 12,157, 1990.
- Brown, D. G., G. R. Wilson, J. L. Horwitz, and D. L. Gallagher, "Self-consistent" production of ion conics on return current region auroral field lines: A time-dependent semikinetic model, *Geophys. Res. Lett.*, **18**, 1841, 1991.
- Brown, D. G., J. L. Horwitz, and G. R. Wilson, Synergetic effects of hot plasma-driven potentials and wave-driven ion heating on auroral ionospheric plasma transport, *J. Geophys. Res.*, **100**, 17,499, 1995.
- Chang, T., and M. André, Ion heating by low frequency waves, in *Auroral Plasma Dynamics*, edited by R. L. Lysak, *Geophys. Monogr. Ser.*, vol. 80, p. 207, AGU, Washington, D. C., 1993.
- Chang, T., and B. Coppi, Lower hybrid acceleration and ion evolution in the subauroral region, *Geophys. Res. Lett.*, **8**, 1253, 1981.
- Chernov, A. A., G. V. Khazanov, and Y. V. Konikov, On the linear theory of the oblique Alfvén and fast magnetosonic waves in the Earth's magnetosphere, *Planet. Space Sci.*, **38**, 823, 1990.
- Comfort, R. H., J. H. Waite Jr., and C. R. Chappell, Thermal ion temperatures from the retarding ion mass spectrometer on DE 1, *J. Geophys. Res.*, **90**, 3475, 1985.
- Cornwall, J. M., Cyclotron instabilities and electromagnetic emission in the ultra low frequency and very low frequency ranges, *J. Geophys. Res.*, **70**, 61, 1965.
- Cornwall, J. M., Micropulsations and the outer radiation zone, *J. Geophys. Res.*, **71**, 2185, 1966.
- Cornwall, J. M., F. V. Coroniti, and R. M. Thorne, Turbulent loss of ring current protons, *J. Geophys. Res.*, **75**, 4699, 1970.
- Cornwall, J. M., F. V. Coroniti, and R. M. Thorne, Unified theory of SAR-arc formation at the plasmopause, *J. Geophys. Res.*, **76**, 4428, 1971.
- Decreau, P. M. E., C. Beghin, and M. Parrot, Global characteristics of cold plasma in the equatorial plasmasphere region as deduced from the GEOS-1 mutual impedance probe, *J. Geophys. Res.*, **87**, 695, 1982.
- Erlandson, R. E., L. J. Zanetti, T. A. Potemra, L. P. Block, and G. Holmgren, Viking magnetic and electric field observations of Pc 1 waves at high latitudes, *J. Geophys. Res.*, **95**, 5941, 1990.
- Fraser, B. J., J. C. Samson, Y. D. Hu, R. L. McPherron, and C. T. Russel, Electromagnetic ion cyclotron waves observed near the oxygen cyclotron frequency by ISEE 1 and 2, *J. Geophys. Res.*, **97**, 3063, 1992.
- Galeev, A. A., Plasma turbulence in the magnetosphere with special regard to plasma heating, in *Physics of the Hot Plasma in the Magnetosphere*, edited by B. Hultquist and L. Stenflo, p. 251, Plenum, New York, 1975.
- Gamayunov, K. V., E. N. Krivorutsky, and G. V. Khazanov, Hydrodynamic description of magnetosphere plasma with due regard to the wave activity of Alfvén and fast magnetosonic waves, *Planet. Space Sci.*, **39**, 1097, 1991.
- Gamayunov, K. V., E. N. Krivorutsky, A. A. Veryaev, and G. V. Khazanov, Saturation of Alfvén oscillations in the ring current region due to generation of lower hybrid waves, *Planet. Space Sci.*, **40**, 477, 1992.
- Ganguli, S. B., and P. J. Palmadesso, Plasma transport in the auroral return current region, *J. Geophys. Res.*, **92**, 8673, 1987.
- Ganguli, S. B., and P. J. Palmadesso, Generalized fluid model of plasma outflow processes in the topside ionosphere, *Adv. Space Res.*, **8**, 69, 1988.
- Gendrin, R., General relationships between wave amplification and particle diffusion in a magnetoplasma, *Rev. Geophys.*, **19**, 171, 1981.
- Gendrin, R., Wave particle interactions as an energy transfer mechanism between different particle species, *Space Sci. Rev.*, **34**, 271, 1983.
- Gendrin, R., Consequences of hydromagnetic waves on magnetospheric particle dynamics, *Space Sci. Rev.*, **42**, 515, 1985.
- Gorbachev, O. A., Y. V. Konikov, and G. V. Khazanov, Quasilinear heating of electrons in the Earth's plasmasphere, *Pure Appl. Geophys.*, **127**, 545, 1988.
- Gorbachev, O. A., G. V. Khazanov, K. V. Gamayunov, and E. N. Krivorutsky, A theoretical model for the ring current interaction with the Earth's plasmasphere, *Planet. Space Sci.*, **40**, 859, 1992.
- Gringauz, K. I., Plasmasphere and its interaction with the ring current, *Space Sci. Rev.*, **34**, 245, 1983.
- Gringauz, K. I., Structure and properties of the Earth plasmasphere, *Adv. Space Res.*, **5**, 391, 1985.
- Iyemori, T., and K. Hayashi, Pc 1 micropulsations observed by Magsat in the ionospheric F region, *J. Geophys. Res.*, **94**, 93, 1989.
- Kennel, C. F., and H. E. Petschek, Limit on stably trapped particle fluxes, *J. Geophys. Res.*, **71**, 1, 1966.
- Khazanov, G. V., and A. A. Chernov, The effect of Alfvén and fast magnetosonic turbulence on the heating of electrons and kinetic coefficients of the plasmasphere (in Russian), *Ionos. Invest.*, **44**, 100, 1988.
- Khazanov, G. V., T. E. Moore, E. N. Krivorutsky, J. L. Horwitz, and M. W. Liemohn, Lower hybrid turbulence and ponderomotive force effects in space plasmas subjected to low-frequency waves, *Geophys. Res. Lett.*, **23**, 797-800, 1996.
- Konikov, Y. V., O. A. Gorbachev, G. V. Khazanov, and A. A. Chernov, Hydrodynamical equations for thermal electrons taking into account their scattering on ion cyclotron waves in the outer plasmasphere of the Earth, *Planet. Space Sci.*, **37**, 1157, 1989.
- Kozyra, J. U., L. H. Brace, T. E. Cravens, and A. F. Nagy, A statistical study of the subauroral electron temperature enhancement using Dynamics Explorer 2 Langmuir probe observations, *J. Geophys. Res.*, **91**, 11,270, 1986.
- Labelle, J., and R. A. Treumann, Poynting vector measurements of electromagnetic ion cyclotron waves in the plasmasphere, *J. Geophys. Res.*, **97**, 13,789, 1992.
- LaBelle, J., R. A. Treumann, W. Baumjohann, G. Haerendel, N. Schopke, G. Paschmann, and H. Luhr, The duskside plasmapause/ring current interface: Convection and plasma wave observations, *J. Geophys. Res.*, **93**, 2573, 1988.
- Lee, N. C., and G. K. Parks, Ponderomotive acceleration of ions by circularly polarized electromagnetic waves, *Geophys. Res. Lett.*, **23**, 327, 1996.
- Li, X., and M. Temerin, Ponderomotive effects on ion acceleration in the auroral zone, *Geophys. Res. Lett.*, **20**, 13, 1993.
- Liemohn, H. B., Cyclotron resonance amplification of VLF and ULF whistlers, *J. Geophys. Res.*, **72**, 39, 1967.

- Lin, J., J. L. Horwitz, G. R. Wilson, C. W. Ho, and D. G. Brown, A semikinetic model for early stage plasmasphere refilling; 2, Effects of wave-particle interactions, *J. Geophys. Res.*, **97**, 1121, 1992.
- Lin, J., J. L. Horwitz, G. R. Wilson, and D. G. Brown, Equatorial heating and hemispheric decoupling effects on inner magnetospheric core plasma evolution, *J. Geophys. Res.*, **99**, 5727, 1994.
- Mauk, B. H., Electromagnetic wave energization of heavy ions by the electric "phase bunching" process, *Geophys. Res. Lett.*, **9**, 1163, 1982.
- Musher, S. L., A. M. Rubenchik, and B. I. Sturman, Collective effects associated with low hybrid heating of plasma, *Plasma Phys.*, **20**, 1131, 1978.
- Olmelchenko, Y. A., V. D. Shapiro, V. I. Shevchenko, M. Ashour-Abdalla, and D. Schriver, Modified lower hybrid fan instability excited by precipitating auroral electrons, *J. Geophys. Res.*, **99**, 5965, 1994.
- Omura, V., M. Ashour-Abdalla, R. Gendrin, and K. Quest, Heating of thermal helium in the equatorial magnetosphere: A simulation study, *J. Geophys. Res.*, **90**, 8281, 1985.
- Retterer, J. M., T. Chang, and J. R. Jasperse, Ion acceleration by lower hybrid waves in the supauroral region, *J. Geophys. Res.*, **91**, 1609, 1986.
- Shapiro, V. D., and V. I. Shevchenko, Astrophysical plasma turbulence, I, *Sov. Sci. Rev. E. Astrophys. Space Phys.*, **6**, 425, 1988.
- Shapiro, V. D., V. I. Shevchenko, G. I. Solov'ev, V. P. Kalinin, R. Bingham, R. Z. Sagdeev, M. Ashour-Abdalla, J. Dawson, and J. J. Su, Wave collapse at the lower-hybrid resonance, *Phys. Fluids B*, **5**, 3148, 1993.
- Singh, N., Refilling of a plasmaspheric tube—Microscopic plasma processes, in *Modeling Magnetospheric Plasma*, *Geophys. Monogr. Ser.* vol. 44, edited by T. E. Moore and J. H. Waite Jr., p. 87, AGU, Washington, D. C., 1988.
- Singh, N., and D. G. Torr, Effects of ion temperature anisotropy on the interhemispheric plasma transport during plasmaspheric refilling, *Geophys. Res. Lett.*, **17**, 925, 1990.
- Tanaka, M., Simulations of heavy ion heating by electromagnetic ion cyclotron waves by proton temperature anisotropies, *J. Geophys. Res.*, **90**, 6459, 1985.
- Young, D. T., S. Perraut, A. Roux, C. de Villedary, R. Gendrin, A. Korth, G. Kremser, and D. Jones, Wave-particle interactions near Ω_{He^+} observed on GEOS 1 and 2, I, Propagation of ion cyclotron waves in the He^+ rich plasma, *J. Geophys. Res.*, **86**, 6755, 1981.

J. L. Horwitz and E. N. Krivorutsky, Department of Physics and Center for Space Plasma and Aeronomic Research, The University of Alabama in Huntsville, Huntsville, AL 35899.

G. V. Khazanov and T. E. Moore, Space Sciences Laboratory, NASA Marshall Space Flight Center, Code ES83, Huntsville, AL 35812. (e-mail: Khazagv@ssl.msfc.nasa.gov)

M. W. Liemohn, Space Physics Research Laboratory, University of Michigan, Ann Arbor, MI 48109.

(Received March 15, 1996; revised July 19, 1996; accepted July 23, 1996.)