

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public report of burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS	
EVIDENCE FOR INCREASED BETA-ADRENORECEPTOR RESPONSIVENESS INDUCED BY 14 DAYS OF SIMULATED MICROGRAVITY IN HUMANS			PE-62202F PR-7755 TA-B5 WU-01	
6. AUTHOR(S)				
Victor A. Convertino, Jill L. Polet, Keith A. Engelke, G. Wyckliffe Hoffler, Lynda D. Lane, and C. Gunnar Blomqvist				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION	
Armstrong Laboratory (AFMC) Aerospace Medicine Directorate Clinical Sciences Division, Physiological Research Branch 2507 Kennedy Circle Brooks AFB, TX 78235-5117			AL/AO-JA-1996-0041	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING/MONITORING	
Life and Biomedical Sciences & Application Division Mail Stop ULR National Aeronautics Space Administration Headquarters Washington, DC 220546-0001				
11. SUPPLEMENTARY NOTES				
Armstrong Laboratory Technical Monitor: Dr. Victor A. Convertino				
12a. DISTRIBUTION/AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
Approved for public release; Distribution is unlimited.				
13. ABSTRACT (Maximum 200 words)				
<p>We studied hemodynamic responses to alpha- and beta- receptor agonists in 8 healthy men (38 ± 2 yrs) before and after 14 days of 6 degree head-down tilt (HDT) to test the hypothesis that increased adrenergic responsiveness is induced by prolonged exposure to microgravity. Immediately following a 30-min baseline period, a steady-state infusion of isoproterenol (ISO) was used to assess beta 1- and beta 2-adrenergic responsiveness. ISO was infused at three graded constant rates of 0.005, 0.01 and 0.02 ug/kg/min. After heart rate and blood pressure had been allowed to return to baseline levels following ISO infusion, graded infusion of phenylephrine (PE) was used to assess responsiveness of alpha 1-vascular receptors. PE was infused at three graded constant rates of 0.25, 0.50 and 1.00 ug/kg/min. Each infusion interval for both drugs was 9 min. During the infusions, constant monitoring of beat-to-beat blood pressure and heart rate was performed and leg blood flow was measured with occlusion plethysmography at each infusion level. The slopes calculated from linear regressions between ISO and PE doses and changes in heart rate, blood pressure, and leg vascular resistance for each subject were used to represent alpha- and beta- adrenoreceptor responsiveness. Fourteen days HDT increased the slopes of heart rate (1056 ± 107 to 1553 ± 83 beats/ug/kg/min; $P = 0.014$) and vasodilation (-469 ± 111 to -1446 ± 309 PRU/ug/kg/min; $P = 0.0224$) to ISO infusion. There was no alteration in blood pressure or vascular resistance responses to PE infusion after HDT. Our results provide evidence that microgravity causes selective increases in beta 1- and beta 2-adrenergic responsiveness without affecting alpha 1-vascular responses.</p>				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
autonomic function; sympathetic; adrenoreceptors; heart rate; blood pressure; baroreflex; vascular resistance			24	
17. SECURITY CLASSIFICATION OF REPORT			18. PRICE CODE	
Unclassified				
19. SECURITY CLASSIFICATION OF THIS PAGE			20. LIMITATION OF ABSTRACT	
Unclassified			UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev 2-88) Prescribed by ANSI Std Z-39-18
298-102 COMPUTER GENERATED

Evidence for increased β -adrenoreceptor responsiveness induced by 14 days of simulated microgravity in humans

VICTOR A. CONVERTINO, JILL L. POLET, KEITH A. ENGELKE,
G. WYCKLIFFE HOFFLER, LYNDIA D. LANE, AND C. GUNNAR BLOMQVIST
*Physiology Research Branch, Clinical Sciences Division, Brooks Air Force Base, Texas 78235;
Biomedical Operations and Research Office, National Aeronautics and Space Administration,
Kennedy Space Center, Florida 32899; Department of Physiology, University of Florida,
Gainesville, Florida 32611; and University of Texas Southwestern Medical Center, Dallas, Texas 75235*

Convertino, Victor A., Jill L. Polet, Keith A. Engelke, G. Wyckliffe Hoffler, Lyndia D. Lane, and C. Gunnar Blomqvist. Evidence for increased β -adrenoreceptor responsiveness induced by 14 days of simulated microgravity in humans. *Am. J. Physiol.* 278 (Regulatory Integrative Comp. Physiol. 42): R93–R99, 1997.—We studied hemodynamic responses to α - and β -receptor agonists in eight healthy men before and after 14 days of 6° head-down tilt (HDT) to test the hypothesis that increased adrenoreceptor responsiveness is induced by prolonged exposure to simulated microgravity. Steady-state infusions of isoproterenol (Iso) at rates of 0.006, 0.01, and 0.02 $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ were used to assess β_1 - and β_2 -adrenoreceptor responsiveness. Infusions of phenylephrine (PE) at rates of 0.25, 0.50, and 1.00 $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ were used to assess responsiveness of α_1 -vascular adrenoreceptors. Slopes calculated from linear regressions between Iso and PE doses and changes in heart rate, blood pressure, and leg vascular resistance (occlusion plethysmography) for each subject were used as an index of α - and β -adrenoreceptor responsiveness. HDT increased the slopes of heart rate ($1,066 \pm 107$ to $1,553 \pm 83$ beats $\mu\text{g}^{-1}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$; $P = 0.014$) and vasodilation (-469 ± 111 to $-1,446 \pm 309$ peripheral resistance units $\mu\text{g}^{-1}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$; $P = 0.0224$) to Iso infusion. There was no alteration in blood pressure or vascular resistance responses to PE infusion after HDT. Our results provide evidence that simulated microgravity causes selective increases in β_1 - and β_2 -adrenoreceptor responsiveness without affecting α_1 -vascular adrenoreceptor responses.

autonomic function; sympathetic activity; heart rate; blood pressure; baroreflex; vascular resistance

ALTERATIONS IN autonomic function are commonly exhibited in individuals who have been exposed to microgravity environments. These include the loss of aortic, carotid, and cardiopulmonary baroreflex reserve to buffer against fluctuations in arterial and central venous blood pressures (6, 7, 9, 14). These changes in autonomic reflex control have resulted in excessive cardioacceleration and limitations in peripheral resistance responses during return to ambulation in terrestrial gravity. There is evidence that changes in central nervous system integration of afferent stimuli (12), reduced baseline vagal tone (9), and/or elevated sympathetic activity (28) may contribute as underlying mechanisms.

Increased sensitivity of end-organ responses to neuroendocrine stimuli resulting from prolonged exposure to the relative inactivity of microgravity has recently been hypothesized (28). This notion is based on the inverse relationship between circulating norepinephrine (NE)

and β -adrenoreceptor sensitivity. β -Adrenoreceptor activity is reduced in individuals who have elevated plasma NE as a result of regular exposure to upright posture (18) and physical exercise (4). In contrast, adrenoreceptor hypersensitivity has been reported in patients with dysautonomias in which circulating catecholamines are absent or reduced (25). Taken together, these studies and the observation that circulating plasma NE can be reduced during spaceflight (18) and in ground-based simulations of microgravity (7, 16, 23) prompt the suggestion that adrenoreceptor hypersensitivity may be a consequence of the adaptation to spaceflight.

We conducted an experiment to study the effects of prolonged exposure to a ground-based analog of microgravity (6° head-down tilt (HDT)) on various regulatory functions in human subjects (8–12). As part of that study, we tested the hypothesis that adaptation to simulated microgravity increases cardiovascular responses to adrenoreceptor agonists and that this adaptation is associated with reduced levels of circulating NE.

METHODS

Subjects. After being informed of all procedures and risks, eight healthy, normotensive, nonsmoking men with mean (\pm SE) age of 38 ± 2 yr, height of 183 ± 2 cm, and weight of 81 ± 3 kg gave their written consent to serve as subjects for this investigation. All experimental procedures and protocols were approved by the Human Research Review Boards of the National Aeronautics and Space Administration (NASA)-Kennedy Space Center, NASA-Ames Research Center, and Brooks Air Force Base. Selection of subjects was based on results of a screening evaluation composed of a detailed medical history, physical examination, blood chemistry analysis, urinalysis, chest X-ray, and electrocardiogram (ECG). Subjects were made familiar with all laboratory personnel, procedures, and protocols during an orientation session conducted before the study.

General protocol. The experimental protocol comprised 4 days of ambulatory control followed by 16 days of 6° HDT and 2 days of post-HDT reambulation. During HDT, the subjects were continuously monitored by staff nurses to ensure that they remained head-down without interruption and that no physical exercise was performed by the subjects between pre- and post-HDT measurements.

During the 22-day experimental period, subjects lived 24 h/day in the Human Research Facility at NASA-Ames Research Center and followed the same controlled diet. The average daily caloric intake was 2,600–2,800 kcal (45% carbohydrate, 38% fat, 17% protein). Dietary sodium and potassium were held constant at ~ 120 and 60–80 meq/day,

respectively. Fluid intake was ad libitum; the average was $1,844 \pm 17$ ml/day. The photoperiod was 16 h light to 8 h dark, with lights on at 0700. The 16-day HDT period was chosen because this represents the projected minimum duration of future Extended Duration Orbiter space missions. The 6° head-down position was chosen because actual flight changes in some cardiovascular responses are closely simulated by this ground-based analog (5). Each subject underwent an adrenoreceptor response test on the third day before HDT and on day 14 of HDT. In addition, antecubital venous blood samples were drawn for determination of circulating NE and plasma volume.

Measurements of adrenoreceptor responsiveness. Each subject was moved to a quiet room on a gurney specially designed with 6° HDT, and intravenous catheters were placed in the antecubital vein of each arm while the subject was supine. All infusions were performed through the intravenous catheter placed in the left arm using an Auto-Syringe infusion pump, and administration rates were achieved by appropriate combinations of infusion rate and agonist concentrations. The catheter placed in the right arm was used to draw blood samples for analysis of plasma NE. Instrumentation was completed during a 30-min period to establish a resting baseline. Baseline measurements included plasma volume, plasma norepinephrine, heart rate (ECG), blood pressure (Finapres), and leg blood flow (LBF) (occlusion plethysmography). After baseline measurements, graded infusions of β - and α -adrenoreceptor agonists were performed with isotonic saline as a vehicle. The total volume infused was <50 ml. A recovery period of at least 30 min was allowed between the two agonist infusion protocols to allow hemodynamic measurements to return to preinfusion baseline levels. Before and during both infusion protocols, constant monitoring of beat-to-beat blood pressure and heart rate was performed. In addition, leg blood flows were measured at each infusion level. A blood sample (20 ml) for plasma NE determination was drawn at the end of the third infusion level for each drug.

β -Adrenoreceptor responsiveness. Immediately after the 30-min baseline period, infusions of isoproterenol (Iso) were used to assess the responsiveness of β_1 - and β_2 -adrenoreceptors. Iso was infused at three graded constant rates of 0.005, 0.01, and 0.02 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$. Each infusion interval was 9 min in duration to establish steady state and allow adequate time for all measurements. The protocol and dosages of Iso (and phenylephrine (PE), see below) were adopted from those used during spaceflight experiments on NASA's Space Life Sciences (SL6)-1 and SL6-2 missions. Appropriate dosages for these adrenergic agonists were determined by laboratory experience to produce safe but significant physiological responses. An elevation of heart rate by 25 beats/min above resting baseline was the predetermined end point for test termination. However, all β -adrenoreceptor responsiveness test protocols were completed. Linear regression relationships were then constructed relating the increase in heart rate and the decrease in leg vascular resistance to the dose of Iso. The slopes describing the linear stimulus-response relationship between the dose of Iso and heart rate and leg vascular resistance provided a measure of the functional responses of β_1 - and β_2 -adrenoreceptors, respectively.

α_1 -Adrenoreceptor responsiveness. After heart rate and blood pressure had been allowed to return to baseline levels following Iso infusions, graded infusion of the α_1 -adrenoreceptor agonist PE was used to assess the responsiveness of these vascular receptors. PE was infused at three graded constant rates of 0.25, 0.50, and 1.00 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$. As in the case of the Iso infusion test, each infusion interval was 9 min in duration. An elevation of systolic blood pressure of 20 mmHg

above or reflex reduction of heart rate 20 beats/min below resting baseline were predetermined and points for test termination. One pre-HDT and two post-HDT tests were terminated during the third stage of infusion using these criteria. The response of α_1 -adrenoreceptors was assessed by relating the PE dose with the increment in mean arterial pressure and reduction in leg vascular resistance. The relationships between PE doses and blood pressure and leg vascular resistance were linear, and the slopes describing these relationships were used to represent an index of α_1 -adrenoreceptor responsiveness.

Heart rate and blood pressure. Continuous heart rate was recorded using a standard three-lead ECG. A photoplethysmographic Finapres finger cuff blood pressure monitoring device was used to provide continuous beat-by-beat measurement of peripheral systolic and diastolic arterial blood pressures. Finapres recordings were verified by blood pressure measured during each stage of infusion with a sphygmomanometer. Heart rate and blood pressure responses were saved as a digital record for subsequent data analysis. Mean arterial pressure was calculated as systolic pressure plus twice diastolic divided by three.

Leg vascular resistance. LBF was measured using venous occlusion plethysmography employing a dual loop mercury-in-Silastic strain gauge placed around the left leg at the point of maximal calf circumference to determine changes in venous volume. Venous outflow from the leg was prevented by the placement of a cuff around the thigh just above the knee using an occlusion pressure of +40 mmHg. Arterial occlusion to reduce blood flow to the foot was applied by an ankle cuff inflated at a pressure of +250 mmHg. After ankle cuff inflation for 1 min, venous occlusion was initiated for 10 s, followed by its release for 10 s for six sequential occlusions. The relative change (percent) in strain gauge length over 10 s was quantified as a volume of blood per unit time, i.e., LBF. The 10-s occlusions were repeated during the final 2 min of drug infusion at each stage, and the average of the six measurements represented the LBF for that drug dose. An index of leg vascular resistance was calculated by dividing mean arterial pressure by average LBF during the final 2 min of each drug infusion and expressed as peripheral resistance units ((PRU); in $\text{mmHg} \cdot \text{l}^{-1} \cdot \text{min}$).

Plasma measurements. Plasma concentrations of NE were measured by high-performance liquid chromatography (Waters). NE was extracted by absorbing plasma samples onto alumina. After washing of the absorbed alumina with a dilute buffer solution, NE was eluted from the alumina when treated with an acidic solution. 3,4-Dihydroxybenzylamine (DHBA) was used as an internal standard, and extraction efficiency of NE and DHBA was based on the extraction of known standards. After extraction, the samples were assayed using a Waters 712 Wisp to inject the samples onto a reverse-phase C_{18} column. A Waters 460 electrochemical detector was used to determine the concentration of NE in the samples. The within-assay coefficient of variation (CV) was 1.4%, and between-assay CV was 3.8%.

Plasma volume was determined by a dilution technique using sterile solutions of Evans blue dye (New World Training, DeBary, FL) previously described and reported (8). Total circulating plasma NE was calculated as the product of plasma volume and plasma NE concentration.

Statistics. Standard descriptive statistics were performed on each of the response variables of interest, with results presented as means \pm SE. Standard paired *t*-test statistics were used to compare mean slopes of the dose-response relationships between drug infusions and heart rate, blood pressure, and vascular resistance before and after HDT and




Fig. 1. Dose-response relationships between isoproterenol (Iso) and heart rate (top) and log vascular resistance (bottom) before (O) and after (●) head-down tilt (HDT). Linear regressions are calculated from mean values. For heart rate, the linear equation for mean pre-HDT data is $y = 954 \cdot x - 57.4$ ($r^2 = 0.992$) and for mean HDT day 14 is $y = 1,429 \cdot x + 68.0$ ($r^2 = 0.990$). For log vascular resistance, the linear equation for mean pre-HDT data is $y = -471 \cdot x + 41.6$ ($r^2 = 0.784$) and for mean HDT day 14 is $y = -1,429 \cdot x + 65.7$ ($r^2 = 0.971$). bpm, Beats/min; PRU, peripheral resistance units.

for comparison of baseline heart rate, blood pressure, plasma NE concentrations, and total circulating NE before and after HDT. Repeated-measures analysis of variance was performed to determine differences between measurements in blood pressures and NE between pre- and post-HDT and across drug infusions. A least-significant difference (LSD) post hoc test was conducted to determine statistical differences across the drug treatments. The null hypothesis was rejected when $P < 0.05$.

RESULTS

Baseline measurements. Mean body weight was reduced from 81.0 ± 3.4 kg before to 79.5 ± 3.3 kg after HDT ($t = 4.446$, $df = 7$, $P = 0.0015$). Baseline heart rate and log vascular resistance were elevated by day

14 of HDT compared with pre-HDT (Fig. 1). Plasma volume was decreased by 16%, from $3,759 \pm 154$ ml before HDT to $3,159 \pm 114$ ml after HDT (5). Baseline plasma NE was reduced from 174 ± 11 pg/ml before HDT to 139 ± 6 pg/ml after HDT ($t = 2.878$, $df = 7$, $P = 0.0237$). As a result of the reduction in plasma volume, total circulating NE was dramatically lowered from 655 ± 49 ng before HDT to 434 ± 15 ng after HDT ($t = 4.685$, $P = 0.0022$).

Adrenoreceptor responsiveness. Fourteen days of HDT increased the average slope of the individual subject dose-response relationships between Iso and heart rate by 47%, from $1,058 \pm 107$ beats $\cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ before HDT to $1,553 \pm 83$ beats $\cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ on day 14 of HDT ($t = 3.235$, $df = 7$, $P = 0.0144$). Figure 1 (top) represents the regressions calculated from the mean (\pm SE) heart rates at each Iso level before and after HDT. Similarly, the average slope of the individual subject dose-response relationships between Iso and log vascular resistance increased by threefold from -469 ± 111 PRU $\cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ before HDT to $-1,446 \pm 309$ PRU $\cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ on day 14 of HDT ($t = 2.919$, $df = 7$, $P = 0.0224$). Figure 1 (bottom) represents the regressions calculated from the mean (\pm SE) log vascular resistances at each Iso level before and after HDT. Mean arterial pressure was unchanged during the Iso infusions and increased during PE infusions, but there were no differences in blood pressure responses to the infusions from before to after HDT (Tables 1 and 2).

The dose-response relationship between PE and log vascular resistance shifted upward on the response axis (Fig. 2, top) with HDT, and the average dose-response relationships between PE and mean arterial pressure before and after HDT were superimposed (Fig. 2, bottom). The average pre-HDT slope calculated from the individual subject dose-response relationships between PE and log vascular resistance (24.7 ± 5.9 PRU $\cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) was not altered ($t = 0.370$, $df = 7$, $P = 0.7224$) by HDT (27.0 ± 3.4 PRU $\cdot \mu\text{g}^{-1}$).

Table 1. Blood pressure responses at baseline and three levels of isoproterenol infusion before and after 14 days of HDT

	Baseline	Isoproterenol, $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$		
		0.005	0.010	0.020
Systolic blood pressure, mmHg				
Before HDT	126 \pm 2	127 \pm 3	127 \pm 4	132 \pm 3
After HDT	124 \pm 3	127 \pm 3	132 \pm 3	134 \pm 5
Diastolic blood pressure, mmHg				
Before HDT	76 \pm 1	70 \pm 1	66 \pm 2	67 \pm 2
After HDT	74 \pm 2	74 \pm 2	72 \pm 2	73 \pm 3
Mean arterial pressure, mmHg				
Before HDT	92 \pm 1	89 \pm 1	89 \pm 2	88 \pm 2
After HDT	91 \pm 2	91 \pm 2	92 \pm 2	94 \pm 3

Values are means \pm SE; HDT, head-down tilt. Analysis of variance revealed no statistical differences.

Table 2. Blood pressure responses at baseline and three levels of phenylephrine infusion before and after 14 days of HDT

	Baseline	Phenylephrine, $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$		
		0.25	0.50	1.00
Systolic blood pressure, mmHg				
Before HDT	127 \pm 2	132 \pm 3	134 \pm 3	139 \pm 3
After HDT	131 \pm 1	133 \pm 1	139 \pm 3	143 \pm 3
Diastolic blood pressure, mmHg				
Before HDT	77 \pm 2	79 \pm 3	80 \pm 3	84 \pm 3
After HDT	77 \pm 2	77 \pm 2	79 \pm 2	84 \pm 2
Mean arterial pressure, mmHg				
Before HDT	94 \pm 2	97 \pm 3	98 \pm 2	102 \pm 3
After HDT	95 \pm 1	96 \pm 2	99 \pm 2	104 \pm 2

Values are means \pm SE; analysis of variance revealed no statistical differences.

$\text{kg}^{-1} \cdot \text{min}^{-1}$). The average slopes calculated from the individual subject dose-response relationships between PE and mean arterial pressure before ($11.9 \pm 3.2 \text{ mmHg} \cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) and after ($12.6 \pm 2.3 \text{ mmHg} \cdot \mu\text{g}^{-1} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) HDT were not statistically distinguishable ($t = 0.165$, $df = 7$, $P = 0.8734$).

NE responses. Both plasma concentration and total circulating plasma NE during baseline rest before HDT ($174 \pm 11 \text{ pg/ml}$ and $655 \pm 49 \text{ ng}$, respectively) were greater ($t = 2.878$, $df = 7$, $P = 0.0237$ and $t = 4.685$, $df = 7$, $P = 0.0022$, respectively) than after HDT ($139 \pm 7 \text{ pg/ml}$ and $411 \pm 15 \text{ ng}$, respectively). Under all infusion conditions, both plasma concentration and total circulating plasma NE were lower [$F(1,7) = 6.67$, $P = 0.0363$] on day 14 of HDT compared with pre-HDT (Fig. 8). Compared with preinfusion baseline, plasma NE was increased by Iso infusion and decreased by PE infusion [$F(2,14) = 28.15$, $P = 0.0001$] both before and after HDT ($\text{LSD} \leq 67 \text{ pg/ml}$, $P < 0.05$).

DISCUSSION

We measured plasma NE, heart rate, blood pressure, and peripheral vascular responses during graded infusion of cardiac and vascular adrenoceptor agonists in eight healthy men before and after 14 days of 6° HDT to test the hypothesis that adaptation to simulated microgravity leads to adrenoceptor hypersensitivity. The major finding of this study was that HDT led to substantial increases in the heart rate and vasodilatory responses to a β -adrenergic agonist but had little effect on the vasoconstrictive response to α -adrenergic stimulation. The results of the present study also substantiated that both circulating concentrations and total content of NE were dramatically reduced during HDT. Our data may be the first to provide evidence that microgravity may cause selective increases in β_1 - and β_2 -adrenoceptor responsiveness in healthy humans that were associated with reduced circulating NE without affecting α_1 -vascular responses.

Tachycardia is a well-documented phenomenon associated with return to the upright posture following exposure to actual or simulated microgravity. Elevation in postflight heart rate has been attributed to hypovolemia (3), elevated catecholamines (1, 6, 12, 23, 18), and increased aortic baroreceptor responsiveness (9). Our data support the notion that increased responsiveness of cardiac β_1 -adrenergic receptors represents an additional mechanism that may contribute to postspace-flight tachycardia.

In addition to chronotropic effects, the alteration in cardiac adrenoceptor responsiveness was associated with frequent occurrence of junctional or nodal arrhythmias during agonist infusion in some of our subjects. This is particularly intriguing in light of observations that altered autonomic function was postulated as a




Fig. 2. Dose-response relationships between phenylephrine (PE) and log vascular resistance (top) and mean arterial pressure (bottom) before (○) and after (●) HDT. Linear regressions are calculated from mean values. For log vascular resistance, the linear equation for mean pre-HDT data is $y = 20.3 \cdot x + 43.0$ ($r^2 = 0.988$) and for mean HDT day 14 is $y = 22.4 \cdot x + 57.0$ ($r^2 = 0.991$). For mean arterial pressure, the linear equation for mean pre-HDT data is $y = 7.7 \cdot x + 94.4$ ($r^2 = 0.978$) and for mean HDT day 14 is $y = 10.4 \cdot x + 94.2$ ($r^2 = 0.974$).




Fig. 3. Plasma concentrations of norepinephrine (NE) during baseline rest and at the end of Iso and PE infusions before (○, broken lines) and after (●, solid lines) 14 days of HDT. HDT reduced plasma NE under all drug conditions ($P = 0.0063$). Compared with preinfusion baseline, plasma NE was increased by Iso infusion and decreased by PE infusion ($P = 0.0001$) both before and after HDT (least-significant difference ≤ 57 pg/ml, $P < 0.05$).

possible mechanism underlying changes in junctional rhythm frequently observed in astronauts during the US Skylab missions (17). There were no junctional rhythms in any of our eight subjects during adrenergic agonist infusion tests conducted before HDT. However, after 14 days of HDT, Iso induced junctional rhythms in three subjects, with premature ventricular contractions (PVCs) occurring in one of those subjects and premature atrial contractions (PACs) occurring in another. PE infusion was associated with junctional rhythm for 2 minutes with one interpolated beat in one subject, junctional rhythm and PVCs in one subject, bradycardia of 35 beats/min with erratic rhythm with escape beats in one subject, and PACs in one subject. These observations indicate that increased responsiveness of cardiac adrenoceptors resulting from exposure and adaptation to microgravity may represent a mechanism for increased risk of cardiac arrhythmias during and after a space mission.

Elevated baseline vasoconstriction and peripheral resistance were evident in our subjects by increased leg vascular resistance (Figs. 1, bottom, and 2, top) and were well-documented responses during orthostatic challenges after adaptation to actual or simulated microgravity (2, 7, 15). It is unclear whether the increased vascular responses to Iso and maintained vascular responses to PE after HDT were affected by the elevated baseline peripheral vasoconstriction. Consequently, we could not verify with our techniques whether the peripheral responses to adrenergic agonists reflected actual alterations in β_1 - and α_1 -adrenoceptors. Physiologically, our finding of increased vascular β_2 -responses following HDT that were greater than α_1 -responses in the presence of lower circulating NE is similar to the relationship between β_2 - and α_1 -vascular adrenoceptors observed in patients with Bradbury-Eggleston syndrome who also have dramatically depressed plasma NE (24). β_2 -Adrenergic hypersensitivity in the absence of a change in α -vascular

adrenoceptor response to adrenergic agonists may present a significant consequence to blood pressure regulation following return from spaceflight. It is clear that the capacity for peripheral vasoconstriction is an important determinant of orthostatic performance following spaceflight because astronauts who successfully finished 10 min of standing after 9- to 14-day missions had significantly higher total peripheral resistance than astronauts who could not complete the stand challenge (2). Because vascular β_2 -adrenoceptors elicit vasodilation compared with vascular constriction mediated by α_1 -adrenoceptors, the overall effect of greater β_2 -responsiveness in the absence of changes in α_1 -responses could produce a lesser vasoconstrictive effect, especially under a condition of increased sympathetic discharge during standing after return to the upright posture (30). This hypothesis is supported by the observation that normal reductions in blood flow to inactive muscle and visceral tissues during exercise did not occur in rats exposed to HDT (22). The potential to limit orthostatically induced elevations in peripheral resistance could compromise the capacity of the cardiovascular system to maintain adequate arterial blood pressure and cerebral perfusion during postspaceflight standing.

The mechanism of increased β -Adrenoceptor responsiveness observed after HDT in the present study is unclear but may be associated with lowered sympathetic discharge (23). This hypothesis is supported in the present study by the dramatic reduction in total circulating NE at day 14 of HDT in our subjects. There is evidence from other investigations that NE is reduced during exposure to actual (18) and ground-based simulation of (6, 16, 21, 23) microgravity. β -Adrenoceptor activity is reduced in individuals who have elevated plasma NE as a result of regular exposure to upright posture (13) and physical exercise (4), but is accentuated in subjects exposed to 10 days of HDT (21) and patients with dysautonomias in which circulating catecholamines are absent or reduced (23, 25). It is therefore possible that reduction of orthostatic and physical work stresses in a microgravity environment could be responsible for chronically reduced secretion of NE during spaceflight, leading to increased responsiveness of β -adrenoceptors and greater tachycardia and vasodilation to sympathomimetic stimulation.

Elevated circulating thyroid hormone has caused increased numbers of cardiac β -adrenoceptors in rats (31), and hyperthyroid state is associated with low levels of plasma catecholamines in humans (20). Plasma thyroxine concentration was elevated from 7.0 ± 0.3 $\mu\text{g}/100$ ml preflight to 8.7 ± 0.5 $\mu\text{g}/100$ ml postflight in the nine crewmembers who participated in NASA's three Skylab space missions (19), and serum triiodothyronine was elevated throughout 54 days of bedrest in human subjects (29). Although circulating thyroid hormone that caused increased numbers of cardiac β -adrenoceptors in an animal model was much greater (threefold) than normal baseline levels (31) compared with 18–24% elevation observed in humans during actual or simulated exposures to microgravity, the

possibility of thyroid hormone as a contributing mechanism to increased cardiac β -adrenoreceptor responsiveness observed following HDT should be considered for investigation.

We chose to use steady-state rather than bolus infusion of the agonists to measure important vascular resistance data that could not be obtained by a distribution-phase method. The use of steady-state infusion could influence our interpretations if HDT altered *Iso* or *PE* clearance by the liver. We are unaware of any published data that indicate that liver function is altered by HDT or spaceflight. On the other hand, if liver metabolism of the agonists is similar to that of the endogenous catecholamines, then the interpretation of the effects of HDT on adrenoreceptor function should be appropriate.

Another limitation to the use of systemic steady-state infusion in human subjects is the presence of compensatory baroreflex responses to adrenergic stimulation. This is complicated by alterations in functions of arterial and cardiopulmonary baroreflex control of heart rate and peripheral vascular resistance induced by exposure to microgravity or its ground-based analogs (6, 7, 9, 14). The chronotropic response to equal elevations in systemic arterial pressure is increased by aortic baroreceptor stimulation (9) and reduced by carotid baroreceptor stimulation (6, 7) in subjects who have been exposed to HDT. During *Iso* infusion in the present experiment, arterial pressures were not altered, suggesting that it is unlikely that the increased heart rate (β_1) responsiveness was influenced by altered arterial baroreflex function. It could be argued that the elevated adrenoreceptor responsiveness following HDT observed in the present study may be underestimated because circulating NE was significantly attenuated under all baseline and infusion conditions.

Peripheral vascular adrenergic responsiveness in a hypovolemic state is heavily influenced by the cardiopulmonary baroreflex control of vascular resistance. HDT decreased plasma volume and increased the vasoconstrictive response to changes in central venous pressure (7). Although not measured during the agonist infusions in the present study, *Iso* infusions at doses greater than those used in our investigation have failed to alter central venous pressure (26, 27). It is therefore unlikely that changes in cardiac and vascular resistance responses to carotid, aortic, or cardiopulmonary baroreceptor control could explain alterations in β_1 - and β_2 -responsiveness observed after HDT in the present study.

PE is known to increase both arterial and central venous pressure (9). A hypertensive stimulus to arterial and cardiopulmonary baroreceptors would be expected to reflexly induce vasodilation by sympathetic withdrawal because peripheral vascular resistance is increased following exposure to HDT during hypotensive stimulation (7). This could lead to an underestimation of the α_1 -response observed in the present study. In addition, the arterial hypertension induced by *PE* would be expected to elicit carotid and aortic baroreceptor-mediated vasodilation. The potential impact of these

arterial baroreflexes on the peripheral α_1 -response is unclear because the effect of microgravity on arterial baroreceptor control of vascular resistance is unknown.

The reduction in plasma volume may also complicate the interpretation of dose-response relationships after HDT because the concentration of the adrenergic agonists might be expected to increase by 16% if the doses of drugs were not adjusted. Because the dose-response relationships are linear, it could be suggested that 16% of the β -receptor responses was accounted for by a dilution effect. Although we did not measure the concentration of *Iso* or *PE* in the plasma, we attempted to adjust for vascular volume reduction by calculating the dose of each drug by body weight, which decreased at the end of HDT in proportion to the decrease in plasma volume (8). In any event, the increase in β_1 - and β_2 -adrenoreceptor responses by 47 and 208%, respectively, indicated that these changes occurred despite the possibility of a slightly higher concentration of agonists due to hypovolemia following HDT.

Perspectives

The heightened β -adrenergic response following HDT in the present study may provide partial explanations for a number of physiological consequences, including orthostatic hypotension and instability, that have been commonly exhibited in individuals who have been exposed to microgravity environments. Numerous factors have been identified as potential contributing mechanisms, including hypovolemia and reduced capacity of baroreflexes to buffer against changes in blood pressure. Although hypovolemia contributes to post-spaceflight tachycardia and hypotension, it is an insufficient explanation because attempts at fluid replacement have not eliminated orthostatically induced cardioacceleration (3, 28). A sharp elevation in the amounts of NE discharged on standing after exposure to microgravity (30) could produce a more tachycardic and less vasoconstrictive character in the presence of hyperresponsive adrenoreceptors. Enhanced elevation in heart rate in a setting of hypovolemia and lower vasoconstriction is a well-known stimulus for the Bezold-Jarisch reflex and vasovagal syncope. All of the subjects in the present study experienced earlier presyncope symptoms and orthostatic hypotension after HDT (11, 12). Therefore, in addition to hypovolemia and impaired baroreflex function, our data support the previously proposed hypothesis (23) that increased responsiveness of β -adrenoreceptors can be induced by prolonged exposure to microgravity and could contribute to postspaceflight orthostatic intolerance.

The authors thank Drs. Jean Varmikos and Charles Wade for their assistance with this project; Deo O'Hara and her staff at the National Aeronautics and Space Administration-Ames Human Research Facility; Don Doerr, Art Maples, and Sandy Reed for engineering support; Dick Triandifile and Barry Black for their technical assistance with data collection; and the subjects for their cheerful cooperation during the prolonged head-down tilt periods.

This research described herein was sponsored by the National Aeronautics and Space Administration administered under Contracts NAS10-10285, NAG 2-408, and W-181568. K. A. Engelke was supported by a Florida Space Grant Research Consortium Fellowship.

Opinions, interpretations, conclusions, and recommendations are those of the authors and are not necessarily endorsed by the US Air Force.

Address for reprint requests: V. A. Convertino, Physiology Research Branch, Clinical Sciences Division, AL/AOCY, 2507 Kennedy Circle, Brooks AFB, TX 78232.

Received 7 November 1996; accepted in final form 13 February 1997.

REFERENCES

- Blomqvist, C. G., and H. L. Stone. Cardiovascular adjustments to gravitational stress. In: *Handbook of Physiology. The Cardiovascular System. Peripheral Circulation and Organ Blood Flow*. Bethesda, MD: Am. Physiol. Soc., 1983, sect. 2, vol. III, pt. 2, chapt. 28, p. 1025-1068.
- Buckley, J. C., Jr., L. D. Lane, B. D. Levine, D. E. Watenough, S. J. Wright, W. E. Moore, F. A. Gaffney, and C. G. Blomqvist. Orthostatic intolerance after spaceflight. *J. Appl. Physiol.* 61: 7-18, 1986.
- Bunge, M. W., J. B. Charles, and P. C. Johnson. Cardiovascular deconditioning during space flight and the use of saline as a countermeasure to orthostatic intolerance. *Aviat. Space Environ. Med.* 58: 985-990, 1986.
- Butler, J., M. O'Brien, K. O'Malley, and J. G. Kelly. Relationship of β -adrenoreceptor density to fitness in athletes. *Nature* 298: 60-62, 1982.
- Convertino, V. A. Exercise and adaptation to microgravity environments. In: *Handbook of Physiology. Environmental Physiology*. Bethesda, MD: Am. Physiol. Soc., 1996, sect. 4, vol. II, chapt. 28, p. 815-843.
- Convertino, V. A., D. F. Doerr, D. L. Eklberg, J. M. Fritsch, and J. Vernikos-Danellis. Head-down bedrest impairs vagal baroreflex responses and provokes orthostatic hypotension. *J. Appl. Physiol.* 68: 1458-1464, 1990.
- Convertino, V. A., D. F. Doerr, D. A. Ludwig, and J. Vernikos. Effect of simulated microgravity on cardiopulmonary baroreflex control of forearm vascular resistance. *Am. J. Physiol.* 266 (Regulatory Integrative Comp. Physiol. 35): R1962-R1969, 1994.
- Convertino, V. A., K. A. Engelke, D. A. Ludwig, and D. F. Doerr. Restoration of plasma volume after 16 days of head-down tilt induced by a single bout of maximal exercise. *Am. J. Physiol.* 270 (Regulatory Integrative Comp. Physiol. 39): R3-R10, 1996.
- Crandall, C. G., K. A. Engelke, V. A. Convertino, and P. B. Raven. Aortic baroreflex control of heart rate following 16 days of simulated microgravity exposure. *J. Appl. Physiol.* 77: 2134-2139, 1994.
- Crandall, C. G., K. A. Engelke, J. A. Pawelczyk, P. B. Raven, and V. A. Convertino. Power spectral and time based analysis of heart rate variability following 16 days simulated microgravity exposure in humans. *Aviat. Space Environ. Med.* 65: 1105-1109, 1994.
- Engelke, K. A., D. F. Doerr, and V. A. Convertino. Effect of a single bout of exhaustive exercise on integrated baroreflex function after 16 days of head-down tilt. *Am. J. Physiol.* 269 (Regulatory Integrative Comp. Physiol. 38): R614-R620, 1995.
- Engelke, K. A., D. F. Doerr, C. G. Crandall, and V. A. Convertino. Application of acute maximal exercise to protect orthostatic tolerance following 16 days of simulated microgravity. *Am. J. Physiol.* 271 (Regulatory Integrative Comp. Physiol. 40): R837-R847, 1996.
- Feldman, R. D., L. E. Limbird, J. Nadeau, G. A. Fitzgerald, D. Robertson, and A. J. J. Wood. Dynamic regulation of leukocyte β adrenergic receptor agonist interactions by physiological changes in circulating catecholamines. *J. Clin. Invest.* 72: 164-170, 1983.
- Fritsch, J. M., J. B. Charles, B. S. Bennett, M. M. Jones, and D. L. Eklberg. Short-duration spaceflight impairs human carotid baroreceptor-cardiac reflex responses. *J. Appl. Physiol.* 73: 684-671, 1992.
- Gabrielsson, A., P. Norak, R. Vidarbak, and O. Hendriksen. Effect of microgravity on forearm subcutaneous vascular resistance in humans. *J. Appl. Physiol.* 79: 434-438, 1995.
- Goldstein, D. S., J. Vernikos, C. Holmes, and V. A. Convertino. Catecholaminergic effects of prolonged head-down bed rest. *J. Appl. Physiol.* 78: 1023-1029, 1995.
- Hoffler, G. W., R. L. Johnson, A. E. Nicogossian, S. A. Bergman, and M. M. Jackson. Vectorcardiographic results from Skylab Medical Experiment M092: lower body negative pressure. In: *Biomedical Results of Skylab*, edited by R. S. Johnston and L. F. Dietlein. Washington, DC: US Government Printing Office, 1977, p. 313-323 (NASA SP-377).
- Leach, C. S., S. I. Altchuler, and N. M. Cistrou-Trevino. The endocrine and metabolic response to space flight. *Med. Sci. Sports Exerc.* 15: 432-440, 1983.
- Leach, C. S., and P. C. Rambaut. Biomedical responses of the Skylab crewmen: an overview. In: *Biomedical Results of Skylab*, edited by R. S. Johnston and L. F. Dietlein. Washington, DC: US Government Printing Office, 1977, p. 204-216 (NASA SP-377).
- Leroy, G. S., and I. Klein. Catecholamine-thyroid hormone interactions and the cardiovascular manifestations of hyperthyroidism. *Am. J. Med.* 88: 642-646, 1990.
- Mason, H. J., Trueman, and F. Baloch. Response of adrenergic receptors to 10 days head-down tilt bedrest. *Acta Physiol. Scand.* 144, Suppl. 604: 61-65, 1992.
- McDonald, K. S., M. D. Delp, and R. H. Fitts. Effect of hindlimb unweighting on tissue blood flow in the rat. *J. Appl. Physiol.* 72: 2210-2218, 1992.
- Robertson, D., V. A. Convertino, and J. Vernikos. The sympathetic nervous system and the physiologic consequences of spaceflight: a hypothesis. *Am. J. Med. Sci.* 306: 126-132, 1994.
- Robertson, D., A. S. Holmster, E. L. Carey, C.-S. Tung, M. R. Goldberg, and R. M. Robertson. Increased vascular β adrenergic responsiveness in autonomic dysfunction. *J. Am. Coll. Cardiol.* 3: 850-856, 1984.
- Robertson, D., S. E. Perry, A. S. Holmster, R. M. Robertson, and I. Baggio. Dopamine- β -hydroxylase deficiency: a genetic disorder of cardiovascular regulation. *Hypertension* 18: 1-8, 1991.
- Roth, C. F., A. D. Flanagan, and R. Mason-McCormick. Role of β -adrenergic agonists in the control of vascular capacitance. *Can. J. Physiol. Pharmacol.* 68: 575-585, 1990.
- Tabrizchi, R., S. L. Lam, and C. C. Y. Pang. Possible equilibration of portal venous and central venous pressures during circulatory arrest. *Am. J. Physiol.* 264 (Heart Circ. Physiol. 33): H269-H281, 1993.
- Vernikos, J., and V. A. Convertino. Advantages and disadvantages of hydrocortisone or saline load in preventing post-spaceflight orthostatic hypotension. *Acta Astronautica* 33: 259-266, 1994.
- Vernikos-Danellis, J., C. S. Leach, C. M. Winget, P. C. Rambaut, and P. B. Raven. Thyroid and adrenal cortical rhythmicity during bed rest. *J. Appl. Physiol.* 33: 644-648, 1972.
- Whitson, P. A., J. B. Charles, W. J. Williams, and N. M. Cistrou. Changes in sympathoadrenal response to standing in humans after spaceflight. *J. Appl. Physiol.* 79: 428-433, 1995.
- Williams, L. T., R. J. Leftowits, A. M. Watanabe, D. R. Hathaway, and H. R. Sack, Jr. Thyroid hormone regulation of β -adrenergic receptor number. *J. Biol. Chem.* 262: 2787-2799, 1977.