

**NH<sub>3</sub>, H<sub>2</sub>S, AND THE RADIO BRIGHTNESS TEMPERATURE SPECTRA  
OF THE GIANT PLANETS**

THOMAS R. SPILKER

*Jet Propulsion Laboratory / California Institute of Technology  
Pasadena, CA, USA 91109*

Recent radio interferometer observations of Neptune enable comparisons of the radio brightness temperature ( $T_B$ ) spectra of all four giant planets. This comparison reveals evidence for fundamental differences in the compositions of Uranus' and Neptune's upper tropospheres, particularly in their ammonia (NH<sub>3</sub>) and hydrogen sulfide (H<sub>2</sub>S) mixing ratios, despite those planets' outward similarities. The tropospheric abundances of these constituents yield information about their deep abundances, and ultimately about the formation of the planets from the presolar nebula (Atreya *et al.*, 1995).

Figures 1, 2, 3, and 4 show the  $T_B$  spectra of Jupiter, Saturn, Uranus, and Neptune, respectively, from 0.1 to tens of cm wavelength. The data shown are collected from many observers. Data for Jupiter, Saturn, and Uranus are those cataloged by de Pater and Massie (1985), plus the Saturn Very Large Array (VLA) data by Grossman *et al.* (1989). Figure 3, Uranus, shows only data acquired since 1973. Before 1973 Uranus'  $T_B$  increased steadily as its pole moved into view, causing significant scatter in those data. Neptune data at  $>1$  cm, all taken at the VLA, are collected from de Pater and Richmond (1989), de Pater *et al.* (1991), and Hofstadter (1993). For a variety of reasons, such as susceptibility to source confusion, single-dish data at those wavelengths are much noisier than the more reliable VLA data and have been ignored. Single-dish data by Griffin and Orton (1993) shortward of 0.4 cm are shown, along with the Owens Valley Radio Observatory (interferometer) datum at 0.266 cm by Muhleman and Berge (1991).

Spectra of Jupiter, Saturn, and Neptune share certain gross characteristics. In each spectrum,  $T_B$  at 1.3 cm is ~120-140 K, less than ~30 K different from that at 0.1 cm. All three spectra show a break in slope at or near 1.3 cm, with  $T_B$  increasing fairly rapidly with wavelength longward of 1.3 cm. Visible and IR spectroscopy show that NH<sub>3</sub>, whose strong inversion spectrum peaks at ~1.3 cm, is an important tropospheric species at Jupiter and Saturn. Its signature on the Jovian radio spectrum is obvious, causing the prominent "hole" at 1.3 cm. At Saturn it is more subdued but is the source of that spectrum's change in slope at 1.3 cm. Radiative transfer models of Jupiter and Saturn with near-solar deep NH<sub>3</sub> abundances agree well with the data (*e.g.*, de Pater, 1990).

Uranus'  $T_B$  spectrum does not fit this pattern.  $T_B$  is ~175 K at 1.3 cm, ~80 K warmer than at 0.1 cm and much warmer than the other three planets. The data indicate a nearly linear increase in  $T_B$  with  $\log(\lambda)$  over the entire range shown in figure 3, requiring no break in slope near 1.3 cm. Notably, there is a distinct difference between Uranus' and Neptune's spectra: at ~20 cm, and also at 0.1-0.4 cm, Uranus'  $T_B$  are quite similar to Neptune's, but in the intervening 1-10 cm range Uranus averages 30-55 K warmer than Neptune. Gulkis *et al.* (1978) first showed that Uranus radiative transfer models with near-solar NH<sub>3</sub> deep abundances predict  $T_B$  at cm wavelengths that are much too cold. Their model using an NH<sub>3</sub> abundance about 1% of solar fit the data better, though far




Figure 1. Jupiter's radio brightness temperature spectrum




Figure 2. Saturn's radio brightness temperature spectrum

from perfectly. They offered one possible cause for the apparent  $\text{NH}_3$  depletion: a superabundance of  $\text{H}_2\text{S}$  could react out most of the  $\text{NH}_3$ , forming  $\text{NH}_4\text{SH}$  as first discussed by Wildt (1937). At the time most researchers assumed that essentially all nitrogen in giant planet tropospheres is in the highly reduced form. In that case the solar




Figure 3. Uranus' radio brightness temperature spectrum




Figure 4. Neptune's radio brightness temperature spectrum

abundance of sulfur made  $\text{H}_2\text{S}$  seem the most likely candidate to deplete  $\text{NH}_3$ , despite the lack of direct observational evidence of  $\text{H}_2\text{S}$  at Uranus or any of the other giant planets. This nondetection is expected even if sulfur abundances are approximately solar, since models predict the  $\text{NH}_4\text{SH}$ -forming reaction would restrict  $\text{H}_2\text{S}$  to depths inaccessible to visible or IR spectroscopy (Atreya and Romani, 1985).




Figure 5. Neptune's radio brightness temperature spectrum, with predictions of various models superposed. See text for model descriptions.

Some researchers suggest a large  $\text{H}_2\text{S}$  superabundance at Neptune also. Problems fitting radiative transfer models to cm data prompted de Pater *et al.* (1991) to invoke  $\text{NH}_3$ -depleting  $\text{H}_2\text{S}$  at Neptune, and to suggest that  $\text{H}_2\text{S}$  might contribute significantly to the total opacity. Recently DeBoer and Steffes (1994) (hereafter "DBS") made laboratory measurements of cm  $\text{H}_2\text{S}$  opacities and found them a factor of two larger than Van Vleck-Weisskopf predictions. Based on this they suggest  $\text{H}_2\text{S}$  may be the major source of cm opacity in Neptune's upper troposphere, and reinterpret Lindal's (1992) Voyager 2 radio occultation data. Lindal assumed all opacity at the 6.3 bar level, the deepest probed, was due to  $\text{NH}_3$  and derived a volume mixing ratio of  $5 \times 10^{-7}$ . DBS assume all opacity there is due to  $\text{H}_2\text{S}$  and derive a mixing ratio of  $1.7 \times 10^{-4}$ . Based on that result, they use radiative transfer models with  $\text{H}_2\text{S}$  above the  $\text{NH}_4\text{SH}$  cloud to generate predicted spectra they compare to the Neptune  $T_B$  data (both single-dish and VLA). Agreement between the predicted spectra and the superior VLA data is rather poor.

Figure 5 duplicates Figure 4 except it includes results from various radiative transfer models. Models 1-4 are after DBS, with 30 times solar  $\text{H}_2\text{O}$  and  $\text{CH}_4$ ;  $\text{NH}_3$  and  $\text{H}_2\text{S}$  abundances, respectively, are 0.5 solar and 15 times solar for model 1, solar and 18 times solar for model 2, twice solar and 25 times solar for model 3, and solar and 6 times solar for model 4. Models 1-3, whose spectral results are identical shortward of 6 cm, yield  $1.7 \times 10^{-4}$   $\text{H}_2\text{S}$  above the  $\text{NH}_4\text{SH}$  cloud, while model 4 yields Lindal's  $5 \times 10^{-7}$   $\text{NH}_3$ . Model 5 is after de Pater and Richmond (1989), using an  $\sim 2\%$  solar  $\text{NH}_3$  mixing ratio ( $3 \times 10^{-6}$ ) throughout the atmosphere, limited by saturation, and no  $\text{H}_2\text{S}$ . Model 6, by the author, uses approximately solar  $\text{NH}_3$  ( $2 \times 10^{-4}$ ) and no  $\text{H}_2\text{S}$  to demonstrate that Neptune models with uniformly near-solar tropospheric  $\text{NH}_3$  mixing ratios are not consistent with the observed spectrum, so some form of  $\text{NH}_3$  depletion (not necessarily  $\text{H}_2\text{S}$ ) is needed.

Only the models with  $\text{NH}_3$  above the  $\text{NH}_4\text{SH}$  cloud reproduce Neptune's  $T_B$  dip at cm wavelengths. Model 5, with more  $\text{NH}_3$  than model 4, provides the best fit; slightly

more  $\text{NH}_3$  would provide a better fit, further decreasing  $T_B$  longward of 1 cm. This does not conflict with Lindal's result, since he states  $\text{NH}_3$  is probably still saturated at the deepest datum. Due to the  $\text{H}_2\text{S}$  spectrum's simple  $f^2$  dependency longward of 0.4 cm, models dominated by  $\text{H}_2\text{S}$  above the  $\text{NH}_4\text{SH}$  cloud (DBS models 1-3) deviate  $<10$  K from a straight line on the plot, quite unlike the data. Reproducing the  $T_B$  dip with such an absorber requires a relatively thin tropospheric layer near the 120-130 K level ( $\sim 5$  bar pressure level) with a much larger absorber mixing ratio than adjacent layers, a situation more appropriate to the upper stratosphere than the troposphere. Without a mechanism to maintain such a tropospheric layer, it is unlikely that the observed cm opacity in Neptune's upper troposphere is primarily due to  $\text{H}_2\text{S}$ . Neptune's radio spectrum appears to require  $\text{NH}_3$ , or another species with an opacity peak near 1-2 cm, in the upper troposphere.

Applying the DBS models 1-3 to Uranus leads to a different conclusion for that planet. Upper tropospheric T-P (temperature-pressure) relations for Uranus and Neptune are very similar: at equal pressures, their temperatures differ by  $\sim 5\text{K}$  at most from well above their




Figure 6. Uranus' radio brightness temperature spectrum, with predictions of  $\text{H}_2\text{S}$ -dominated models of Neptune by DBS (solid lines). See previous page for model descriptions.

tropospheres to the deepest level probed by radio occultation (Lindal *et al.*, 1987; Lindal, 1992). Also, recent work by Killen and Flasar (1995) show that the cm spectra of the giant planets are largely insensitive to T-P profiles, but instead depend most strongly on the relative humidities of condensible absorbers. Given a fixed set of constituent abundance profiles, a model using Neptune's T-P profile will yield a  $T_B$  spectrum quite similar to one produced using Uranus' T-P profile. The dissimilarity of the two planets' observed radio spectra makes it highly unlikely they have similar constituent profiles. Figure 6 shows the result of using DBS'  $\text{H}_2\text{S}$ -dominated models of Neptune as first approximations to such models for Uranus. The models fit Uranus' observed spectrum much better than Neptune's, suggesting that tropospheric constituents whose cm opacities have  $f^2$  dependencies, such as  $\text{H}_2\text{S}$ , could produce Uranus' observed radio  $T_B$  spectrum.

This does not rule out the presence of  $\text{NH}_3$ , but it points to other possibilities. Thus, while Neptune seems to need a small (relative to solar) but nontrivial amount of  $\text{NH}_3$  in its upper troposphere, Uranus does not.

Recent work by Atreya *et al.* (1995) suggests that a significant fraction of Uranus' and Neptune's nitrogen could be in the form of  $\text{N}_2$  instead of  $\text{NH}_3$ . Thus a range of nitrogen and sulfur abundances, both not greatly different from solar, could yield the observed depletion of  $\text{NH}_3$ . Further radiative transfer modeling can better define the limits of such abundances, which currently are poorly constrained. Insufficient data are available at this time to state with certainty the precise mechanisms for  $\text{NH}_3$  depletions at Uranus and Neptune. Whatever the mechanisms, comparing the two planets' radio  $T_B$  spectra shows that the resulting absorber mixing ratio profiles in their upper tropospheres are quite different.

#### Acknowledgement

The research described in this paper was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration.

#### REFERENCES

- Atreya, S.K., and P.N. Romani (1985) Photochemistry and clouds of Jupiter, Saturn, and Uranus, in *Recent Advances in Planetary Meteorology*, G.E. Hunt ed., Cambridge Univ. Press, Cambridge, 17-68.
- Atreya, S.K., S.G. Edgington, D. Gautier, and T.C. Owen (1995) Origin of the Major Planet Atmospheres: Clues from Trace Species, this publication.
- DeBoer, D.R., and P.G. Steffes (1994) Laboratory Measurements of the Microwave Properties of  $\text{H}_2\text{S}$  under Simulated Jovian Conditions with an application to Neptune, *Icarus* **109**, 352-366.
- de Pater, I., and S.T. Massie (1985) Models of the Millimeter-Centimeter Spectra of the Giant Planets, *Icarus* **62**, 143-171.
- de Pater, I., and M. Richmond (1989) Neptune's Microwave Spectrum from 1 mm to 20 cm, *Icarus* **80**, 1-13.
- de Pater, I. (1990) Models of the Millimeter-Centimeter Spectra of the Giant Planets, *Ann.Rev.Astr. Astrophys* **28**, 347-399.
- de Pater, I., P.N. Romani, and S.K. Atreya (1991) Possible Microwave Absorption by  $\text{H}_2\text{S}$  Gas in Uranus' and Neptune's Atmospheres, *Icarus* **91**, 220-233.
- Griffin, M.J., and G.S. Orton (1993) The Near-Millimeter Brightness Temperature Spectra of Uranus and Neptune, *Icarus* **105**, 537-547.
- Grossman, A.W., D.O. Muhleman, and G.L. Berge (1989) High-Resolution Microwave Images of Saturn, *Science* **245**, 1211-1215.
- Gulkis, S., M.A. Janssen, and E.T. Olsen (1978) Evidence for the Depletion of Ammonia in the Uranus Atmosphere, *Icarus* **34**, 10-19.
- Hofstadter, M.D. (1993) Microwave Imaging of Neptune's Troposphere, *Bull.Am.Astr.Soc.* **25** No. 3, 1077.
- Killen, R.M., and F.M. Flasar (1995) Microwave Sounding of the Giant Planets, *Icarus*, in press.
- Lindal, G.F., J.R. Lyons, D.N. Sweetnam, V.R. Eshleman, D.P. Hinson, and G.L. Tyler (1987) The Atmosphere of Uranus: Results of Radio Occultation Measurements With Voyager 2, *JGR* **92** No. A13, 14987-15001.
- Lindal, G.F. (1992) The Atmosphere of Neptune: An Analysis of Radio Occultation Data Acquired With Voyager 2, *Astron.J* **103** No. 3, 967-982.
- Muhleman, D.O., and G.L. Berge (1991) Observations of Mars, Uranus, Neptune, Io, Europa, Ganymede, and Callisto at a Wavelength of 2.66 mm, *Icarus* **92**, 263-272.