

NASA/TM—1998-207915



# Effect of Gravity on Surface Tension

M.M. Weislogel  
Lewis Research Center, Cleveland, Ohio

M.O.J. Azzam and J.A. Mann  
Case Western Reserve University, Cleveland, Ohio

National Aeronautics and  
Space Administration

Lewis Research Center

---

August 1998

Available from

NASA Center for Aerospace Information  
7121 Standard Drive  
Hanover, MD 21076  
Price Code: A02

National Technical Information Service  
5287 Port Royal Road  
Springfield, VA 22100  
Price Code: A02

# EFFECT OF GRAVITY ON SURFACE TENSION

M.M. Weislogel  
National Aeronautics and Space Administration  
Lewis Research Center  
Cleveland, Ohio 44135

M.O.J. Azzam and J.A. Mann  
Department of Chemical Engineering  
Case Western Reserve University  
Cleveland, Ohio 44106

The surface tension of a liquid-fluid pair has been viewed as a thermodynamic property of the system principally dependent on temperature and, to a lesser extent, pressure (refs. 1 and 2). Recent theoretical work (refs. 3 and 4) has introduced the likelihood of an additional body force dependency, in our case gravity,  $g$ , which, though perhaps small, may manifest itself at a measurable level. The principal provision enhancing such a measurement would be that the gravity vector be aligned perpendicular to the interface. This is because any variance of surface tension with gravity would be attributable to compressibility effects in the interfacial region, owing directly to an increase/decrease in interfacial thickness; the latter being a dimension of the interface which increases dramatically near the liquid-vapor critical point.

The following is a description of a simple experiment using laser light scattering heterodyne spectroscopy to determine the effects of orientation with respect to earth gravity ( $g_0$ ) on the surface tension of a liquid-vapor interface. The experimental technique is well established (refs. 5 to 9) and measures the frequency and damping of thermal fluctuations at the surface (thermal ripples) from which the surface tension is calculated through the dispersion equation.<sup>1</sup> A schematic of the optical path is shown in figure 1. In addition to the potential for increased accuracy, this approach eliminates any gravity dependence on the measurement technique itself, a unique quality when contrasted to other techniques where gravity is essential to the measurement (refs. 11 and 12).

To provide an ability to vary the gravitational field in a terrestrial laboratory, a capillary tube approach was taken in which the tube size was selected for the test fluid which yielded a meniscus that could be inverted (gravity vector opposite in direction to the density gradient), flattened, and stabilized without the heavier fluid falling out. The values as measured in the inverted configuration ( $-g_0$ ) were then compared to measured values of the same fluid in the normal configuration ( $+1g_0$ , gravity vector in direction of density gradient).

Doubly distilled water (in air) was chosen as the test fluid due to its high surface tension. This permitted the use of a relatively large diameter capillary tube (up to 12.7 mm) and also reduced the ratio of incident beam diameter to characteristic surface length ( $d/l$ ). The tube diameter selected for this experiment was 9.52 mm.

The tube was made of aluminum and the critical surfaces were machined on an airbearing lathe to a mirror finish. The tube was then fitted to a semi-flexible teflon tube which passed through a volume control fitting and connected the aluminum tube to the fluid reservoir (fig. 2).

The teflon tubing and fittings were cleaned ultrasonically in a mild soap solution and were then rinsed with acetone and/or ethanol and allowed to air dry. The capillary tube was treated separately with an optical cleaning solution to avoid marring the critical surfaces. The assembled components were then flushed with large quantities of distilled water. To monitor possible contamination of the test liquid, Du Noüy ring surface tension measurements were made regularly of the water in the reservoir and the water as drained from the capillary during the flush procedure. In every case these values were  $72.5 \pm 1.1$  dyne/cm. Surface tension values measured of the water left stagnant in the lines for long periods were also consistent at 72.5 dyne/cm.

After extensive flushing the capillary tube was aligned vertically in position for light scattering, either inverted or in the "normal" configuration. Once stabilized, the surface was "flattened" by adjusting the volume of the liquid in the capillary tube. The flatness of the surface was gauged by observing the reflected/scattered light in the far field until it became collimated with a circular symmetric shape. This flat surface condition

---

<sup>1</sup>When capillary waves move on a surface both the surface tension and the gravity field provide a restoring force. The latter force enters the theory of the spectroscopic approach through a pressure term of the form  $\Delta \rho g \zeta$ , where  $\Delta \rho$  is the density difference across the interface and  $\zeta$  is the surface elevation. Thus  $g$  appears as a parameter in the dispersion equation ref. (10) under which the assumptions of continuum, linear behavior hold.

could not be maintained for longer than 10 sec due to gradual, though slight, changes in the fluid volume (due to possible thermal expansion, evaporation, and/or pressure leaks) resulting in slight changes in surface curvature. However, for such spectroscopic measurements, only 1 to 5 sec are necessary for the collection of quality data. The data sampling time used for these measurements was 3 sec.

Table I presents the results of the measurements of the surface tension under gravity fields  $\pm 1g_0$ . The difference is small (0.34 dyne/cm), and when compared to the uncertainties in the measurements it is reasonable to conclude that the difference between the two cases is within the experimental uncertainty. It should be emphasized that both measurements at  $\pm 1g_0$  yielded surface tensions which are higher than the commonly accepted values found in the literature. This does not affect the fact that the measurement at  $-1g_0$  was slightly higher than the one made at  $+1g_0$  since everything in both experiments remained the same except for the orientation. Further tests could be conducted along this line on fluids nearer to critical point conditions where the effect of gravity on the surface tension would be amplified. Such information could prove valuable for future space experimentation where  $g$  approaches zero.

#### REFERENCES

1. L.A. Turkevich and J.A. Mann, *Langmuir*, **6**, p. 445 (1990).
2. J.S. Rowlinson, B. Widom, *International Series of Monographs on Chemistry*, **8**, Oxford Science Publications (1989).
3. R.E. Salvino, *Physical Review A*, **41**, p. 4236 (1990).
4. U. Marini Bettolo Marconi and B.L. Gyorffy, *Phys. Rev. A*, **41**, p. 6732 (1990).
5. S. Hard, Y. Hamnerius and O. Nilson, *J. Appl. Phys.*, **47**, p. 2433 (1976).
6. J.A. Mann and R.V. Edwards, *Rev. Sci. Instrum.*, **55**(5), p. 727 (1984).
7. R.V. Edwards, R.S. Sirohi, J.A. Mann, L.B. Shih and L. Lading, *Applied Optics*, **21**(19), 3555 (1982).
8. D. Langevin and J. Mennier, *Photon Correlation Spectroscopy and Velocimetry*, (Plenum Press, New York, 1976).
9. M.O.J. Azzam, Ph.D. dissertation (Case Western Reserve University, 1992).
10. J.A. Mann, *Surface and Colloid Science*, **177**, eq. 114 (Plenum Press, New York, 1984).
11. A.W. Adamson, *Physical Chemistry of Surfaces*, (Interscience Publishers Inc., 1960).
12. C.A. Miller and P. Neogi, *Interfacial Phenomena* (Marcel Dekker Inc., New York, 1985).

TABLE I.—MEASURED SURFACE TENSION  
(USING LIGHT SCATTERING TECHNIQUE)  
OF WATER/AIR SYSTEM AT 22C, UNDER  
A GRAVITY FIELD OF  $\pm 1g_0$ . (ALL  
MEASUREMENTS PERFORMED  
AT ROOM TEMPERATURE  $23.5 \pm .1$  C).

Surface tension, dyne/cm		
Test condition	Measured	Literature
+1 $g_0$	73.12 $\pm$ 0.41	72.44
-1 $g_0$	73.46 $\pm$ 0.23	72.44




Figure 1.—The optical path from the laser to the photodetector.




Figure 2.—Experimental setup for measuring the surface tension of water at gravity fields of  $+1g_0$  and  $-1g_0$  in a capillary tube. ( $-1g_0$  configuration shown).

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1998	3. REPORT TYPE AND DATES COVERED Technical Memorandum	
4. TITLE AND SUBTITLE Effect of Gravity on Surface Tension			5. FUNDING NUMBERS WU-962-24-00-00	
6. AUTHOR(S) M.M. Weislogel, M.O.J. Azzam, and J.A. Mann				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Lewis Research Center Cleveland, Ohio 44135-3191			8. PERFORMING ORGANIZATION REPORT NUMBER E-11189	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA TM-1998-207915	
11. SUPPLEMENTARY NOTES M.M. Weislogel, NASA Lewis Research Center (presently with TDA Research, Inc., 12345 West 52nd Avenue, Wheat Ridge, Colorado 80033); M.O.J. Azzam and J.A. Mann, Case Western Reserve University, Department of Chemical Engineering, Cleveland, Ohio 44106. Responsible person, M.M. Weislogel, organization code 6712, (303) 940-2320.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified - Unlimited Subject Categories: 23, 34, and 35 This publication is available from the NASA Center for AeroSpace Information, (301) 621-0390.			12b. DISTRIBUTION CODE Distribution: Nonstandard	
13. ABSTRACT (Maximum 200 words) Spectroscopic measurements of liquid-vapor interfaces are made in $\pm 1g_0$ environments to note the effect of gravity on surface tension. A slight increase is detected at $-1g_0$ , but is arguably within the uncertainty of the measurement technique. An increased dependence of surface tension on the orientation and magnitude of the gravitational vector is anticipated as the critical point is approached.				
14. SUBJECT TERMS Surface tension; Ripplon, Spectroscopy; Interfacial tension; Gravity			15. NUMBER OF PAGES 9	
			16. PRICE CODE A02	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT	