

FINAL REPORT

Grant Number: NAG5-2477

Project Title: "The Structure of the Broad-Line Region in Well-Studied AGNs" *P4*

Investigators: Bradley M. Peterson, Ohio State University (PI)
Gary J. Ferland, University of Kentucky

Award Period: 1994 February 1 - 1997 July 31

Total Award Amount: \$ [REDACTED]

Institution: The Ohio State University Research Foundation
1960 Kenny Road
Columbus, OH 43210

ABSTRACT

Large amounts of high quality UV and optical data have been obtained in massive multi-wavelength monitoring campaigns on a small number of active galactic nuclei, and these data are changing our understanding of the central engines in these sources in a fundamental way. Preliminary analyses have shown that more comprehensive approaches will be necessary to make full use of these data. We propose to undertake a complete set of photoionization equilibrium calculations with a state-of-the-art computer code in order to determine the radial structure of the broad-line region in a way that is consistent with the emission-line fluxes, profiles, and transfer functions.

Project Description

I. Introduction

The broad emission lines in the ultraviolet and optical spectra of AGNs provide an important probe of both the mass and radiated spectrum of the central engines in these luminous sources. Through analysis of the broad-line spectrum, it is possible to determine something about the nature of the otherwise unobservable spectrum shortward of the Lyman limit. Furthermore, the dynamics of the gas are almost certainly determined by some combination of radiation pressure and gravity. It is possible to determine the distribution and kinematics of the broad-line region (BLR) clouds by observing the response of the broad emission lines to continuum variations; the structure of the BLR can be unraveled by observing how the BLR “reverberates” in response to continuum variability (Blandford & McKee 1982). Measurement of the continuum and emission-line light curves consequently provides a tool of tremendous potential importance for probing the very central regions of AGNs. Indeed, determination of the size and kinematics of the BLR can result in unambiguous measurement of the mass of the central engine and lead to the identification of some of the most important physical processes in this region.

The structure and kinematics of the BLR can be solved for formally by inversion of the transfer equation

$$L(v, t) = \int_{-\infty}^{\infty} \Psi(v, \tau) C(t - \tau) d\tau, \quad (1)$$

where $C(t)$ is the continuum light curve, $L(v, t)$ is the emission-line flux at line-of-sight velocity v at time t , and Ψ is a geometry-dependent “transfer function” (Blandford & McKee 1982), which can be thought of as the response of the emission line to a δ -function continuum outburst as seen by external observer. The aim of reverberation mapping is to invert the transfer equation to solve for $\Psi(v, t)$ and thus infer the BLR geometry and velocity field. Obviously, successful inversion of the transfer equation requires accurate determination of $C(t)$ and $L(v, t)$. Even with modest amounts of data, it is possible to extract limited information about the transfer function by cross correlation of $C(t)$ and $L(t)$, since convolution of eq. (1) with $C(t)$ shows that the cross-correlation function is the convolution of the transfer function with the continuum autocorrelation function.

As described in our original proposal, our intent has been to carry out a more thorough analysis of existing variability data than has been undertaken to date. Our ultimate goal is to determine the nature and origin of the broad-line clouds, since this may well provide the key element in explaining the AGN phenomenon.

II. Project Summary

Virtually all of the funds requested through the ADP program were used to provide funding for a postdoctoral research associate (Dr. Ignaz Wanders) to work on analysis of the existing space-based and related ground-based data AGNs that have been monitored with *IUE*, *HST*, and other observatories.

The first major result of the project was determination of the C IV transfer function for NGC 5548 (Wanders et al. 1995). The transfer function is consistent with an anisotropic

continuum source illuminating clouds in randomly inclined circular Keplerian orbits. Recognition that the continuum source might be beamed led to a full consideration of how different orientations might lead to a variety of different one-dimensional (i.e., velocity-independent) transfer functions and line profiles (Goad & Wanders 1996).

We then followed up on this work with a detailed study of emission-line profile variability (Wanders & Peterson 1996; Peterson et al. 1998). We have found that emission-line profile variations are unrelated to reverberation effects — profile variability is due to geometrical or physical changes in the BLR on time scales of several months (the dynamical time scale). We believe that the observed profile variations can be interpreted through the anisotropic continuum model: in this model, the profile variations are due to stochastic effects, specifically to individual emission-line clouds moving in and out of the beamed continuum. The argument for this is that profile changes seem to occur only rather rarely and a given profile state seems to persist for a long time, longer than the light travel-time across the BLR. However, when profile changes occur, they can occur fairly abruptly. The strongest argument against this scenario is that stochastic effects are unlikely to be responsible for profile variability if the number of clouds is very large, 10^{7-11} in various traditional estimates. However, we have argued earlier that these usual arguments about cloud numbers are generally specious (Peterson 1994). The absence of profile structure on small velocity scales (500 km s^{-1} or less) indicates that the individual line-emitting clouds must have a supersonic component (such as a wind, perhaps), since the individual clouds must emit their radiation over a large line-of-sight velocity range, something like 10^3 km s^{-1} rather than the 10 km s^{-1} thermal width.

The work initiated under this grant is being continued and additional refereed publications describing more recent results are being prepared for submission under support for subsequent proposals.

References

- Blandford, R.D., and McKee, C.F. 1982, ApJ, 255, 419
 Goad, M., and Wanders, I. 1996, ApJ, 469, 113
 Peterson, B.M. 1994, in *Reverberation Mapping of the Broad-Line Region in Active Galactic Nuclei*, ed. P.M. Gondhalekar, K. Horne, and B.M. Peterson (San Francisco: Astronomical Society of the Pacific), p. 1.
 Peterson, B.M., Pogge, R.W., and Wanders, I. 1998, in *Structure and Kinematics of Quasar Broad Line Regions*, ed. C.M. Gaskell, W.N. Brandt, D. Dultzin-Hacyan, M. Dietrich, and M. Eracleous (San Francisco: ASP), in press.
 Wanders, I., Goad, M.R., Korista, K.T., Peterson, B.M., Horne, K., Ferland, G.J., Koratkar, A.P., Pogge, R.W., and Shields, J.C. ApJL, 453, L87 (1995).
 Wanders, I. and Peterson, B.M. 1996, ApJ, 466, 174. Erratum: ApJ, 477, 990 (1997).

PAPERS PUBLISHED BASED ON NASA SUPPORT
THROUGH GRANT NAG5-2477

1. "Emission-Line Profile Variations in the AGN NGC 5548", I. Wanders and B.M. Peterson, *Bulletin of the American Astronomical Society*, **26**, 1500 – 1501 (1994).
2. "Aperture Effects and Limitations on the Accuracy of Ground-Based Spectrophotometry of Active Galactic Nuclei", B.M. Peterson, R.W. Pogge, I. Wanders, S.M. Smith, and W. Romanishin, *Publications of the Astronomical Society of the Pacific*, **107**, 579 – 589 (1995).
3. "The Geometry and Kinematics of the Broad-Line Region in NGC 5548 from HST and IUE Observations", I. Wanders, M.R. Goad, K.T. Korista, B.M. Peterson, K. Horne, G.J. Ferland, A.P. Koratkar, R.W. Pogge, and J.C. Shields, *Astrophysical Journal (Letters)*, **453**, L87 – L90 (1995).
4. "The Effect of a Variable Anisotropic Continuum Source Upon Broad Emission-Line Profiles and Responses" M. Goad and I. Wanders, *Astrophysical Journal*, **469**, 113 – 130 (1996).
5. "A Long-Term Study of Emission-Line Profile Variability in NGC 5548", I. Wanders and B.M. Peterson, *Astrophysical Journal*, **466**, 174 – 190 (1996). Erratum: *Astrophysical Journal*, **477**, 990 (1997).
6. "Multiwavelength Observations of Short Time-Scale Variability in NGC 4151. IV. Analysis of Multiwavelength Continuum Variability", R. Edelson et al. (91 authors, including B.M. Peterson), *Astrophysical Journal*, **470**, 364 – 377 (1996).
7. "Steps Toward Determination of the Size and Structure of the Broad-Line Region in Active Galactic Nuclei. XI. Intensive Monitoring of the Ultraviolet Spectrum of NGC 7469", I. Wanders, B.M. Peterson, et al. (57 authors), *Astrophysical Journal Supplement*, **113**, 69–88 (1997).
8. "Emission-Line Profile Variability". B.M. Peterson, R.W. Pogge, and I. Wanders, in *Structure and Kinematics of Quasar Broad Line Regions*, ed. C.M. Gaskell, W.N. Brandt, D. Dultzin-Hacyan, M. Dietrich, and M. Eracleous (San Francisco: ASP), in press.