


NASA/CR-1998-208737
ICASE Report No. 98-49



Effects of Helicity on Lagrangian and Eulerian Time Correlations in Turbulence

*Robert Rubinstein
ICASE, Hampton, Virginia*

*Ye Zhou
ICASE, Hampton, Virginia
and
IBM, Yorktown Heights, New York*

*Institute for Computer Applications in Science and Engineering
NASA Langley Research Center
Hampton, VA*

Operated by Universities Space Research Association



National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23681-2199

Prepared for Langley Research Center
under Contract NAS1-97046

November 1998

Available from the following:

NASA Center for AeroSpace Information (CASI)
7121 Standard Drive
Hanover, MD 21076-1320
(301) 621-0390

National Technical Information Service (NTIS)
5285 Port Royal Road
Springfield, VA 22161-2171
(703) 487-4650

EFFECTS OF HELICITY ON LAGRANGIAN AND EULERIAN TIME CORRELATIONS IN TURBULENCE *

ROBERT RUBINSTEIN[†] AND YE ZHOU[‡]

Abstract. Taylor series expansions of turbulent time correlation functions are applied to show that helicity influences Eulerian time correlations more strongly than Lagrangian time correlations: to second order in time, the helicity effect on Lagrangian time correlations vanishes, but the helicity effect on Eulerian time correlations is nonzero. Fourier analysis shows that the helicity effect on Eulerian time correlations is confined to the largest inertial range scales. Some implications for sound radiation by swirling flows are discussed.

Key words. helicity, time correlations

Subject classification. Fluid Mechanics

1. Introduction. The theory of turbulent time correlations might be said to have begun with Kraichnan's demonstration [1] that the DIA response equation is infrared divergent when evaluated on a Kolmogorov inertial range. This calculation demonstrated the nonlocality of Eulerian time correlations. In later Lagrangian modifications [2] of DIA, the corresponding integrals converge; this confirmed the intuitive expectation [3] that Lagrangian time correlations are local in scale.

A simplification of Kraichnan's calculations was proposed by Kaneda [4] who observed that short time Taylor series expansions using the Navier-Stokes equations could produce useful information without explicitly invoking a specific closure scheme. These expansions express two-time correlations in terms of single-time correlations. Later, Kaneda and Gotoh [5] posed the question of the effect of helicity on Lagrangian and Eulerian time correlations. The present report proposes to complete this calculation using Kaneda's method and to note some possible applications.

2. Helical Turbulence. In the classic account of the kinematics of isotropic turbulence, Batchelor [6] noted the possibility that the turbulent random velocity field can lack mirror symmetry, and called this condition '*skew isotropy*.' Subsequent work used the term '*helicity*' instead. In terms of the single-time velocity correlation $Q_{ij}(\mathbf{k})$ defined by

$$(1) \quad \langle v_i(\mathbf{k}, t) v_j(\mathbf{k}', t) \rangle = Q_{ij}(\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}')$$

a helical random field is such that $Q_{ij}(\mathbf{k}) \neq Q_{ij}(-\mathbf{k})$, although necessarily $Q_{ij}(\mathbf{k}) = Q_{ji}(-\mathbf{k})$. Therefore, the correlation tensor is not symmetric: $Q_{ij}(\mathbf{k}) \neq Q_{ji}(\mathbf{k})$.

We will assume time-stationary, homogeneous, isotropic turbulence. For helical turbulence, the single-time correlation function is

$$(2) \quad Q_{ij}(\mathbf{k}) = Q(k) P_{ij}(\mathbf{k}) + i \epsilon_{imj} k_m H(k)$$

*This research was supported by the National Aeronautics and Space Administration under Contract No. NAS1-97046 while the authors were in residence at the Institute for Computer Applications in Science and Engineering (ICASE), NASA Langley Research Center, Hampton, VA 23681-2199.

[†] Institute for Computer Applications in Science and Engineering, NASA Langley Research Center, Hampton, VA 23681

[‡] Institute for Computer Applications in Science and Engineering, NASA Langley Research Center, Hampton, VA 23681 and IBM Research Division, T.J. Watson Research Center, P.O. Box, 218, Yorktown Heights, NY 10598.

Note that the *helicity spectrum* $H(k)$ contributes neither to the total energy of the velocity field nor more generally to any of the Reynolds stress components. A nonvanishing helicity spectrum is therefore consistent with the symmetry of the Reynolds stress tensor.

Define total helicity H by

$$(3) \quad H = \langle \mathbf{u} \cdot \nabla \times \mathbf{u} \rangle = \int d\mathbf{k} \, 2k^2 H(k)$$

An elementary calculation [7] shows that total helicity is an inviscid invariant of the equations of motion. Helicity has a geometric significance in terms of the linking of vortex lines: if the vortex lines never link, the total helicity vanishes, but nonzero helicity indicates nontrivial linking of vortex lines. [7]

Although initial speculation that helicity plays a dynamic role in three-dimensional turbulence analogous to enstrophy in two-dimensional turbulence has not proven fruitful, interest in the possibility that helicity impedes energy transfer remains. Moffat and Tsinober [7] provide a recent review of helicity effects in turbulence.

3. Analysis. First note that

$$(4) \quad \langle u_p \frac{\partial u_i}{\partial x_q} \rangle = \frac{1}{6} H \epsilon_{piq}$$

Let us begin by calculating the effect of helicity on Lagrangian single-point two-time correlations.¹ Let

$$(5) \quad u'_i(t) = \frac{Du_i}{Dt}$$

denote the convective derivative, so that

$$(6) \quad u'_i(t) = -\frac{\partial p}{\partial x_i}$$

Then the Lagrangian two-time correlation can be expanded in the Taylor series

$$(7) \quad \langle u_i(t) u_i(0) \rangle_L = \langle u_i u_i \rangle + \langle u'_i u_i \rangle t + \langle u''_i u_i \rangle \frac{1}{2} t^2 + \dots$$

where in Eq. (7) and subsequently, a velocity without a time argument indicates evaluation at time $t = 0$. In homogeneous turbulence, the linear term vanishes in view of Eq. (6), and stationarity in time implies

$$(8) \quad \langle u''_i u_i \rangle = -\langle u'_i u'_i \rangle = -\langle \frac{\partial p}{\partial x_i} \frac{\partial p}{\partial x_i} \rangle$$

It is convenient to write the last correlation as the Fourier transform

$$(9) \quad \begin{aligned} \langle \frac{\partial p}{\partial x_i} \frac{\partial p}{\partial x_i} \rangle &= \int d\mathbf{k} \, k^{-2} \int \delta(\mathbf{k} - \mathbf{p} - \mathbf{q}) \delta(-\mathbf{k} - \mathbf{p}' - \mathbf{q}') d\mathbf{p} d\mathbf{q} d\mathbf{p}' d\mathbf{q}' \times \\ &\quad \langle p_j q_i p'_n q'_m u_i(\mathbf{p}) u_j(\mathbf{q}) u_m(\mathbf{p}') u_n(\mathbf{q}') \rangle \\ &= \int d\mathbf{k} \, k^{-2} k_j k_i k_m k_n \int \delta(\mathbf{k} - \mathbf{p} - \mathbf{q}) \delta(-\mathbf{k} - \mathbf{p}' - \mathbf{q}') d\mathbf{p} d\mathbf{q} d\mathbf{p}' d\mathbf{q}' \times \\ &\quad \langle u_i(\mathbf{p}) u_j(\mathbf{q}) u_m(\mathbf{p}') u_n(\mathbf{q}') \rangle \end{aligned}$$

Assuming the quasi-normal closure for fourth-order single-time correlations,

$$(10) \quad \langle \frac{\partial p}{\partial x_i} \frac{\partial p}{\partial x_i} \rangle = 2 \int d\mathbf{k} \, k_i k_j k_m k_n k^{-2} \int \delta(\mathbf{k} - \mathbf{p} - \mathbf{q}) d\mathbf{p} d\mathbf{q} \, Q_{im}(\mathbf{p}) Q_{jn}(\mathbf{q})$$

¹We are grateful to Prof. Y. Kaneda for this argument.

On substituting Eq. (2) in Eq. (10), the symmetry of the wavevector factors in Eq. (10) implies that helicity makes no contribution to the second order expansion of the Lagrangian time correlation.

In the corresponding calculation for Eulerian correlations, let

$$(11) \quad \dot{u}_i = \frac{\partial u_i}{\partial t}$$

The obvious analogs of Eqs. (7)-(8) apply. Then

$$(12) \quad \langle \dot{u}_i \dot{u}_i \rangle = \langle u_p \frac{\partial u_i}{\partial x_p} u_q \frac{\partial u_i}{\partial x_q} \rangle + 2 \langle u_i \frac{\partial p}{\partial x_i} \rangle + \langle \frac{\partial p}{\partial x_i} \frac{\partial p}{\partial x_i} \rangle$$

The second correlation on the right side vanishes by homogeneity, and it has just been shown that there is no helicity contribution to the the last term. Expanding the first correlation by quasi-normality,

$$(13) \quad \begin{aligned} \langle u_p \frac{\partial u_i}{\partial x_p} u_q \frac{\partial u_i}{\partial x_q} \rangle &= \langle u_p \frac{\partial u_i}{\partial x_p} \rangle \langle u_q \frac{\partial u_i}{\partial x_q} \rangle + \langle u_p u_q \rangle \langle \frac{\partial u_i}{\partial x_q} \frac{\partial u_i}{\partial x_q} \rangle \\ &\quad + \langle u_p \frac{\partial u_i}{\partial x_q} \rangle \langle u_q \frac{\partial u_i}{\partial x_p} \rangle \\ &= \frac{2}{3} k \varepsilon / \nu - \frac{1}{6} H^2 \end{aligned}$$

The first term in Eq. (13) is the non-helical contribution to the time correlation and the second term shows the effect of helicity. Note that helicity decreases the turbulent frequency, hence it increases the Eulerian time scale.

This calculation can be refined somewhat by evaluating the same effect in wavevector space. Substituting Kolmogorov forms in Eq. (2),

$$(14) \quad Q_{ij}(\mathbf{k}) = C_K \varepsilon^{2/3} k^{-5/3} P_{ij}(\mathbf{k}) / 4\pi k^2 + C_H \varepsilon_H^{2/3} i \epsilon_{imj} k_m k^{-7/3} / 4\pi k^3$$

Then

$$(15) \quad H = 6 C_H \varepsilon_H^{2/3} k_0^{-1/3}$$

where $2\pi k_0^{-1}$ is the integral scale of the turbulence.

The quadratic term in Kaneda's expansion is

$$(16) \quad \langle \ddot{u}_i(\mathbf{k}) u_j(-\mathbf{k}) \rangle = -\frac{1}{4} \int \delta(\mathbf{k} - \mathbf{p} - \mathbf{q}) d\mathbf{p} d\mathbf{q} P_{imn}(\mathbf{k}) P_{mrs}(\mathbf{p}) Q_{rj}(\mathbf{k}) Q_{ns}(\mathbf{q})$$

We will investigate how this integral diverges in the limit $\mathbf{q} \rightarrow 0$, isolating the helical contribution. In this limit, the contribution due to helicity to the integrand in Eq. (16) is

$$(17) \quad \begin{aligned} P_{imn}(\mathbf{k}) P_{mrs}(\mathbf{p}) Q_{rj}(\mathbf{k}) Q_{ns}(\mathbf{q}) &= -P_{imn}(\mathbf{k}) \{ P_{mrs}(\mathbf{k}) - q_r P_{ms}(\mathbf{k}) \\ &\quad - q_s P_{mr}(\mathbf{k}) + k_r q_a P_{ams}(\mathbf{k}) k^{-2} + k_s q_a P_{amr}(\mathbf{k}) k^{-2} \} \times \\ &\quad \epsilon_{rj\alpha} \epsilon_{ns\beta} k_\alpha q_\beta \varepsilon_H^{4/3} (kq)^{-10/3} / 16\pi^2 k^2 q^2 \end{aligned}$$

Then a simple calculation shows that

$$(18) \quad \begin{aligned} &-P_{imn}(\mathbf{k}) \{ P_{mrs}(\mathbf{k}) - q_r P_{ms}(\mathbf{k}) - q_s P_{mr}(\mathbf{k}) \\ &\quad + k_r q_a P_{ams}(\mathbf{k}) k^{-2} + k_s q_a P_{amr}(\mathbf{k}) k^{-2} \} \epsilon_{rj\alpha} \epsilon_{ns\beta} k_\alpha q_\beta \\ &= q_r \beta k_{n\alpha} \epsilon_{rj\alpha} \epsilon_{ni\beta} - k_{s\alpha} q_r \beta \epsilon_{rj\alpha} \epsilon_{is\beta} \\ &= -2o_i o_j \end{aligned}$$

where

$$(19) \quad o_i = \epsilon_{ipq} k_p q_q$$

The integral with respect to \mathbf{q} in Eq. (16) therefore diverges as

$$(20) \quad \int d\mathbf{q} q^{-10/3+2} \sim k_0^{-1/3}$$

consequently, for the helical contribution to the second order term,

$$(21) \quad \langle \ddot{u}_i(\mathbf{k}) u_j(-\mathbf{k}) \rangle \sim P_{ij}(\mathbf{k}) k^2 \epsilon_H^{2/3} k^{-10/3} \epsilon_H^{2/3} k_0^{-1/3} = P_{ij}(\mathbf{k}) E_H(k) H$$

Integration of this result over k recovers the H^2 dependence in Eq. (13) of the helical contribution to the single-point correlation. Note the non-local dependence on the total helicity, which is analogous to the non-local dependence of Eulerian time correlations on the total rms velocity fluctuations.

Define the Eulerian frequency scale by

$$(22) \quad \omega_E^2 = - \langle \ddot{u}_i(\mathbf{k}) u_i(-\mathbf{k}) \rangle / Q(k)$$

Then adding the usual sweeping contribution gives the complete Eulerian frequency scale to second order

$$(23) \quad \omega_E = [C_D^2 (Vk)^2 - C'_H H \frac{\epsilon_H^{2/3}}{\epsilon_H^{2/3} k^{1/3}}]^{1/2} = C_D V k \{1 - \frac{C'_H}{2C_D^2} \frac{H(\epsilon_H/\epsilon)^{2/3}}{V^2} k^{-5/3}\}$$

Again, the frequency is decreased by helicity. Note that the effect of helicity is confined to the largest inertial range scales.

4. Conclusion. This calculation reveals an additional dynamic distinction between Eulerian and Lagrangian time correlations. The agreement to second order of the Taylor series expansion of helical and non-helical Lagrangian time correlation functions suggests that the effect of helicity on Lagrangian time correlations is relatively weak, whereas helicity strongly alters Eulerian correlations.

The dependence of turbulent energy transfer on Lagrangian correlations [2, 3] might seem to suggest a relatively small effect of helicity on turbulent energy transfer in steady-state flows. However, closures suggest that both spatial and temporal correlations influence energy transfer, so that this analysis is certainly consistent with some connection between helicity and energy transfer. These connections are discussed in [7, 8]. It is possible that helicity could stabilize turbulence and play a role in the sustainment of large-scale structures, as suggested by Yokoi and Yoshizawa. [9]

Although less theoretical interest is naturally attached to Eulerian time correlations, they arise in problems including wave propagation through turbulence sound radiation by turbulence. In the sound radiation problem, the Eulerian space-time correlation function appears in the turbulent sound source in Lighthill's theory. [10] The effect of helicity on radiated sound is a redistribution of acoustic energy radiated by the largest inertial range scales to lower frequencies. In helical flows, like swirling jets, helicity may also greatly alter the large structures in the flow [9] and consequently modify the sweeping velocity. The combination of both helicity effects may therefore cause significant modification of sound radiation from swirling flows.

5. Acknowledgments. Our discussions with Prof. Y. Kaneda have been very helpful and are gratefully acknowledged.

REFERENCES

- [1] R.H. KRAICHNAN, *The structure of turbulence at very high Reynolds number*, J. Fluid Mech. **5**, (1959), p. 497.
- [2] R.H. KRAICHNAN, *Lagrangian-history closure approximation for turbulence*, Phys. Fluids **8**, (1965), p. 575.
- [3] R.H. KRAICHNAN, *Kolmogorov's hypotheses and Eulerian turbulence theory*, Phys. Fluids **7**, (1964), p. 1723.
- [4] Y. KANEDA, *Lagrangian and Eulerian time correlations in turbulence*, Phys. Fluids A **5**, (1993), p. 2835.
- [5] Y. KANEDA AND T. GOTOH, *Lagrangian velocity autocorrelation in isotropic turbulence*, Phys. Fluids A **3**, (1991), p. 1924.
- [6] G.K. BATCHELOR, *The theory of homogeneous turbulence*, Cambridge University Press, (1953).
- [7] H.K. MOFFAT AND A. TSINOBER, *Helicity in laminar and turbulent flow*, Ann. Rev. Fluid Mech. **24**, (1992), p. 281.
- [8] R.H. KRAICHNAN, *Helical turbulence and absolute equilibrium*, J. Fluid Mech. **59**, (1973), p. 745.
- [9] N. YOKOI AND A. YOSHIKAWA, *Statistical analysis of the effects of helicity in inhomogeneous turbulence*, Phys. Fluids A **5**, (1993), p. 464.
- [10] M.J. LIGHTHILL, *On sound generated aerodynamically: I. General theory*, Proc. Roy. Soc. **A211**, (1952), p. 1107.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE November 1998	3. REPORT TYPE AND DATES COVERED Contractor Report		
4. TITLE AND SUBTITLE Effects of Helicity on Lagrangian and Eulerian Time Correlations in Turbulence		5. FUNDING NUMBERS C NAS1-97046 WU 505-90-52-01		
6. AUTHOR(S) Robert Rubinstein Ye Zhou				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute for Computer Applications in Science and Engineering Mail Stop 403, NASA Langley Research Center Hampton, VA 23681-2199		8. PERFORMING ORGANIZATION REPORT NUMBER ICASE Report No. 98-49		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Langley Research Center Hampton, VA 23681-2199		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA/CR-1998-208737 ICASE Report No. 98-49		
11. SUPPLEMENTARY NOTES Langley Technical Monitor: Dennis M. Bushnell Final Report Submitted to Physics of Fluids				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified Unlimited Subject Category 34 Distribution: Nonstandard Availability: NASA-CASI (301)621-0390		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Taylor series expansions of turbulent time correlation functions are applied to show that helicity influences Eulerian time correlations more strongly than Lagrangian time correlations: to second order in time, the helicity effect on Lagrangian time correlations vanishes, but the helicity effect on Eulerian time correlations is nonzero. Fourier analysis shows that the helicity effect on Eulerian time correlations is confined to the largest inertial range scales. Some implications for sound radiation by swirling flows are discussed.				
14. SUBJECT TERMS helicity; time correlations		15. NUMBER OF PAGES 10		16. PRICE CODE A03
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	