

11

Quaternary Research,
in press

Final
copy

3/4/98

11 4/1
11 4/1

**Antiphasing Between Rainfall in
Africa's Rift Valley and North America's Great Basin**

Wallace S. Broecker
Lamont-Doherty Earth Observatory
Palisades, NY 10964

Dorothy Peteet
Lamont-Doherty Earth Observatory
Palisades, NY 10964

and
NASA's Goddard Institute for Space Studies
New York, NY 10025

Irena Hajdas
ITP ETH Hoenggerberg
CH-8093 Zürich, Switzerland

Jo Lin
Department of Geography
University of California
Berkeley, CA 94720

Elizabeth Clark
Lamont-Doherty Earth Observatory
Palisades, NY 10964

In press

Quaternary Research

Abstract

The beginning of the Bølling-Allerød warm period is marked in Greenland ice by an abrupt rise in $\delta^{18}\text{O}$, an abrupt drop in dust rain, and an abrupt increase in atmospheric methane content. The surface waters in the Norwegian Sea underwent a simultaneous abrupt warming. At about this time, a major change in the pattern of global rainfall occurred. Lake Victoria (latitude 0°), which prior to this time was dry, was rejuvenated. The Red Sea, which prior to this time was hypersaline, freshened. Lake Lahontan, which prior to this time had achieved its largest size, desiccated. Whereas the chronologic support for the abruptness of the hydrologic changes is firm only for the Red Sea, in keeping with evidence obtained well away from the northern Atlantic in the Santa Barbara basin and the Cariaco Trench, the onset and end of the millennial-duration climate events were globally abrupt. If so, the proposed linkage between the size of African closed basin lakes and insolation cycles must be reexamined.

Johnson *et al.* (1996) reported an astounding phenomenon. Based on sediment cores and seismic profiles across the deep portion of equatorial Africa's Lake Victoria, they concluded that prior to about 12,800 ^{14}C years ago, this lake was dry. The evidence comes from a soil encountered in piston cores at the base of the late-glacial-Holocene limnic sequence. The sound reflection from the sediment-soil interface allows its extent to be traced across the entire deep portion of the Lake Victoria basin. Whereas it has long been known that the rift valley lakes experienced dry conditions during peak glacial time (Haberyan and Hecky, 1987; Gasse *et al.*, 1989; Taylor, 1990), the Victoria results provide a means of quantifying the magnitude of this drying.

We say astounding, because in order to dry out this lake, an enormous change in climate is required. To see this, we must consider today's water budget (Table 1). Currently, the amount of water entering the lake from the rivers that feed it is comparable to the amount overflowing into the White Nile from Victoria (Piper *et al.*, 1986). Based on rainfall at stations around the lake (Fig. 1), it is estimated that four times as much water is added to the lake by direct rainfall than from rivers. Taken together, these three sets of measurements suggest that evaporation from the lake surface must nearly match rainfall onto the lake (i.e., ca. 1.6 m/yr). Indeed, independent estimates of evaporation rate are consistent with this conclusion (Crul, 1995). The central 20% of the lake's basin is deeper than 60 m. It is in this portion that the coring and seismic surveys conducted by Johnson *et al.* (1996) were concentrated (Fig. 1). Clearly, in order to dry the lake, its size would have to be reduced by at least a factor of ten. Assuming no change in evaporation rate, this would require that the input of water to the lake be reduced tenfold. However, as today only 9% of the rain water falling onto the land portion of the Victoria drainage basin is discharged into the lake, the fractional reduction in rainfall need not be nearly as large as that in lake size. For example, at the -60 m level, 80% of the former lake area would have become land. This complicates the problem of estimating the paleo-rainfall rate. The relationship between the steady state size of the lake (A) and the rate of rainfall (R) and the rate of evaporation (E) over the lake is given by the following equation:

$$EA = fR (D - A) + RA$$

where D is the area of the drainage basin and f is the fraction of rainfall onto the land portion of the drainage basin which runs off into the lake. Solving for R, this equation becomes:

$$R = \frac{E(A / D)}{f + (1 - f)(A / D)}$$

Assuming that the evaporation rate over the lake and fraction of runoff from the land portion of the drainage basin were equal to their present day values (i.e., 1.6 and 0.09 m/yr, respectively), then this equation becomes

$$R = \frac{0.42(A / A^\circ)}{0.09 + 0.24(A / A^\circ)}$$

where A° is the present size of the lake (i.e., 0.26 D). This relationship is displayed graphically in Figure 2. As can be seen, in order to dry up, the rainfall rate would have to be reduced by at least a factor of four (i.e., to no more than 0.4 m/yr). Of course, with drier conditions, it might be expected that the evaporation rate over the lake would rise and that the fraction of runoff from the land portion of the basin would fall. However, to the extent the tropics cooled during glacial time, evaporation rates would have been reduced. Unfortunately, there is no way to assess how large these changes would be. Nevertheless, it is clear that in order to dry up, the rainfall in the Victoria drainage basin must have been far lower than at present.

The documentation by Johnson *et al.* (1996) of dry conditions in the equatorial portion of the African rift zone prior to 12,800 years ago is consistent with previous studies (Table 2). Taylor (1990) and Bonnefille *et al.* (1990) provide pollen evidence for dry conditions during the period 21,000 to ca. 12,000 ^{14}C years ago. Roberts *et al.* (1993) present evidence for dry conditions prior to 12,700 yr B.P. in the Lake Magadi basin (2°S). Gasse *et al.* (1989) document low stands of Lake Tanganyika (3° to 9°S) from 22,000 to 13,000 ^{14}C yr ago, with a wet phase following 13,000 ^{14}C yr ago. Although we would like to believe that the studies by Johnson *et*

al. (1996) and Gasse *et al.* (1989) indicate that the transition from wet to dry conditions close to 12,800 years ago was abrupt, we must admit that firm supporting evidence is lacking. Needed are radiocarbon dates for terrestrial macrofossils from basal sediments at various water depths in Lake Victoria. Only if all these ages group close to 12,800 ^{14}C yr B.P. could it be stated with confidence that conditions went from dry to comparable to today's in a short interval of time. Such a series of dates would also firm up the apparent tie between the timing of this wetting and that of the onset of the Bølling-Allerød in northern Europe about 12,550 ^{14}C yr ago (Table 3).

The dry interval in Africa's equatorial rift valley corresponds to that during which hypersaline conditions prevailed in the Red Sea (Deuser and Degens, 1969; Hemleben *et al.*, 1996) as indicated by the absence of planktonic foraminifera (Fig. 3) and by thick secondary aragonite coatings on the pteropod shells (Fig. 4). As shown by Bijma (1991), planktonic foraminifera growth is highly impaired at salinities above 44 gm/L. In this case, the end of this period of hypersalinity was abrupt. It occurred close to 13,000 ^{14}C yr B.P. (Fig. 4). One could postulate that the abruptness of this change merely reflects the point at which a more gradual end to the period of dry conditions brought the salinity down to the point where foraminifera could survive. However, in this case, the match between the timing of Lake Victoria's rejuvenation and the Red Sea's passage through a salinity threshold would have to be written off as a coincidence. To us, this match suggests a rapid change in regional climate.

Having maintained a continuing interest in the history of another similar large intermountain water body, i.e., Lake Lahontan, located in the western part of the Great Basin (36-42°N), we were struck by the asymmetry of its late-glacial wetness record with that of Lake Victoria. During the interval 21,000 to 13,000 ^{14}C yr B.P., Lake Lahontan and other of the lakes in the Great Basin (i.e., Bonneville, Searles, and Russell) were far larger than today's small remnants (Benson *et al.*, 1990, 1995). Very close to the time that Victoria was rejuvenated, Lake Lahontan experienced a dramatic desiccation that reduced its area close to that occupied by its late Holocene remnants (Table 4). The area of these lakes in the year 1900 (i.e., before diversions

of its feed rivers for irrigation use) was about one tenth that of Lake Lahontan at its maximum extent (Benson, *et al.*, 1990).

While not precisely defined, the desiccation of Lake Lahontan was well underway no later than 12,300 ^{14}C yr ago and it began no earlier than about 13,800 ^{14}C yr ago. The lower age limit is firm because it is based on radiocarbon dating of terrestrial wood samples from packrat middens in small rock shelters 90 m below the highest level of the lake (Table 5). Studies dating back to that of Russell (1885) conclude that the largest size achieved by the lake occurred close to the end of the last pluvial interval. Radiocarbon dates for CaCO_3 associated with the highest shorelines of these lakes average about 12,800 yr, and range from 12,200 to 13,800 yr B.P. (Table 5, Fig. 5). While consistent with the interpretation that this high stand was brought to an end by a sudden desiccation, several red flags must be raised. First, a reservoir correction must be made to account for the difference between the $^{14}\text{C}/\text{C}$ for lake ΣCO_2 and that for atmospheric CO_2 . As discussed by several authors (Broecker and Walton, 1959; Benson, 1993), this correction could be anywhere from 200 to 700 years. If the proper correction were toward the high end of this range, when made, it would bring most of these radiocarbon ages below the firm limit set by the packrat midden wood ages. But, another factor could swing the pendulum back toward older ages; namely, exchange with younger CO_2 or HCO_3^- or the deposition of secondary CaCO_3 likely reduced the ^{14}C age of all these samples. In hindsight, the early measurements on bulk uncleaned tufa made by Broecker and Orr (1958) clearly document such contamination. Later attempts by Benson (1981) and by Lin *et al.* (1996) using dense and acid-leached tufa give the ^{14}C ages summarized in Figure 5. However, there is still no guarantee that all the contamination was entirely removed. The fact that several quite different types of material (tufa, shell, and organic matter encased in tufa) yield similar ages suggests that contamination is under control. If the proper reservoir correction is at the low end of the permissible range (i.e., ~200 yr), one could take the ages at face value and conclude that the lake stood within 20 m of its highest level between 13,800 and 12,400 yr ago.

However this conclusion presents a major problem. Uranium series ages on high shoreline tufas yielding ^{14}C ages of $\sim 12,500$ yr are close to 16,500 cal yr for both Lahontan and Searles (Lin *et al.*, 1996). These calendar ages translate to about 13,700 yr on the radiocarbon time scale (Bard *et al.*, 1992). For Lake Lahontan, the ^{230}Th age is based on a well-constrained isochron (Lin *et al.*, 1996). For Searles Lake, the $^{230}\text{Th}/^{232}\text{Th}$ ratio measured in a tufa from the highest shoreline is so large that the isochron approach is not required (Lin, 1996; Lin *et al.*, in press). It is possible, of course, that uranium has been lost from the tufas over their lifetime. This explanation seems unlikely because it would be fortuitous that the loss affected the Lahontan and the Searles tufas by the same amount. Thus, in the absence of any satisfactory explanation for the large range in the ^{14}C ages for high shoreline samples and for the discordance between the ^{230}Th and ^{14}C ages on identical tufas, we can conclude only that the desiccation of Lake Lahontan occurred sometime between 13,700 and 12,300 ^{14}C yr ago. While this range brackets the ^{14}C age for the end of Lake Victoria's period of desiccation and that for the end of the Red Sea's hypersalinity period, it leaves open the question as to whether these events were synchronous.

We were interested to learn from B.D. Allen and R.Y. Anderson (personal communication, 1997) of the University of New Mexico that Lake Estancia desiccated shortly after $12,490 \pm 60$ ^{14}C yr B.P. (based on AMS measurements on thoroughly cleaned hand-picked ostracods). This playa, located southeast of Albuquerque, has deflation blowouts in which the sedimentary record for the last ca. 50,000 yr is well displayed, allowing a more-detailed history of pluvial conditions to be reconstructed than is possible for any other Great Basin lake.

We would like to believe that this evidence points to an abrupt change in the pattern of global rainfall. Of course, were this conclusion based on the records for just two places on the globe, it could be passed off as a coincidence. However, the time of this change is tantalizingly close to that of the abrupt onset of the northern Atlantic basin's Bølling-Allerød warm interstade. This event is one of the most prominent features in the Greenland ice core records and also in the pollen records from small lakes north of the Alps (Lotter, 1991). Its radiocarbon age is 12,550 yr (Table 3). Furthermore, pollen changes in eastern North America (Peteet *et al.*, 1990, 1993;

Maenza-Gmelch, 1997a, 1997b; Kneller and Peteet, 1998), and changes in the beetle assemblage in the Rocky Mountains (Elias, 1996) and of pollen in the Chilean lake district (Lowell *et al.*, 1995) occurred at about this time. Also, meltwater event #1 revealed in the Barbados coral record occurred on the heels of this event (Fairbanks, 1989). As proponents of abrupt and globally synchronous climate changes, we take these similarities in timing to indicate, as is the case for the abrupt onset and demise of the Younger Dryas cold event, that the climate change at the onset of the Bølling-Allerød interstade was a global event. Nevertheless, we must admit that at present the accuracy of relative dating is no better than ± 500 yr.

It should be noted that we take a different view of Africa's pluvial history than do many authors. Following the lead of Kutzbach and Street-Perrott (1985), it has become common practice to relate the changes discussed here to the cycle in the strength of the monsoon caused by the orbitally induced cycles in seasonality over Africa and Asia. The fact that the transition from dry to wet conditions occurred during the rise toward a maximum in northern hemisphere seasonality fits well with this explanation. On the other hand, the timing of this transition also seems to fit neatly into the record of abrupt events in the northern Atlantic basin. The frequency of these events is far too high to be attributable to Milankovitch forcing, so the question is: did rainfall in equatorial Africa follow seasonality or did it respond abruptly to a switch in the climate system's mode of operation? Whereas there is no doubt that, when averaged over thousands of years, the Earth's climate has been paced by Milankovitch cycles, the climate record appears to be telling us that, rather than preceding in smooth sinusoids, climate has responded by jumping from one mode of operation to another (Broecker and Denton, 1989). Thus, whereas Kutzbach and his followers may be correct in concluding that the wetting of equatorial Africa was brought about by the increasing of seasonality, it is our view that the wetting did not smoothly follow the rise in seasonality but came mainly in a single jump that reflected a reorganization of the Earth's climate system at the onset of the Bølling-Allerød warm interval.

This conclusion is consistent with the records of $\delta^{18}\text{O}$ (Dansgaard *et al.*, 1989), dust fall (Taylor *et al.*, 1993) and the atmospheric methane content (Chappellaz *et al.*, 1993) from the Greenland ice cores and of conditions in the Santa Barbara basin (Behl and Kennett, 1996) and the Cariaco Trench (Hughen *et al.*, 1996). Indeed, the onset of the Bølling-Allerød warm interval appears to have been marked by an abrupt reorganization of the Earth's climate system.

TABLE 1
Vital statistics and water budget for Lake Victoria

(based on Crul, 1995)

<i>Lake Area</i>	0.69 x 10 ⁵ km ²
Dry Drainage Area	1.94 x 10 ⁵ km ²
TOTAL AREA	2.63 x 10 ⁵ km ²
<i>Lake Volume</i>	2.76 x 10 ³ km ³
Mean Depth	40 m
Maximum Depth	79 m
<i>Inputs to lake</i>	
Rain	1.66 m/yr
River	0.42 m/yr
TOTAL	2.08 m/yr
<i>Losses from lake</i>	
Evaporation	1.59 m/yr
Outflow	0.49 m/yr
TOTAL	2.08 m/yr
<i>H₂O residence time</i>	~20 yr

TABLE 2
Radiocarbon dates ^{to} related the boundary between the late-
glacial, cold-dry interval and the succeeding wet-warm period
in equatorial Africa

Location	Radiocarbon age	
	yr B.P.	Reference
Lake Kivu	13,700 - 12,500	Hecky and Degens, 1973
Lake Victoria	13,750 ± 200	Kendall, 1969
Lake Albert	13,340 ± 190	Livingstone, 1975
Kawasenkoko Swamp	13,835 ± 120	Hamilton, 1982
Kaisungor Swamp	12,650 ± 100	Coetzee, 1967
Sacred Lake	14,050 ± 360	Coetzee, 1967
Laboot Swamp	13,776 ± 80	Hamilton, 1976
Lake Mahoma	12,700 ± 120	Hamilton, 1982
Muchoya Swamp	12,890 ± 130	Morrison, 1968
Muchoya Swamp	13,900 ± 130	Taylor, 1990
Kamiranzovu Swamp	13,575 ± 130	Hamilton, 1982
Lake Tanganyika	12,710 ± 120	Gasse <i>et al.</i> , 1989
Kashiru Swamp	13,250 ± 200	Bonnefille <i>et al.</i> , 1990
Lake Magadi	12,840 ± 120	Roberts <i>et al.</i> , 1993

TABLE 3
Radiocarbon ages related to the onset of the Bølling-Allerød warm interval in northern Europe*

Location	Material	Radiocarbon Age	
		(yr B.P.)	Reference
Ballybetagh, Ireland	Twig	12,540 ± 80	Cwynar and Watts, 1989
Neuchatel, Switzerland	Ter. macro-fossils	12,490 ± 95	Schwalb <i>et al.</i> , 1994
Lobsigensee, Switzerland	Ter. macro-fossils	12,300 to 12,800	Ammann and Lotter, 1989
Zurichsee, Switzerland	Ter. macro-fossils	12,600 ± 200	Lister, 1988
Holzmaar, Germany	Ter. macro-fossils	12,560 ± 80	Zolitschka, 1996
Lago di Monticchio, Italy	Betula seeds	12,540 ± 130	Watts <i>et al.</i> , 1996

*The radiocarbon ages for this transition all lie in the range 12,550 ± 100 yr B.P.

TABLE 4
Change in area of lakes in the African equatorial rift valley
and in the North American Great Basin

and in the North American Great Basin			
Lake	Area of present-day lake	Area ~14,000 ¹⁴ C yr ago	$\frac{A_{14,000}}{A_{\text{present}}}$
	(km ²)	(km ²)	
Africa			
Victoria	69,000	<7000	< $\frac{1}{10}$
North America *			
Lahontan	2420	22,300	-9
Bonneville	5140	51,300**	-10
Searles	315	1725	-5
Russell	190	790	-4

*See Benson *et al.*, 1990.

**Area before overflow and downcutting of outlet (Currey, 1988).

TABLE 5

Location	Elevation (m)	Material	Radiocarbon age (yr B.P.)	Reference
Lahontan Post Highstand Desiccation				
Fishbone Cave	1235	<i>Equus</i>	12,280 ± 180	Benson and Thompson, 1987
Crypt Cave	1240	Juniper	12,350 ± 180	"
Crypt Cave	1240	Juniper + dung	12,240 ± 180	"
Lahontan Basin (Highest Level ~1330 m)				
Pyramid Lake	1321	Tufa	12,540 ± 190	Benson, 1981
"	1321	"	12,570 ± 190	"
"	1325	"	12,610 ± 180	"
"	1326	"	12,770 ± 190	"
"	1312	"	13,820 ± 200	"
"	1331	"	12,850 ± 190	"
"	1303	"	12,890 ± 190	"
"	1324	"	13,050 ± 190	"
"	1312	"	13,130 ± 190	"

TABLE 5 (continued)

Location	Elevation (m)	Material	Radiocarbon age (yr B.P.)	Reference
Pyramid Lake	1311	Tufa	13,430 \pm 200	Benson, 1981
"	1311	"	13,550 \pm 200	"
"	1330	"	13,060 \pm 100	Lin <i>et al.</i> , 1996
"	1330	"	13,150 \pm 100	"
"	1330	"	12,200 \pm 100	"
"	1330	"	12,850 \pm 100	"
"	1327	"	12,320 \pm 100	"
"	1327	"	12,200 \pm 100	"
"	1327	"	12,980 \pm 100	"
"	1327	"	12,690 \pm 100	"
"	1327	"	12,830 \pm 95 *	"
"	1327	"	12,790 \pm 110	"
"	1327	"	12,260 \pm 100	"
"	1328	Organic Matter +	12,870 \pm 120	Lin, 1996
"	1311	Shell	13,260 \pm 200	Benson, 1981
Black Rock Desert	1306	Tufa	13,810 \pm 600	Benson and Thompson, 1987

TABLE 5 (continued)

Location	Elevation (m)	Material	Radiocarbon age (yr B.P.)	Reference
"	1332	"	12,850 ± 600	"
Walker Lake	1318	"	12,240 ± 160	Benson, 1981
"	1327	"	12,280 ± 160	"
"	1324	"	12,340 ± 160	"
Walker Lake	1324	Tufa	12,690 ± 160	Benson and Thompson, 1987
"	1330	"	13,300 ± 190	Benson, 1981
"	1330	"	13,300 ± 190	"
"	1330	"	13,300 ± 180	"
Carson Sink	1311	"	12,310 ± 150	Benson and Thompson, 1987
"	1323	"	12,980 ± 540	"
"	1327	"	12,700 ± 95	Lin, 1996
"	1325	Shell	13,110 ± 110	"
"	1325	"	13,280 ± 110	"
Searles Lake (Highest stand 695 m)				
Navy Road	695	Tufa	12,430 ± 90	Lin, 1996

TABLE 5 (continued)

Location	Elevation (m)	Material	Radiocarbon age (yr B.P.)	Reference
Lake Russell (Highest stand 2155 m)				
Monocraters	2155	Tufa	12,910 ± 70	Benson <i>et al.</i> , 1990
Black Point	2144	"	12,770 ± 70	"

Table 5 Caption

Radiocarbon ages for the high stands of the lakes (Lahontan, Searles and Russell) in the western Great Basin range from 12,200 to 13,800 ^{14}C years. Almost all these dates are on beach-rock deposits (tufa). Only two are on shell and one is on organic matter encapsulated in the tufa. A reservoir correction estimated to be somewhere between 200 and 700 years has not been included in the table. Uranium series dating of tufas from the high shoreline of Lahontan and Searles give age close to 17,000 years or ~14,000 years on the radiocarbon scale (Bard *et al.*, 1992). Terrestrial organic matter from pack-rat middens in the Winnemucca caves located 90 meters below the highest shoreline of Lake Lahontan demonstrate that drying was well underway by 12,300 ^{14}C years ago. Hence, at least some and perhaps all of the ^{14}C ages have been shifted downward by the incorporation of younger CaCO_3 .

Figure Captions

FIG. 1. Map showing the depth contours (in m) for Lake Victoria, the locations of piston cores studied by Johnson *et al.* (1996), and the seismic profiling tracks along which the soil horizon formed during the dry period was mapped. Also shown are the mean monthly rainfall rates (histograms, mm/month) and the mean annual rainfall (in boxes, m/yr) for eight stations (Crul, 1995). Shown on the lower right are the radiocarbon dates obtained by Johnson *et al.* (1996) for core V95-2P which document that the dry period came to a close about 12,800 ^{14}C yr ago.

FIG. 2. Steady-state size of Lake Victoria as a function of the rate of rainfall. The assumption is made that the rate of evaporation over the lake (i.e., 1.6 m/yr) and the fractional runoff of rain falling on the dry portion of the drainage basin (i.e., 0.09) remain unchanged. The rainfall required to maintain the lake just below its outflow level is 1.25 m/yr. Assuming that in order to dry up entirely the area of the lake must be reduced to less than 10% of its present area, the

rainfall during the period of desiccation could have been no more than 0.4 m/yr (i.e., less than a quarter of the present rate).

FIG. 3. Oxygen isotope record for planktonic foraminifera from a Red Sea sediment core, KL11 (Hemleben et al., 1996). The exceptionally large glacial to interglacial $\delta^{18}\text{O}$ swings are attributed to hypersalinity during glacial times when much of the water flowing in from the Indian Ocean was lost to evaporation. The absence of planktonic foraminifera in the 80 cm-long-section of the core corresponding to marine isotope stage 2 suggests that the interval of hyperaridity lasted ca. 10,000 yr.

FIG. 4. A. Oxygen and carbon isotope records from pteropods from the Red Sea sediment cores (Deuser and Degens, 1969). The open circles are for measurements on pteropods free of secondary coatings and the closed circles for those on pteropods heavily encrusted with secondary aragonite. B., C. ^{14}C ages (corrected for a 400-yr age of surface-water) versus depth in two Red Sea cores. In both, the transition from hypersaline to present conditions occurred about 13,000 ^{14}C yr ago.

FIG. 5. Histogram of radiocarbon ages (Table 2) for samples collected within 20 m of the highest shorelines of Lake Lahontan (open), Searles Lake (filled), and Lake Russell (shaded). Except for the squares marked S (for shell) and OM (for organic matter encapsulated in the tufa), the results are all on tufa CaCO_3 . Note that ^{230}Th dates (transformed to the ^{14}C scale) suggest that the raw ^{14}C ages (i.e., without reservoir correction) range up to 1800 years too young. At least the youngest of the tufas must be contaminated with younger carbon, as their ^{14}C ages overlap firm ^{14}C ages on woody terrestrial plants from packrat middens 90 m below the highest shoreline of Lake Lahontan.

REFERENCES

- Ammann B., Lotter, A. F. (1989). Late-Glacial radiocarbon- and palynostratigraphy on the Swiss Plateau. *Boreas* **18**, 109-126.
- Bard, E., Fairbanks, R. G., Arnold, M., and Hamelin, B. (1992). $^{230}\text{Th}/^{234}\text{U}$ and ^{14}C ages obtained by mass spectrometry on corals from Barbados (West Indies), Isabela (Galapagos) and Murua (French Polynesia). In "The Last Deglaciation: Absolute and Radiocarbon Chronologies" (E. Bard and W. S. Broecker, Eds.), NATO ASI Series I: Global Environmental Change **2**, 103-110.
- Behl, R. J., and Kennett, J. P. (1996). Brief interstadial events in the Santa Barbara basin, NE Pacific, during the past 60 kyr. *Nature* **379**, 243-246.
- Benson, L. V. (1981). Paleoclimatic significance of lake-level fluctuations in the Lahontan Basin. *Quaternary Research* **16**, 390-403.
- Benson, L. V., and Thompson, R. S. (1987). Lake-level variation in the Lahontan Basin for the past 50,000 years. *Quaternary Research* **28**, 69-85.
- Benson, L. V., Currey, D. R., Dorn, R. I., Lajoie, K. R., Oviatt, C. G., Robinson, S. W., Smith, G. I., and Stine, S. (1990). Chronology of expansion and contraction of four Great Basin lake systems during the past 35,000 years. *Palaeogeography, Palaeoclimatology, Palaeoecology* **78**, 241-286.
- Benson, L. (1993). Factors affecting ^{14}C ages of lacustrine carbonates: Timing and duration of the last Highstand Lake in the Lahontan Basin. *Quaternary Research* **39**, 163-174.
- Benson, L., Kashgarian, M., and Rubin, M. (1995). Carbonate deposition, Pyramid Lake subbasin, Nevada: 2. Lake levels and polar jet stream positions reconstructed from radiocarbon ages and elevations of carbonates (tufas) deposited in the Lahontan basin. *Palaeogeography, Palaeoclimatology, Palaeoecology* **117**, 1-30.

- Bijma, J. (1991). *On the Biology of Tropical Spinose Globigerinidae (Sarcodina, Foraminiferida) and its Implications for Paleoecology*. PhD Thesis from Rijksuniversiteit Groningen, 242 pp., Hans-Joachim Köhler, Druck & Reprografie, Tübingen.
- Bonnefille, R., Roeland, J. C., and Guiot, J. (1990). Temperature and rainfall estimates for the past 40,000 years in equatorial Africa. *Nature* **346**, 347-349.
- Broecker, W. S., and Orr, P. C. (1958). Radiocarbon chronology of Lake Lahontan and Lake Bonneville. *Geological Society of America Bulletin* **69**, 1009-1032.
- Broecker, W. S. and Walton, A. F. (1959). The geochemistry of C14 in fresh-water systems. *Geochimica et Cosmochimica Acta* **16**, 15-38.
- Broecker, W. S., and Denton, G. H. (1989). The role of ocean-atmosphere reorganizations in glacial cycles. *Geochimica et Cosmochimica Acta* **53**, 2465-2501.
- Chappellaz, J., Blunier, T., Raynaud, D., Barnola, J. M., Schwander, J., and Stauffer, B. (1993). Synchronous changes in atmospheric CH₄ and Greenland climate between 40 and 8 kyr BP. *Nature* **366**, 443-445.
- Coetzee, J. A. (1967). Pollen analytical studies in East and Southern Africa. *Paleoecology of Africa* **3**, 1-146.
- Crul, R. C. M. (1995). *Limnology and Hydrology of Lake Victoria*, UNESCO Publishing, Paris, France, 79 pp.
- Currey, D. R. (1988). Isochronism of final Pleistocene shallow lakes in the Great Salt Lake and Carson Desert regions of the Great Basin. In "Abstract Program of American Quaternary Association Tenth Biennial Meeting" 117.
- Cwynar, L. C., and Watts, W. A. (1989). Accelerator-mass spectrometer ages: Late-Glacial events at Ballybetagh, Ireland. *Quaternary Research* **31**, 377-380.
- Dansgaard, W., White, J. W. C., and Johnsen, S. J. (1989). The abrupt termination of the Younger Dryas climate event. *Nature* **339**, 532-533.

- Deuser, W. G., and Degens, E. T. (1969). O^{18}/O^{16} and C^{13}/C^{12} ratios of fossils from the hot-brine deep area of the central Red Sea, In "Hot Brines and Recent Heavy Metal Deposits in the Red Sea" (E. T. Degens and D. A. Ross, Eds.), pp. 336-347, Springer-Verlag, New York.
- Elias, S. A. (1996). Late Pleistocene and Holocene seasonal temperatures reconstructed from fossil beetle assemblages in the Rocky Mountains. *Quaternary Research* 46, 311-318.
- Fairbanks, R. G. (1989). A 17,000-year glacio-eustatic sea level record: Influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation. *Nature* 342, 637-642.
- Gaillard, M. J., and Moulin, B. (1989). New results on the Late-Glacial and environmental of the Lake Neuchatel (Switzerland). *Eclogae Geologicae Helvetiae* 82, 203-218.
- Gasse, F., Lédée, Massault, M., and Fontes, J.-C. (1989). Water-level fluctuations of Lake Tanganyika in phase with oceanic changes during the last glaciation and deglaciation. *Nature* 342, 57-59.
- Haberyan, K. A. and Hecky, R. E. (1987). The Late Pleistocene and Holocene stratigraphy and paleolimnology of Lakes Kivu and Tanganyika. *Palaeogeography, Palaeoclimatology, Palaeoecology* 61, 169-197.
- Hamilton, A. C. (1976). The significance of patterns of distribution shown by forest plants and animals in tropical Africa for the reconstruction of upper Pleistocene paleo-environments: a review. *Paleoecology of Africa* 9, 63-97.
- Hamilton, A. C. (1982). *Environmental History of East Africa: A Study of the Quaternary*, Academic Press, New York.
- Hecky, R. E., and Degens, E. T. (1973). Late Pleistocene-Holocene chemical stratigraphy and paleolimnology of the Rift Valley lakes of Central Africa *Technical Report* Woods Hole Oceanographic Institution, Cambridge, MA, 93 pp.
- Hemleben, C., Meischner, D., Zahn, R., Almogi-Labin, A., Erlenkeuser, H., and Hiller, B. (1996). Three hundred eighty thousand year long stable isotope and faunal records from the Red Sea: Influence of global sea level change on hydrography. *Paleoceanography* 11, 147-156.

- Hughen, K. A., Overpeck, J. T., Peterson, L. C., and Trumbore, S. (1996). Rapid climate changes in the tropical Atlantic region during the last deglaciation. *Nature* **380**, 51-54.
- Johnson, T. C., Scholz, C. A., Talbot, M. R., Kelts, K., Ricketts, R. D., Ngobi, G., Beuning, K., Ssemmanda, I., McGill, J. W. (1996). Late Pleistocene desiccation of Lake Victoria and rapid evolution of Cichlid fishes. *Science* **273**, 1091-1093.
- Kendall, R. L. (1969). An ecological history of the Lake Victoria basin. *Ecological Monographs* **39**, 121-176.
- Kneller, M., and Peteet, D. M. (1998). Late-glacial climate changes from a central Appalachians pollen and macrofossil record. *Quaternary Research*, in press.
- Kutzbach, J. E., and Street-Perrott, F. A. (1985). Milankovitch forcing of fluctuations in the level of tropical lakes from 18 to 0 kyr BP. *Nature* **317**, 130-134.
- Lehman, S. J. and Keigwin, L. D. (1992). Sudden changes in North Atlantic circulation during the last deglaciation. *Nature* **356**, 757-762.
- Lin, J. C.-F. (1996). "U-Th, ^{14}C and Sr Isotopic Studies of Late Pleistocene Hydrological Events in Western Great Basin, Nevada and California." Unpub. Ph.D. dissertation, Columbia University, New York.
- Lin, J. C., Broecker, W. S., Anderson, R. F., Hemming S., Rubenstone, J., and Bonani, G. (1996). New $^{230}\text{Th}/\text{U}$ and ^{14}C ages from Lake Lahontan carbonates, Nevada, USA, and a discussion of the origin of initial thorium. *Geochimica et Cosmochimica Acta* **60**, 2817-2832.
- Lin, J. C., Broecker, W. S., Hemming, S. R., Hajdas, R., Phillips, F., Smith, G. I., Kelley, M., and Bonani, G. (1997). Significance of U-Th and ^{14}C age determinations in Searles Lake sediments for late glacial high frequency hydrological events in the western Great Basin, USA. *Quaternary Research*, in press.
- Livingstone, D. A. (1975). Late Quaternary climate change in Africa. *Annual Review of Ecology and Systematics* **6**, 249-280.
- Lister, G. S. (1988). A 15,000 year isotopic record from Lake Zurich of deglaciation and climate change in Switzerland. *Quaternary Research* **29**, 129-141.

- Lotter, A. F. (1991). Absolute dating of the Late-Glacial period in Switzerland using annually laminated sediments. *Quaternary Research* 35, 321-330.
- Lowell, T. V., Heusser, C. J., Andersen, B. G., Moreno, P. I., Hauser, A., Heusser, L. E., Schlüchter, C., Marchant, D. R., Denton, G. H. (1995). Interhemispheric correlation of Late Pleistocene glacial events. *Science* 269, 1541-1549.
- Maenza-Gmelch, T. E. (1997a). Vegetation, climate, and fire during the late-glacial-Holocene transition at Spruce Pond, Hudson Highlands, southeastern New York, USA. *Journal of Quaternary Science* 12, 15-24.
- Maenza-Gmelch, T. E. (1997b). Late-glacial - early Holocene vegetation, climate, and fire at Southerland Pond, Hudson Highlands, southeastern New York, USA. *Canadian Journal of Botany* 75, 431-439.
- Morrison, M. E. S. (1968). Vegetation and climate in the uplands of southwestern Uganda during the Later Pleistocene period, 1. Muchoya Swamp, Kigezi District. *Journal of Ecology* 56, 363-384.
- Peteet, D. M., Vogel, J. S., Nelson, D. E., Southon, J. R., Nickman, J. R., and Heusser, L. E. (1990). Younger Dryas climatic reversal in northeastern USA? AMS ages for an old problem. *Quaternary Research* 33, 219-230.
- Peteet, D. M., Daniels, R. A., Heusser, L. E., Vogel, J. S., Southon, J. R., and Nelson, D. E. (1993). Late-glacial pollen, macrofossils, and fish remains in northeastern USA - the Younger Dryas oscillation. *Quaternary Science Reviews* 12, 597-612.
- Piper, B. S., Plinston, D. T., and Sutcliffe, J. V. (1986). The water balance of Lake Victoria. *Hydrological Sciences Journal* 31, 25-38.
- Roberts, N., Taleb, M., Barker, P., Damnati, B., Icole, M., and Williamson, D. (1993). Timing of the Younger Drya event in East Africa from lake-level changes. *Nature* 366, 146-148.
- Russel, I. C. (1885). "Quaternary History of Lake Lahontan, A Quaternary Lake of Northern Nevada" USGS Monograph 11.

- Schwalb, A., Lister, G. S., and Kelts, K. (1994). Ostracode carbonate DO18- and DC13-signatures of the hydrological climatic changes affecting Lake Neuchatel, Switzerland since the latest Pleistocene. *Journal of Paleolimnology* **11**, 3-19.
- Talbot, M. R., and Livingstone, D. A. (1989). Hydrogen index and carbon isotopes of Lacustrine organic matter as lake level indicators. *Palaeogeography, Palaeoclimatology, Palaeoecology* **70**, 121-137.
- Taylor, D. M. (1990). Late Quaternary pollen records from two Ugandan mires: evidence for environmental change in the Rukiga Highlands of southwest Uganda. *Palaeogeography, Palaeoclimatology, Palaeoecology* **80**, 282-300.
- Taylor, K. C., Hammer, C. U., Alley, R. B., Clausen, H. B., Dahl-Jensen, D., Gow, A. J., Gundestrup, N. S., Kipfstuhl, J., Moore, J. C., and Waddington, E. D. (1993). Electrical conductivity measurements from the GISP2 and GRIP Greenland ice cores. *Nature* **366**, 549-552.
- Watts, W. A., Allen, J. R. M., Huntly, B., and Friz, S. C. (1996). Vegetation history and climate of the last 15,000 years at Laghi di Monticchio, Southern Italy. *Quaternary Scientific Review* **15**, 113-132.
- Zolitschka, B. (1996). *Palaoklimatische Bedeutung Laminierter Sedimente*, Habilitationsschrift, Potsdam.




Fig. 1




Fig. 2




Fig. 3






Fig. 4




Fig. 5