

**Intercomparison of Global Precipitation Products: The Third Precipitation  
Intercomparison Project (PIP-3)**

**Robert F. Adler  
Laboratory for Atmospheres, NASA Goddard Space Flight Center,  
Greenbelt MD**

**Christopher Kidd  
Department of Geography, University of Birmingham**

**Grant Petty  
Department of Meteorology, University of Wisconsin**

**Mark Morrissey  
College of Geosciences, University of Oklahoma**

**H. Michael Goodman  
Global Hydrology and Climate Center, NASA Marshall Space Flight Center**

**(Submitted to Bulletin American Meteorological Society/September 2000)**

**Corresponding author:**

**Robert F. Adler  
Laboratory for Atmospheres  
Mail Code 912  
NASA Goddard Space Flight Center  
Greenbelt, MD 20771**

**Ph: 301-614-6290  
Email: Robert.Adler@gsfc.nasa.gov**

## **Abstract**

A set of global, monthly rainfall products has been intercompared to understand the quality and utility of the estimates. The products include 25 observational (satellite-based), four model and two climatological products. The results of the intercomparison indicate a very large range (factor of two or three) of values when all products are considered. The range of values is reduced considerably when the set of observational products is limited to those considered quasi-standard. The model products do significantly poorer in the tropics, but are competitive with satellite-based fields in mid-latitudes over land. Over ocean, products are compared to frequency of precipitation from ship observations. The evaluation of the observational products point to merged data products (including rain gauge information) as providing the overall best results.

## 1. Introduction

Over the past six years several intercomparisons of satellite precipitation algorithms, such as the first and second Precipitation Intercomparison Projects (PIP-1 and PIP-2), and AIP (Algorithm Intercomparison Project) -1, -2 and -3 have aided the development and use of global satellite precipitation products. A summary of results from the AIP program is given by Ebert et al. (1996). The PIP-1 project, which is closest in form to the current PIP-3 being discussed, is discussed in Barrett et al. (1994). The PIP-2 intercomparison which focused on instantaneous estimates based on passive microwave observations is summarized by Smith et al. (1998). The current PIP-3 follows the success of these previous efforts, but puts increased emphasis on evaluation of quasi-standard, satellite-based, global, monthly precipitation fields.

The PIP-3 project was sponsored by NASA through the WetNet Project and was endorsed by the Global Precipitation Climatology Project (GPCP) of the WCRP/GEWEX Program. This article summarizes the results of the PIP-3 Workshop, held at College Park, MD, and the pre-workshop and post-workshop analysis carried out with the submitted data sets. Sixty scientists attended the workshop representing numerous organizations involved in precipitation analysis from both an observational and modeling perspective. Additional information on the project and the workshop and detailed results can be found at the PIP-3 homepage address of <http://ghrc.msfc.nasa.gov/pip3>. A compact disc (CD) of the results, data sets, images, etc. is also available.

## 2. PIP-3 Objective and Approach

The objective of PIP-3 is to determine the utility of the current quasi-standard global, monthly precipitation products to the climate modeling and diagnostic community and the potential improvement expected with the latest satellite algorithms. The user community needs recommendations on accuracy and usefulness for a variety of applications including global model validation and climate monitoring and diagnostics. The algorithm community needs information on the future requirements of the user community. PIP-3 was designed to produce an evaluation of the current products and facilitate the exchange of information on future directions.

The monthly, global rain totals and rainfall frequencies submitted by the participants were evaluated against surface validation data sets, including an atoll gauge data set, ocean precipitation frequency, and land gauge data sets.

A full year (1992) was analyzed to test annual cycle retrieval. January and July of 1991, 1992, and 1993 were included to allow evaluation of interannual variations. August 1987 from the PIP-1 period was also evaluated to seek evidence of algorithm improvement during the past five years. Products using Special Sensor Microwave / Imager (SSM/I), geosynchronous infrared, Microwave Sounding Unit (MSU), TIROS Operational Vertical Sounder (TOVS) data, merged analysis schemes and composite microwave algorithms were included as well as prototype Tropical Rainfall Measuring Mission (TRMM) and Earth Observing System (EOS) microwave algorithms. Precipitation fields calculated from General Circulation Models (GCMs) were also included in the comparison.

The evaluation statistics were kept fairly simple and consist mainly of bias, root mean square error, and correlation versus the validation data. The satellite-based products were

examined with regard to their overall reasonableness (e.g., rainfall maxima in the right place and of reasonable intensity), freedom from artifacts (e.g., unnatural coastline precipitation features) and the statistical comparison to the validation data.

### 3. Description of Products and Validation Data Sets

Table 1 summarizes the thirty-one products. The twenty-five observational products, four model-based products and the two climatologies were intercompared with each other and the validation data sets in terms of monthly rainfall statistics during 1992, interannual variations among the January's and July's of 1991, 1992 and 1993 and the frequency of precipitation over the ocean (monthly and annual statistics).

The twenty-five observational products were divided into two groups for certain aspects of the analysis. The Quasi-Standard (Q-S) products were identified as those already in use by the modeling/diagnostic community, available for long, multi-year periods and readily available from archives, etc. These products tended to be the more mature products among the observational submissions. The remaining observational products were categorized as Experimental (EXP). All but one of the Experimental products were based on SSM/I data alone and typically were produced especially for the PIP-3 activity. Seven SSM/I-based products and one other product were ocean only estimates. Of the nine Quasi-Standard (Q-S) products, three were SSM/I-based, one each based on Microwave Sounding Unit (MSU) data, TIROS Operational Vertical Sounder (TOVS) data, and geosynchronous IR data and three were merged estimates using a combination of satellite observations or a combination of satellite and surface gauge observations.

The four model products included calculated fields from the reanalysis efforts at the European Centre for Medium-range Weather Forecasting (ECMWF), the National Centers for Environmental Prediction (NCEP) and the NASA Goddard Space Flight Center (GSFC) and a climatological average of the Atmospheric Model Intercomparison Project (AMIP) climate models (Lau et al.,1996).

The descriptions of the submitted products can be found on the noted web site or project CD, or through the references in Table 1.

The validation data set used in the PIP-3 study was accumulated from a number of sources.

Land areas:

i) the Global Precipitation Climatology Centre (GPCC) gauge product (Rudolf, 1994) was taken as the base validation data set. The raw data product (as opposed to the climatological-corrected product) was used. However, there were some notable areas that have little data. Outside Western Europe the number of gauges is sparse, even in countries such as the US and Australia where gauge coverage is known to be good.

ii) the Surface Reference Data Centre (SRDC) (Huffman et al., 1997) data set was seen as the most accurate of the data sets. Each 2.5 degree box used numerous gauges to generate the rainfall estimate. Unfortunately, it was also the least comprehensive, being only available for a few 2.5 degree boxes.

iii) supplementary gauge data was sought for USA, Australia and South Africa to fill in some of the voids in the GPCC data set. The gauge data from these areas was interpolated and mapped to the 2.5 degree resolution. In addition, to boost the number of gauges in the

tropical region, data from the Amazon region was incorporated from the Amazon River Basin Precipitation data set at the Oak Ridge National Laboratory. The land gauge data sets were merged on the basis of the SRDC product having top priority, followed by the supplementary gauge data where the number of gauges exceeded that of the GPCC data, and lastly, the GPCC data set.

A subset of this data base was chosen for the validation of the algorithm products in order to achieve a representative geographic data set. The selection was based upon the number of gauges available per 2.5 degree box, by the number of boxes within each climatic region, and the proximity to other boxes: Figure 1 shows the distribution of the validation boxes chosen.

The interannual validation data set was based upon selected areas of four contiguous 2.5 x 2.5 degree boxes in order to reduce the noise from both the validation and algorithm data. Areas were chosen as representative samples of the different climatic regimes. These areas can be seen in Fig. 1 as the three groups of four outlined boxes in the U.S. and Australia.

#### Ocean:

Oceanic validation data, especially gauge data, is very limited. For the PIP-3 study, atoll raingauge data from the Comprehensive Pacific Rainfall Data Base (CPRDB; Morrissey et al., 1995) were used. Data from the atolls were collected, quality controlled and mapped by the Environmental Verification and Analysis Center (EVAC) at the University of Oklahoma. The data were grouped into three regions reflecting the seasonal characteristics of the rainfall data, namely north of 5° N, 10° S - 5° N and south of 10° S. Figure 1

shows the locations of the atoll validation grid boxes. For the interannual comparisons the sum of the 10° S - 5° N boxes was used (as shown by the outline in Fig. 1).

#### Frequency of Precipitation:

Data from the Comprehensive Ocean-Atmosphere Data Set (COADS; see Petty 1995) was used as validation data over the oceans, and was prepared by one of the authors (Petty). The COADS data set, comprised of ship observations of present and past weather, were used to determine the occurrence, or frequency, of precipitation. Due to the sparse nature of the observations in certain parts of the globe, data from the period 1958 to 1991 were used to generate an average, and therefore should be treated as a climatological average of the frequency of precipitation. The fractional-time-precipitating was derived from ship reports falling within a latitude-longitude window centered on the grid box in question. The dimensions of the window were chosen so as to achieve an adequate statistical sample without unnecessarily smoothing real gradients in rainfall distributions. Two sets of validation data were generated, one using all the COADS data with observations reporting all precipitation, except drizzle, and another set reporting all precipitation except drizzle and snow. The latter data set was included on the basis that estimates of precipitation from the passive microwave sensor would not include drizzle and snowfall.

#### 4. Intercomparison of Monthly Rainfall Totals

The global, monthly rainfall total maps for 1992 were examined and intercompared in a number of ways and against the validation data sets over the Western Pacific Ocean atolls and over land. They were also examined for artifacts and for reasonableness over areas where no validation data sets exist, for example in the mid-latitude oceans. Four examples (of the different product types) of monthly maps for July 1992 are shown in Fig. 2. All



four examples display the main features of a July precipitation map. The Inter-Tropical Convergence Zone (ITCZ) stretches across the Pacific and Atlantic Oceans and northern South America. The Asian summer monsoon is producing rainfall maxima over India, Indochina and adjoining water areas. Northern Australia is in its dry season. In the tropics, the four example maps show very similar patterns and similar magnitudes. In mid-latitudes oceanic maxima are evident, with varying intensities. For example, the three maxima (the top, right panel (model)) at approximately 40°S east of Africa, in the central Pacific and east of South America are evident in the experimental and quasi-standard examples, but with different magnitudes. The noisiness of the Experimental product is due to the limited sampling with the low-orbit satellite. A similar noisy pattern is evident in the quasi-standard example at ocean latitudes above 40°.

#### 4.1 Zonal annual totals over water and land

Zonal averages of the annual (1992) total over the ocean of each of the 25 observational products indicate a wide variation among the products, both in the tropics and in middle and high latitudes. The mean value and range of values at each latitude are plotted in Fig. 3a. All the products generally capture the tropical maximum, the sub-tropical minima and the mid-latitude maxima. However, among all the observational products the peak value in the Inter-Tropical Convergence Zone (ITCZ) at 8°N varies from 1300 to 3200 mm for the annual total. This large variability among the estimates is also evident in mid-latitudes with values ranging from 900 to 1800 mm in the Northern Hemisphere maximum, with additional outliers above and below those values. In very high latitudes, for example at 60°S, the range becomes an order of magnitude, going from 100 to 1000 mm.

At first this large variability among the estimates is disconcerting. However, if the set of products is limited to the Quasi-Standard data products, the range of values decreases significantly. This effect can be seen in Fig. 3b, which shows the standard deviation among the estimates as a function of latitude for both the 25 observational estimates and the subset of eight Q-S products. This decrease in the variability as we go from all to the Quasi-standard products mainly reflects the maturity of the products. In addition, there is some interdependence among the Q-S products because of merged products using some of the same input fields. Many of the products in the Experimental group were based on early versions of retrieval algorithms and due to errors, or perhaps a lack of tuning, some of these products produced values outside the range of reasonableness. These facts point to the need for the user community to exercise caution in selecting products with which to work.

Fig. 4a compares the average of all the observational products with the two climatologies. The tropical peak in the Legates/Wilmott climatology is significantly larger than the observational products or the Jaeger climatology. In fact, the zonal totals also indicate that the Legates/Wilmott climatology has higher values in the ITCZ as compared to all the Q-S products. This difference is mainly related to the large peak found in the climatology in the east-central Pacific Ocean in the ITCZ during the Northern Hemisphere summer. None of the observational products support the existence of this feature, although they are looking at only one year of data. In the dry, subtropic zone in the Southern Hemisphere oceans the Legates/Wilmott (L/W) climatology also carries significantly higher values than all the observational products. The difference here is due mainly to the lesser westward extent from the South American and African coasts of the subtropic minima in the climatology as compared to the satellite estimates. In mid-latitudes (poleward of  $40^{\circ}$ ) the mean of all the observational products is significantly less than the climatologies. The observational mean

is dominated by SSM/I-based products which seem to have a tendency for underestimation at these latitudes, as is discussed in a later section.

Still higher than the L/W climatology in the ITCZ is the ECMWF reanalysis model result with a peak annual total of 3100 mm (Fig. 4b). Between 0° and 10°S the ECMWF model result is significantly larger than the Q-S products, being 2500 mm and nearly double the mean of the Q-S products at 5°S. This difference in zonal total is related to very strong precipitation features in the ECMWF calculations in the central Pacific Ocean and over the Indian Ocean. The satellite estimates do not support the magnitude of the maxima, especially in the Indian Ocean. In mid-latitudes the models generally agree with the shape and magnitude of the climatologies and the mean of the Q-S products.

The zonally-averaged totals over land show similar results with the tropical peak at the Equator ranging from 1300 to 2800 mm, but with a smaller range when only “mature” products were considered. Products with errors related to misidentifying desert surface as rain and products with other artifacts were identified with the zonal annual averages.

#### 4.2 Validation of estimates with Western Pacific Ocean atoll data set

The Western Pacific Ocean atoll rainfall data set (Morrissey, et al.1995) was used to compare with the monthly rainfall totals from the products. Fig. 5 displays bar graphs of the statistical results for each of the products using monthly totals for each of the months of 1992. Of the twenty-five observational products, nineteen had a negative bias compared to the atoll data. When limited to the Quasi-standard products, the data set had eight out of nine products having a negative bias, with the ratio being 0.88. This result is in contrast to that of a previous intercomparison (AIP-3) in the Western Pacific Ocean which used

surface-based radar data from the TOGA-COARE program as its validation (Ebert, 1998). There a very large majority of products (including many of the same retrieval schemes represented here) had a positive bias. Although these two validation data sets are in different locations and use different measurement methodologies, this difference needs to be addressed before we can be confident of the absolute magnitude of the rainfall estimates over the tropical oceans.

As a group, the Q-S products have a higher correlation, a lower root mean square error (RMSE) and a smaller bias than the Experimental products. The models as a group have a reasonable bias, but very low correlations, indicating that they are less accurate than the observations in portraying the spatial and temporal variations of the monthly fields on these scales over this portion of the tropical ocean. Of the standard products the NOAA Merged Product (*nmg*) has the best statistics; however, it uses the atoll information in its merger process, so these statistics may not reflect what would be the results in other locations where there are no atolls. The venerable *gpi* has the highest correlation of any satellite-information-only product, reemphasizing the importance of sampling. Although the *gpi* also has a very small bias error against the atolls, the zonally-averaged, oceanic results again point to its limitations in sub-tropical regions (a large positive bias), as is well know to its users. Two of the Q-S products, the *tov* and *nmi* have relatively large biases.

Of the Experimental products, the OLR Precipitation Index (*opi*) also exhibits very good statistics, although it has been derived by correlating against the *nmg* product where gauges are used. These validation statistics may therefore not be indicative of the accuracy of the product in locations without gauges. Of the SSM/I-based products (Experimental or Quasi-Standard) the *buc* and *gem* have the lowest RMSE, with a number of other retrievals close behind. The SSM/I-based products seem to have an upper limit to the correlation of about 0.75 because of the sampling limitation of the polar orbit, even though

almost all of the SSM/I products used data from two polar orbiting satellites, therefore maximizing the sampling.

#### 4.3 Validation of estimates over land

Validation over land for the months of 1992 was carried out using the raingauge data sets previously described. Products that directly incorporated raingauge information (*gpm* and *nmg*) performed best statistically among all the products. Even the Experimental (EXP) product *opi*, which is derived through comparison with the *nmg* provides very good statistics. However, caution must be used in evaluation of these statistics. Because we are validating the products in locations where the raingauge information is of high quality and plentiful, these statistics may be overly optimistic as to how these products perform in general, especially in areas of poor raingauge coverage or quality. One example of the impact of the raingauge data can be seen by comparing the *gps* and *gpm* products. The *gps* is a merged satellite data product, whereas *gpm* additionally incorporates the gauges. In the tropical belt (30 ° N - 30° S) the *rms* drops from 56 to 30 mm when gauges are added, nearly a 50% reduction.

The comparison of combined 12 month statistics also indicates that the climatologies are competitive with the satellite-based and model products. This result is related to the use of only one calendar year for this comparison and the large variance in the rainfall data set related to the climatological spatial and seasonal patterns. Only the products incorporating the raingauge data for the particular year had significantly better validation statistics than the best of the climatologies.

In the tropics over land (30°N - 30°S) the Q-S products and the climatologies as groups have the best RMSE, with the models generally having a positive bias and a larger RMSE (see Fig. 6). Of the observational products without influence from gauge information, the *gps* has the lowest RMSE. Over land the *gps* is a merger of the *nmi* and *gpi* and gives an RMSE of 58 mm as compared to the RMSE of 91 mm for the *nmi* and 77 mm for the *gpi*. In this region five observational products not affected by gauge information have ratios between 0.80-1.20 and correlations greater than 0.75. These are *bup*, *cuf*, *cul*, *pur*, *plc*, *gps*. Of the SSM/I products the PIP-1 composite (*plc*) has the lowest RMSE of 61 mm.

In mid-latitudes (30°N - 60°N) the models and even the climatologies outperformed the observational products, except for those using raingauge information (see Fig. 7). The models had small biases and high correlations, with the ECMWF model (*ecm*) having the best RMSE (31 mm) of all products not using gauge information. In this region, when the same ratio and correlation criteria as in the previous paragraph are used, no non-gauge products meet the criteria. If the criteria are loosened, to 0.75-1.25 for the ratio and greater than 0.65 for the correlation, then *buc* and *bus* emerge.

When the latitude boundaries are expanded to encompass the entire 60°N-60°S region (not shown) the *buc*, *bup*, *pur*, *gps* and *plc* products meet the original criteria. It should be remembered that the SSM/I-based products do not typically have estimates over portions of the winter hemispheres over land due to cold land/snow contamination and that the statistics shown are for a matched set, using only those points where all the algorithms produced estimates.

In a footnote related to sampling the relatively poor showing by the *nmi* product in the tropics over land (RMSE of 91 mm) is due mainly to the use of only one satellite in this Q-

S product. The other SSM/I-based products used two satellites in producing their 1992 estimates. When *nmi* was run with two satellites after the workshop the RMSE for the tropical land area was reduced to 71 mm, which is comparable with most of the other SSM/I products. This very significant difference emphasizes the importance of sampling in the production of monthly precipitation estimates.

In general, these statistical results for 1992 over land indicate that in the tropics a number of observational algorithms produce good results, but that over mid-latitude land results degrade, both absolutely and relative to the model calculations. This is especially true for the microwave-based products. The addition of gauge information greatly increases the accuracy of the products.

#### 5. Interannual results

An evaluation of product performance related to the estimation of inter-annual variations was performed using the Januarys and Julys of 1991, 1992 and 1993. However, due to the near total absence of SSM/I data from January 1991, the difference fields between January 1991 and January 1992 were eliminated from the statistical comparison, leaving one January difference field and two July difference fields. The SSM/I products used for the interannual comparisons were produced using data from only one DMSP satellite. Example interannual difference maps between January 1992 and January 1993 are shown in Fig. 8. The strong El Niño in 1992 produced a strong mid-Pacific Ocean maxima in January 1992 and that is reflected in the interannual difference fields. All four example fields have very similar patterns over the tropical Pacific Ocean and to a lesser degree elsewhere in Fig. 8. Subtle differences are evident over the Indian Ocean and in mid-latitudes.

Statistical results over the Western Pacific Ocean region using the atoll data set indicate that as a group the Q-S products performed well with correlations in the 0.75-0.80 range and relatively low RMSE, with only the PIP-1 composite (*p1c*) doing poorly (see Fig. 9). This low correlation for the *p1c* product may be due more to the one satellite sampling in the interannual exercise. The other Q-S products either have the better sampling related to the geosynchronous satellites or multiple polar-orbiters, such as with the *tov* and *msu* products. Among the EXP products the results are very variable with the *opi* and a few of the SSM/I products having reasonable correlations, approaching or equaling the validation statistics of the Q-S products. However, those SSM/I products that have relatively good interannual statistics with the atoll data are generally not among the best on the monthly statistics discussed earlier. Thus, taking into account both the annual and interannual statistics over the atolls, there is no clearly superior product among the SSM/I-based entrants.

The models did poorer than the observational products as a group in terms of the correlation (0.4-0.6), as expected, with the *ecm* (ECMWF) results having the highest model correlation. The *ecm* also has the highest rmse, indicating that this product is not reproducing the magnitude of the interannual differences. In the 1992 monthly statistics of the previous section, *ecm* was roughly comparable or a little worse than the other models. Thus, combining the information in both the monthly and interannual statistics leads again to the fact of no clear distinction among the model products.

Over land areas in the Tropics (30°N-30°S) the products directly using gauges easily do best (see Fig. 10). Again it should be stressed that these statistics are for areas where the validation data, and therefore the input data to these products are the best. Again the Q-S products generally provide the best answers with better correlations and RMSE's than the experimental products or models. Among the Q-S products that do not include gauge



information, the *gpi* has a high correlation, but an RMSE about the same as *gps* and *tov*. The PIP-1 composite product (*plc*) does relatively poorly in terms of both correlation and RMSE. In fact all the SSM/I-based products have poorer statistical results than the Q-S products (other than *plc*). The *opi* product has interannual statistics in this region roughly equivalent to the satellite-only Q-S products, a drop-off compared to its better relative statistics on the monthly rain totals. The model products have lower correlations and higher RMSE's than most of the observational products.

In mid-latitudes (30°-60°N) over land the interannual statistics (Fig. 11) indicate the models approximately matching the quality of the observational estimates, with the exception of those that include the raingauge data. The median model correlation of 0.65 is equal to or better than all the values for the SSM/I-based products and approximately matches that of the non-gauge Q-S products and the *opi*. This interannual result is similar to that found with the monthly statistics in mid-latitudes.

Figure 12 shows examples of interannual change during July in four example areas. Examples from January are not used because of the lack of SSM/I data in January 1991. In this exercise data from 4-6 adjacent or nearby boxes were combined to reduce the algorithm and validation random error. In Fig 12a the three year July variation over the six boxes of the atoll data set between 5°N and 10°S is shown for the pertinent products that produced results for two or three of the months. Most of the observational products reproduce the interannual changes in the atoll gauges, especially the relatively large change from July 1992 to July 1993, which is partially related to the end of the 1992 ENSO event. The three model calculations do not fare as well, although the *ecm* reproduces the tendencies correctly, but not the absolute values of the precipitation.

Two nearby areas over land in summer are shown in Figs. 12b and 12c. In Fig. 12b results for a 5° by 5° latitude-longitude area (37.5°-42.5°N, 87.5°-92.5°W) centered over the state of Missouri in the United States show a large increase in rain leading up to the heavy flooding in 1993 with mean July rainfall in the area of over 300mm. The observational products as a whole do not reproduce the interannual change correctly. The microwave-based products all overestimate the magnitude of the change, sometimes very substantially. This may be due to the microwave algorithms mis-identifying wet ground or standing water as falling precipitation. Infrared-based products, such as *gpi* and *opi* do better than the microwave products, but seem to saturate, and do not identify the increase from 1992 to 1993. The three models, on the other hand, reproduce the raingauge results in this location very well. A different pattern of interannual changes is shown in Fig. 12c for an area to the southeast covering part of the state of Mississippi (32.5°-37.5°N, 85°-90°W). With much smaller magnitudes (note the scale difference in the diagrams) and a sharp decrease between 1992 and 1993, the observational products overall do a very good job in this area, while two of the models indicate poor results.

Over southeast Australia (32.5°-37.5°N, 145°-150°E) in the cool season (July), Fig. 12d indicates that most of the satellite products capture the interannual variation qualitatively, but a number underestimate the magnitude of the precipitation. The *gpi* matches the validation nearly perfectly, while the *opi* seems to underestimate the magnitude of the changes. The models also capture the time change, but underestimate the precipitation.

The four examples shown are not all encompassing. However, the results indicate that both for the observational products and the models as groups, and for individual products, the interannual results are very variable in quality and therefore these products should be

used with caution when assessing interannual change. The results over land also indicate that including gauge information, where available, is critical.

## 6. Frequency of oceanic precipitation

Validation of monthly rainfall totals over oceans is limited to the atoll data set as representative of the open ocean. This limits the geographic scope of a validation exercise to a portion of the Western Pacific Ocean. In order to obtain measures of how well the precipitation products in this intercomparison were reproducing the observed precipitation distributions over the oceans generally; the product producers were asked to provide estimates of precipitation frequency in addition to monthly rainfall totals. These estimates could then be compared to the precipitation frequencies derived by Petty (1995). Examples of maps of the annual frequency of precipitation from the submitted products and from the COADS data are shown in Fig. 13. The upper left panel in Fig. 13 shows the COADS climatology from Petty (1995). The main tropical and extratropical maxima and minima are evident with some boxes in the central Pacific ITCZ reaching 15-18%. Elsewhere in the tropics there are significant areas with rain frequencies of 8-10% in the climatology. In mid-latitudes of the Northern Hemisphere, the COADS data indicate maxima of 8-10%, increasing to 14% at 60°N in the Atlantic. The two experimental observational products shown reproduce the main features with significantly different magnitudes (from each other) in both the tropics and mid-latitudes. For example, in the central North Pacific one technique shows 2-4%, while the other product has 8-10% over a large area. This difference is typical of differences shown among the SSM/I-based products. The example model product in the lower, right panel of Fig. 13 has very large frequencies in the tropics and frequencies in mid-latitudes more comparable to the validation data and the other products shown. This difference in the model product between tropics and mid-latitudes is partially due to the convective nature of the precipitation in the tropics compared to the

widespread precipitation in mid-latitudes. This difference and the relatively coarse mesh of the model may help produce the initial result in Fig. 13.

The frequency of precipitation comparison was very useful in delineating the reasonableness of the precipitation patterns and in some cases the accuracy of the products. However, the quantitative usefulness of the comparison is muted by the sensitivity of the results to the rainfall rate threshold used in defining the rainfall frequency both in the satellite and model products and the COADS surface observations. For this study the frequency of precipitation from the COADS data was defined as light precipitation and heavier (eliminating drizzle) at the station and at the time of observation. The product producers were advised to use 0.5 mm/h as a threshold, if they had that flexibility in their product.

The elimination of drizzle from the COADS frequency of precipitation values may produce an underestimate of the actual precipitation frequency, when compared to the threshold of 0.5 mm/h, especially in middle and high latitudes. This can be seen in Fig. 14 where a scatter plot of human observations of precipitation (excluding drizzle) are compared with raingauge measurements of 0.5 mm/h or greater over the United Kingdom. The results indicate a bias of approximately 3%, with the human observation (equivalent to COADS) higher than the raingauge values (presumably comparable to the precipitation products in the intercomparison). Use of a raingauge threshold of 0.2 mm/h eliminates most of the bias. Therefore, this sensitivity of the results to the rainrate threshold and the magnitude of the bias should be kept in mind when evaluating the following results.

The results indicate a very large variability among all the products in the general oceanic precipitation frequency. Some of the large variation stems from various satellite and model footprint (or grid resolution) sizes and in different precipitation rate thresholds used. The

most homogeneous data set is that coming from the SSM/I-based algorithms, although among this subset there was still considerable variation (see Fig. 15). The zonally-averaged, annual (for 1992) peak in the ITCZ shows values ranging from 4% to 13% among the SSM/I products, with the COADS climatology indicating 8%. The SSM/I results are approximately equally distributed above and below the COADS value. In the Northern Hemisphere sub-tropical minimum the COADS climatology shows 3%, while the majority of SSM/I values are lower. Moving poleward from 30°N the COADS values increase continuously from 4% at 30N to 11% at 60°N. Many of the SSM/I algorithms produce lower estimates of the precipitation frequency throughout this zone, although a few compare favorably with the COADS up to 45N, where they peak and then decrease rapidly, producing a very pronounced underestimation. The possible ~3% high bias in the COADS estimates would reduce the difference, but the latitudinal profile would still not be similar. This middle and high latitude underestimate is probably related to a failure to detect frozen precipitation and the relatively light rain at these latitudes as can be inferred by examining the second, lower COADS curve in high latitudes which excludes frozen precipitation. A few of the SSM/I products (*rss*, *pur*, *gpf*) do reasonably well up to 50°N in this comparison.

Among the non-SSM/I satellite products there is a general overestimate of the precipitation frequency. Both the *tov* and *msu* products have an ITCZ zonally-averaged annual peak of about 30%, a Northern Hemisphere subtropical minimum of 8% and 45N value of 18%, all more than double the COADS values. On the other hand, both of these products show an underestimate in total rainfall in relation to the atoll data, indicating a relatively small rainrate in the raining areas of these products. These large values of rainfall frequency may partially be related to the larger footprint (relative to SSM/I) of *tov* and *msu* retrievals and/or the threshold used by the product producers. The *gpi* also has high values of

frequency, especially in the ITCZ, because of the use of an infrared  $T_b$  threshold of 235K, which gives the best correlation with rainfall occurrence over large areas and long times, but produces a “cold cloud” area usually much larger than the rain area.

Two of the model-based products (*geo* and *nep*) submitted frequency of precipitation information. The zonal-averaged precipitation frequencies from these two models are given in Fig. 16, along with the COADS estimates. The model estimates represent the occurrence of precipitation somewhere in the relatively (to SSM/I observations) large grid box, so that the model estimates might be expected to be an overestimate compared to the COADS frequency. The model-based precipitation frequencies exhibit very distinct differences from the COADS numbers and the satellite estimates. The two models both have significant positive biases in the tropical rain areas and negative biases in mid-latitudes. The zonally-averaged, annual peak in the N.H. ITCZ for both the models is about twice the COADS value of 8%. In the S.H. there is a secondary peak at approximately 5°S with the *geo* frequency again being about twice the COADS value (5%); however the *nep* value is significantly higher (15%), about three times the COADS value. This large overestimate in the *nep* values, relative to both the COADS and the *geo* values is related to an eastward extension of the South Pacific Convergence Zone (SPCZ) into the central South Pacific Ocean and a western extension of the Indian Ocean maximum toward Africa that do not exist in either the COADS climatology or the satellite frequency estimates. The *geo* model product has a very weak eastern Pacific maximum, while the *nep* nearly fails to produce the Atlantic ITCZ.

In terms of geographic distribution of the frequency features most of the SSM/I-based products reproduce the locations of the maxima and their relative magnitudes (see Fig. 13). The western and eastern Pacific Ocean features are reproduced, as well as their seasonal movements. The advance and retreat of the Asian monsoon can be easily traced in the

rainfall frequency diagrams. In mid-latitudes the feature agreement is somewhat worse due to the apparent inability of the algorithms to capture the increase in precipitation frequency with latitude, although even somewhat subtle features, such as the maximum off the west coast of Canada and Alaska in winter is clearly defined.

In terms of seasonal variation in the 30-45N region the COADS data show a distinct wintertime maximum that is only clearly reproduced in a few of the SSM/I-based products, namely *pur* and *tam* (see Fig. 18). Some other products fail to clearly retrieve this seasonal variation, or in a few cases even place the maximum in the summer. In the Southern Hemisphere mid-latitudes the annual frequency map from the COADS data show rainfall frequency maxima located southeast of Africa and South America and in the mid-Pacific Ocean where COADS data is sufficient for analysis. The satellite-based products that best reproduce both the location of the features and the approximate magnitude of the features are *gpf*, *pur* and *rss*. The *gme* product also reproduces the location well, but greatly overestimates the magnitude.

For seasonal variation both model products capture the annual variation in mid-latitudes with a peak precipitation frequency in the winter of the respective hemispheres, although both consistently underestimate the frequency. In the tropics the models show the general overestimate, as stated before, but the *geo* product has a small overestimate in the N.H. winter, but a large overestimate in the N.H. summer, primarily related to an apparent large overestimate of precipitation in the Western Pacific Ocean. The twelve-month correlation statistics, which indicate how well the products are delineating the spatial and seasonal variations over the ocean, show that in general the models have much lower correlations than the observational products in the tropics, but have correlations better than many, but not all of the satellite products in the 30-60N zone. Between the two models the

correlations show approximate equality in the mid-latitude zone, with the *ncp* having higher correlations in the tropics.

The quantitative comparison of the SSM/I-based precipitation frequency products with the COADS data indicate some clear differentiation among the products. An examination of the annual average frequencies in the tropics (30°N-30°S) finds eight products (*cuf*, *cul*, *dlr*, *gpf*, *iow*, *rss*, *nmi*, *tam*) that have a bias ratio of between 0.70-1.30 and a correlation at or above 0.8. Expanding the zone to 45°N and 45°S reduces the list to *gpf*, *pur*, *nmi.*, *rss*, and *tam* . A table of results for the 45°N-45°S region is given in Fig. 17. Again expanding the area to 60N-60S, one is forced to loosen the requirements for the correlation to 0.70. Again the five products meeting these criteria are *gpf*, *pur*, *nmi.*, *rss*, and *tam*, with the highest correlation (0.81) held by *pur*. One must remember that even as we expand the latitudinal zone the preponderance of the data set is in the tropics. Restricting the latitude band to 30-60N to examine the mid-latitude numbers, we find the highest correlation belongs to *pur*.

Similar results, but with much poorer correlation coefficients, is evident if we use the monthly statistics instead of the annual means described in the last paragraph. However, as mentioned previously most products did not reproduce the seasonal cycle in the 30-60°N band and this shows up clearly in the correlation statistics, where *pur* had a significantly higher correlation than any other SSM/I-based product.

Among the non-SSM/I products, the *msu* does well in terms of correlation coefficient in the tropics (30°N-30°S), equaling the best SSM/I value there. However, in the expanded latitude zones it loses significant ground to the SSM/I values. The *tov* correlations are



lower than the leading SSM/I-based values in all latitude zones. The *gpi*, surprisingly, also does poorly compared to almost all the SSM/I values in the 30°N-30°S zone, with a correlation of 0.60 for the monthly statistics.

## 7. General Conclusions

The intercomparison of the many observational and model-based precipitation products in this effort results in a plethora of images and statistics. However, because of the limitations of the validation data in terms of both coverage and quality and because of products often performing well in only certain locations or situations, there is great difficulty in unambiguously pointing to a certain product as “best” in terms of a monthly precipitation total over most of the globe. However, general conclusions can be drawn from the intercomparison results and recommendations to both the producer and user communities can be made.

The intercomparison reveals a very large range of estimates among the products. Even the zonally-averaged, annual total field has a factor of two to three between the smallest and largest values, depending on the latitude. The range of values is reduced considerably when the observational products are limited to those from the Quasi-Standard set. This generally better performance by the Q-S products is also evident in both the atoll and land monthly validation and even the inter-annual results. The model-based precipitation fields do significantly poorer than the observational fields in the Tropics, but are competitive with the satellite-based fields in mid-latitudes over land. The inter-annual statistics do not necessarily follow the monthly statistics in terms of which products perform well, at least among the SSM/I-based products. The frequency of precipitation intercomparison was

very worthwhile in terms of gaining a better picture of how oceanic algorithms performed over a wide range of climatological zones. In terms of SSM/I-based products, the COADS-based frequency information clearly helps to diagnose the performance of the products.

Some general conclusions from the intercomparison are as follows. This intercomparison clearly establishes the value of the merged analysis products that incorporate information from two or more satellite sources and blend in the raingauge data. Over land these products demonstrate superior statistics because of the incorporation of the gauge data. However, because the over-land evaluation centers on areas of plentiful gauge data, further evaluation of the products is needed using data exclusion tests to determine the error characteristics in areas of little or no gauge information. Even without the gauges there is evidence that the merger of microwave and geosynchronous IR data produce a better product. A second general conclusion is that the Quasi-Standard products generally outperform the Experimental group as a whole. This result is closely related to the maturity of the products. The Q-S products are, for the most part, products that have undergone substantial testing over a period of time, including in other intercomparisons. Many of the EXP products were being produced for the first time and will no doubt perform better in the future.

Results from the Tropical Rain Measuring Mission (TRMM), launched in late 1997, should be a major source of new information on tropical rain totals, structure of tropical rain systems and how current rain estimation techniques compare with the improved information from TRMM.

## 8. Recommendations to the User and Producer Communities

The PIP-3 Workshop and the related intercomparison activities resulted in an excellent review of the status of global precipitation analysis on a monthly time scale and, as always, raised a number of questions as to the large variability and reliability of estimates in certain regions and the validity of the validation or comparison *in-situ* data sets. The workshop and associated analysis did arrive at some general conclusions which are related to the following recommendations to the user and producer communities.

Recommendations:

- 1) For the period 1987-present, the user community should focus their use of global, monthly precipitation fields on the merged analysis products combining information from SSM/I data, geostationary IR data and raingauge data. Users should note possible limitations in these products, for example, a high latitude low bias over oceans (related to SSM/I estimates) and some observed artifacts in interannual fields. The over ocean values in these products are critically dependent on the SSM/I-based estimates, thereby emphasizing the need to use accurate, validated frequency of rain and rain amount information from the SSM/I algorithms.
- 2) The apparent high latitude ocean low bias in most (not all) SSM/I precipitation total and frequency fields should be a focus of research attention with the objective of development of an approach which agrees more closely with the presumed correct precipitation frequency climatologies. Mechanisms for development of validation data sets in middle and high latitude oceans should also be pursued.
- 3) Continued research and analysis should be done on the use of MSU, TOVS, OPI and other data sources for potential use in global precipitation analysis in the 1979-1987 period before SSM/I.

4) A product or products providing information on the variability among all (or a subset) of the observational products should be developed and analyzed to judge its utility as a measure of the unanimity of our estimates as a function of location and season. In the absence of high quality validation data this approach would give indications where estimates agreed and disagreed.

5) The user community requires global precipitation fields at finer space and time scales for diagnostics, model validation (including diurnal variations) and assimilation into models. Requirements: ~6 hours, 100 km for global coverage.

6) For future intercomparisons all products that include gauges explicitly should have a non-gauge version in order to better intercompare satellite-only products. In addition frequency of precipitation should be included along with precipitation total with each submitted product so that the COADS frequency information may be used for direct comparison with all the products.

#### Acknowledgements:

The authors wish to thank the following for providing special validation data sets: Beth Ebert of Bureau of Meteorology, Australia; Johan Koch, National Weather Service, South PAfrica; and Oak Ridge National Laboratory. The analysis was supported by NASA HQ (James Dodge). The authors especially want to thank the contributors for all their efforts and inputs to the intercomparison effort.

References:

- Adler, R. F., G. J. Huffman, and P. R. Keehn, 1994: Global tropical rain estimates from microwave-adjusted geosynchronous IR data. *Remote Sensing Reviews*, **11**, pp. 125-152.
- Arkin, P. A. and B. N. Meisner, 1987: The relationship between large-scale convective rainfall and cold cloud over the Western Hemisphere during 1982-1984. *Mon. Wea. Rev.*, **115**, pp. 51-74.
- Barrett, E. C., R. F. Adler, K. Arpe, P. Bauer, W. Berg, A. Chang, R. Ferraro, J. Ferriday, S. Goodman, Y. Hong, J. Janowiak, C. Kidd, D. Kniveton, M. Morrissey, W. Olson, G. Petty, B. Rudolf, A. Shibata, E. Smith, and R. Spencer, 1994: The first WETNET precipitation intercomparison project: interpretation of results. *Remote Sensing Reviews*, **11**, pp. 303-373.
- Bauer, P. and Schuessel, P., 1993: Rainfall, total water, ice water and water-vapor over sea from polarized microwave simulations and SSM/I data. *Journal of Geophysical Research*, **98**(D11), pp. 20737-20759.
- Chiu, L., A. Chang and J. Janowiak, 1993: Comparison of monthly rain rates derived from GPI and SSM/I using probability distribution functions. *J. Appl. Meteor.*, **32**, pp. 323-334.
- Ebert, E. E., 1996: *Results of the 3<sup>rd</sup> Algorithm Intercomparison Project (AIP-3) of the Global Precipitation Climatology Project (GPCP), Revision 1*. Bureau of Meteorology Research Centre, Melbourne, Australia, 199 pp.

Ebert, Elizabeth E., Michael J. Manton, 1998: Performance of Satellite Rainfall Estimation Algorithms during TOGA COARE. *Journal of the Atmospheric Sciences*: **55**(9), pp. 1537–1557.

Ferraro, R. R. and G. F. Marks, 1995: The development of SSM/I rain rate retrieval algorithms using ground based radar measurements. *J. Atmos. Oceanic Technol.*, **12**, pp. 755-770.

Haferman, J. L., Anagnostou, E., Tsintikidis, D., Krajewski, W. F., and Smith, T. F., 1996: Physically Based Satellite Retrieval of Precipitation using a 3D Passive Microwave Radiative Transfer Algorithm, *Journal of Atmospheric and Oceanic Technology*, **13**, pp. 832-850.

Huffman, G. J., R. F. Adler, P. Arkin, A. Chang, R. Ferraro, A. Gruber, J. Janowiak, A. McNab, B. Rudolf, and U. Schneider, 1997: The Global Precipitation Climatology Project (GPCP) Combined Precipitation Data Set. *Bull. Amer. Meteor. Soc.*, **78**(1), 5-20.

Jaeger, L., 1976: Monatskarten des Niederschlags fur die ganze Erde. *Ber. Dtsch. Wetterdienstes*, **139**, 33pp.

Kalnay, E., et al., 1996: The NCEP/NCAR 40-year reanalysis project. *Bull. Amer. Meteor. Soc.*, **77**, 437-471.

Kidd, C., 1998: On Rainfall Retrieval using Polarization-Corrected Temperatures. *International Journal of Remote Sensing*, **19**(5), pp. 981-996.

Kniveton, D. R., Barrett, E. C. Dodge, J. C. 1996: The Development of Composite Algorithms for Global Rainfall Estimation data from DMSP SSM/I. *Int. Journal of Remote Sensing*, **18**(3), pp. 517-534.

Kummerow, C., W. S. Olson, and L. Giglio, 1996: A simplified scheme for obtaining precipitation and vertical hydrometeor profiles from passive microwave sensors. *IEEE Trans. Geosci. Remote Sensing*, **34**, pp. 1213-1232.

Lau, K. M., J.H. Kim, and Y. Sud, 1996: Intercomparison of Hydrologic Processes in AMIP GCMs. *Bull. Amer. Met. Soc.*, **77**, pp. 2209-2227.

Legates, D., and C. J. Willmott, 1990: Mean seasonal and spatial variability in gauge-corrected, global precipitation. *Int. J. Climatol.*, **10**, 111-127.

Liu and Curry, 1996: Large-scale cloud features during January 1993 in the North Atlantic Ocean as determined from SSM/I and SSM/T2 observations. *J. Geophys. Res.*, **101**, pp. 7019-7032.

Morrisey, M.L., Shafer, M.A., Postawko, S.E. and Gibson, B. (1995). Pacific Raingauge Data, *Water Resources Research*, **31**, pp. 2111-2113

Petty, G. W., 1994a,b: Physical retrievals of over-ocean rain rate from multichannel microwave imagery. Parts I and II. *Meteorol. Atmos. Phys.*, **54**, pp. 79-122.

Petty, G.W. (1995). Frequencies and Characteristics of Global Oceanic Precipitation from Shipboard Present-Weather Reports. *Bulletin of the American Meteorological Society*, **76**(9), pp. 1593-1616.

Rudolf, B., H. Hauschild, W. Rueth and U. Schneider (1994): Terrestrial Precipitation Analysis: Operational Method And Required Density Of Point Measurements. In: Global Precipitations and Climate Change (Ed. M. Desbois, F. Desalmond), NATO ASI Series I, Springer-Verlag, **26**, pp. 173 - 186.

Schubert, S., R. Rood, and J. Pfaendtner, 1993: An assimilated data set for earth science applications, *Bull. Atmos. Meteor. Soc.*, **74**, pp. 2331-2342.

Smith, E. A., J. E. Lamm, R. Adler, J. Alishouse, K. Aonashi, E. Barrett, P. Bauer, W. Berg, A. Chang, R. Ferraro, J. Ferriday, S. Goodman, N. Grody, C. Kidd, D. Kniveton, C. Kummerow, G. Liu, F. Marzano, A. Mugnai, W. Olson, G. Petty, A. Shibata, R. Spencer, F. Wentz, T. Wilheit, and E. Zipser, 1998: "Results of the WetNet PIP-2 Project", *J. Atmos. Sci.*, **55**, pp. 1483-1536.

Smith, D. M., Kniveton, D. R., Barrett, E. C., 1998: A Statistical Modeling Approach to Passive Microwave Rainfall Retrieval. *J. Appl. Meteor.*, **37**(2), pp. 135-154.

Smith, E. A., X. Xiang, A. Mugnai, and G. J. Tripoli, 1994: Design of an inversion-based precipitation profile retrieval algorithm using an explicit cloud model for initial guess microphysics. *Meteorol. Atmos. Phys.*, **54**, pp. 53-78.

Spencer, R. W., 1993: Global oceanic precipitation from the MSU during 1979-92 and comparisons to other climatologies. *J. Climate*, **6**, pp. 1301-1326.

Susskind, J., P. Piraino, L. Rokke, L. Iredell, and A. Mehta, 1997: Characteristics of the TOVS Pathfinder Dataset. *Bull. Am. Met. Soc.*, **78**(7), pp. 1449-1472.



Tiedtke, M., 1993: Representative of Clouds in Large-Scale Models, *Mon. Wea. Rev.*, **121**(11), pp. 3040-3061.

Wentz, F. J., and R. Spencer, 1998: SSM/I Rain Retrievals within an Unified All-Weather Ocean Algorithm, Tech. Rpt. 011596, Remote Sensing Systems, Santa Rosa, CA 95404. *Journal of Atmospheric Sciences*, PIP-2 Special Issue, **55**(9), pp. 1613-1627

Wilheit, T. T., et al., 1997: A satellite technique for quantitatively mapping rainfall rates over the ocean, *J. Appl. Meteor.*, **16**, pp. 551-560.

Wilheit, T., A. Chang and L. Chiu, "Retrieval of monthly rainfall indices from microwave radiometric measurements using probability distribution function," *J. Atmos. Oceanic Tech.*, **8**, pp. 118-136.

Xie, P., and P. A. Arkin, 1996: Global Monthly Precipitation Estimates from Satellite-Observed Outgoing Longwave Radiation. *J. Climate*, **11**(2), pp. 137-164.

Xie, P., and P. A. Arkin, 1996: Analysis of Global Precipitation Using Gauge Observations, Satellite Estimates and Numerical Model Predictions. *J. Climate*, **9**, pp. 840-858.

## Table Caption

Table 1. List of precipitation products along with short code for each product, source of data for each product, land/ocean coverage for each product and reference.

## Figure Captions

Fig. 1. Location of  $2.5^\circ$  boxes of validation data. The shaded boxes were used in the validation of monthly amounts and the sources of data are described in the text. The outlined boxes were used for validation of interannual changes.

Fig. 2. Four examples of monthly maps of estimates of precipitation for July 1992 representing different types of products.

Fig. 3. Zonal totals of precipitation for 1998. (a) the mean, maximum and minimum from the product data set as a function of latitude. (b) the standard deviation of the zonal mean among the observational products and among the Quasi-Standard products.

Fig. 4. Zonal totals of precipitation for 1998. (a) the mean of the observed products compared to climatologies. (b) results from the four model products.

Fig. 5. Statistical results for each of the products for monthly rain over the atoll region of the Western Pacific Ocean. The results are grouped by product type.

Fig. 6. Statistical results for each of the products for monthly rain over land in the Tropics from 30°N to 30°S . The results are grouped by product type.

Fig. 7. Statistical results for each of the products for monthly rain over land in mid-latitudes from 30°N to 60°N . The results are grouped by product type.

Fig. 8. Four examples of monthly maps of interannual change of precipitation from July 1992 to July 1993 representing different types of products.

Fig. 9. Statistical results for each of the products for interannual change of precipitation over the atoll region of the Western Pacific Ocean. The results are grouped by product type.

Fig. 10. Statistical results for each of the products for interannual change of precipitation over land in the Tropics from 30°N to 30°S . The results are grouped by product type.

Fig. 11. Statistical results for each of the products for interannual change of precipitation over land in the Tropics from 30°N to 60°N . The results are grouped by product type.

Fig. 12. Examples of interannual change in precipitation in four areas (one ocean, three land). Over land the areas are 5° by 5° latitude/longitude boxes shown by the outlined areas in Fig. 1. Over water the area is the sum of six 2.5° boxes in the outlined area in the Western Pacific in Fig. 1.

Fig. 13. Four examples of maps of estimates of precipitation frequency over ocean for 1992 representing different types of products and the climatology based on COADS data.

Fig. 14. Comparison of the frequency of human observation of precipitation (excluding drizzle) to the frequency of raingauge measured rainrates of 0.5 mm/h, or greater.

Fig. 15. Comparison of the zonal average of precipitation frequency over ocean for the SSM/I products as compared to the COADS climatology.

Fig. 16. Comparison of the zonal average of precipitation frequency over ocean for 1992 for two model products as compared to the COADS climatology.

Fig. 17. Statistical results for each of the products for frequency of precipitation over ocean for 1992 from 45°S to 45°N . The results are grouped by product type.

Fig. 18. Seasonal variation of precipitation frequency over the ocean between 30°N to 45° for 1992 as estimated by various products and the COADS data and compared to the COADS climatology.

Table 1. List of precipitation products along with short code for each product, source of data for each product, land/ocean coverage for each product and reference.

Algorithm Description	Algo code	Algo class	Source data				Coverage			Reference
			SSMI	IR	Gauge	Model	Other	Land	Ocean	
Jaeger Climatology	jae	C	-	-	-	-	X	X	X	Jaeger, 1976
Legates/Willmott Clim.	lnw	C	-	-	-	-	X	X	X	Legates, 1990
Bristol (Comb.algorithm)	buc	E	X	-	-	-	X	X	X	Ebert, 1996
Bristol (PCT)	bup	E	X	-	-	-	X	X	X	Kidd, 1998
Bristol (Self calibrating)	bus	E	X	-	-	-	X	X	X	Smith, 1996
Colorado (F)	cuf	E	X	-	-	-	X	X	X	Ferriday, 1994
Colorado (L&C)	cul	E	X	-	-	-	X	X	X	Liu, 1996
DLR	dlr	E	X	-	-	-	-	X	X	Bauer, 1993
Florida State	fsu	E	X	-	-	-	-	X	X	Smith, 1994
Goddard (GPROF)	gpf	E	X	-	-	-	X	X	X	Kummerow, 96
Goddard (GSCAT)	gsc	E	X	-	-	-	X	X	X	Adler, 1994
Goddard (mod. emission)	gme	E	X	-	-	-	-	X	X	Chiu, 1993
Iowa	iow	E	X	-	-	X	-	X	X	Haferman, 1996
NCAR	ncr	E	X	-	-	-	-	X	X	Sheu
NOAA (OPI)	opi	E	-	X	-	-	-	X	X	Xie, 1996
Purdue	pur	E	X	-	-	-	-	X	X	Petty, 1994
Remote Sensing System	rss	E	X	-	-	-	-	X	X	Wentz, 1998
Texas A&M	tam	E	X	-	-	-	-	X	X	Wilheit, 1997
AMIP models	amp	M	-	-	-	X	-	X	X	Lau, 1996
ECMWF model	ecm	M	-	-	-	X	-	X	X	Tiedtke, 1993
GEOS model	geo	M	-	-	-	-	X	X	X	Schubert, 1993
NCEP model	ncp	M	-	-	-	X	-	X	X	Kalnay, 1996
Goddard (Emission)	gem	Q	X	-	-	-	-	X	X	Wilheit, 1997
Goddard (TOVS)	tov	Q	-	-	-	-	X	X	X	Susskind, 1997
GPCP (Merged satellite)	gps	Q	X	X	-	-	-	X	X	Huffman, 1997
GPCP (Satellite/Gauge)	gpm	Q	X	X	X	-	-	X	X	Huffman, 1997
Marshall MSU	msu	Q	-	-	-	-	X	X	X	Spencer, 1993
Marshall PIP-1	plc	Q	X	-	-	-	-	X	X	Kniveton, 1996
NOAA GPI	gpi	Q	-	X	-	-	-	X	X	Arkin, 1987
NOAA (satellite/gauge)	nmg	Q	X	X	X	X	-	X	X	Xie, 1996
NOAA (SSM/I)	nmi	Q	X	-	-	-	-	X	X	Ferraro, 1995





Figure 1



Climatology (lnw)



Model (ecm)



Experimental (buc)



Quasi-standard (gpm)

July 1992




Figure 2





Figure 3







Figure 4

# Statistics Plots : Atoll Region (12 months)




Figure 5

# Statistics Plots : Land Areas 30S-30N (12 months)




Figure 6

# Statistics Plots : Land Areas 30N-60N (12 months)




Figure 7




Figure 8

Statistics Plots : Atoll Region (all) (interannual)




Figure 9

Statistics Plots : Land Areas 30S-30N (interannual)




Figure 10

Statistics Plots : Land Areas 30N-60N (interannual)






Figure 11




a) Atolls (172.5-180.0E, 10.0S-5.0N, July)



b) Missouri (97.5-92.5W, 37.5-42.5N, July)



c) Mississippi (85.0-90.0W, 32.5-37.5N, July)



d) Australia (145.0-150.0E, 37.5-37.5S, July)




Figure 12




Figure 13




Figure 14

# Global Oceanic Latitudinal Profile (1992) SSM/I Products




Figure 15

# Global Oceanic Latitudinal Profile (1992) Models




Figure 16

# Statistics Plots : COADS 45S-45N (annual)




Figure 17

# COADS Seasonal plots : 30N-45N




Figure 18