

Discovery of a New Nearby Star

B. J. Teegarden*, S. H. Pravdo[†], K. Covey[‡], O. Frazier[‡], S. L. Hawley[‡], M. Hicks[‡], K. Lawrence[‡], T. McGlynn*, I. N. Reid^{||}, and S. B. Shaklan[‡],

**Laboratory for High Energy Astrophysics, NASA/Goddard Space Flight Center, Greenbelt, MD 20771, US*

†Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

‡University of Washington, Dept. of Astronomy, Box 351589, Seattle, WA 98195, USA

|| Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA

We report the discovery of a nearby star with a very large proper motion of 5.06 ± 0.03 arcsec/yr. The star is called SO025300.5+165258 and referred to herein as HPMS (high proper motion star). The discovery came as a result of a search of the SkyMorph¹ database, a sensitive and persistent survey that is well suited for finding stars with high proper motions. There are currently only 7 known stars with proper motions > 5 arcsec/yr. We have determined a preliminary value for the parallax of $\pi = 0.43 \pm 0.13$ arcsec. If this value holds our new star ranks behind only the Alpha Centauri system (including Proxima Centauri) and Barnard's star in the list of our nearest stellar neighbours. The spectrum and measured tangential velocity indicate that HPMS is a main-sequence star with spectral type M6.5. However, if our distance measurement is correct, the HPMS is underluminous by 1.2 ± 0.7 mag.

Proper motion surveys have historically revealed a large fraction of our nearest stellar neighbours. The usual method is to search plate pairs typically taken a decade or more apart for relative displacements. However, stars with large proper motions can get lost in the background. The SkyMorph database of NEAT² (Near Earth Asteroid

Tracking) data is ideally suited to search for such objects. NEAT is a NASA-funded project intended to perform a comprehensive survey of near-earth asteroids. Beginning in 1995, several dedicated telescopes have been employed to repeatedly take CCD exposures of the sky mostly in the region within $\pm 25^\circ$ of the celestial equator (excluding galactic latitudes $< \sim 20^\circ$). The instantaneous fields-of-view have varied over the years with telescope and CCD camera, but generally covered areas of 2-3 deg^2 . Our survey, thus far, has included data from May 1996 – Dec 2001 from two 1-m telescopes located on Maui. The total sky coverage is $\sim 10000 \text{ deg}^2$, and within the region of coverage there are typically 20 exposures at any given point. The revisit time ranges from \sim hours to years. With such a large number of exposures over a ~ 6 yr period, the Skymorph database is an ideal place to search for objects with very large proper motion.

We have searched the Skymorph database for objects with unfiltered CCD magnitudes in the range 11.0-18.0 having proper motions $> 0.25 \text{ arcsec/yr}$. Candidate objects were visually confirmed by comparison with Digital Sky Survey images using the overlay feature of the SkyView Program³. A total of 3154 objects were found of which 63 had proper motions $> 1.0 \text{ arcsec/yr}$. ~ 400 of these objects are previously uncatalogued and a paper on these is in preparation⁴. The HPMS had the largest proper motion of all the detected objects: $5.06 \pm 0.03 \text{ arcsec/yr}$. To further refine the HPMS proper motion determination we have included additional observations from the USNO A2.0 Catalogue⁵ and the 2nd Guide Star Catalogue⁶. The measured parameters of our fast-moving star are given in Table 1. We derive from our observations a tangential velocity of $52.9 \pm 11.7 \text{ km/s}$. This is consistent with the velocity dispersion of main sequence red dwarfs of $\sigma \sim 50 \text{ km/s}$.⁷

Figure 1 shows the motion of HPMS across the sky with images from the Palomar Sky Surveys and SkyMorph. The parallax was estimated by fitting the position of HPMS on 43 SkyMorph images (many averaged in sets of three) taken from Julian Dates

2450779-2452498 (November 26, 1997- August 11, 2002) to a reference frame defined by 12 nearby stars in the field. Since the HPMS is quite red, a significant correction for differential chromatic refraction (DCR) was required. The DCR correction was both estimated from first principles and fit to the data taken over one night with varying hour angles. We chose a value for the DCR correction that minimized the scatter in the position on single nights. Figure 2 shows the result of the fit to parallax of the HPMS position with the NOVAS⁸ model for proper motion and parallax. The result for the parallax is 0.43 ± 0.13 arcsec. The 1-sigma error is larger than we would like because the camera pixel sizes in the measurements, 1 and 1.4 arcsec, are not ideal for this purpose. More precise observations with cameras designed for this purpose are underway.

A spectrum of the HPMS taken on 11 July, 2002 using the Apache Point 2.5 m telescope is shown Figure. 3. For comparison the spectra of the M6.5 V star GJ1111 and the M8 V star VB10 are also shown. The match to GJ1111 is good. We identify the HPMS as a main-sequence dwarf with spectral type M6.5 V based upon detailed examination of the CaH2 ($\lambda 7042-7046$, $\lambda 6814-6846$), CaH3 ($\lambda 7042-7046$, $\lambda 6960-6990$), and TiO5 ($\lambda 7042-7046$, $\lambda 7126-7135$) spectral features⁹. The relative strengths of the CaH and TiO bands show no evidence that the star is significantly metal-poor.

The HPMS object was observed photometrically on the night of 7 August, 2002 at the JPL Table Mountain Observatory 0.6-meter telescope using broad band UBVRI Bessel filters and the facility CCD imaging system. Multiple UBVRI exposures of 15 Landolt (1992) stars ranging in color and air mass were also measured. With this data both linear and color-corrected extinction curves for each filter were computed. The nominal extinction coefficients and small (~ 1 mMag) color coefficients indicated a good photometric night. The color-corrected extinction curves were used to compute the measured magnitudes listed in Table 1.

The HPMS is also in the soon-to-be-released 2MASS all-sky point source catalog (2MASS J02530084+1652532)¹⁰, with magnitudes of $J = 8.39 \pm 0.03$, $H = 7.88 \pm 0.04$ and $K_s = 7.59 \pm 0.05$. The photometric measurements are summarized in Table 1. Combining these data with the trigonometric parallax given above allows estimation of the absolute magnitude. Despite the strong spectroscopic similarity to GJ 1111, the derived values are significantly fainter. For example, we derive $M_J = 11.58 \pm 0.35$ for the HPMS, as compared with $M_J = 10.40 \pm 0.03$ for GJ 1111. If our trigonometric parallax is accurate, then the HPMS is underluminous relative to a normal M6.5 disk dwarf by 1.2 ± 0.7 magnitudes. This result might be expected for an extreme metal-poor subdwarf, such as LHS 1826 (Gizis & Reid, 1997, PASP, 109), but, as noted above, there is no indication that HPMS is metal-poor. If the star has the same effective temperature as GJ 1111, then the low luminosity implies that the HPMS has a radius only 60% that of GJ 1111, or $\sim 0.68 R_{\text{Jupiter}}$, violating electron degeneracy.

Given these results, we have estimated a spectrophotometric parallax to HPMS. Dahn et al (2002, AJ) have shown that spectral type is well correlated with M_J for ultracool dwarfs. Using their linear relation, we derive $M_J = 10.59 \pm 0.25$, corresponding to a distance of 3.6 ± 0.4 parsecs. At this distance, which is almost identical to that of GJ 1111, HPMS is the 17th nearest system (and 27th nearest star) to the Sun.

Table 2 summarizes the properties of the 10 nearest known stars including distances and proper motions. Given our parallax value, the HPMS ranks 3rd in distance among stellar systems (or 5th among individual objects) and 8th in proper motion. (We emphasize again that the parallax value is preliminary and that the distance ranking could change.). In the visual band it is the faintest star in the list, which is part of the reason it was not discovered in previous surveys. Since the NEAT survey has only covered the region $\pm 25^\circ$ in declination it is entirely possible that other faint nearby objects remain to be discovered.

Table 1. HPMS Parameters

RA (J2000)*	42.24369°
DEC (J2000)*	16.89200°
Epoch*	1989.84
Magnitudes (B,V, R, I, J, H, K _s) [†]	17.21 ± 0.11, 15.40 ± 0.08, 13.26 ± .06, 10.66 ± 0.03, 8.39 ± 0.03, 7.88 ± 0.04, 7.59 ± 0.05
Spectral Type	M6.5 V
Proper Motion	5.06 ± 0.03 arcsec/yr
Position Angle	138.1 ± 0.3°
Parallax (π)	0.43 ± 0.13 arcsec
Distance (1/ π)	2.3 (+1.0,-0.5) pc
Tangential Velocity	52.9 ± 11.7 km/s

* From 2nd Guide Star Catalogue.

† B,V,R,I are from this work. J, H, K_s are from the 2MASS Catalogue.

Table 2. Nearest Known Stars*

Name	D(pc)	PM [†] (arcsec/yr)	M(V) [‡]	Sp. Type
Proxima Centauri	1.30	3.81	15.49	dM5 e
Alpha Centauri A	1.34	3.69	4.38	G2 V
Alpha Centauri B	1.34	3.69	5.71	K0 V
Barnard's Star	1.83	10.31	13.23	M5 V
HPMS (this work)	2.3 (+1.0,-0.5)	5.06 ± .03	18.5 ± 0.7	M6.5 V
Wolf 359	2.39	4.70	16.56	M6
Gl 411	2.52	4.81	10.48	M2 Ve
Luyten 726-8 A	2.63	3.37	15.47	dM5.5 e
Luyten 726-8 B (UV Ceti)	2.63	3.37	15.60	dM5.5 e
Sirius A	2.63	1.33	1.47	A1 V
Sirius B	2.63	1.33	11.34	DA2

* All values in this table (with the exception of the HPMS) were either taken directly or derived from the Preliminary Version of the Third Catalogue of Nearby Stars¹¹.

† Proper motion.

‡ Absolute visual magnitude.

Acknowledgments

We thank D. Kirkpatrick for his reduction of the HPMS 2MASS data. We also thank N. Zacharias, S. Levine, and C. Dahn for useful discussions. This research was performed in part by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. The results presented in this paper are based in part on data obtained using the NASA-JPL camera at the Oschin Telescope, Palomar Observatory, California Institute of Technology. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. SkyMorph is funded by NASA's Applied Information Systems Program.

The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to B. Teegarden (e-mail: bonnard@lheamail.gsfc.nasa.gov).




Figure 1. Images from the Palomar Sky Surveys and SkyMorph (lower right) show the motion of HPMS south and east across the sky from 1951 until the current epoch.




Figure 2. The fit to the parallax of HPMS results in a value of 0.43 ± 0.13 arcsec. The errors per measurement that produce the good fit shown are 0.28 arcsec. The contribution to this error from photon-counting statistics is ~ 0.1 arcsec. The remainder is due to uncalibrated DCR or other unknown systematic errors.




Figure 3. The spectrum of HPMS compared with two standard stars. Based upon this comparison and detailed analysis of spectral features HPMS is a main sequence dwarf with spectral type M6.5.

¹. The SkyMorph database is accessed via the Worldwide Web at:

<http://skyview.gsfc.nasa.gov/skymorph/>.

² Pravdo, S. H. et al., The Near-Earth Asteroid Tracking (NEAT) Program: An Automated System for Telescope Control, Wide-Field Imaging, and Object Detection, *Astron. J.*, **117**, 1616-1633 (1999).

-
- ³ McGlynn, T., Scollick, K., White, N., SkyView: The Multi-Wavelength Sky on the Internet, McLean, B.J. et al., *New Horizons from Multi-Wavelength Sky Surveys*, Kluwer Academic Publishers, IAU Symposium No. 179, p465 (1996).
- ⁴ Pravdo, S. H. et al. 2002, in preparation.
- ⁵ . Monet, D. G., The 526,280,881 Stars in the USNO-A2.0 Catalog, *A. A. S.*, **193**, 12003 (1998).
- ⁶ Morrison, J.E. and McLean, G., The Second Generation Guide Star Catalog, *AAS DDA meeting #32*, #06.03 (2001).
- ⁷ Binney, J. and Merrifield, M., *Galactic Astronomy*, Princeton Univ. Press, Princeton, p. 627 (1998).
- ⁸ G. H. Kaplan et al., Mean and apparent place computations in the new IAU system. III - Apparent, topocentric, and astrometric places of planets and stars, *Astron. J.* **97**, 1197-1210 (1989).
- ⁹ Reid, I. N., Hawley, S.L., & Gizis, J. E., The Palomar/MSU Nearby-Star Spectroscopic Survey. I. The Northern M Dwarfs - Bandstrengths and Kinematics, *Astron. J.*, **110**, 1838-1859 (1995).
- ¹⁰ Kirkpatrick, D., private communication.
- ¹¹ Gliese, W. and Jahreiss, H., *Preliminary Version of the Third Catalog of Nearby Stars*, Astron. Rechen-Institut, Heidelberg (1991).