


NASA/CP—2002—211842



A DRAFT TEST PROTOCOL FOR DETECTING POSSIBLE BIOHAZARDS IN MARTIAN SAMPLES RETURNED TO EARTH

Edited by:

*John D. Rummel
NASA Headquarters
Washington, D.C.*

*Margaret S. Race
SETI Institute
Mountain View, California*

*Donald L. DeVincenzi
NASA Ames Research Center
Moffett Field, California*

*P. Jackson Schad
NASA Headquarters
Washington, D.C.*

*Pericles D. Stabekis
The Windermere Group
Washington, D.C.*

*Michel Viso
CNES Headquarters
Paris, France*

*Sara E. Acevedo
SETI Institute
Mountain View, California*

National Aeronautics and
Space Administration

Ames Research Center
Moffett Field, California 94035-1000

October 2002

Acknowledgements

The Editors wish to acknowledge the contributions of the diverse group of scientific experts who participated in the development of this Draft Protocol. We thank both the scientists who participated in the Workshop Series and the members of the Oversight and Review Committee. Their contributions are represented here as fully as possible. Any errors that may have crept in are the responsibility of the Editors alone.

Available from:

NASA Center for AeroSpace Information
7121 Standard Drive
Hanover, MD 21076-1320
301-621-0390

National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
703-605-6000

PREFACE

This document provides the final version of a *Draft Test Protocol for Detecting Possible Biohazards in Martian Samples Returned to Earth*. This Draft Protocol was developed through an iterative process of discussion and review during the Mars Sample Handling Protocol Workshop Series, as well as afterwards. The table below is a chronological list of key workshops, reviews, and publications that led to the development of the Draft Protocol, and gives the terminology used in this document to refer to earlier versions. The final reports from the Workshops are cited in Appendix B, and contain full documentation and details of the sub-group discussions at each Workshop. The discussions from Workshops 1 through 3 led to a consensus that was reached during Workshop 4, resulting in the first complete protocol (denoted below as the “Completed Working Draft Protocol”). That document underwent review and revision by a special Oversight and Review Committee (see Appendix C), and a reading by the NASA Planetary Protection Advisory Committee. This “final” version of the Draft Protocol resulted from their critical reading and revisions, and supercedes all earlier versions. It is anticipated that this Draft Protocol will be subject to extensive further review and debate prior to development of any final protocol for use in receiving and testing samples from Mars.

Terminology Used	Date/Location	Report Citation or Annotation
Workshop 1 Final Report	March 2000, Bethesda, MD	<i>Race and Rummel, 2000</i>
Workshop 2 Final Report	October 2000, Bethesda, MD	<i>Race et al., 2001a</i>
Workshop 2a Final Report	November 2000, Rossllyn, VA	<i>Bruch et al., 2001</i>
Workshop 3 Final Report	March 2001, San Diego, CA	<i>Race et al., 2001b</i>
Penultimate Working Draft Protocol	May 2001	First compilation of the developing protocol from recommendations of Workshops 1, 2, 2a, and 3
SSB/COMPLEX Report: The Quarantine and Certification of Martian Samples	May 2001 Advance Copy	<i>SSB 2002</i>
Workshop 4 Final Report	June 2001, Arlington, VA	<i>Race et al., 2002.</i>
Completed Working Draft Protocol	June 2001	A consensus working draft resulting from the entire Workshop Series — published in WS 4 Final Report (see <i>Race et al., 2002</i> , Appendix A, page 71.); submitted to the ORC for comment and review.
Oversight & Review Committee (ORC) review process Oct-Nov 2001	12 November, 2001, ORC Meeting, Rockefeller University New York, NY	Review of the Completed Working Draft Protocol.
<i>A Draft Test Protocol for Detecting Possible Biohazards in Martian Samples Returned to Earth</i>	October 2002	<i>Rummel et al., 2002</i> (this document); the final version of the Draft Protocol incorporating comments and recommendations from the ORC.

TABLE OF CONTENTS

23	TABLE OF CONTENTS		
24			
25	Preface		i
26	Table of Contents		iii
27	Draft Test Protocol		
28	Introduction to the Draft Protocol		1
29	Why a Draft Protocol?		4
30	Containment in the Sample Receiving Facility (SRF) and Elsewhere.....		5
31	“Sterilization” of Martian Samples		9
32	Criteria for Release		14
33	Overview of the Draft Protocol		17
34	Physical/Chemical Processing		19
35	Future P/C Research and Development Needs		37
36	Life Detection Testing		40
37	Future LD Research and Development Needs		54
38	Biohazard Testing		54
39	Future BH Research and Development Needs		71
40	Facility Requirements		72
41	Future Research and Development Needs		75
42	Environmental and Health Monitoring and Safety		75
43	Future Research and Development Needs		82
44	Personnel Management Considerations in Protocol Implementation		83
45	Future Considerations.....		91
46	Contingency Planning for Different Protocol Outcomes		91
47	Maintaining and Updating the Protocol		97
48	Appendices		
49	A. MSHP Workshop Series Basic Assumptions		105
50	B. References		107
51	C. MSHP Oversight and Review Committee		109
52	D. MSHP Leadership Group		113
53	E. Glossary of Terms and Acronyms		115

54 **A DRAFT TEST PROTOCOL**
55 **FOR DETECTING POSSIBLE BIOHAZARDS**
56 **IN MARTIAN SAMPLES RETURNED TO EARTH**
57

58 **Introduction to the Draft Protocol**

59 In anticipation of missions to Mars that will involve the return of samples, it is
60 necessary to prepare for the safe receiving, handling, testing, distributing, and
61 archiving of martian materials here on Earth. Previous groups and committees
62 have studied selected aspects of sample return activities, but a specific protocol
63 for handling and testing of returned samples from Mars must still be developed.
64

65 For upcoming Mars sample return missions, NASA is committed to following the
66 recommendations developed by the Space Studies Board (SSB) of the National
67 Research Council (NRC) in its report on sample handling and testing [SSB 1997].

68 In particular, the NRC recommended that:

69 *a) “samples returned from Mars by spacecraft should be contained and*
70 *treated as potentially hazardous until proven otherwise,” and b) “rigorous*
71 *physical, chemical, and biological analyses [should] confirm that there is no*
72 *indication of the presence of any exogenous biological entity.”*
73

74 To develop and refine the requirements for sample hazard testing and the criteria
75 for subsequent release of sample materials from precautionary containment, the
76 NASA Planetary Protection Officer convened the Mars Sample Handling Protocol
77 (MSHP) Workshop Series from March 2000 to June 2001. The overall objective of
78 the Workshop Series was to produce a Draft Protocol by which returned martian
79 sample materials could be assessed for biological hazards and examined for
80 evidence of life (extant or extinct), while safeguarding the samples from possible
81 terrestrial contamination. In addition to U.S. and international participants invited by
82 NASA, significant participation and support by French scientists were provided in
83 all aspects of the Workshops and protocol development through arrangement with
84 the Centre National d’Études Spatiales (CNES).

85 The stated objective for the Workshop Series was:

86 *“For returned Mars samples, develop a recommended list of comprehensive*
87 *tests, and their sequential order, that will be performed to fulfill the NRC*
88 *recommendations that ‘rigorous analyses determine that the materials do*
89 *not contain any biological hazards.’”*

90
91 Throughout the Workshop Series, these analyses were anticipated to comprise
92 not only a series of tests to detect a possible living entity (‘life detection’), but also
93 tests to look for biological activity, even if a living entity were not detected
94 (‘biohazard testing’).¹ Therefore the Workshop Series was designed to devise a
95 protocol that could rigorously analyze returned martian sample materials to
96 determine that those materials are free from biohazards and/or extraterrestrial life-
97 forms, and are therefore safe to be released from containment in their native state
98 for further scientific research. To accomplish this, Workshop Series participants
99 focused on a variety of questions that had to be addressed about the protocol to
100 meet the Series’ objective (see Appendix A). This Draft Protocol is intended to
101 incorporate the answers developed to those questions.

102

103 To keep the Workshop Series focused, a set of basic assumptions (see Appendix
104 A) was given to the participants at each of the Workshops to guide and constrain
105 their deliberations. Subsequent to the failure of the Mars Surveyor 1998 missions,
106 these assumptions were subject to some modification during the re-planning
107 process that NASA and its international partners undertook (i.e., the change of the
108 return date from ‘2007’ to ‘in the next decade’ in Assumption #2). However, none of
109 the modifications affected the basic premises under which the Workshop
110 participants undertook their task. These assumptions are consistent with the
111 plans of NASA and its international partners as of the publication of this report

1. This two-pronged approach is consistent with the Space Studies Board’s recommendations for returned martian samples [SSB 1997, p. 27]: “The initial evaluation of samples returned from Mars will focus on whether they pose any threat to the Earth’s biosphere. The only potential threat posed by returned samples is the possibility of introducing a replicating biological entity of non-terrestrial origin into the biosphere. Therefore, the initial evaluation of potential hazards should focus on whether samples contain any evidence of organisms or biological activity.”

112 (October 2002), and are expected to remain current despite the inevitable program
113 delays and likelihood of future changes.

114

115 In addition to the development of this Draft Protocol through the NASA-led
116 Workshop Series, the SSB was asked by NASA in early 1999 to develop
117 recommendations for the quarantine and certification of martian samples—both
118 as an input to the NASA Workshop Series, and as recommendations to NASA to
119 be assessed in their own right. The SSB report [*SSB 2002*] was released in
120 preliminary form in May 2001, just prior to Workshop 4. Thus participants of
121 Workshop 4 had access to an Advance Copy of the SSB report during their review
122 of the Penultimate Working Draft Protocol. Therefore, both the completed Working
123 Draft Protocol (as published in the Workshop 4 final report [*Race et al., 2002*]) and
124 this final version of the Draft Protocol reflect, to a great degree, an examination of
125 the findings and recommendations of the Space Studies Board study.²

126

127 This document is the first complete presentation of the Draft Protocol for Mars
128 sample handling that meets planetary protection needs, and represents a
129 consensus that emerged from the work of sub-groups assembled during the five
130 Workshops of the Series.³ Over the course of the Workshops, participants
131 converged on a conceptual approach to sample handling as well as on specific
132 analytical requirements. Further discussions identified important issues
133 remaining to be addressed, including research and development necessary for
134 optimal protocol implementation. This Draft Protocol also incorporates the review
135 comments of an Oversight and Review Committee (see Appendix C) that
136 examined the Completed Working Draft subsequent to the end of the Workshop
137 Series.

138

-
2. See Appendix B for a complete list of workshops and reports contributing to this Draft Protocol.
 3. The final reports from the Workshops in the Series [*Race and Rummel, 2000; Race et al., 2001a, 2001b, and 2002; Bruch et al. 2001*] contain full documentation and details of the sub-group discussions that fed into this final version of the Draft Protocol.

139 Why a 'Draft Protocol'?

140 What is reported here is termed a 'Draft' Protocol because it is intended to be just
141 that. While it is a responsibility of NASA's Planetary Protection Officer [NASA 1999]
142 to prescribe "standards, procedures, and guidelines applicable to all NASA
143 organizations, programs, and activities" to achieve the policy objectives of NASA's
144 planetary protection program, including ensuring that Earth is "protected from the
145 potential hazard posed by extraterrestrial matter carried by a spacecraft returning
146 from another planet or other extraterrestrial sources," (in this case, Mars), it is
147 neither practical nor useful for this Draft Protocol to be developed into a final form
148 at this time. The final protocol that will guide the process of assessing the martian
149 samples should owe much to new knowledge about Mars that will be gained in
150 robotic exploration on Mars leading up to the sample return mission, as well as
151 detailed information available only on the sample return mission itself. In addition,
152 the final protocol should take into account the *specific* nature of the receiving facility
153 that is developed for the initial processing and testing of the returned samples, as
154 well as the requirements and abilities of the *specific* instrumentation and
155 personnel selected to undertake the challenging task of testing the samples while
156 protecting Earth from possible hazards, and preserving the scientific value of the
157 sample return undertaking. It is anticipated that the final protocol will receive its
158 final review at or about the time the first samples leave the martian surface.


159
160 Meanwhile, this Draft Protocol is intended to provide a proof-of-concept model of
161 the final protocol, demonstrating one approach (and more importantly, a sufficient
162 approach) to testing returned Mars samples for possible biohazards or biological
163 activity of martian origin. This Draft Protocol has been developed to provide a
164 sequential series of tests that can be applied to martian samples to provide data
165 that can be used to make decisions about the release of unsterilized samples
166 from containment—either wholly or partially—while allowing for an earlier release
167 of samples subjected to a decontamination process ("sterilization") to ensure they
168 are safe for analyses outside of containment.

169 **Containment in the Sample Receiving Facility and Elsewhere**

170 In order to preserve the scientific value of returned martian samples under safe
171 conditions and avoid false indications of life within the samples, the capability is
172 required for handling and processing Mars samples while preventing their
173 contamination by terrestrial materials (i.e., cleanroom conditions, technical criteria
174 TBD) and while maintaining strict biological containment. This requirement is a
175 major challenge in the design of what will be described here as a Sample
176 Receiving Facility (SRF).⁴ To some degree, the cleanroom requirement is likely to
177 constrain the working space inside an SRF even more than might normally be
178 experienced in a “typical” Biosafety Level 4 (BSL-4) facility of similar size. An SRF
179 will require combining technologies currently found in maximum containment
180 microbiological laboratories (e.g., BSL-4, BSL-3)⁵ with those used in cleanrooms
181 to preserve the pristine nature of rare samples. Such an integrated facility is not
182 currently available anywhere. Some of the challenges of providing such a facility
183 may be alleviated through a design and development process that will include
184 mock-ups of containment/cleanroom combinations whose efficacy can be tested
185 thoroughly (see Figure 1 for some options). Some of the overall facility constraints
186 may be lessened through the use of multiple containment facilities to accomplish
187 different aspects of the protocol, especially where material (as opposed to
188 biological) contamination constraints can be relaxed. It is anticipated that samples
189 may be shipped among appropriate containment facilities wherever necessary
190 under procedures developed in cooperation with the U.S. Centers for Disease
191 Control and Prevention, the U.S. Department of Transportation, and appropriate
192 international authorities. Nonetheless, it is envisaged that all samples initially

-
4. A variety of names have been used in reference to the place where returned samples will be handled and tested initially (e.g. Sample Receiving Facility (SRF), the Quarantine Facility, the Mars receiving laboratory, primary containment facility, quarantine facility, etc.). A recent NRC report [SSB 2002] has used “Quarantine Facility,” but it is more useful in this report to use the generic SRF. The actual name and location(s) of the facility or facilities where the protocol will be executed is TBD. Use of these facilities beyond the receipt of martian samples may be anticipated.
 5. “BSL” levels are a North American convention. European equivalents will be considered and described as necessary in implementation of the final protocol.

193 returned from Mars will be placed in a single SRF and held there through the
 194 preliminary examination phase (i.e., “Preliminary Evaluation,” as envisaged in
 195 Figure 2 on page 18), and for those subsequent steps compatible with SRF
 196 design and capacity.



197
 198 Figure 1. Top and Center: Simple options for the combination of a biological
 199 containment facility with a cleanroom. Arrows show gas flow (*via* leakage) caused
 200 by pressure differentials in the spaces shown. Gray areas are potentially
 201 contaminated by any organisms the Mars samples might contain. Bottom: A more
 202 complex arrangement with double walls separating workers from samples, and in
 203 which the gases from the workers and the samples both are exhausted through the
 204 space between the walls (and in the case of the gases from the personnel, to the
 205 outside atmosphere). *From SSB 2002.*

206 BSL-4 is required for work with dangerous and exotic agents that pose a high risk
207 to the individual of aerosol-transmitted laboratory infection and life-threatening
208 disease. The unknown nature of any possible biohazard in returned martian
209 samples demands, at least initially, this most stringent containment presently
210 afforded to the most hazardous biological entities known on Earth. In the
211 biomedical community, biohazard testing is a pathway towards gradual
212 “decontainment” of dangerous and/or exotic bioagents, when supported by
213 experimental evidence. Decisions about the appropriate biosafety level for a
214 particular bioagent can be made when sufficient data are obtained to support
215 either the need for continued work at a high level of containment, or allowance to
216 conduct work at a lower level.

217

218 Generally, lower biosafety levels are assigned to bioagents with less human
219 virulence. If sufficient data are gathered to rule out concerns about human
220 virulence and infection, a decision could later be made to allow subsequent work
221 at a lower containment level during tests investigating possible environmental
222 effects. A lower level of containment would potentially enhance sample access
223 within the scientific community while still providing adequate biosafety conditions
224 under existing biosafety guidelines and regulations.

225

226 In addition to satisfying both biosafety and cleanliness needs, the SRF will need to
227 provide different types of laboratory environments for carrying out the various
228 aspects of protocol testing. During the Workshop Series, the new term ‘Planetary
229 Protection Level’ (PPL) was developed for the purpose of categorizing and
230 describing the different combinations of containment and cleanliness conditions
231 required within the SRF for different testing needs. Although details of various PPL
232 designations will require further definition, it is possible to anticipate a number of
233 laboratory conditions that may be required during the protocol testing. The four
234 PPLs are described in the following text and in Table 1:

235

- 236 ● PPL- α – for incoming samples and archived samples; maximum
 237 biocontainment and cleanliness; maintains samples in an inert gas
 238 environment and Mars-like conditions (TBD).⁶
- 239 ● PPL- β – maintains maximum biocontainment and protection for workers
 240 and the environment; maximum cleanliness, but allows exposure to
 241 ambient terrestrial conditions.
- 242 ● PPL- γ – maintains maximum biocontainment with moderate cleanliness
 243 and ambient terrestrial conditions (i.e., for animal testing scenarios).
- 244 ● PPL- δ – maintains BSL-3-Ag containment conditions, with less
 245 emphasis on cleanliness, and ambient terrestrial conditions.⁷
- 246

PPL-type	Biocontainment	Cleanliness	'Ambient' Conditions	Used For:
PPL- α	Maximum (BSL-4)	Maximum	Mars-like (pristine); <i>Although at 1 atm w/inert gas environment.</i>	Incoming container and materials; some preliminary tests; sample bank/storage; some Life Detection
PPL- β	Maximum (BSL-4)	Maximum	Earth-like	Life Detection; some Physical/Chemical; TBD
PPL- γ	Maximum (BSL-4)	Moderate	Earth-like	Some Biohazard testing, some Physical/Chemical processing, and animal testing
PPL- δ	Strict BSL-3-Ag	Ambient	Earth-like	Some Biohazard testing; 'post-release' tests TBD

247
 248
 249
 250
 251

Table 1. Anticipated laboratory conditions and PPL categories. Note: Levels of cleanliness associated with each PPL are TBD and should be defined explicitly well in advance of sample return.

-
6. It is anticipated that only the primary SRF will be required to have PPL- α conditions. If other facilities beyond the SRF are used as part of the protocol testing, they will be certified for conducting particular tests or studies at the appropriate PPL conditions.
7. PPL- δ provides a level of containment for the samples that allows investigators to work in a laboratory situation providing protection to personnel through an engineered environment with HEPA filtered air entering and leaving the area, containment of water and/or waste to the laboratory, and protection through personnel protective equipment consistent with U.S. BSL-3 Agriculture and French P4 standards. It was recommended that the BSL-3-Ag facilities used should be designed to accommodate large instruments, rather than miniaturizing the instruments to fit into a pre-existing lab.

253 It is important to note that, regardless of cleanliness requirements or ambient
254 conditions, all initial testing will be done under maximum biocontainment
255 equivalent to United States BSL-4 [CDC-NIH, 1993]. In addition, Biohazard
256 testing will not require the extreme cleanliness levels to be used for initial
257 sample processing, or certain Physical/Chemical or Life Detection tests. The
258 majority of Biohazard tests will be done in PPL- γ . If the results of the initial Life
259 Detection and Biohazard tests are all negative, it may be appropriate to conduct
260 some subsequent tests under less strict containment conditions. The first step
261 in downgraded containment for untreated samples has been designated as
262 PPL- δ , which is equivalent to BSL-3-Ag.⁸

263

264 **“Sterilization” of Martian Samples**

265 Recognizing that a species' adaptation to physiological stress may evolve through
266 natural selection, it is expected that possible extant life on Mars could be able to
267 survive extremely hostile conditions. Surface temperatures at the equator of Mars
268 range from -100°C during the martian winter to 20°C during the martian summer.
269 Mars is extremely dry; the partial vapor pressure of water on the surface is
270 approximately 0.1 bar. The martian atmosphere is 95% CO_2 and provides no
271 protection against exposure to 200-300 nanometer ultraviolet light, which may
272 generate strong oxidants in the surface material. It is believed that organic
273 compounds on the surface of Mars are subject to oxidation by this UV-induced
274 photochemistry. Since this combination of conditions cannot be found on Earth, it
275 is unlikely that a single terrestrial species will be found that can serve as a
276 surrogate for a putative martian organism when evaluating methods for sterilizing
277 martian samples. There are terrestrial environments, however, that are sufficiently
278 similar to the martian environment to allow the isolation of species that exhibit
279 extreme resistance to a subset of the conditions (e.g., desiccation, radiation, or

8. PPL- δ applies at the point in the protocol where samples do not require atmospheric isolation and may be moved to outside laboratories with suitable facilities for further testing. In general, level 3 biosafety laboratories (BSL-3) abide by different standards within the U.S. and Europe. For clarity, the U.S. standard for BSL-3-Ag will be used.

280 cold) to be encountered on Mars. As an item for further research, it is anticipated
281 that an effort will be made to identify and characterize terrestrial species from
282 environments as similar as possible to those on Mars, and that these species will
283 be used to validate sterilization processes.

284

285 In the context of this Draft Protocol and the relevant NRC reports [SSB 1997; SSB
286 2002], the term “sterilization” is used to connote the decontamination process that
287 will be used to ensure that the samples are safe for analyses outside of
288 containment. It is possible, though very unlikely, that martian organisms are not
289 carbon based, and martian biology could conceivably be based on other elements
290 (e.g., Si, N, P, O, H, S, Al, B). But overall, it should be noted that the chemical
291 elements on Mars and the forces holding molecules together are the same as on
292 Earth. If there were a life-form on Mars based on other than carbon-containing
293 molecules, the energies holding such molecules together would not be much
294 different than those for proteins and polynucleotides. Hence, bond breakage by
295 heat or gamma radiation should be similar for Earth and Mars life-forms, and
296 sterilization conditions for Earth microorganisms should eradicate
297 microorganisms of similar size from Mars. There is no absolutely optimal
298 approach to decontamination under these circumstances, but enough is known
299 about the relationships among organism size, repair mechanisms, and
300 survivability, that the maximum survivability of any martian organisms can be
301 estimated with some confidence.

302

303 Whether we assume that life on Mars is based on the same building blocks as
304 terrestrial life, or on other covalently bonded complex molecules, only two methods
305 of sterilization are considered viable options at present—dry heat and gamma
306 radiation, either alone or in combination. These methods will penetrate the
307 sample and, therefore, provide the highest level of assurance that putative
308 organisms will be destroyed. It is recognized that the application of heat, and in
309 some cases gamma irradiation, will modify the geological properties of the

310 sample. Within reason, every effort should be made to develop and implement a
311 method of sterilization that protects the scientific integrity of the sample.

312
313 Many of the key parameters measured by geochemists are unaffected by sterilizing
314 representative geological samples with gamma radiation [Allen et al., 2000].
315 Gamma photons from ^{60}Co (1.17 – 1.33 MeV) in doses as high as 30 Mrads do
316 not induce radioactivity in rock and mineral samples. Such doses also produce no
317 measurable changes in isotopic compositions, elemental compositions, or
318 crystallographic structures. The only detectable effects are changes in albedo,
319 color, and thermoluminescence in selected minerals. Isotopic and elemental
320 compositions will not be affected regardless of gamma dose. Sterilization at
321 doses significantly above 30 Mrads may induce changes in crystallographic
322 structure (*caveat*: research required) and dose-dependent changes in albedo,
323 color and thermoluminescence may affect sample science. On balance, if
324 samples returned from Mars require biological sterilization, exposure to gamma
325 rays may provide a feasible option.

326
327 For the development of a final protocol for use with martian samples, a program of
328 research should be initiated to determine the effects of varying degrees of
329 treatment by heat and by gamma irradiation on organic compounds in rocky
330 matrices, and also on microscopic morphological evidence of life. This research
331 should be started well in advance of the return of the Mars samples, so that the
332 decontamination process can be designed to allow data obtained from analyses
333 of sterilized samples to be interpreted with minimal ambiguity and maximum utility
334 for the scientific purposes intended. Research should also be conducted to
335 determine the efficacy of various supercritical fluids and commonly used organic
336 solvents in killing model microorganisms, allowing the possibility that solvent
337 extracts might be safe to remove from containment without the damage to
338 dissolved biomarker compounds that would be caused by heat or ionizing
339 radiation. Whether decontamination is systematically achieved by any supercritical
340 fluids used in making extracts is a matter that must be investigated further, prior to

341 the removal of any such samples from the SRF. Also critical will be the
342 atmospheric conditions (gas mix, humidity) under which irradiation conditions are
343 qualified for use. Lethality of irradiation is enhanced by the presence of oxygen,
344 whether from O₂, H₂O, or other sources.

345

346 The aim of a sterilizing process is to reduce the risk of significant adverse effects
347 of samples distributed to the scientific community. The sterilization levels will be
348 defined to be such that the likelihood of adverse effects, given exposure to
349 humans, animals, and the environment, is less than 10⁻⁶. A suggested process
350 for sterilization consists of irradiation with gamma rays at temperatures up to
351 approximately 105°C [Bruch et al., 2001, page 5]. This procedure has the
352 advantage of being able to kill all known terrestrial organisms, while doing
353 minimal damage to the non-biologic constituents of the Mars samples.

354

355 The survival rate of a large number of terrestrial organisms exposed to ⁶⁰Co
356 gamma rays has been determined as a function of dosage, dose rate, and
357 temperature. There are no terrestrial organisms known whose probability of
358 survival is >10⁻⁶ at a dose of 20 Mrads at room temperature. Nonetheless,
359 populations of organisms may require higher doses to ensure that the probability
360 of finding any survivor is <10⁻⁶. The survival rate at a given total dose decreases
361 with increasing temperature during irradiation. For example, the sensitivity of dry
362 T1 bacteriophage to inactivation by X-rays increases, or the D₃₇ decreases by
363 approximately ten-fold between 60 and 105°C [Pollard 1953].

364

365 Protocol "Sterilization" Conditions A large number of geochemical tests will be
366 carried out in the SRF upon arrival of the samples. These tests will likely include
367 X-ray tomography to determine loci of cracks and other separations where life-
368 forms most likely would be, and total organic carbon (TOC), which permits a limit
369 on the density of carbon-based organisms to be calculated.

370

371 Irrespective of the chemical basis of any life-form, a confidence level of sterilization
372 can be provided with only two assumptions: 1) any reproducing life-form must be
373 based on macromolecules (i.e., polymers) with interatomic covalent bonds (not
374 crystal lattices), and 2) since all such bonds have similar strength, destroying
375 these bonds destroys the life-form.

376

377 Evidence shows that (at or near room temperature) 55 Mrads of radiation will
378 destroy almost all known bacteria, viruses, spores, and prions (i.e., the causative
379 agent in Scrapie) by 1 million-fold. Using 100 Mrads would give a 10-fold safety
380 margin. If worst-case estimates are used (10^6 – 10^{12} organisms/gram of martian
381 sample and a tiny target, such as a virus) sterilization would require 400 Mrads.
382 Even after this higher dose, most geologic studies may still be accomplished. This
383 amount of radiation could be safely reduced if the irradiation were carried out at
384 elevated temperature (e.g., 105°C), and/or if the TOC (or equivalent for non-carbon-
385 based organisms) is low enough to rule out large numbers of organisms being
386 present in the sample.

387

388 If martian organisms returned to Earth are similar to terrestrial organisms, a dose
389 of 20 Mrads at 105°C should reduce their number to $<10^{-6}$ of their initial number
390 (but not necessarily kill them all). It is not clear, however, that martian organisms
391 should be similar to terrestrial organisms; it is possible that they could be much
392 more resistant to gamma radiation. A good deal is known about the relationship
393 between the size and the biochemistry of terrestrial organisms and their
394 resistance to gamma radiation. For example, it has been shown that smaller
395 organisms tend to survive higher radiation doses, but the strategies used by
396 microorganisms to increase their resistance to radiation are not well understood.
397 It might, therefore, be a useful exercise to explore hypothetical possibilities for the
398 evolution of martian organisms adapted to the much higher radiation fluxes to
399 which they would be subjected naturally, compared to terrestrial microbes. The
400 radiation dose at various temperatures required to reduce the probability of the
401 survival of even a *single* organism below 10^{-6} per sample could then be estimated

402 and could become the basis of irradiation protocols for the sterilization of returned
403 Mars samples. In particular, tests should be made against radiation-tolerant
404 species like *Deinococcus radiodurans*, which possesses amazing radiation repair
405 capabilities [Daly 2000]. In such tests, it will be important to consider the
406 destruction of both the smallest and most hardy known Earth organisms, as well
407 as the destruction of non-living surrogates (such as viruses and viroids) that can
408 serve to provide effective sterilization doses for martian organisms that may be
409 smaller—as small as conceivably possible (see SSB 1999). Such surrogates also
410 can provide for the eventuality that, if Earth life and putative Mars life are somehow
411 related, the sterilization conditions will provide effective protection against martian
412 virus- or viroid-like entities that may be potentially hazardous.

413

414 **Criteria For Release**

415 As part of the charge to the recent NRC study of *The Quarantine and Certification of*
416 *Martian Samples* [SSB 2002], the Committee on Planetary and Lunar Exploration
417 (COMPLEX) was asked to study “What are the criteria that must be satisfied before
418 martian samples can be released from the facility?” The Committee’s
419 recommendations were weighed extensively in the derivation of the release criteria
420 given here. For the most part, their recommendations are incorporated in spirit, if
421 not in specific wording. Departures from the Committee’s report were the subject
422 of Workshop Series discussions, and were addressed in the review of the
423 Oversight and Review Committee. The departures are most obvious where the
424 NRC Committee made recommendations that were not fully consistent with their
425 own assumptions. An example of this is given in a footnote to the NRC report [SSB
426 2002, p. ES-5], which states that, “The word ‘life,’ when used in the context of
427 martian life, should always be understood to mean ‘Life as we know it,’ to allow for
428 the possibility of life-forms distinctly outside our terrestrial experience.” This is an
429 important footnote, but it has been noted that not all of the Committee’s release
430 criteria (for example, ‘no carbon equals no hazard’) were consistent with this
431 possibility. Additionally, COMPLEX’s recommendations place a heavy emphasis
432 on “sterilization” of Mars samples as a key to their release—yet the report states in

433 a number of places that the effects of sterilizing doses of heat and/or gamma
 434 radiation on the geochemical and biological signals the samples may carry are
 435 not known. Overall, the release criteria listed below are slightly more stringent, as
 436 well as somewhat more comprehensive, than those recommended by COMPLEX.

437
 438 Table 2 gives the basic overview of the questions that need to be answered prior to
 439 the release of unsterilized samples from the SRF. These questions will be asked
 440 of a representative sub-sample of the material returned from Mars.

441

Item	Question	Strategy
1	Is there anything that looks like a life-form?	Microscopy; beam synchrotron or other non-destructive high-resolution analytic probe, particularly one that would allow testing unsterilized (yet still contained) samples outside main facility.
2	Is there a chemical signature of life?	Mass spectrometer and/or other analytical measurement systems (to be used in containment) that would identify biomolecules, chiral asymmetry, special bonding, etc.
3	Is there any evidence of self-replication or replication in terrestrial living organism?	Attempts to grow in culture, in cell culture, or in defined living organisms.
4	Is there any adverse effect on workers or the surrounding environment?	Microcosm tests; medical surveillance of workers and monitoring and evaluation of living systems in proximity of receiving facility to ensure no release or exposure associated with operations of SRF.

442
 443 Table 2. Sequence of questions and possible strategies for decisions about release of
 444 sample material from containment.

445
 446

448 In any event, only evidence of measurable biohazards or active martian life-forms
 449 or their biomaterials should be regarded as relevant criteria for deciding whether
 450 to release any unsterilized samples (the specific release criteria are TBD).

451 Depending on results of Life Detection and Biohazard tests, remaining portions of
 452 samples will either be released for allocation outright, or sterilized and then
 453 released for allocation. Hence, the following criteria are intended to govern the
 454 release of samples evaluated using this Draft Protocol:

455 Protocol Release Criteria

- 456 ● No solid sample shall be released from containment in the Mars receiving
457 laboratory until it or its parent sample undergoes preliminary examination,
458 baseline description, cataloguing, and any necessary repackaging.
- 459 ➤ Samples to be used for Life Detection procedures or to be released from
460 containment will be screened for radioactivity and potential chemical
461 hazards.
- 462 ➤ Additionally, samples to be used for Biohazard testing will be screened
463 for known toxicity to bacterial and eukaryotic cells.
- 464 ● Samples containing any active martian form of life, *be it hazardous or not*,
465 will be kept under appropriate level of containment, or be thoroughly
466 sterilized before release.
- 467 ● Samples providing indications of life-related molecules, including proteins,
468 nucleic acids, or molecular chirality, will require more extensive testing,
469 including additional Biohazard testing, prior to their release.
- 470 ● Samples may be released if they are first subjected to a sterilizing process
471 involving heat, radiation, or a combination of these agents, to ensure they
472 are safe for analyses outside of containment. A sample that is 'safe' is
473 stipulated to be free of any viable self-replicating entities or entities able to
474 be amplified.
- 475 ● Samples may be released if Biohazard testing does not yield evidence of
476 live, extraterrestrial, self-replicating entities, or of harmful effects on
477 terrestrial life-forms or environment under Earth-like conditions.
- 478 ➤ Biohazard testing will involve assays for: 1) replication in media with
479 various organic and inorganic carbon sources, including enriched media
480 (liquid/solid), and sparse media appropriate to photo- or chemo-
481 autotrophs; 2) effect/growth on various cell cultures; 3) effect/growth on
482 whole organisms (i.e., murine/specified rodent; plant); and, 4) effect on
483 the ecosystem level.
- 484 ➤ Basic Biohazard testing will be required even in the absence of evidence
485 of organic carbon in a sample returned from Mars.
486


487 **Overview of the Draft Protocol**

488 The Draft Protocol has one basic purpose—to ensure that a representative set of
489 sub-samples undergoes sufficient testing to evaluate them against the release
490 criteria. Samples must be characterized, categorized, and analyzed to ensure that
491 they can be sorted according to a procedure providing ‘statistical relevance’ to any
492 sub-sampling (whether homogenized or pre-sorted for ‘biologically interesting
493 features’), within a reasonable time using a minimal amount of sample. Early
494 results in the Biohazard testing will need to be screened to ensure that potentially
495 chronic effects are not overlooked. The tests themselves should be performed in
496 an order that takes into account the relative harm posed by a potential biohazard
497 (e.g., to humans, animals, environments) and takes into consideration a variety of
498 routes of exposure and infection. Samples must be tested for biomolecules
499 (known or suspected), for other organic compounds, and for non-carbon evidence
500 of an active metabolism being present (e.g., alterations of sulfur, iron, or other
501 compounds). Life Detection and Biohazard testing partially overlap, and both will
502 depend on the processing of the samples and data from the Physical/Chemical
503 processes to evaluate their results and how to interpret them.

504

505 The Draft Protocol has three main segments: Physical/Chemical (P/C)
506 processing, Life Detection (LD) testing, and Biohazard (BH) testing. Figure 2 is a
507 simplified overview of how these segments are related. In this protocol, P/C
508 processing refers to all of the analytical testing and sample description that will be
509 accomplished prior to materials being tested for signs of life, or in support of
510 various forms of life and biohazard detection. LD testing is also mainly analytical
511 and descriptive. LD testing seeks signs of life in either morphology, chemistry, or
512 cultivation, as well as detecting a life-form in a manner that may be informed by
513 hypotheses about what signs of life a martian biota might leave. BH testing seeks
514 to challenge test sample materials against a variety of model systems to see if the
515 sample contains any hazardous properties that can be shown to be the result of a
516 self-replicating entity contained within the sample. BH testing should be as free as
517 possible from assumptions about the putative nature of a martian life form.

OVERVIEW: DRAFT MARS SAMPLE RETURN PROTOCOL



518
519
520

Figure 2. A simplified overview of the Draft Protocol showing the 3 main segments: Physical/Chemical processing, Life Detection, and Biohazard testing.


521 The overall process is as follows: the sample(s) will be removed from the Sample
522 Return Canister (SRC) under maximum biocontainment in gloveboxes containing
523 an inert gas atmosphere and housed within a combination cleanroom/biosafety
524 lab. After initial documentation, samples will undergo preliminary characterization,
525 splitting, and detailed examination using a variety of different methodologies.
526 Ultimately, data from LD and BH testing will be used to determine whether to
527 release materials from biocontainment. All sample materials not selected for
528 further testing will be archived in sealed containers in an inert atmosphere
529 glovebox within the lab for future scientific purposes. The Draft Protocol also
530 addresses issues related to facilities, personnel management, monitoring,
531 contingency planning, decision making, protocol review, implementation, and
532 approval processes.

533

534 **Physical/Chemical Processing**

535 The overall objective for P/C processing is to specify information about the
536 samples required to enable effective LD and BH testing, and curation. The focus is
537 on sample characteristics that could be determinative in understanding the results
538 of any *in vitro* and *in vivo* testing that may be required, as well as on information
539 needed for sample preservation purposes. P/C processing includes actions
540 affecting the returned samples between the time the SRC arrives in the SRF and
541 the time sample aliquots are apportioned for LD and BH tests. P/C processing
542 under this protocol should include only those actions required in support of
543 planetary protection and future sample utilization. Figure 3 outlines the proposed
544 P/C processing, which draws heavily from protocols proposed or used by others.⁹

9. This Draft Protocol is based on a framework developed at the first Workshop in this Series [Race and Rummel, 2000, p.14-19], and on an earlier report by MSHARP [Carr et al., 1999], which are, in turn, based on protocols developed at Johnson Space Center for handling and processing Apollo lunar samples, Antarctic meteorites, and cosmic dust. During the Workshop Series, modifications to the Draft Protocol were suggested by various sub-groups [Race et al., 2001a, 2001b, 2002], and many of those have been included here resulting in several significant differences from the framework developed in Workshop #1. In general, the proposed Draft Protocol is consistent with the requirements and conditions set forth by the Space Studies Board [SSB 1997], the MSHARP Committee [Carr et al., 1999], an earlier workshop on sample quarantine protocols [DeVincenzi et al., 1999], and CAPTEM [Neal, 2000].



545
546
547
548
549

Figure 3. The Physical/Chemical processing will occur in four sequential stages leading into the Life Detection and Biohazard testing. The numeric annotations refer to numbered sections of text below, which elaborate on the proposed P/C steps.

550 Principles The selected steps and investigations in the P/C processing tracks are
551 motivated by the following principles, as functions of the SRF: know what the
552 returned samples are; preserve sample integrity; document everything; anticipate
553 that different types of samples (e.g., gases, fines, rocks, and cores) require
554 different treatment; recognize that all data obtained in the P/C processing must
555 serve later scientific investigations; use the minimum sample possible; and
556 provide real-time guidance and adjustment to the process. These principles,
557 initially outlined by the report of the Mars Sample Handling and Requirements
558 Panel (MSHARP) [Carr et al., 1999], have been endorsed by all the Mars Sample
559 Handling Protocol Workshops [Race and Rummel, 2000; Race et al., 2001a;
560 Bruch et al., 2001; Race et al., 2001b; Race et al., 2002].

561

562 The first two principles (know the sample; preserve sample integrity) are, to some
563 extent, inconsistent because every characterization method or action on the
564 returned samples will affect them in some regard. This inconsistency has been
565 addressed in two ways. First, all characterization procedures in P/C processing
566 are nominally non-contact and non-destructive—all the sample mass remains in
567 the same physical and chemical state after each analysis. Second, most of the
568 returned sample is subjected to only minimal investigations, while only a
569 representative portion of the sample is subjected to more specific (and potentially
570 sample-altering) analyses. The P/C processing and screening methods, except
571 for weighing, involve sample interactions with electromagnetic radiation, principally
572 near-visible wavelengths (near ultraviolet, visible, and near infrared). Several
573 methods use X-rays to probe the samples, but it was recognized that X-rays can
574 (at some dosages) affect biological/organic systems.

575

576 This Draft Protocol attempts a compromise between the desire to affect only a
577 small proportion of the returned sample by planetary protection testing, and the
578 need to assure safety by testing all portions of all samples. A range of strategies
579 have been advocated to deal with the sample testing issue, from “characterize
580 everything with all available non-destructive methods,” to “store most of the

581 sample uncharacterized, and do only the minimum with the rest” (see discussions
582 in: Carr, et al., 1999, p. 37; Race and Rummel, 2000, p. 18; Race et al., 2001a,
583 p. 35; and Race et al., 2001b, p. 34). Here it is stipulated that it will be essential to
584 examine all the returned material in at least a minimal fashion to: confirm
585 spacecraft operations in sample transfer from Mars to the Sample Return
586 Canister; correlate returned samples with documentation developed by the
587 mission on Mars; and provide enough data to make informed choices about
588 samples for LD/BH analyses. Examining all returned materials in at least a
589 minimal fashion will help avoid a worst case scenario where an obviously
590 biogenic sample could be stored unexamined and only discovered after nominal
591 LD/BH tests were completed.

592

593 Documentation All treatments and actions with the returned samples need to be
594 documented fully. Without a high level of documentation, it would be impossible to
595 establish which samples are representative or particularly interesting, and to
596 indicate what had been done to which sample during processing.

597

598 Different Samples It is clear that the different types of samples will require different
599 processing techniques. Gases and bulk fines samples are expected to be
600 inherently homogeneous to some level, and will require only minimal processing
601 to derive characteristic and representative samples. However, solid materials are
602 anticipated to be potentially heterogeneous and more extensive study and real-
603 time decisions about their processing will be required.

604

605 Minimum Sample Mass The amount and size of returned Mars samples will be
606 small, and it will be desirable to subject sample materials to a great range of
607 biological, physical, and chemical tests. Thus, by necessity, each test on a
608 returned sample must use the minimum mass consistent with achieving the
609 scientific goal of the test.

610

611 Real-Time Adjustments – Oversight Committee Provisions must be made to
612 adjust the P/C processes in response to changing technology and mission
613 specifics, to monitor the processes in progress, and to adjust them in real-time to
614 fit the actual returned samples [Carr et al., 1999, pp. 7, 9]. This Draft Protocol is
615 being written more than 10 years before the nominal return of Mars samples to
616 Earth. We do not know the spacecraft configuration, the types of martian samples
617 that will be collected, their return configuration, and the exact nature of planetary
618 protection measures. Similarly, we cannot anticipate all of the advances in
619 instrumentation and analytical methods that are likely between now and the time of
620 sample return.

621

622 It is likely that the returned samples will not be exactly as we imagine them now,
623 and may include materials that are complex (e.g., breccias) or unusual
624 (e.g., a possible stromatolite fossil). Treatment of these types of samples must be
625 sample-specific, and cannot be defined in advance. Thus, there must be a
626 mechanism such as an SRF oversight committee to adjust the final protocol to fit
627 the actual samples.

628

629 Assumptions In preparing the P/C portion of the Draft Protocol, the mission profile
630 and constraints outlined in the initial *Assumptions* of the Workshop Series [see
631 *Appendix A*] were adopted. It is worth reiterating here a few of the key assumptions
632 which hold particular relevance to physical chemical processing: the SRCs will be
633 received at the SRF free of exterior contamination with Mars materials, intact, and
634 with no breaches of containment (see page 96); the returned samples will include
635 gas, fines material (bulk regolith), and solids; the total mass of all samples is
636 expected to be ~ 500 to 1000 grams.

637

638 Overview of Physical/Chemical Processing Physical and chemical processing
639 comprises the priority actions taken concerning the returned Mars samples
640 between arrival of the SRC at the SRF, and initial examination for hazards and the
641 LD/BH testing of fines and solids. These anticipated steps in P/C processing are

642 shown schematically in Figure 3, which is based on portions of Figures 6-2 and
643 6-3 of Carr et al. (1999), Figure 2 on page 18 of Race and Rummel (2000), and the
644 narrative of Race et al. (2001a). The numeric annotations in Figure 3 refer to
645 similarly numbered sections of text below, which elaborate on the proposed P/C
646 processing steps in narrative form.

647

648 P/C processing can be divided into three phases in roughly sequential order:

- 649 ● Pre-processing, before preliminary examination of the samples;
- 650 ● Preliminary examination and screening of gas, fines, and solids, to permit
651 informed choices about samples for later detailed testing, banking, or
652 curation; and,
- 653 ● Sub-division of samples selected for Life Detection and Biohazard tests.

654

655 Following P/C processing, Life Detection and Biohazard testing will begin. Those
656 processes may require information developed during preliminary examination and
657 screening, and may also require subsequent and more detailed information of a
658 physical or chemical nature; these additional analyses are not included here as
659 they are contingent upon the results of the Life Detection and Biohazard testing.

660

661 The steps of preliminary examination and screening were judged to be different for
662 three types of samples: gases, homogeneous particulate samples, and
663 inherently inhomogeneous samples like rocks, rock cores, and regolith cores.

664 Each of these sample types will follow a different track through preliminary
665 examination and screening as described in the text below and shown on Figure 3
666 as the 'Gases Track,' 'Solids Track,' and 'Fines Track.'

667

668 Pre-processing Samples

- 669 ● 1.0 Pre-Processing Steps. Pre-processing steps outlined here are those
670 between arrival of the SRC at the SRF, and initial examination of gas, fines,
671 and solids. Pre-processing steps refer to cleaning and decontaminating the
672 exterior of any containers holding samples, as well as the initial steps in

673 each of the gases-, fines-, and solids-tracks involving opening containers
674 and removal of samples.

675 ● *1.1 Clean and Decontaminate Exterior of SRC.* It is imperative that the
676 exterior of any sample return containers or vessel(s) carry no terrestrial
677 microbes, and are organically clean. (It is assumed that the exterior of the
678 SRC is not contaminated with martian materials.)¹⁰ If these states are not
679 achieved, all subsequent analyses for life or biohazard are severely
680 compromised. Actual methods of cleaning and decontamination are to be
681 determined. An interesting new method to be considered is laser ablation of
682 the SRC exterior.

683 Procedures for opening sample containers are mission specific as to
684 number, types, and contents of containers. At a minimum, we assume that
685 some solid materials with surrounding gas will be in the container(s). It is
686 recommended that the gas be extracted for separate treatment, and that the
687 solid samples be contained thereafter in an inert gas, such as dry nitrogen.

688 ● *1.2 Extract Head Gas and Back-fill.* The returned solid samples will arrive
689 on Earth with some gas surrounding them. Presumably, this “head gas”
690 would consist originally of martian atmosphere. By the time of arrival on
691 Earth, the gas might have been affected by chemical and physical reactions
692 with the solids (rock and soil), by out-gassing from the solids (especially if
693 the temperature rises above 25°C during return), and possibly by biological
694 activity in the sample. This gas may contain information important to
695 understanding the thermal, chemical, and biological histories of the solid
696 returned samples. Therefore, extraction and analysis of the head gas is a
697 high priority.

698 In this step of pre-processing, the head gas would be extracted from the
699 SRC, and the SRC back-filled with a chemically unreactive gas to ambient
700 “room” pressure. Exact procedures for extraction and back-filling will
701 depend on the SRC design and construction, but might (for instance)
702 include puncturing the SRC at an intentional thin point, extracting the head

10. It should be noted that planetary protection requirements will exist for a Mars Sample Return (MSR) Project to assure that the sample return container(s) is(are) intact and free of exterior contamination with Mars materials when delivered to the Sample Receiving Facility. Compliance with these requirements is the responsibility of the MSR Project Office and, therefore, not a function to be included in this protocol, which begins at the point of opening that clean and intact container.

703 gas to a pre-determined vacuum pressure, and refilling the SRC with dry
704 clean N₂ gas. The extracted head gas would be processed as set forth
705 below (see 2.0 – 2.2 *Gases Track*).

706 Three issues related to gases were identified for further consideration and
707 possible research: 1) the effects of vacuum and non-martian gas on the
708 chemical properties of the sample; 2) the effects of vacuum and non-
709 martian gas on any live martian biota; and 3) the effects of extraction on gas
710 isotope ratios.

711 For the first issue, experience with curation of the *Apollo* lunar samples has
712 shown that few geochemical and other inorganic investigations are
713 materially affected by holding and processing the samples in dry N₂ gas at
714 1 bar. Of course, the lunar samples originated at hard vacuum on the Moon.
715 It is not clear what changes might be wrought on returned Mars samples
716 (possibly containing clays or other hydrous materials) by first vacuum
717 pumping, and then immersion in dry N₂ gas; further research is required in
718 this area.

719 For the second issue, there is reason for the returned solid samples to be
720 treated under an atmosphere as near to martian as possible, i.e., both to
721 preserve key geochemical signatures [*Neal, 2000, p. 22492ff*], and to
722 maintain possible microorganisms in their native environment. It is
723 unknown whether live martian organisms could be killed by removal of
724 0.006 bars of CO₂ and then immersion in 1 bar of N₂, and there may not be
725 comparable terrestrial biota to test. Some samples eventually will be
726 subjected to higher pressures merely because the biota of BH tests would
727 not survive in martian atmosphere. On the other hand, there are serious
728 problems in sample handling and geochemistry that would be caused by
729 immersing the samples in a model martian atmosphere. Sample handling
730 and LD/BH testing at reduced pressure (the near vacuum of 0.006 bars
731 CO₂) present severe problems. Sample handling under vacuum was
732 attempted during the *Apollo* program with lunar samples, and was found to
733 be extremely difficult, expensive and contaminating (e.g., mercury or oil from
734 vacuum pumps). Similarly, back-filling the sample container with a relatively
735 reactive gas like CO₂ would change the isotopic nature of the sample.
736 Terrestrial carbon and oxygen will exchange with the sample and
737 compromise biological and geochemical inferences from these two stable

738 isotope systems. This is an area of future research and discovery. One
739 possible approach would be to backfill the SRC and perform sample
740 handling and examination, where possible, under 1 bar of dry N₂ gas with
741 0.006 bars of CO₂ added. This might satisfy the constraints of easy sample
742 handling, while being consistent with the desire to not kill live martian
743 organisms, if any, and should be considered for the final protocol.

744 For the third issue, it is known that the elemental and isotopic ratios of a
745 gas sample can be fractionated during transfer from one reservoir to
746 another. With the head gas in contact with the abundant surface area of the
747 returned samples, fractionation could become a serious potential problem.

748

749 Gases Track

- 750 ● *2.0 Gases Track.* Gas withdrawn from the SRC, the “head gas,” will be
751 processed by filtering and subsequently split for Life Detection and
752 Biohazard testing and would be available relatively rapidly for other
753 investigations [*Race and Rummel, 2000, p. 17*].
- 754 ● *2.1 Filter to <TBD Nanometers.* During or after removal of the head gas
755 from the SRC, the gas should be filtered to remove particles [*Race and*
756 *Rummel, 2000, p. 17*]. The purpose of filtering the head gas is to remove
757 objects that could reasonably constitute viable organisms, or that might
758 present biohazards. The size of objects passing the filter is to be
759 determined. Sizes suggested by sub-groups in the Workshop Series have
760 ranged from <0.5 μm [*Race et al., 2001a, p. 34*] to <0.02 μm [*Race et al.,*
761 *2001b, p. 27*], both of which are realizable with current technology (currently,
762 some methods are rated to remove particles larger than 0.003 μm). It is not
763 clear if filtering could change the chemical or molecular composition of the
764 head gas, for instance by preferential adsorption of heavy noble gases or by
765 catalysis of reactions; this also requires additional research.
- 766 ● *2.2 Distribute in Sealed Containers.* Filtered head gas should be released
767 from the SRF and distributed in sealed containers. Unlike the returned solid
768 samples (rock, regolith, etc.), a returned gas sample is only useful for
769 investigation if it is contained. Typically, a gas sample like this would be
770 placed in a glass bulb, which would then be sealed by melting the stem of
771 the bulb. Containment at PPL-α or PPL-β levels is inherent in the

772 combination of filtration and this procedure. The filtered gas will be available
773 for immediate allocation from the SRF without further processing or
774 sterilization.¹¹

775

776 Solids Track

777 ● *3.0 Solids Track.* After removal and filtering of the SRC head gas, the
778 remaining returned samples would be solids of various types, i.e., regolith
779 samples, rocks, rock cores, soil cores, and fines. The specifics of this solid
780 sample set are to be determined during mission design. These solid
781 samples will be processed through two separate tracks, *Solids Track (3.0)*
782 and *Fines Track (4.0)*, for basic documentation, further preliminary testing,
783 and selection for subsequent LD and BH tests.

784 Some principles of this P/C process are worth restating here. The P/C
785 process is a method to obtain the minimum data needed to characterize the
786 samples adequately and to permit selection of suitable samples for LD/BH
787 tests. The remaining samples will be preserved and made available for
788 subsequent investigations and analyses. The samples will be changed as
789 little as possible from their original state.

790 The martian samples will only be touched by or come in contact with a
791 limited set of materials under controlled temperature, pressure, humidity,
792 and atmospheric conditions. Pristine lunar samples are touched only by
793 stainless steel, aluminum, and Teflon™; these might also be suitable for
794 returned Mars samples. Neal cites the considerations, from a geochemical
795 perspective, for choices of materials for sample handling and suggests
796 several types [Neal, 2000]. Whether these materials are appropriate for
797 returned martian samples should be determined through additional
798 research with Mars simulants prior to sample return.

799 The temperature of processing is TBD, and will depend in great part on
800 technical mission constraints. The implicit assumption here has been that
801 the temperature of processing will be between 0°C (273K) and ambient

11. To date, no decisions have been made about when and under what conditions sample materials will be eligible for release from containment at the SRF. Ultimately, it is likely that decisions about what is done with sample materials will be made after review by an appropriate international scientific oversight committee at the SRF in consultation with NASA's Planetary Protection Officer and other responsible officials.

802 (~298K), for which the protocols and experience with the *Apollo* samples
803 are relevant. On the other hand, it will be important from geochemical and
804 biological perspectives to maintain the returned sample at its ambient
805 martian temperature, ~240K [Carr et al., 1999; Neal, 2000]. This
806 temperature may not be possible within mission constraints, and there
807 appears to be no compelling reason to process at temperatures
808 significantly below those experienced by the samples during their transit to
809 Earth. It is not clear, at this point, what problems and attendant costs would
810 be associated with sample curation and processing at sub-freezing
811 temperatures.

812 It is suggested that an atmosphere of 1 bar of unreactive gas be used in
813 processing, curation, and back-filling of the SRC. The steps outlined below
814 assume that processing and curation will take place under 1 atmosphere of
815 a pure unreactive gas (e.g., N₂). It is not known whether this gas would
816 present problems for the LD and BH testing procedures. The composition
817 and pressure of the atmosphere has implications for biological and
818 geochemical testing, and is an area of concern (see sections 1.2, 5.0, and
819 “Future Research”). It must be recognized that a requirement for processing
820 at low pressure, like the atmosphere of the martian surface (0.006 atm),
821 would have significant implications for the design and cost of a SRF.

- 822 ● 3.1 *Open SRC and Remove Samples*. The SRC must be opened to
823 retrieve and remove the solid samples. The procedures for opening the
824 SRC and removing the samples are to be determined and will depend
825 largely on the design of the SRC.
- 826 ● 3.2 *Preliminary Examination and Documentation*. As part of the P/C
827 processing, *Preliminary Examination and Documentation* includes the
828 minimal investigations deemed critical to an understanding of the nature of
829 the returned sample, and to support initial biohazard investigations [Race
830 and Rummel, 2000, pp. 14, 17; Race et al., 2001a, p. 37].

831 The first material-hazard investigation is a measurement of sample
832 radioactivity. Some forms of ionizing radiation can penetrate the curation
833 barriers between the returned sample and human processors. The
834 purpose is not to measure abundances of indigenous radioisotopes
835 (e.g., ²³⁸U), nor cosmogenic radioactivities (e.g., ²⁶Al), but rather to
836 determine whether radiation levels associated with the samples could pose

837 a threat to workers at the SRF. Biohazard radioactivity can be measured on
838 the bulk returned sample (safety level TBD), and need not be measured on
839 individual samples unless the bulk presents a radiation biohazard. Only
840 gamma radiation need be measured, as beta and alpha radiation will not
841 penetrate the barriers between the returned samples and human
842 processors. Based on prior experience with martian materials in
843 meteorites, it is considered unlikely that returned martian samples will
844 present a radiation safety hazard.

845 Imaging provides the first critical documentation of the returned sample
846 [*Race and Rummel, 2000, p. 17*]. Imaging at this stage serves multiple
847 objectives: verification of mission success; correlation of specific samples
848 with images of them taken on Mars and their sources; documentation of
849 physical effects of transport to Earth (e.g., fracturing, disaggregation);
850 preliminary identification of rock types; and measurement of sample
851 volumes. It is anticipated that the returned samples would be imaged at a
852 high spatial resolution (TBD, perhaps ~0.1 millimeter per pixel), over a
853 range of perhaps seven to nine different wavelengths TBD, with at least
854 three or four in the visible. These data will be critical to understanding the
855 nature of the returned sample, and in processing and selection of samples
856 for Life Detection and Biohazard tests.

857 The sample masses should be measured at this stage, and each time a
858 sample is cleaned, split, or allocated. Measurement of mass is important
859 as a mission requirement, for sample tracking and curation, and in
860 allocating suitable samples for LD/BH testing. For instance, it is likely that a
861 given mass of martian material would be returned to Earth as a mission
862 requirements, and weighing at this stage will determine if that mission
863 requirement has been fulfilled.

864 ● **3.3 Separate Rock Fragments and Cores From Fines.** At this stage of
865 processing, the solid samples would be separated into larger and smaller
866 fragments. The larger samples would include drill cores, whole rocks, and
867 rock fragments or rocklets (equivalent to the *Apollo* “coarse-fines”).¹² The

12. The terminology used to refer to small rocky materials has varied from workshop to workshop in this Series. The terms rock fragments, rocklets, and pebbles have been used to identify a general class of solid material that is distinct from fines, larger rocks, or rock cores. In addition to determining cut-off sizes at some later date, it will be necessary to use consistent terminology in all parts of the protocol.

868 smaller samples would include unconsolidated regolith, atmospheric dust,
869 and dust generated by coring operations. This separation is necessary
870 because the larger fragments cannot be treated as homogeneous
871 powders, and must be examined individually for Life Detection and
872 Biohazard analyses. It is possible that the regolith samples will include
873 small rocks and rocklets, comparable to the case with the lunar regolith
874 samples returned by the *Apollo* missions. As with *Apollo*, the small rocks
875 and rocklets would be separated from the finer material, cataloged, and
876 curated individually throughout subsequent processing and analyses.
877 The cut-off size for rock fragments or rocklets remains to be determined.
878 The standard cut-off size in the soil science community is greater than
879 2 millimeters. Sub-groups in the Workshop Series have suggested sizes
880 ranging from greater than 1 millimeter to greater than 2 millimeters, and
881 even "... greater than several millimeters ..." for martian samples [*Race et*
882 *al., 2001a, p. 34; Race and Rummel, 2000, p. 17*]. Decisions about cut-off
883 sizes for different classes of solid materials will be made when the sample
884 is returned and first examined, based on a recommendation of the SRF
885 Oversight Committee (see *Personnel Management Considerations* later in
886 this document).

887 Given the dusty nature of the martian surface, and the likelihood of dust
888 generated during coring, it is anticipated that the surfaces of cores and rock
889 samples will be coated with fine-grained materials. After separation,
890 preliminary examination, and documentation of the returned solid materials,
891 it will be necessary to remove dust from surfaces of the cores, rocks, and
892 rocklets [*Race et al., 2001b, p. 22*]. These fine materials constitute distinct
893 samples of martian material, and will require different processing and
894 curation than the solids (i.e., the fines track). In addition, the fine materials
895 on solids likely will hinder identification and processing of the latter by
896 obscuring their surfaces. Selection of samples for Life Detection and
897 Biohazard assays will require knowledge of the mineralogy, structure, and
898 textures of the samples. The analytical probes available (primarily visual
899 and near-infrared optics) will be unable to operate effectively on dust-
900 covered samples.

901 The exact methods of fines removal are TBD. Suggested methods have
902 included vacuuming the samples, blowing the dust off, a combination of

- 903 vacuuming and blowing, and laser desorption. In all these cases, thought
904 needs to be given to how the fines will be collected after removal. The fines
905 collected from each solid sample would be identified individually, and
906 treated as a separate fines sample within the *Fines Track*, as described in
907 section 4.0 below.
- 908 ● **3.4 Sort to Groups.** After removal of adhering fines, the solid samples
909 should be sorted into groups of similar materials using visual clues and
910 information from *Preliminary Examination* data [Race and Rummel, 2000,
911 p. 17; Race et al., 2001a]. This step assumes that the returned sample will
912 contain several cores and/or multiple millimeter-sized rock fragments
913 (“rocklets”). Criteria for sorting would include size, rock type (including
914 color), grain size, texture, and other readily observable properties. This
915 sorting is an important first step towards selecting representative samples
916 for Life Detection and Biohazard tests [Race et al., 2001a, p. 26].
 - 917 ● **3.5 Pristine Bank.** Samples and sub-samples that are not chosen at this
918 point for *Further Screening* and/or for Life Detection and Biohazard tests will
919 be stored in a *Pristine Sample Bank* [Race and Rummel, 2000, p. 17]. This
920 “bank” will serve as a containment system designed to maintain the
921 physical/ chemical, and biological integrity of samples while they await
922 allocation for other analyses at a later date. According to recommendations
923 by the Curation and Analysis Planning Team for Extraterrestrial Materials
924 (CAPTEM), the “bank” should hold the samples under an inert atmosphere
925 at temperatures below 240K [Neal, 2000]. The pristine solid samples are
926 those that have been affected by no procedures beyond those of preliminary
927 examination, dust removal, and sorting. The pristine bank will serve the
928 critical purpose of preserving a portion of the returned sample for analyses
929 beyond and after the Life Detection and Biohazard assays associated with
930 planetary protection. The pristine bank samples will become the principal
931 resource for all subsequent chemical, geological, physical, and biological
932 analyses on the returned samples.
 - 933 ● **3.6 Further Screening.** At this point, sub-samples of each rock type group
934 sorted previously (see section 3.4 above) would be subjected to additional
935 analyses in support of (and preliminary to) Life Detection and Biohazard
936 tests [Race and Rummel, 2000, p. 14; Race et al., 2001a, p. 37]. The exact
937 analyses needed are to be determined in conjunction with the detailed

938 LD/BH tests (see *Future Research*, below). Whenever possible, selected
939 analyses should emphasize non-destructive methods that are not likely to
940 modify or destroy biological molecules or biohazards, and would not be
941 anticipated to kill or weaken live martian organisms. Once the tests are
942 defined, it will be possible to learn what characteristics of the returned
943 samples might affect or interfere with particular tests, and what data are
944 essential prior to the tests. With this information in hand, the *Further*
945 *Screening* analyses can be tailored to meet the requirements of life and
946 biohazard detection. Given these restrictions and uncertainties, the
947 following screening methods have been suggested:

- 948 ► Multi-spectral imagery of the samples in visible, near-infrared, and/or
949 thermal infrared light can provide identification of the minerals (inorganic
950 chemical compounds) and the presence and distributions of organic
951 matter and water (molecular and bound) in the sample. Raman
952 spectroscopy should be considered here, also, with the caveat that
953 samples can experience significant heating during Raman analysis. For
954 instance, 514.5 nanometer green light from an argon laser is absorbed
955 significantly more than 1064 nanometer infrared light from a Nd:YAG
956 laser. Heating can also be mitigated by distribution of laser power in
957 space and time over the sample. The distributions of minerals on the
958 samples' surfaces will be crucial clues to understanding their internal
959 structures. X-ray diffraction analysis would also be valuable in defining
960 the minerals in the samples (see *Race et al., 2001a*, p. 35ff, for more
961 detail on these methods.)
- 962 ► It is important to know the internal structures of the samples (especially
963 the larger ones), because biogenic material could reasonably be
964 concentrated in cracks and open spaces (analogous to terrestrial
965 endolithic organisms). Building on the multi-spectral imagery,
966 tomographic analyses could provide three-dimensional visualizations of
967 the internal structures of the samples. Among tomographic methods, the
968 most developed at present is X-ray tomography. To provide X-ray
969 tomographic maps of density (i.e., continuum absorption of X-rays) now
970 requires only a bench-top instrument. X-ray tomographic maps for
971 individual elements like carbon require at present the X-ray intensity of a

972 synchrotron light source, and is considered impractical for this *Further*
973 *Screening* step.

974 ➤ Abundances and distributions of major elements and several minor
975 elements will likely be important for sample selection in Life Detection
976 and Biohazard analyses. It is also possible that abundances of certain
977 elements could produce false positives or negatives on Life Detection
978 and Biohazard tests. A likely method for elemental analysis is X-ray
979 fluorescence, a mature technique used routinely in inorganic
980 geochemistry and studies of human bone composition.

981 ➤ It would be very useful at this stage to have bulk analyses for carbon as a
982 guide to sample selection. However, a non-destructive test for bulk
983 carbon that is sufficiently precise, and has low enough detection limits to
984 be useful here, has not been identified; this requires future research.

985 ● *3.7 Selection of Sub-samples.* Representative sub-samples will be
986 selected for Life Detection and Biohazard tests based on data from the
987 *Further Screening* tests (see section 3.6). The remaining unselected
988 samples will be stored in the *Returned Sample Bank* (see section 3.8) for
989 future research access. Additional research will be required to define
990 representative sample and sub-sample criteria for all martian materials in
991 light of a potential for extreme heterogeneity of rock and soil samples, and a
992 concomitant likelihood that putative biohazards may be limited in terms of
993 location. Selected samples will carry forward to the actual Life Detection and
994 Biohazard investigations (see section 5.0).

995 ● *3.8 Returned Sample Bank.* The *Returned Sample Bank*, distinct from the
996 *Pristine Sample Bank* (see section 3.5), is for storage of samples that have
997 experienced the analysis of *Further Screening*, but have not yet been
998 allocated for Life Detection and Biohazard tests. These returned samples
999 should be labeled and kept distinct from the pristine samples, as the former
1000 have had more chance for contamination than the latter.

1001

1002 *Fines Track*

1003 ● *4.0 Fines Track.* Fines samples are those with particle sizes smaller than
1004 some limit TBD; the size limit suggested in the MSHP Workshop Series
1005 was 1 or 2 millimeters [*Race and Rummel, 2000; Race et al., 2001a,*

1006 2001b]. In either case, it is anticipated that fines samples will contain so
1007 many grains, mixed homogeneously, that it will be readily possible to take
1008 representative splits for Life Detection and Biohazard tests. Fines samples
1009 may include materials from a variety of sources: material collected as such,
1010 like dust from a wind-deposited dune; regolith that has had coarser material
1011 removed (see section 3.3); dust filtered out of the SRC headspace gas (see
1012 section 2.1); or particulates removed from surfaces of rocks or cores (see
1013 section 3.3).

1014 ● **4.1 Characterization.** Characterization of fines samples would be limited to
1015 imagery of each bulk fines sample (possibly including multi-spectral
1016 imagery) and weighing of each bulk sample [Race et al., 2001a, p. 35].
1017 There is no need to image or otherwise characterize each individual particle
1018 within a bulk fines sample. Only these minimal analyses are needed to
1019 document each fine sample at this stage in order to select samples or
1020 representative sub-samples for Life Detection and Biohazard assays. Each
1021 fines sample may be subdivided into fragments larger and smaller than
1022 1 millimeter [Race and Rummel, 2000], but the desirability of this further
1023 splitting is an area requiring additional research.

1024 ● **4.2 Split for LD/BH Tests and Banking.** At this point in P/C processing, fines
1025 samples would be selected for Life Detection and Biohazard tests, and split
1026 into representative aliquots. Some aliquots would be carried forward to Life
1027 Detection and Biohazard tests (see section 5.3), and some would be
1028 reserved in the 'Pristine Sample Bank' (see section 3.5). Since additional
1029 chemical analyses will be included as part of the LD/BH testing, no
1030 separate elemental analyses will be conducted on fines at this point in the
1031 P/C processing.

1032 The methods for splitting the fines samples are TBD. Methods used in
1033 typical terrestrial applications (e.g., riffle splitter, or coning-and-quartering),¹³
1034 may not be appropriate or practical here [Race et al., 2001a, p. 14]. First,
1035 these methods will involve considerable contact between and among the
1036 sample, tools, and surfaces, and may be deemed too contaminating.

13. A riffle splitter is a mechanical separation device that is able to split an unconsolidated soil sample into two equal parts that have the same grain size distribution (and presumably composition) as the parent sample. Coning-and-quartering is another commonly-used separation method (as described in Maxwell 1968).

1037 Second, both methods have the potential for considerable loss of sample
1038 through embedding in metal surfaces or electrostatic adhesion to metal
1039 and plastic surfaces. The electrostatic adhesion problem will be
1040 exacerbated in the dry atmosphere of the PPL- α spaces, as has been found
1041 with curation of lunar samples. In fact, neither method is now used for
1042 splitting lunar fines samples. This clearly is another area of required
1043 research.

1044 In this Draft Protocol, it is assumed that a sub-sample of fines is
1045 representative, based on confirmation of an adequate splitting method.
1046 However, it is suggested initially [Race *et al.*, 2001, p. 14] that each sample
1047 of fines be split into multiple sub-samples, each of which should be
1048 analyzed for bulk composition and mineralogy (as under *Further Screening*,
1049 see section 3.6) to determine whether splits are homogeneous. Further
1050 consideration of this issue is needed.

1051

1052 *Preparation for Life Detection and Biohazard Testing*

- 1053 ● *5.0 Samples for Life Detection and Biohazard Testing.* At this point,
1054 samples have been selected for LD/BH tests as well as other P/C analyses.
- 1055 ● *5.1 Split into Representative Sub-samples for LD/BH.* The samples
1056 selected for LD/BH tests will be split into representative sub-samples at this
1057 point. This splitting is necessary to ensure that analyses are performed on
1058 similar materials, and so that the results of one test may be reasonably
1059 correlated with the results of another. Splits chosen for immediate analysis
1060 will proceed to various LD/BH tests (see section 5.3 below). Some splits
1061 will be held in reserve as part of the *Return Sample Bank* as described in
1062 section 5.2. below.
- 1063 ● *5.2 Reserve.* Some splits from section 5.1 will be held in reserve for LD/BH
1064 tests, in anticipation of future needs. Should a test fail or require repetition,
1065 this reserve material would be available. These reserve splits could
1066 reasonably be kept in the '*Return Sample Bank*,' but labeled accordingly.
- 1067 ● *5.3 Parallelism of Tasks.* It is beyond the scope of the P/C procedure to
1068 describe the actual operation of LD/BH analyses and supporting inorganic
1069 analyses. However, they are included on Figure 3 for completeness. It is
1070 anticipated that these three types of tests will be run in parallel, with the

1071 results of each influencing the interpretation and course of the other tests
1072 [Carr et al., 1999, p. 9].

1073

1074 Future P/C Research and Development Needs In the discussions of P/C

1075 processing of the returned martian samples, several areas were identified where

1076 data were not available or could readily be obtained without additional research.

1077 Each research suggestion discussed below is keyed to the particular numbered

1078 text section above, where it is called out:

1079 ● Exactly what analyses and data do the LD/BH analyses require from the P/C
1080 processing? (see sections 3.2, 3.6, and 4.1). The P/C process here reflects
1081 informed judgment about which analyses would be most useful in LD/BH
1082 studies, but it will be very important to know what information about sample
1083 characteristics, or about the particular P/C processing, will be useful when
1084 assessing LD/BH results (for example, to determine possible causes of
1085 false positives or negatives; to document abundances of specific elements
1086 of interest (e.g., arsenic) or minerals (e.g., saponite clay); or to characterize
1087 surface reactivity and constituents (e.g., super-oxidants, etc.).

1088 ● In implementing the final protocol, there must be close collaboration
1089 between biohazard, toxicology, and pathology disciplines on the one hand,
1090 and chemistry, biochemistry, geochemistry, physics, and geophysics, on the
1091 other, to coordinate a truly integrated testing outcome, pursuant to
1092 augmenting which physical sciences data should be ruled in or ruled out in
1093 ultimate interpretations of sub-sample biohazard and/or toxicity testing.

1094 ● Trial-testing initiatives should be developed before the protocol is fully
1095 implemented in a sample return mission. These trials should be
1096 refinements that take into account the prospective chemical and physical
1097 properties of martian soil and rock(s) (and/or use martian surrogates where
1098 applicable), as well as evaluate biohazard containment facility needs.

1099 ● Is there added value in separating each fines sample into grain size
1100 separates [Race and Rummel, 2000, p. 17]? What additional contamination
1101 might be introduced by this procedure? (see section 4.2)

1102

- 1103 ● How can one remove terrestrial contaminants (including organics) from the
1104 exterior of the SRC before it enters PPL- α space? Laser ablation surfacing
1105 was suggested and should be studied (see section 1.1).
- 1106 ● How can one effectively remove and collect dust and other fines from the
1107 surfaces of rocks and rock cores? (see section 3.3) Three suggestions
1108 were vacuuming, blowing with compressed gas, and laser desorption.
- 1109 ● What effects do X-rays have on biological structures and molecules?
1110 Several analytical methods involve interaction of X-rays with the samples
1111 (e.g., XRD, XRF, XR tomography), and it is not known whether these X-ray
1112 doses interacting with Mars samples would affect LD/BH analyses (see
1113 section 3.6).
- 1114 ● How can one analyze a bulk sample for trace or ultra-trace quantities of
1115 carbon, non-destructively and without anticipated deleterious effects on
1116 biological molecules or viable organisms? (see section 3.6)
- 1117 ● Is the chemical composition of the head gas affected by filtration to remove
1118 small particles? (see section 2.1)
- 1119 ● What chemical and physical effects would removal of head gas and
1120 replacement with dry nitrogen have on the returned martian samples? (see
1121 section 1.2)
- 1122 ● What chemical effects would removal of head gas from the returned sample
1123 canister have on the gas itself? (see section 1.2)
- 1124 ● What effects would removal of head gas and replacement with dry nitrogen
1125 have on live martian and any contaminating terrestrial organisms in the
1126 returned martian samples? Would these effects be mitigated if samples
1127 were curated under dry nitrogen with 0.006 bars of CO₂ gas? (see section
1128 1.2)
- 1129 ● What effects would gas with terrestrial carbon and oxygen isotope ratios
1130 have on live martian organism in the returned martian sample? Would live
1131 martian organisms ingest the terrestrial carbon and oxygen, and become
1132 isotopically indistinguishable from terrestrial organisms? (see section 1.2)
- 1133 ● How can one produce representative splits of martian dust and fines
1134 materials without unacceptable contamination or loss of sample? (see
1135 section 4.2)

- 1136 ● How can one confirm that splits of dust or fines material are representative
1137 before Life Detection and Biohazard analyses, or is such confirmation
1138 necessary? (see section 4.2)
- 1139 ● What are the overall requirements and statistical test methods necessary to
1140 ensure that a representative sub-sample of rock and soil material is
1141 available for further LD and BH testing?
- 1142 ● Using artificially constructed Mars simulants, determine whether materials
1143 and conditions recommended by CAPTEM [Neal, 2000] are appropriate for
1144 handling martian samples. (see sections 3.0 and 4.0)
- 1145 ● Petrographic thin sections are enormously valuable in characterizing the
1146 minerals, structures, textures and history of a rock. Can petrographic thin
1147 sections be produced in a manner consistent with the principles of minimal
1148 sample use and minimal contamination of the section material and the
1149 remaining sample? (see section 5.3)

1150

1151 Areas of Concern Several areas of serious or general concern have been raised
1152 during discussions of physical and chemical processing. These issues, listed
1153 below, are significant enough to affect mission design, and SRC and SRF design.

- 1154 ● The validity and significance of Life Detection and Biohazard procedures in
1155 the SRF are strongly dependent on sample collection procedures on Mars,
1156 and thus on spacecraft and mission design. How can the Life Detection and
1157 Biohazard teams influence the designs of sample return spacecraft and
1158 sample collection procedures?
- 1159 ● What if the return sample container is breached or its seal is compromised?
1160 What contingency plans are possible to achieve PPL- α containment and
1161 biosafety? (see *Assumptions*, Appendix A)
- 1162 ● Is measurement of sample mass important as a preliminary
1163 characterization step? Should it be deferred until the “Further Screening”
1164 step? (see sections 3.2 and 3.6)
- 1165 ● How is the head gas to be removed from the SRC without contamination? Is
1166 backfill with non-reactive gas justifiable in terms of possible effects on
1167 martian biology? Would it be adequate or preferable to backfill with 6 mbar
1168 of terrestrial CO₂ and the remainder a non-reactive gas? (see section 1.2)

- 1169 ● What should be done if a unique critical sample is smaller than the nominal
1170 requirements for LD/BH analyses? (see section 3.4)
- 1171 ● What should be done if the requirements for LD/BH testing evolve to
1172 consume an inordinate quantity of returned sample, to preclude other
1173 biological, organic, and inorganic tests that further NASA's other goals?
1174 (see section 5.0)
- 1175 ● Study the effects of sterilization measures that could have significant
1176 adverse effects on biochemical analyses outside of PPL containment [*Race*
1177 and *Rummel, 2000*].
- 1178

1179 **Life Detection Testing**

1180 Introduction The proposed Life Detection (LD) analyses are intended to detect
1181 specific evidence whether life of any kind exists in the sample, or rule out the
1182 presence of such evidence of life.¹⁴ These analyses will use a broad definition of
1183 and criteria for life, and an approach for detecting life, not intended to be limited by
1184 the specific features of life as we know it on Earth. This approach will begin with,
1185 and rely on, 'signatures' of various types that encompass all known terrestrial life,
1186 and that might encompass non-terrestrial life. These signatures structures,
1187 structural and biosynthetic chemistry, isotopic patterns, and geochemical features
1188 that help define the underlying principles of life (see *Biosignatures*, page 45). The
1189 LD tests will take advantage of, but will not be constrained by, knowledge of the
1190 structural and metabolic intricacies of terrestrial life. In particular, the recent
1191 recognition of our limited ability to cultivate terrestrial microbial life¹⁵ emphasizes
1192 the importance of relying on methods beyond *in vitro* cultivation for detecting
1193 extraterrestrial life. Life is likely to be catalytic and carbon-based. The most
1194 parsimonious scenarios for the existence of extraterrestrial life posit the presence
1195 of a prebiotic mix similar to that which existed on the early Earth. The similarity of
1196 Mars to Earth in this regard is anticipated under current models of solar system

14. The final reports from each Workshop contain detailed documentation of the discussions which occurred at those Workshops [*Race and Rummel, 2000; Race et al., 2001a, 2001b, and 2002*].


15. At the time of this writing, only about 1% of known microbes can be readily cultured.

1197 formation. Evolutionary paths different from those that occurred on Earth may have
 1198 led to the generation of slightly different building blocks and polymers. The LD
 1199 methods should be potentially capable of recognizing the products of these variant
 1200 paths, and be capable of recognizing the various known forms of life on Earth.

1201

1202 An overall strategy for LD is illustrated in Figure 4, showing the expected flow of
 1203 materials into the various testing queues to be established for the protocol. This
 1204 strategy, originally developed in the first Workshop of the Series [Race and
 1205 Rummel, 2000], was refined and elaborated upon in the subsequent Workshops
 1206 [Race et al., 2001a; 2001b; and 2002].

1207



1208

1209

1210

Figure 4. Life Detection Process Flowchart.

1211 Table 3 lists what could be considered ‘universal’ properties of life. Many of these
1212 properties are directly measurable, although some of them, such as replication
1213 or evolution, can, in all likelihood, only be inferred. Evidence for only a subset of
1214 these properties in an extraterrestrial specimen might constitute a sign of life
1215 (e.g., evidence for a self-sustaining catalytic system). However, it is the presence
1216 and combination of all of these properties that define life as we know it.
1217

- Life is catalytic
 - + There should be significant deviations from what chemical kinetics predicts
 - + Life modifies its environment
 - + Life consumes energy
 - + Life creates waste products
 - + Life is exothermic
 - + Life uses thermodynamic disequilibria to build and maintain other thermodynamic disequilibria (in open systems or within a “wall”)
- Life is genetic
 - + There will be some system for storing and propagating information
 - + There will be molecular distributions with significant capacity for complexity
- Life replicates and evolves
 - + There will be evidence for replication of structures and complexity
 - + There may be evidence (structural & chemical) of evolution of form& function

1218
1219
1220
1221

Table 3: Universal properties of life, as we know it.

1222 LD Principles General principles to follow in searching for life or biosignatures
1223 (i.e., signs of life) are shown in Table 4 on the next page. These principles guide
1224 the search from the selection of samples to be tested through the application of
1225 analytical methods, as shown above in Figure 4. Analytical methods can be
1226 divided into those that facilitate a wide survey of a representative portion of different
1227 sample types, and those that facilitate a more focussed, but high-resolution,
1228 examination of areas of interest. Survey methods are less destructive of samples,
1229 and include microscopy, broad band fluorescence, surface scanning and
1230 chemistry, tomography, and isotope release experiments. These methods seek

1231 structural and basic chemical signatures, and local inhomogeneities. Higher
1232 resolution methods are generally more destructive, and include mass
1233 spectroscopic methods, combustion, isotope analysis, and electron microprobe
1234 procedures for elemental mapping. These methods seek to characterize
1235 inhomogeneities and more complex structures, and are discussed below in
1236 further detail (see *Sample and Time Requirements*, page 53).¹⁶
1237

- Begin with a broad survey of a portion of different sample types for more general features suggestive of life, then turn to a higher resolution examination of sites with suggestive features for a more complete characterization
- Emphasize structural signatures of life and other inhomogeneities that can be easily detected as a first order task
- Emphasize less destructive methods in the early stages of investigation, since they can guide the use of more definitive but destructive methods
- Start with samples least likely to contain life (e.g., surface fines); if negative, use these as blanks and controls for spiking experiments
- Recognition of life will require the coincidence of multiple independent signatures
- Inactive or “past” life will be treated as potentially active life
- Generalize a carbon-centered methodology to other chemical species
- Use an iterative approach for the Life Detection protocol
- Invest significant time in the design of controls and blanks, as early in protocol development as possible.

1238

Table 4: General principles guiding the search for life.

1239

1240

1241

1243

1244

1245

1246

1247

1248

1249

One factor that may complicate the Life Detection efforts is the difficulty in detecting or interpreting many of these signatures if the life-forms are inactive, or have been for long periods of time (e.g., hibernation or quiescence), or have become fossilized. One of the large challenges in Life Detection is a more complete understanding of the stability of various biosignatures over time and their dependence on continued metabolic activity. Attempts to induce activity and replication are also posited as a means of amplifying potentially detectable

16. An estimate of the amount of sample required for the survey/less-destructive methods is 200 milligrams, and 3 grams total for all tests (see page 53).

1250 biosignatures. Some indicators, either structural and/or chemical, which may
1251 indicate “past” or inactive life should be treated as potential indicators of active life.

1252

1253 One potentially useful strategy for detecting active life-forms is based on replicate
1254 measurements over time. Repeated analyses for any of the biosignatures
1255 described above may reveal changes in the sample due to metabolic activity. The
1256 search for significant changes in these signatures offers an important potential
1257 source of information, and does not require a thorough understanding of the
1258 signature. The probability of life based on a chemical species other than carbon is
1259 low, but cannot be eliminated. With this in mind, carbon centered methodologies
1260 and approaches which dominate our present thinking need to be generalized to
1261 other chemical species whenever possible. An iterative general approach is
1262 recommended for the Life Detection tests, with results obtained by one method or
1263 analysis being used to specify and direct any subsequent use of such methods or
1264 analyses.

1265

1266 There are three possible outcomes of the Life Detection procedures:

1267 1. *Failure to detect any of the biosignatures described above, and absence of*
1268 *any carbon or complex carbon in representative samples.* This result would
1269 lead to proposals for downgrading of the containment level for controlled
1270 distribution.

1271 2. *Clear and overwhelming evidence of living organisms that appear to be of*
1272 *non-terrestrial origin (for example, evidence of motile structures with no DNA*
1273 *or RNA present).* This finding could result in the continued containment of
1274 all unsterilized samples for an indefinite period of time—until the living
1275 organisms are better understood. Biological experimentation and biohazard
1276 assessment would be given highest priority. It must be emphasized that the
1277 most likely source of life detected in the martian specimens is expected to
1278 be terrestrial contamination (introduced just prior to, or following the
1279 spaceflight portion of the mission).

1280 3. *The third and most likely scenario lies between these extremes, where clear*
1281 *evidence of life or its absence is not forthcoming.* An example would be a

1282 situation in which complex carbon-containing compounds are detected in
1283 the sample, but without other evidence of life or biosignatures.

1284
1285 Extraction of Representative Sample It is anticipated that sample material will
1286 differ widely in size and composition. For discussion purposes, a representative
1287 aliquot of approximately 1 gram would be subjected to extraction for further
1288 destructive tests. This initial extract will be made using ultra-clean water.

1289 Mechanical disruption may be necessary, but should be kept to a minimum so as
1290 not to damage cellular structures or potentially viable cells. A fraction of this
1291 aqueous slurry should be designated for organic solvent extraction. Obviously,
1292 future planning on the extraction of a representative sample will be dependent on
1293 mission capabilities and sampling equipment employed.

1294
1295 Biosignatures The signatures and signs of life that are the principal targets of LD
1296 testing may be defined through different prisms, perspectives, and methods.
1297 Broadly-defined signatures offer the greatest opportunities for detecting life that is
1298 unfamiliar to us in its detail; however, broad signatures also carry the greatest
1299 chance for misleading or false-positive findings. In general, the greater the
1300 number of independently-defined signatures that are detected, and the greater the
1301 spatial co-localization of these signatures, the stronger the evidence for life. As a
1302 simple example, self-sustaining catalytic processes should create a localized
1303 overabundance of a discrete set of related compounds. Useful biosignatures may
1304 exist in a variety of types:

- 1305 ● *Morphological.* As we know them, all forms of life are defined by a boundary
1306 (e.g., a wall) that delineates them from the surrounding environment. This
1307 “spatial-physical incongruity” often contains patterns, complexity and
1308 recognizable features (e.g., size, shape, structure, morphological indicators
1309 of replication or specialized features such as attachment and motility
1310 structures, septae, etc.).
- 1311 ● *Structural Chemistry.* Life can be defined by basic chemical features, such
1312 as organic or complex carbon, or by higher-order features, such as
1313 polymers, membranes, and attachment and motility structures. Methods

- 1314 need to be improved for characterization of complex polymers and criteria
1315 developed for interpreting the patterns associated with complex carbon. We
1316 are even less well-informed about the possible structural complexity that
1317 can be incorporated into silica and silica-carbon polymers.
- 1318 ● **Metabolism and Bioenergetics.** The waste products that are released and
1319 the energy expended by all forms of life as we know them can be detected
1320 with physical and chemical methods. Some products are created through
1321 specific enzyme catalyzed reactions, such as the reduction of nitrogen that
1322 can occur from inorganic reactions. Other products are predicted to result
1323 from reactions in the absence of protein-enzymes, such as those involved in
1324 energy and CO₂ reduction. More work is needed to assess the range of
1325 metabolic mechanisms and products that occur on Earth, as well as
1326 theoretical studies of those that might occur in the absence of carbon.
 - 1327 ● **Biosynthetic Mechanisms.** All life has mechanisms to synthesize structural,
1328 metabolic and replicative macromolecules. Carbon-based life on Earth
1329 uses protein-enzymes and, to a limited extent, ribozymes (catalytic RNA).
1330 The synthesis of macromolecules involves a sequence of reactions that
1331 depends on the availability of basic organic components, such as amino
1332 acids for protein synthesis. Such synthetic mechanisms should provide
1333 detectable biosignatures, if they are present. In taking a broader view, we
1334 must consider the possibility of biosynthetic mechanisms and pathways
1335 catalyzed by inorganic metals and minerals in non-protein matrices, or that
1336 are dependent on physical gradients (temperature, pH, Eh, magnetism),
1337 catalytic mineral surfaces, or various energy sources (UV and other forms of
1338 radiation and light). Such mechanisms may exist, but their detection may be
1339 as a consequence of first detecting other signatures of life.
 - 1340 ● **Isotopic Signatures.** All forms of life with which we are familiar fractionate
1341 various elements; thus, fractionation patterns can be indicative of life.
1342 Organisms that express different metabolic capabilities display distinctive
1343 patterns in the fractionation of carbon, nitrogen and sulfur. This might be
1344 particularly important in assessing the possible origins of organic
1345 compounds and various volatiles such as methane, carbon dioxide, and
1346 carbon monoxide, if detected on Mars. While one cannot assume that
1347 extraterrestrial life will fractionate elements in the same manner as
1348 terrestrial life, it is reasonable to assume that local patterns of fractionation

1349 within or at sites of life-forms in the sample will vary from those measured
1350 in the surrounding sample environment. Some isotopes, such as those for
1351 oxygen (detected in carbon dioxide and phosphate), can be indicators of
1352 environmental temperature. There is promising new technology for
1353 measuring carbon isotope fractionation patterns in single organic
1354 molecules and fractionation patterns in transition metals. The latter may be
1355 very important in identifying a biological source for various minerals such as
1356 magnetite.

1357 ● *Geochemical Signatures.* This family of signatures includes findings such
1358 as magnetite, and other minerals out of equilibrium with their normal
1359 distribution in the environment. Redfield-like ratios¹⁷ of key elements
1360 (e.g., C, H, N, O, P, and S) are found in the pigments of terrestrial life, such
1361 as those known to be associated with photosynthesis, and other inorganic
1362 chemical anomalies (e.g., based on iron, sulfur, etc.). When specific
1363 biologically important elements are limited in the environment, there will be
1364 higher concentrations associated with life-forms or colonies of life-forms.
1365 Usually, the limiting element in the environment will limit the extent of growth
1366 and productivity of organisms (known as Liebig's Law of the Minimum).
1367 Some key elements that are limited in terrestrial environments include iron
1368 and molybdenum (essential for nitrogen cycle reactions), and tungsten
1369 (essential for specific enzymes in hyperthermophilic archaea).

1370
1371 *Analytical Methods* Because deep and surface mineral particles are common
1372 micro-environments for microbial life on Earth, the chemical analysis of Mars
1373 samples at a micrometer scale can yield information about the presence of active
1374 or fossil life on Mars. Raman, IR, and fluorescence micro-spectroscopy are
1375 valuable tools to perform non-destructive analysis of mineral matrices and surface
1376 compounds.

1377 ● *Microscopy.* As part of the preliminary examination of returned samples,
1378 light microscopy of fines as well as surfaces of pebbles or rock should be
1379 used to look for obvious signs of cellular structure and mineral deposits
1380 associated with microbial life.

17. The 'Redfield Ratio' describes the ratio of carbon to nitrogen to phosphorous (C:N:P) found in marine organisms.

- 1381 ● *Analysis of Gases in Head Space.* One potentially important analysis for
1382 Life Detection would be to compare a pristine atmospheric sample from
1383 Mars to the gas occupying the head space above collected soil and rock
1384 samples. If a pristine sample is available, the comparison may yield
1385 differences that could be due to chemical interaction of the gas with
1386 samples, or that may be signs of metabolic activity within the specimens.¹⁸
- 1387 ● *Laser Desorption Mass Spectroscopy and Laser Raman.* Laser desorption
1388 mass spectroscopy (LD/MS) is a rapid, non-destructive method for detecting
1389 low levels of organic matter in geological specimens. It has been
1390 successfully used to analyze PAHs in meteorites and interplanetary dust
1391 particles. Minimal sample preparation is required, and small particles as
1392 well as fresh fracture surfaces of larger specimens can be analyzed. In
1393 LD/MS, a 10-40 micron diameter spot is positioned on the specimen,
1394 organic species are thermally desorbed from the outer few microns of the
1395 specimen, they are photo-ionized and directed into a time-of-flight mass
1396 spectrometer. Continuing developments offer the prospect of high selectivity
1397 in detection of specific classes of organic compounds, (e.g., amino acids).
1398 Additionally, recent studies suggest that for organic compound detection
1399 UV-Raman spectroscopy (especially deep UV Raman, ~224 nanometers)
1400 may be 5-7 orders of magnitude more sensitive than longer-wavelength
1401 Raman spectroscopy, and can use a smaller focused light source that is
1402 less sensitive to rough surfaces. At UV wavelengths, the mineral
1403 fluorescence disappears and the signal, even when small, has little or no
1404 noise attached from that source. Automated scanning technology will be
1405 critical for application of these techniques to the maximum amount of
1406 sample. These techniques are limited to surface analysis.
- 1407 ● *3D Tomography.* Given the present state of the art, 3D tomography would
1408 require transport of a specimen outside of maximum containment facilities
1409 to a synchrotron; however, the specimen can remain in a sealed container,
1410 under the equivalent of PPL- α containment conditions. The availability of an
1411 appropriately qualified synchrotron facility capable of applying this method to
1412 detect specific elements within a sample would be of great interest in the

18. Although not a requirement of the protocol *per se*, the desirability of this analysis suggests the importance of collecting separate gas-only samples from the sample collection sites on Mars.

1413 preliminary examination of rock samples that might have heterogeneous
1414 interior structures.

1415 ● *Carbon Analysis.* High priority should be given to quantitative analysis of
1416 carbon, especially organic carbon. Techniques having the greatest
1417 sensitivity should be applied, including progressive heating/oxidation,
1418 coupled to GC/MS. It is anticipated that multiple samples and sites with
1419 suspicious findings from survey methods will be analyzed to detect and
1420 characterize localized organic or inorganic carbon.

1421 ● *Flow Cytometry.* An aliquot of the aqueous slurry will be subjected to flow
1422 cytometry. Flow cytometry will be used to analyze single particles in the
1423 range of 2 to 100 microns in diameter, at rates of tens to hundreds of
1424 thousands of particles per second. Based on initial, non-destructive
1425 characterization of laser light scatter and auto-fluorescence, particles will be
1426 re-analyzed, with or without staining with fluorochromes specific for DNA,
1427 proteins or functional viability assays. During subsequent analysis, at least
1428 four pre-selected sub-populations can be sorted from each sample for
1429 further analysis by other techniques. Positive fractions can be sorted and
1430 directed toward further chemical and biochemical testing.

1431
1432 Cultivation Elaborate forward-contamination controls will be used on the mission,
1433 but it is still possible that viable terrestrial microbes may be detected in returned
1434 Mars samples (either from contamination on the original spacecraft, the sample
1435 container that made a round-trip, or through sample handling contamination). To
1436 rule out possible terrestrial microbial contamination, an aliquot of the sample
1437 should be subjected to the standard microbiological examinations currently used
1438 for planetary protection, as well as other routine methods for detecting and
1439 identifying terrestrial organisms.

1440
1441 In addition to the procedures used to identify any terrestrial contamination, culture
1442 attempts should be made that represent Mars-like conditions. Culture conditions
1443 that would be compatible with martian micro-environments are not well-
1444 understood and the likelihood of success is small (only about 1% of Earth
1445 organisms can readily be cultured), yet attempts should be made to create such

1446 conditions and propagate life-forms. The composition of gases in the martian
1447 atmosphere, including plausible ancient atmospheres, should be replicated,
1448 especially with CO₂ as a carbon source. Given the current extremely dry conditions
1449 on Mars, the degree of sample hydration should be varied. The range may
1450 fluctuate from partially hydrated specimens to totally aqueous conditions. Energy
1451 sources should include light for any possible photosynthetic organisms and pairs
1452 of electron donors and acceptors for chemosynthetic organisms. Mineralogical
1453 information from samples should be integrated into the decisions in media
1454 formulations. Likewise, any organic compounds detected in the samples should
1455 be considered as carbon sources for possible microbial growth. Cultures will be
1456 monitored by simple microscopy as well as through multiple sequential analyses
1457 by GC/MS, LC/MS, micro-calorimetry, nucleic acid amplification, and other
1458 methods.

1459

1460 Distinguishing Earth-based from Mars-based Life If viable cells are found in the
1461 samples, and especially in cultures taken from samples, it will be important to
1462 address the possibility (even likelihood) of terrestrial microbial contamination.
1463 Detected cells will be subjected to phenotypic and genotypic analyses, with
1464 sequence searches against databases containing large numbers of known
1465 terrestrial organisms to quickly identify contaminants (though it is important to
1466 remember that only a small percentage of Earth microbes are currently known).
1467 Because of the harsh conditions on Mars and the relatively small amount of
1468 sample to be returned, the most likely source for familiar complex polymers such
1469 as nucleic acids is from terrestrial contamination. Amplification techniques such
1470 as the polymerase chain reaction (with broad range primers directed against
1471 targets such as rDNA, and with random oligomers) and subsequent sequencing
1472 methods offer a sensitive and rapid means for detecting and characterizing DNA
1473 and RNA (as a marker for terrestrial contamination), and should be applied to the
1474 outbound spacecraft and container surfaces before and after return, as well as to
1475 the samples themselves. Other assays, such as the *Limulus* Amoebocyte Lysate

1476 (LAL) assay, may assist in detecting extremely small amounts of terrestrial
1477 contamination, but are less specific.

1478
1479 It must also be kept in mind that detection of terrestrial contamination in a
1480 specimen does not exclude the possibility that the same specimen also contains
1481 martian life. The presence of terrestrial contamination could compromise the
1482 detection of potential martian life in a number of ways—e.g., if martian life is
1483 closely related to Earth life, or if the “noise” of terrestrial contamination drowns out
1484 the “signal” of Mars life; this is a key reason for requirements to be imposed on the
1485 sample collection mission that will restrict the transfer of terrestrial contamination
1486 to the sample and/or sample container.

1487

1488 Considerations Concerning Controls and Blanks

- 1489 ● Prior to departure, the spacecraft and specimen containers should be
1490 examined, and samples should be archived; witness plates¹⁹ should be
1491 employed.
- 1492 ● Strong consideration should be given to the return of a sample of martian
1493 atmosphere in a separate, but identical container. If collected and stored
1494 under increased pressure, extra aliquots of atmosphere could be used for
1495 replication of martian conditions in other experiments after specimen return.
- 1496 ● Early determination of negative findings for life in low-likelihood martian
1497 samples may allow these samples to be used as negative controls.
- 1498 ● Because negative results are expected in many of the Life Detection
1499 procedures, determinations of assay sensitivity using known specimens of
1500 terrestrial life would aid in the interpretation of these negative results.
- 1501 ● Methods should be validated and evaluated using a wide variety of
1502 terrestrial life-forms.
- 1503 ● Simulants of martian samples and conditions should be refined for protocol
1504 development prior to sample return. Particular attention should be given to
1505 the probability of highly-oxidizing sample surfaces.

19. ‘Witness plates’ are controls for forward contamination, used to monitor the bioload on a spacecraft before launch.

- 1506 ● Exposure of the sample surface to PPL- α conditions will inevitably lead to
1507 deposition of particulate matter from the surrounding enclosure. The
1508 features of this process should be characterized prior to specimen return.
- 1509 ● Questions that yield answers for which a statistical assessment of
1510 confidence can be performed should be identified. Principles to be applied
1511 in order to generate statistically robust findings should be determined.

1512

1513 *Life As We Don't Know It* The possibilities of dealing with “life as we don't know it”
1514 need to be considered seriously, including: a composition devoid of organic
1515 carbon; the unconventional reliance on “non-biological” elements such as Si, Fe,
1516 and Al; structures less than 100 nanometers in diameter; and a composition
1517 based on organic monomers. Of course, it is difficult to evaluate the probability of
1518 encountering forms of life with these features.

1519

1520 Discussions of the possibility of non-carbon based life have had a rich history,
1521 especially in the realm of science fiction.²⁰ Life based on organic monomers has
1522 recently been proposed as a model for the ‘metabolism-first’ scenario for the
1523 origin of life.²¹ According to this model, a set of self-sustained chemical reactions
1524 might be considered ‘living’ if metabolism is considered to be more important than
1525 replication as a fundamental basis of life. Some of these unlikely scenarios might
1526 require alternative laboratory conditions for proper study (e.g., use of inert gases).

1527

1528 Existing theories of the origin of life on Earth suggest that life will arise as a
1529 consequence of chemical and physical principles anywhere prebiotic carbon
1530 compounds accumulate in suitable environments (e.g., water, temperature, etc.)
1531 in sufficient amounts for sufficient time. Although the precise process for life's

20. H.G. Wells, writing in the Pall Mall Gazette in 1894, scolded scientists for thinking of only carbon-based life: “It is narrow materialism that would restrict sentient existence to one series of chemical compounds – and the conception of living creatures with bodies made up of the heavier metallic elements and living in an atmosphere of gaseous sulfur is no means so incredible as it may, at first sight, appear.”

21. Wächtershäuser, G., *Science* 289:1307-1308 (2000).

1532 origins on Earth is not known, it is perceived to have been a progression in
1533 complexity beginning from an original prebiotic mixture, at some stage involving
1534 RNA catalysis, and probably at later stages catalysis by peptides and proteins,
1535 ultimately culminating with the first simple organisms that had a metabolism, the
1536 ability to replicate, and the capability of preserving useful information during the
1537 replication process. The most likely scenario we can conceive of for the
1538 independent development of life on Mars is by a similar process, which if
1539 stochastic, may have deviated from our own terrestrial process and resulted in
1540 different fundamental amino acids or nucleotides used, types of lipids, chirality,
1541 etc. The primary indicator of past or present life of this type would be the finding of
1542 unusual macromolecular assemblages (e.g., peptides or oligonucleotides with
1543 nonstandard amino acids, nonstandard bases, nonstandard linkages). If deviation
1544 occurred only later in the process, we might find Earth-like complex structures
1545 such as recognizable ribosomal RNAs.

1546

1547 It also should be noted that if there is, or has been, life on Mars, it might be related
1548 to life on Earth by descent. If an evolved living organism reached Earth from Mars,
1549 or less likely, reached Mars from Earth, the two life forms should be closely similar
1550 in their biochemistry. They should, for example, use DNA as a genetic molecule
1551 and might have the same genetic code. If two life forms originate and evolve
1552 independently, however, there is no *a priori* reason to expect them to be similar.

1553

1554 Sample and Time Requirements It is estimated that approximately 3 grams of
1555 sample will be required to conduct the proposed preliminary Life Detection tests
1556 on returned martian sample materials.²² As methods mature and new
1557 approaches become available, these sample requirements may change.
1558 Estimates of the time needed for Life Detection are difficult to make. Survey
1559 methods can be completed within weeks-to-months, in some cases. However,

22. Estimates for sample amounts are based on what is necessary to conduct the tests outlined in the Draft Protocol; however, actual amounts may depend on definitions of "representative samples" made at the time samples are returned.

1560 any positive or suspicious findings may impose additional time requirements,
1561 depending on the strength of the findings and the follow-up methods required for
1562 further assessment. For example, enrichment culture experiments as part of the
1563 Life Detection protocol may extend for many months, even though they are not
1564 considered a strong methodology for detecting martian life.²³

1565

1566 Future LD Research and Development Needs

- 1567 ● Miniaturization of many chemical/physical analyses
- 1568 ● Sample registry, for re-interrogating precisely defined sites within the sample
- 1569 ● Micro-calorimetry
- 1570 ● Database development
- 1571 ● Software for "multiple sequential analysis" search logic
- 1572 ● Effect of Mars atmosphere versus inert atmosphere on proposed methods
- 1573 ● Cleaning/cleanroom technologies
- 1574 ● Validation of controls
- 1575 ● 3-dimensional nano-scale structural mapping of specimens
- 1576 ● Characterization of complex compounds based on Si, Al, Fe
- 1577 ● More complete inventory of life on Earth, using molecular methods

1578

1579 **Biohazard Testing**

1580 Introduction The Biohazard testing process is intended to determine if samples
1581 from Mars pose any threat to terrestrial organisms or ecosystems, regardless of
1582 whether the samples are found to contain life-forms or non-replicative hazards. In
1583 this Draft Protocol, it is recognized that potential hazards could take one or more of
1584 a multitude of forms (e.g., toxic, mutagenic, life-cycle altering, hazardous through
1585 genetic recombination, disruptive to ecosystems, capable of biasing phenotypes,
1586 or even behavior). Thus, the spectrum of tests selected is deliberately diverse.

23. Attempts to culture potential microorganisms from Mars samples will be done recognizing that, even on Earth, the vast majority of terrestrial organisms cannot be cultured under known conditions. Bearing this in mind, the length of various culture experiments may be allowed to extend into months even though the likelihood of positive outcomes is extremely low.


1587 Both conventional whole-organism animal and plant *in vivo* testing are planned, in
1588 addition to *in vitro* cellular assays and molecular biology tests (see Figure 5).

1589
1590 In light of the robust nature of emerging molecular, cellular, and conventional
1591 testing procedures, specific methods will be selected later in accordance with
1592 state-of-the-art practices and refinements at the time the final protocol is
1593 implemented [Race et al., 2002]. Selections should take into account evolving test
1594 methods (e.g., toxicogenomics) that are anticipated to replace many current
1595 conventional practices over the coming years. These newer procedures may
1596 ultimately become refined state-of-the-art approaches. In such instances,
1597 advances in testing methodologies that presently await standardization and
1598 validation should allow modifications and refinements to Biohazard testing
1599 adopted for the final protocol applied to samples from Mars.

1600
1601 The proposed tests and procedures for Biohazard testing reflect the current state
1602 of knowledge and practice. It is anticipated that this Draft Protocol will evolve both
1603 in content and implementation as a result of new or improved methodologies or
1604 expanded states of knowledge prior to sample return, and in response to real-time
1605 information about sample materials learned during implementation of the various
1606 processes at the SRF. A sketch of the pathway of experiments for Biohazard
1607 testing is given in Figure 5 and further details of those pathways are in Table 5.
1608 The approach outlined in Table 5 was developed early in the MSHP Workshop
1609 Series [Race et al., 2001a], and refined at subsequent Workshops in the Series
1610 [Race et al., 2001b and 2002]. Throughout the Workshop Series, the development
1611 of a general approach for Biohazard testing, rather than a specific list of tests, was
1612 considered the most useful and responsible approach for deliberations at this
1613 time. [Race and Rummel, 2000; Race et al., 2001a, 2001b, 2002],

1614
1615 The data from Biohazard testing will be used in combination with those from Life
1616 Detection and Physical/Chemical testing to determine what level of containment, if
1617 any, will be required for the further study of the samples. In practical terms,

1618



1619

1620 Figure 5. Proposed Flow Chart for Biohazard testing. The clear region contains tests
 1621 (chiefly for pathogenicity) that should be done in strict containment (PPL- $\alpha/\beta/\gamma$),
 1622 while the shaded region represents similar tests for broader-spectrum biohazards
 1623 done in less strict, but still secure, containment (PPL- δ).
 1624

1625

Test Type	Procedures/Questions	Sample Usage and Time Required
Verification that any potential organisms do not attack biocontainment materials (e.g., Silastic™, rubber, etc.).	Do samples affect test coupons of containment materials at various humidity levels and temperatures?	Sample expended: 1 gram Time: 1 - 3 months?
<p>Input from Life Detection Procedures (discussed separately):</p> <ul style="list-style-type: none"> • If life detected, this would radically change/focus the approach to Biohazard testing by providing focus in terms of conditions for replication, agents that can kill the organism(s), etc. • If no life is detected, still run subsequent tests for toxicity and biohazard. 	<ul style="list-style-type: none"> • Carbon? • Carbon-carbon bonds? • Complex carbon compounds (indicative of metabolic processes)? • Skeletal remains or fossilized remnants? • Indication of live organisms (organelles, membranes, structures on microscopic evaluation)? • Life-like structures? • Living agent (replicates in environment, with co-agent/host, in terrestrial cells)? • Mutual/commensal/parasitic relationship? • Kills cells or organisms? • Kills complex multicellular organisms? • Kills everything? 	Sample expended: TBD Time: TBD
<p>Multi-species infectivity, pathogenicity, toxicity testing.</p> <ul style="list-style-type: none"> • Look at broad host ranges (assuming that any pathogens would not be too host-specific) with well-known and standardized model systems. • Use small organisms in small volumes, allowing for maximum sample conservation. • Initial work all done at BSL-4 biological containment level. 	<p>Sample preparation (rough cut):</p> <ul style="list-style-type: none"> • Crush larger clumps/rocks but do not pulverize particulates. • Filter? • Mix into sterile water. • Chelate heavy metals? • pH buffer? • Use serum for some samples? <p>Heavily irradiate sterilized control samples w/ ⁶⁰Co.</p> <p>Introduce appropriate amount of sample (10 -100 milligrams for statistical relevance) to culture of unicellular organism and cell lines.</p> <p>Inoculate whole organisms (animals as human models) with primary (not passaged) material.</p> <p>Monitor:</p> <ul style="list-style-type: none"> • Cell proliferation, • Cell morphology, • Differential analyses of biochemicals and gene expression • Comparative genomics (any inserted genes in host?) • Reporter assays (?) • etc. 	<p>Sample expended: Three trials plus sterilized control per organism, assuming 100 mg per sample = 1.6 grams.</p> <p>Time: ~ 6 months to allow for passage times.</p>
Negative results with multi-species tests may lead to downgrading to PPL-δ.	<p>The following tests/criteria are proposed:</p> <ul style="list-style-type: none"> • First passage from infectivity analysis (+ or -), but second and subsequent passages all neg. • DNA damage assays (mutagenesis: Ames- test, strand break analysis). • Environmental damage. • Whole plant inoculations. • Diversity of growth conditions extant on Earth (extremophiles, etc.) and other media. <p>Monitor: cell viability, expression of toxic response genes.</p> <p>Negative results on these tests may allow a decision to downgrade to a lower containment level or release.</p>	<p>Sample expended: ~10 - 20 grams (very rough estimate).</p> <p>Time: ~6 months to allow for passage times.</p> <p>Note: There was consensus on the 'first round' (infectivity), but it was also clear that the containment-level determination issues need considerably more analysis and study.</p>
		Total = 15-25 grams

1626
1627

Table 5. An outline of a possible pathway of experiments for Biohazard testing.

1628 Biohazard testing should allow a determination—with a high degree of confidence
1629 and a clear understanding of the conditions of release—of whether the samples
1630 contain any biohazard and whether to distribute sub-samples. A determination
1631 about releasing a sample from containment will be made with careful
1632 consideration of applicable regulatory requirements and will provide a reasonable
1633 assurance that the samples will not put humans or other terrestrial organisms at
1634 risk.

1635

1636 *Biohazard Defined* In general terms, hazards of concern to biological systems
1637 may be caused by materials or entities of biological origin, and by those materials
1638 or entities replicating or being amplified²⁴ toxic and by a biological system. Such
1639 hazards are capable of producing an adverse effect on or significant alteration of a
1640 biological system at the level of individual organisms or ecosystems.²⁵ In the
1641 special case of hazards from returned martian samples, a distinction can be
1642 made between replicating and non-replicating hazards. For the purpose of this
1643 Draft Protocol, a *biohazard* is defined as a hazard that can either replicate or be
1644 amplified by a biological system. In practical terms, replication is a key distinction
1645 between a biohazard (i.e., replicating and potentially contagious) and a simple
1646 toxin or hazard (e.g., a non-replicating substance that can be diluted down below
1647 an initial toxic concentration). Only replicating entities, or entities that are able to be
1648 amplified by a biological system, pose a potential widespread threat. While other
1649 hazardous materials are of concern, the quantities returned from Mars will be
1650 extremely limited, and they thus represent a potential hazard of real significance
1651 only to scientists and others who may be exposed to them.

1652

24. In this context, biohazards are not limited to 'living' entities—and may include biohazards such as viruses that are not living or self-replicating *per se*.

25. In the context of potentially biohazardous extraterrestrial entities, "adverse effects" includes any significant alteration on a biological system, and is not limited to adverse effects that are immediately or acutely toxic.

1653 If the distinction between a biohazard and a non-biological hazard is made, the
1654 level of containment and procedure for distribution of the samples can be
1655 appropriately defined. The existence of either biohazards, which are self-
1656 replicating or able to be amplified by another biological system, or toxic hazards
1657 would require further study and characterization of the nature of the hazard
1658 (e.g., strong chemical oxidizer, radioactive, replicating life-form, etc.) so that
1659 appropriate subsequent containment and/or handling procedures can be
1660 determined and stipulated to avoid potential biological impacts during future
1661 research.

1662
1663 Assumptions About Containment Containment at the SRF will be designed to
1664 provide a range of environmental conditions for the martian samples, while
1665 maintaining them at appropriate biocontainment levels. It is important to
1666 understand the various containment types at the SRF and the anticipated
1667 containment needs during Biohazard testing. Life Detection and
1668 Physical/Chemical tests will seek to characterize the sample materials and
1669 determine if evidence for “life” can be found under conditions that are both Mars-
1670 like and Earth-like. In contrast, Biohazard tests are designed to determine the
1671 effect of martian samples on terrestrial life-forms under Earth-like conditions.
1672 Thus, containment requirements for execution of the Biohazard testing will not
1673 require the same stringent clean room conditions associated with the preliminary
1674 P/C tests, certain Life Detection studies, and ‘banking’ or curation. The appropriate
1675 initial containment level for the Biohazard testing is thus anticipated to be PPL- γ ,
1676 which translates to the maximum BSL-4 biocontainment, but with less demanding
1677 cleanliness restrictions than PPL- α .

1678
1679 The unknown nature of any possible biohazard in returned martian samples
1680 demands, at least initially, the most stringent containment presently afforded to the
1681 most hazardous biological entities known on Earth. If sufficient data are gathered
1682 to rule out concerns about human virulence and infection, a decision could be
1683 made later to allow subsequent work at a lower containment level during tests

1684 investigating possible environmental effects. The Biohazard testing process is
1685 designed to allow for gradual decontainment or adjustment to less stringent
1686 containment levels if justified upon review of accumulated data about the sample
1687 materials during implementation of the Draft Protocol. If the initial Life Detection
1688 and Biohazard tests are all negative, it would be appropriate to conduct
1689 subsequent tests under less strict containment conditions once sample materials
1690 have been shown to be non-biohazardous. In particular, additional geophysical
1691 testing can be done at a reduced level of containment, as well as using selected
1692 biological tests associated with the biohazard analysis. A lower level of
1693 containment would potentially enhance sample access within the scientific
1694 community, while still providing adequate biosafety conditions under existing
1695 biosafety guidelines and regulations.

1696

1697 Biohazard testing will be conducted within containment at the primary receiving
1698 facility or at other secure containment facilities. Since neither all the necessary
1699 scientific expertise, nor all of the high-end scientific instrumentation required, are
1700 located at a single facility, there may be a need to allow samples to be distributed
1701 for study/curation at facilities other than the initial receiving laboratory. The
1702 rationale for the use of multiple containment facilities and the ability to test
1703 unsterilized sample materials outside the primary containment facility depend on
1704 the availability of an adequate means for containing and transporting the samples,
1705 for sterilizing or cleaning the outside of the sample container, and for returning the
1706 remaining samples to the primary containment facility after non-invasive or non-
1707 destructive analyses (e.g., synchrotron analyses). Mobile containers certified at the
1708 appropriate PP level (as distinct from traditional BSL transportation requirements)
1709 should be developed and used for transport of samples between facilities.

1710

1711 Considering that Biohazard testing should yield results within a “reasonable time”
1712 (e.g., most testing completed within approximately 6 to 9 months), the majority of
1713 tests should be started synchronously and conducted in parallel. Nonetheless, the
1714 need to conduct preliminary sample examinations and to work on Life Detection

1715 require that Biohazard researchers proceed with some tests before others.
1716 Common sense and gradual decontainment strategies require tests identifying
1717 deleterious effects on containment equipment before those identifying biohazards
1718 to people, and the latter before identifying biohazards to the environment.

1719

1720 After the equipment-compatibility tests, the types of assays to be accomplished
1721 are prioritized by their likelihood of identifying potential pathogenicity and identifying
1722 any restrictions on the distribution of samples to other laboratories for further
1723 testing. If a possible human pathogen were detected, the strictest of handling
1724 protocols would remain in place. If, in complementary fashion, a pathogen specific
1725 to another host were detected, less stringent handling methods might be
1726 possible. If the only hazard identified were a non-replicating toxic agent (e.g., a
1727 toxic chemical), containment could be less restrictive, and would be definable on
1728 the basis of dose-response characteristics and the nature of the toxicity.

1729

1730 Model Systems for Biohazard Testing Prior to conducting Biohazard tests,
1731 decisions will be needed to identify the exact model systems that will be used for
1732 the specific assays. Working criteria for choosing the models are as follows:

- 1733 ● The models should be relevant to a probable hazard scenario, deliberately
1734 avoiding models that would only be sensitive to an improbable danger
1735 (i.e., very unlikely event, very artificial route, extreme doses, rare species
1736 confined to remote niches, etc.) as such models would be of little relevance
1737 to initial Biohazard testing with Mars samples. The emphasis will thus be
1738 placed on modeling of biological systems likely to be in contact with
1739 samples (e.g., workers, their microbial flora, their pets, insects, life-forms
1740 common to the surrounding of sites of future experimentation with the
1741 samples), via probable routes of exposure (e.g., aerosol, etc.), at probable
1742 (low) doses.
- 1743 ● Subsequent models should be relevant to systems of ecological and/or
1744 economic interest.
- 1745 ● Models should be sensitive, meaningful and, if possible, clear to interpret.
1746 Equivocal answers can needlessly prolong the time required to reach a

- 1747 decision on sample release, and will likely cause samples to be consumed
1748 unnecessarily.
- 1749 ● Models should be robust. Samples are likely to contain complex minerals,
1750 oxidative agents and other elements that should not interfere with its
1751 function.
 - 1752 ● Models should be well documented. Observations and analyses should
1753 identify known behavior of the biological system in the model. Preferably, its
1754 genome should be fully sequenced, and extrapolation to other
1755 species/situations should have been evaluated.
 - 1756 ● Models should provide answers in a reasonably short time.
 - 1757 ● Models should be compatible with handling within the SRF, under
1758 containment. For instance:
 - 1759 ► *Cellular and 'small' models.* Should the model organisms or cells for
1760 Biohazard testing be chosen or developed as of this writing, these would
1761 include:
 - 1762 ◆ wild type, mutant and recombinant yeast bearing special sensitivity to
1763 hazardous materials (e.g., radiation mutants; green and blue
1764 fluorescent protein [GFP and BFP] recombinants to test for
1765 recombinogenicity; etc.);
 - 1766 ◆ human cell lines that are as sensitive to pathogens as standard cell
1767 lines used for Biohazard testing (e.g., a human equivalent to vero E6
1768 cells, as sensitive as BHK-cells to mutagens, etc.);
 - 1769 ◆ bacteria and other microbes associated with people (e.g., *E. coli*,
1770 *Staphylococcus*, *Bacteroides*, *Chlamydomonas*, etc.);
 - 1771 ◆ bacteria found in niches likely to be similar to martian underground
1772 ecosystems (e.g., cold and possibly oxidizing, low-oxygen and with
1773 high radiation levels, etc.);
 - 1774 ◆ relevant algal/planktonic unicellular organisms;
 - 1775 ◆ mammalian (e.g., mouse) egg before re-implantation;
 - 1776 ◆ fish eggs (e.g., Zebrafish, Medaka, etc.);
 - 1777 ◆ models for testing effects on development (e.g., *Neurospora crassa*);
 - 1778 ◆ cells and seeds from *Arabidopsis* and rice;

- 1779 ♦ complete *C. elegans*; and,
1780 ♦ complete *Drosophila melanogaster* (likely a flightless variant).
1781 ➤ *Larger organism models.* For tests in which whole organisms are
1782 required, model organisms would include:
1783 ♦ *Arabidopsis* and rice at different stages of development;
1784 ♦ zebrafish and medaka;
1785 ♦ bird eggs; and,
1786 ♦ a variety of types of mice (e.g., germ-free, humanized, wild type,
1787 mutant, recombinant, immunosuppressed, knockout), whether
1788 reimplanted, newborn, or pregnant.
1789 ➤ *Ecosystem-level models.* For tests of multi-species systems, stable,
1790 replicable, laboratory-scale ecosystem models need to be developed
1791 and tested. Microbial mats may form a promising basis for such a
1792 model.
1793

1794 *Verification of Containment Materials Integrity* As a first order of business, a set of
1795 preliminary tests is required for materials used in containment equipment. It is
1796 important to verify that sample materials or potential organisms growing from
1797 them do not attack rubber, Silastic™, and other bio-containment materials. For
1798 example, ten 10-milligram samples would be taken for each seal/containment
1799 material (e.g., latex, Silastic™, Plexiglas™, cyanoacrylate, epoxy, etc.). ‘Coupons’
1800 (i.e., small, regular samples) of each material would be incubated with martian
1801 sample material at a few different humidity levels, bounding those actually to be
1802 used for sample curation, and including liquid water. Test vessels for these
1803 experiments (i.e., primary containment) should be extremely non-reactive, such as
1804 refractory metals (e.g., titanium). For this example, if ten materials are tested, a
1805 total of one gram (or less) of martian sample would be expended.

1806
1807 At regular intervals (over weeks to months), the sample coupons should be
1808 monitored for degradation using optical methods, mechanical tests, and chemical
1809 analyses. ‘Failure’ criteria would be defined in terms of parameters that would

1810 compromise containment, such as outright consumption, pitting/erosion, pinhole
1811 formation, substantial changes in bulk chemical or mechanical properties, etc.
1812 The results would be used to provide a high level of confidence that the samples
1813 could be kept in storage vessels made of the tested materials without risk of
1814 inadvertent release.

1816

1817 Pathogenicity Testing These Biohazard tests, which have a specific focus on
1818 determining adverse effects on humans, will be done in PPL-γ (containment:
1819 BSL-4; environment: normal terrestrial). Toxic effects on cultured cells and
1820 microorganisms should be anticipated due to the chemical (mineral) composition
1821 of the Mars samples. Appropriate controls (terrestrial or meteoritic) must be run
1822 and interpreted. It is assumed that toxic effects, if any, should diminish rapidly in
1823 sub-culturing ('passaging') experiments, since a replicating agent or one able to
1824 be amplified would not be involved in a toxic response *per se*.

1825

1826 Since fines can be considered 'homogeneous' and can be sub-sampled as a
1827 single category in a statistically relevant way, Biohazard testing should begin with
1828 fines. Whether and when other materials should undergo the full array of
1829 Biohazard testing will be based on the results of initial P/C screening and
1830 processing.

1831

1832 Tests will involve exposing model organisms to the martian sample material.
1833 Specific cell and tissue systems should be used for Biohazard testing, as noted
1834 above in the "model" discussion and below in the discussion of each test. It is
1835 envisaged that a large amount of the cell culture work will be accomplished
1836 robotically using existing or new technologies.

1837

1838 The following specific initial exposure tests [*Race et al., 2001a*] should be
1839 included, based on the knowledge available should it be carried out today:

- 1840 ● Human cell lines and primary cell cultures, with particular emphasis on
1841 epithelial cells (e.g., skin, lung, gut). All cells will be observed for abnormal

1842 growth (e.g., cytopathic effect, morphological changes, genetic response to
1843 stress, integration into host genome, co-growth [mycoplasma-like], and
1844 mutation rates). Cells can be checked for transformation (growth on soft
1845 agar). Both supernatant and homogenized cell pellets should be passaged,
1846 typically twice each week for 3 months. Other replicate cultures must be
1847 observed for 1-2 weeks to look for delayed effects. Cell cultures (and
1848 concentrated medium) should be examined, as well, by electron
1849 microscopy to search for microorganisms that may have replicated without
1850 causing abnormal changes in the cells being cultured.

- 1851 ● Mouse cells should also be tested in similar fashion, with “culture-
1852 adapted” material being injected into mice; three mouse systems should
1853 be employed (i.e., wild-type, SCID, and SCID-Hu).
- 1854 ● Microbial systems to be tested should include *Chlamydomonas* (stress
1855 response), *S. aureus*, yeast, and *E. coli*. In addition, microorganisms that
1856 grow in high salinity should also be considered.

1857

1858 Subsequent Pathogenicity Testing and Possible Decontainment Subsequent
1859 testing should be designed to accommodate a variety of test systems and
1860 representative organisms from different biological domains and ecologically and
1861 economically important phyla. If the initial Biohazard tests (above) and Life
1862 Detection tests are all negative, it may be appropriate to conduct these
1863 subsequent tests under less strict containment conditions (e.g., PPL- δ). In
1864 particular, additional P/C testing, as well as some additional Biohazard tests, can
1865 be conducted at a reduced level of containment using the following models:

- 1866 ● Secondary mammalian cell culture systems.
- 1867 ● Plant cell systems (*Arabidopsis*) and whole-plant growth experiments.
- 1868 ● Additional microbes (e.g., nanobacteria, cyanobacteria, thermophiles,
1869 anaerobes, gram-positive bacteria) and microbial systems (e.g., various
1870 temperature ranges, pH ranges, salinity).
- 1871 ● Other species, such as *Drosophila melanogaster* (e.g., wingless
1872 mutants), worms (*C. elegans*), and amphibian and bird eggs. Horizontal
1873 and vertical transmission studies should be done. All animal species

1874 should be observed for behavior change, toxic and teratogenic effects,
1875 and pathological changes.

1876
1877 Additional experiments can employ a variety of techniques to test for biologically
1878 active compounds, micro-arrays (for proteins), etc.

1879

1880 Broader-Spectrum Biohazard Tests Beyond strict pathogenicity testing, the
1881 Biohazard tests that should be completed include:

- 1882 ● *Direct culture.* This is also part of the Life Detection testing process; any
1883 cultured organism which cannot be clearly identified as terrestrial will be
1884 subjected to further Biohazard studies.
- 1885 ● *Exposure of cellular and 'small' models.* Unicellular organisms, or very
1886 small animals can be used with a limited amount of sample, i.e., ~10-1000
1887 micrograms per test. These tests would be based on exposing the
1888 organisms to the sample and using some form of signal readout, such as
1889 gene expression.
- 1890 ● *Molecular and biological tests (altered levels of proteins and metabolites).*
1891 Rapid progress is being made in developing chip-based, as well as other,
1892 methods that allow one to measure the level of particular proteins or
1893 metabolites in a biological sample. Within the next five years, driven by the
1894 demand of genomics research and drug development, these techniques
1895 are likely to become broadly available. It is difficult to make specific
1896 recommendations at this time before standardized procedures are
1897 established. It is expected, however, that the comparative measurement of
1898 proteins and metabolites associated with the biological response to
1899 infection or toxic exposure will become part of the biohazard assessment
1900 procedure.
- 1901 ● *Genetic testing.*
 - 1902 ➤ *Mutagenesis Assays.* Another possible approach is mutagenesis
1903 assays that look at genetic changes over several rapid reproductive
1904 cycles. Typically, bacteria are used (e.g., the Ames test for mutagenicity
1905 uses *E. coli*). The consensus is that these tests will be problematic in
1906 that mutagenesis results tend to be oversensitive and controls would be

1907 difficult to realize. A related assay type is teratogenicity, but these require
1908 breeding animals, and, thus, can require more time (for some species)
1909 than other assay types.

1910 ➤ *DNA Damage.* Assessment of DNA damage should include the
1911 measurement of mutation frequency, recombination frequency, and the
1912 occurrence of DNA strand breaks. Standardized methods are available
1913 to carry out each of these measurements, for example, genetic reversion
1914 assays for DNA mutation, transposon rearrangement assays for
1915 recombination, and terminal transferase assays for strand breaks. Such
1916 approaches, focusing on general measures of DNA damage, are likely
1917 to be more fruitful than highly specific measurements of DNA damage,
1918 such as comparative sequencing or the measurement of a particular
1919 type of DNA damage.

1920 ➤ *Altered Gene Expression.* Techniques are available for measuring the
1921 relative expression level of almost any gene under various conditions.
1922 For purposes of biohazard assessment, however, it would be preferable
1923 to narrow the focus to genes that are expressed at a significantly altered
1924 level in response to infection or toxic exposure. Testing for altered gene
1925 expression due to toxic exposure is being refined as “toxicogenomics,”
1926 and is anticipated to reach a sophisticated level of standardization by the
1927 time the selection of methods is made for the final protocol.

1928 ● *Whole organisms.* This approach includes ingestion/inhalation/injection of
1929 samples by living organisms with subsequent monitoring of physiologic
1930 functions, behavior, gene expression, inflammatory cascade (e.g., cytokine
1931 levels), etc. Hosts can include animals, plants, and modified organisms
1932 (such as SCID mice, xenograft systems, etc.). Another key aspect of this
1933 approach is the ability to evaluate the infectivity of the potential organisms to
1934 other organisms via passage, and in subsequent generations. The benefits
1935 of this approach to whole organism testing include: direct measurement of
1936 physiologic effects; ability to handle multi-organ interactions in toxicity;
1937 inherent inclusion of complex host characteristics (tough to execute with
1938 cell-based and other assays); and, the possibility of detecting infectivity (if
1939 hosts are appropriate for replication).

1940 Nonetheless, some significant drawbacks exist, including: the difficulty in
1941 seeing long-term effects; it would be impossible to cover all possible

1942 organisms (many terrestrial pathogens are very host-specific); large
1943 samples may be required; tests may be confounded by the presence of
1944 inorganic materials; and, results may depend on the mode of introduction of
1945 sample to test organisms (terrestrial pathogens have specific routes of
1946 infection). A major drawback of this approach is that it requires more
1947 sample, i.e., ~100-5000 micrograms per test. Approaches/organisms
1948 include:

- 1949 ➤ Exposure by direct contact and/or aerosol—*Arabidopsis* and rice at
1950 different stages of development;
- 1951 ➤ Exposure to the sample by routes to be determined (e.g., water
1952 solution, etc.)—Zebrafish and Medaka;
- 1953 ➤ Injection with powdered sample—bird eggs (notably embryonated
1954 chicken eggs); and,
- 1955 ➤ Exposure of a variety of types of mice (such as: germ free,
1956 humanized, wild-type, mutant, recombinant, newborn, pregnant,
1957 immunosuppressed, reimplanted), to the sample as an aerosol, by
1958 intraperitoneal injection, or *per os*. There may also be genetic
1959 designer knockout mice exposures included, which could alleviate
1960 some of the above mentioned drawbacks.

1961 The selection of particular species for whole-organism Mars sample testing
1962 should be based upon (i) state-of-the-art methodology and practices at the
1963 time of the mission and (ii) expert opinion about the suitability and
1964 applicability of employing certain species over other disqualified
1965 candidates. NASA will keep abreast of research developments in whole
1966 organism testing, as well as cultivate and maintain strong liaison
1967 relationships with national and international scientific experts to assure that
1968 appropriate state-of-the-art methods and practices are ultimately employed
1969 and followed.

- 1970 ● *Ecosystems*. Multi-organism population testing is important because
1971 potential biohazard effects may only manifest within the complex
1972 interactions present in ecosystems. The development of microarrays for
1973 analyzing RNA from soil and water will allow both bacterial community
1974 structure and function to be followed in microcosms. Although the
1975 development of reproducible test microcosms will require further research

1976 and development, such assays could be sensitive, fast (on the order of a
1977 week), and include environmental genomics monitoring capabilities.
1978 Microcosm tests could allow monitoring for 'global' characteristics
1979 (e.g., system metabolism, biochemical profile of solid/liquid/gas phases,
1980 etc.), as well as for specific parameters associated with subtle or complex
1981 changes in community structure and function. Additional research will be
1982 required to develop these comprehensive and effective tests.

1983
1984 Sample Size Two different approaches were used to estimate the amount of
1985 sample required for analysis. The first was based on a pre-sorting of the sample
1986 that assumed that 'relevant' biologically interesting sub-samples would be used.
1987 Under this assumption, the amount of sample to be used is dictated by:

- 1988 ● the relevance of the dose being modeled,
- 1989 ● the amount with which the model biological system can be physically
1990 dosed,
- 1991 ● the sample preparation procedure,
- 1992 ● the number of tests to be conducted, and
- 1993 ● the total time Biohazard testing should take.

1994
1995 With this approach, the crudely estimated sample consumption for Biohazard
1996 testing was ten grams.

1997
1998 The second approach did not assume a particular sorting of 'relevant' samples,
1999 but instead used simple statistical methods. Using Earth soil as a crude
2000 reference, a conservative calculation suggested that 15–25 grams of sample
2001 should suffice. These two estimates were quite close despite the very different
2002 approaches used to arrive at them.

2003
2004 Ruling out biohazards in one sample will not allow for extrapolation to other
2005 samples. It will remain a case-by-case task, at least for a considerable period.
2006 This applies even when sub-sampling returned materials. One consideration is

2007 whether samples should be 'homogenized' prior to Biohazard testing. Such a
2008 homogenization is inadvisable because of the loss of information it represents.
2009 For example, sedimentary rocks (which may be in the minority) are more likely to
2010 harbor signs of life than igneous rocks. In addition, since surface conditions may
2011 be toxic to organisms, homogenization with deeper sample components may not
2012 be advisable.

2013

2014 In general, small sample sizes will be required to conserve the returned
2015 specimens, so biological assays that require small quantities are highly
2016 desirable. Examples include cell-based assays (requiring as little as 100
2017 microliters of total fluid volume, making milligram samples potentially adequate)
2018 or the use of small organisms, such as *Arabidopsis* and *C. elegans*.

2019

2020 It was noted that the amount of material needed for destructive testing (consumed)
2021 in biohazard assessments must be determined in consultation with biostatisticians.
2022 Regardless of what starting assumptions are made, the statistics of
2023 sampling will apply, and confidence in 'hazard exclusion' statements can only be
2024 made in the form of "no hazard exists at a concentration greater than X per gram."

2025

2026 Time Needed The time to conduct Biohazard testing was estimated to be twice
2027 the time to conduct the slowest test. It was estimated that most of the results
2028 would be acquired within 90 days, but that 4 to 6 months would be a good
2029 estimate for the completion of the bulk of the testing on the initial samples,
2030 including opportunities to conduct tests on subsequent generations of whole
2031 organisms involved in the testing. As an example, it was estimated that all
2032 Biohazard testing necessary to downgrade the samples from BSL-4 to BSL-3-Ag
2033 would take approximately 6 months, while another 6 months would be required to
2034 downgrade the sample to a lower level of containment or release, as appropriate.

2035

2036 Comments on Controls Control samples clearly are needed for all of the above
2037 experiments. Methods for generating control samples (e.g., dealing with oxidants,

2038 iron, etc.—these contaminants could greatly confound bioassays and not be
2039 modified by some sterilization methods such as high-level irradiation) must be
2040 developed.

2041
2042 Irradiated samples, while somewhat modified, apparently are suitable for much of
2043 the geologic investigations of interest, and along with simulants can be used as
2044 controls. Interestingly, “clean” in terms of geology can mean knowing that certain
2045 elements such as lead are present in concentrations in the parts-per-trillion range.
2046 The important point here is that typical biological containment systems are not
2047 designed with such cleanliness (e.g., molecular/atomic) in mind. A practical
2048 impact of this is that containment/handling equipment and materials should be
2049 characterized in terms of trace concentrations of elements that may be irrelevant
2050 biologically, but damaging to geological and other scientific analyses.

2051
2052 One additional point is that there is a need for pre-launch controls to help rule out
2053 terrestrial contamination. Swab samples, etc., from the assembly and launch
2054 phases and test facility should be taken periodically for two years before mission
2055 launch. This will be a vital piece of the process to establish positive and negative
2056 controls. Negative controls can also be generated at the time of analysis by
2057 treating samples with DNAses, proteases, etc., to subtract out any terrestrial or
2058 Mars biomarkers, so that effects of Mars soil on subsequent assays can be
2059 evaluated.

2060

2061 Future BH Research and Development Needs Further efforts need to be
2062 undertaken to perfect many steps in the final protocol, including:

- 2063 ● A sub-sampling procedure needs to be developed and validated so as to
2064 provide statistical relevance and innate conservatism. This is essential to
2065 ensure that the Biohazard testing is capable of determining the safety of the
2066 samples. Without an effective representative sub-sampling strategy, testing
2067 of the entire sample may be necessary, and untested samples may need to
2068 be kept in containment indefinitely.

- 2069 ● Specific models for use in Biohazard tests have to be chosen or developed.
2070 Each one of them should be validated with terrestrial mimics of martian soil
2071 (possibly with meteoritic minerals from Mars) used “as-is,” or spiked with
2072 known agents to provide a positive control in Biohazard testing.
- 2073 ● Relevant, robust, and reproducible methods of sample preparation and
2074 sample delivery must be developed to ensure the Draft Protocol can be
2075 accomplished effectively.
- 2076 ● The selection of optimal cell and culture systems for use in biohazard and
2077 toxicology assays will be critical. Prior to protocol implementation, research
2078 is needed to select optimum cell and/or molecular assays for BH testing.
- 2079 ● All assay refinements should take into account biohazard containment
2080 issues in their design and implementation. Moreover, it is likely that NASA
2081 will need to coordinate these refinements, and any attendant research
2082 developments, with the toxicology and infectious disease programs at the
2083 National Institutes of Health (NIH), the U.S. Army Medical Research Institute
2084 of Infectious Diseases (USAMRIID), and the Centers for Disease Control
2085 and Prevention (anticipating forthcoming funding increases to integrate
2086 extensive research into infectious diseases and bioterrorism issues). NASA
2087 also must stay abreast of developments in toxicogenomics at the NIH and
2088 in industry, a new field anticipated to replace conventional toxicology
2089 methods over the next five years.

2090

2091 **Facility Requirements**

2092 The size and scope of the facility required to complete the elements of this Draft
2093 Protocol will depend on whether all protocol functions and activities (e.g., sample
2094 receiving and processing, physiochemical characterization, Life Detection studies,
2095 and Biohazard testing) will be conducted at a single SRF or if some elements will
2096 be distributed to secondary labs beyond the SRF. In either case, based on
2097 experience following receipt of lunar samples, the primary SRF should be
2098 designed to be expandable and allow great flexibility in switching functions as
2099 needed. In particular, the SRF should be able to support investigator-driven
2100 research, both to accomplish science objectives that should be addressed prior to
2101 release of unsterilized samples, and to accommodate initial work following the

2102 possible discovery of extraterrestrial life, if necessary. The primary SRF should be
 2103 designed to allow continuous and long-term operation in addition to
 2104 accomplishing its primary goal of receiving the Mars samples and implementing
 2105 the final protocol. There also should be a backup PPL- α facility to contain a subset
 2106 of the initial samples for banking purposes.

2107
 2108 The various elements of the Draft Protocol and appropriate levels of containment
 2109 for completing them are depicted in Figure 6. From a planetary protection
 2110 perspective, these functions can be performed at any facility that meets the
 2111 containment requirements, but as of this writing, no facilities exist which meet PPL-
 2112 α or PPL- β requirements, and only a handful worldwide meet PPL- γ . Similarly, no
 2113 specific test or instrument is precluded from use during the completion of the
 2114 protocol if that test or measurement can be accomplished or placed in
 2115 containment.

2116

2117

TYPE OF TESTS	CONTAINMENT TYPE				
	PPL- α *	PPL- β	PPL- γ	PPL- δ	Other Labs
Physical/Chemical					
Life Detection					
Biohazard					{Fossil}
					
* Simulated martian environment					

2118

2119

2120

Figure 6. Sequential containment requirements by test category.

2121 Regardless of how the final protocol functions are distributed, all ancillary facilities
2122 must meet the same containment guidelines and standard operating procedures
2123 (for items such as personnel monitoring, security assessment, chain of custody
2124 tracking for samples, etc.). There are advantages of utilizing a single facility, at
2125 which the samples are received and all functions up to PPL- γ are performed before
2126 some materials are transferred to PPL- δ facilities to complete the testing. These
2127 advantages include a streamlined management and advisory structure, decreased
2128 sample volume for testing, fewer personnel to monitor for potential exposure,
2129 consolidation of appropriate experts at a single site, and diminished transportation
2130 and logistics concerns. Significantly, this approach assures that the samples are in
2131 the fewest number of facilities practicable, should special actions be necessary if
2132 they are found to contain life or a biohazard. Likewise, there are disadvantages to
2133 building a single large facility instead of a smaller one to be used in combination
2134 with other, existing facilities. Potential disadvantages include increased cost and
2135 complexity, a possible decreased breadth of instrumentation that can be
2136 accommodated, potential delays in recruitment of personnel or complications for
2137 personnel visiting from international partners, and the lack of a second
2138 containment laboratory for the corroboration of test results.

2139

2140 In the final analysis, the facilities required to implement this Draft Protocol, or its
2141 successors, should be the minimum set needed to accomplish the required
2142 planetary protection and science requirements for Mars sample handling in
2143 containment. A variety of facility strategies can be pursued, depending on the
2144 availability of personnel and resources among the partners pursuing a Mars
2145 sample return mission. Further studies of this issue are required, since several of
2146 those strategies can provide for protocol completion as well as the optimal
2147 availability of the samples for scientific studies at the earliest possible time
2148 consistent with Earth safety.

2149

2150 **Future Research and Development Needs** Additional facility-related tasks that
2151 should be addressed in further work include:

- 2152 ● Completely define the PPL containment guidelines and any qualifying or
2153 disqualifying site-related criteria;
- 2154 ● Continue to work with the appropriate agencies and groups²⁶ to explore
2155 containment issues, options, and requirements regarding the refinements
2156 that will be necessary over the coming years to design or retrofit the
2157 appropriate and applicable biohazard containment facility;
- 2158 ● Develop a self-contained structure that could be placed inside of a BSL-4
2159 laboratory, and, as a composite, meet PPL- α containment requirements
2160 (this structure should be able to use robotics to handle the specimens);
- 2161 ● Develop a comprehensive list of equipment, and the required facility
2162 accommodations, for all proposed tests in the Draft Protocol;
- 2163 ● Develop systems needed for some Life Detection testing under simulated
2164 martian environmental conditions, while maintaining PPL- α/β containment;
2165 and,
- 2166 ● Develop cooperative agreements with appropriate BSL-3 and BSL-4
2167 laboratories that can provide experience to NASA personnel prior to the
2168 receipt of Mars samples, or that may act as PPL- δ laboratories thereafter.

2169

2170 **Environmental and Health Monitoring and Safety**

2171 Procedures for monitoring the health and safety of the personnel of the SRF and
2172 the environment in and around the SRF (as well as at secondary sites if used)
2173 must be developed and implemented as part of the final protocol. These will
2174 require a consideration of monitoring over time and an assessment of how long to
2175 continue monitoring, beginning prior to the arrival of Mars samples and continuing
2176 during work on the samples at the SRF and at secondary sites, and for some time
2177 thereafter.

2178

26. Appropriate agencies such as: NIH, USAMRIID, and CDC in the U.S. and Institut National de la Santé et de la Recherche Médicale (INSERM) in France.

2179 Assumptions

- 2180 ● The actual risks associated with the Mars samples are unknown.
- 2181 ● The greatest potential risk is biological. Additionally, the potential existence
2182 of "life as we don't know it," although considered remote, must be
2183 acknowledged and addressed in testing.
- 2184 ● The potential primary exposures will be limited to a small group of trained
2185 professionals in the SRF until more information about the nature of the
2186 specimens is available.
- 2187 ● A high level of security for the SRF and the samples will be maintained as
2188 part of the PPL designation.

2189

2190 Recommended Principles for Development of a Monitoring Program for SRF

2191 Whenever possible, the monitoring plan should use existing regulations and
2192 standards. Since international teams will be working on the Mars samples, the
2193 regulatory standards from participating countries should be reviewed and
2194 considered when developing the final monitoring plan. When considering existing
2195 regulatory standards, the strictest standards, as appropriate for the anticipated
2196 hazards, should apply. Exemptions from existing regulations may be necessary.
2197 For example, differences in the protection of medical information between the
2198 participating countries may be in conflict. The first principle for personnel
2199 monitoring and safety must be to provide optimal protection from anticipated
2200 hazards for the individuals working with Mars samples. Because of the unique
2201 nature of the potential hazards, additional controls beyond those routinely used for
2202 hazard monitoring may be required. The monitoring plan should be designed to
2203 maintain a balance between the estimated risks to individuals, the environment,
2204 and the general population, and the personal and practical impositions of the
2205 monitoring program. The monitoring plan should allow for cross-correlation of the
2206 data from the Life Detection and Biohazard testing with the data from the
2207 monitoring of the SRF personnel and environment, and allow for subsequent
2208 modification of either set of tests.

2209

2210 Potential Hazards Five categories of potential hazards to personnel were
2211 considered: physical hazards, potential chemical hazards from non-biological
2212 toxins, biological hazards, psychological hazards, and loss of containment itself.
2213 The physical hazards include predominantly radiation from the Mars samples
2214 (which is expected to be negligible) and hazards associated with equipment within
2215 the SRF. The potential chemical hazards are predominantly from non-biological
2216 toxins. Any biological hazards will clearly be the most difficult to monitor.
2217 Psychological hazards may arise for personnel working under PPL conditions,
2218 although the psychological risk perception will be far greater for the general public
2219 than for committed risk-taking workers, if generally less immediate. Finally,
2220 ensuring that there is no loss of containment is a significant part of the monitoring
2221 program.

2222

2223 Recommendations for Monitoring

- 2224 ● *Physical Hazard Monitoring (Radiation and Equipment)*. Radiation is a
2225 standard hazard with well-established protocols for protection, handling,
2226 and monitoring. To confirm the expectation that the Mars samples will not
2227 present a radioactivity hazard, a radioactivity measurement should be one of
2228 the initial measurements conducted during the Physical/Chemical
2229 assessments (though technically it is part of the Biohazard testing). The
2230 measurement should be at a level appropriate to assess a biohazard risk,
2231 and need not assess the absolute level of radioactivity present. Standard
2232 radiation safety protocols should be in place prior to the arrival of the Mars
2233 samples, but if the radioactivity level does not represent a biohazard,
2234 monitoring for radioactivity can be discontinued (unless required for
2235 equipment used in the SRF). If a biohazardous level of radioactivity is
2236 detected in the Mars samples, the radioactivity monitoring program would
2237 be continued. Other risks from equipment or facilities can be addressed by
2238 the use of standard procedures, training, and maintenance.
- 2239 ● *Chemical Hazard Monitoring*. A chemical hazard from the Mars samples
2240 would be most likely caused by non-biological, non-replicating toxins, if
2241 present. The presence of toxins will be assessed early in
2242 Physical/Chemical testing. If an unusual substance or chemical is

- 2243 identified, specific monitoring methods for that substance can be designed.
2244 The substance could also be used as a marker for Mars sample breach of
2245 containment monitoring in the SRF and the environment.
- 2246 ● **Monitoring of Containment.** Standard methods for monitoring of
2247 containment can be adapted for use in implementing the PPLs, and can be
2248 used to define a breach of containment or potential personnel exposure. If a
2249 breach occurs within the SRF it can be corrected by standard procedures,
2250 and personnel exposures can be assessed. If a breach occurs to the
2251 environment outside the SRF, a standard procedure should be developed to
2252 assess possible consequences to the environment and/or to humans.
2253 Procedures for handling a breach of the SRF due to different causes
2254 (e.g., leak, disaster, security breach, etc.) should be considered in the
2255 development of the plans for handling a breach.
 - 2256 ● **Monitoring of the Environment.**
 - 2257 ► **Before Mars Sample Arrival.** An assessment of the environment around
2258 the SRF should be made prior to the arrival of the Mars samples.
2259 Environmental monitoring should be implemented in compliance with
2260 the applicable and appropriate regulatory requirements, and in
2261 consultation with relevant U.S. and international agencies. The
2262 environmental assessment should survey the pre-existing conditions,
2263 and include an assessment of the water, air, flora, and fauna. This
2264 survey will likely be accomplished as part of the Environmental Impact
2265 Statement (or Environmental Assessment) required by the U.S. National
2266 Environmental Policy Act and that will be done prior to building the SRF.
2267 During the survey, sentinel species (including microbes, insects, plants,
2268 and animals) can be identified for use as baseline organisms for
2269 monitoring of environmental changes. Consideration should be given to
2270 including some of the same organisms, or closely related organisms, in
2271 Biohazard testing. In case changes in the environment around the SRF
2272 are noted after arrival of the Mars samples, the Biohazard testing results
2273 could assist in determining if the changes are related to the Mars
2274 samples. Environmental monitoring may also include surveillance of
2275 humans in the nearby population, if the facility's location warrants it. If so,
2276 NASA will use attendant, sensitive risk communication practices in
2277 implementation of all public health surveillance initiatives.

- 2278 ➤ *During Mars Sample Handling at the SRF.* Once the Mars samples are
2279 in the SRF, environmental monitoring can focus on the identified sentinel
2280 species and any novel components of the Mars samples, if identified. It
2281 also will be useful to track and record basic weather conditions in the
2282 area of the SRF as part of baseline data. In the event of a breach to the
2283 outside or any unusual occurrences or observations around the SRF,
2284 these data could prove useful in demonstrating either positive or
2285 negative correlation with actual or alleged impacts from SRF operations.
2286 Also, if routine monitoring reveals changes in the environment,
2287 procedures could be undertaken to assess whether an undetected
2288 breach has occurred. SRF personnel would assist with investigating the
2289 cause of the environmental change to establish whether it is related to
2290 the SRF and Mars samples. In the event of a breach, procedures should
2291 be followed to re-establish containment and clean up any detected
2292 contamination.
- 2293 ➤ *After Completion of Life Detection/Biohazard Testing.* The required level
2294 of continued environmental monitoring should be reassessed based on
2295 the outcome of the Mars sample testing protocols. Consideration should
2296 be given to the requirements for maintaining security and containment
2297 within the SRF to assure the proper transition to the long-term curation of
2298 the Mars samples.
- 2299 ● *Monitoring of the SRF Personnel.*
- 2300 ➤ *Before Mars Sample Arrival.* A process of certification for people who will
2301 work in the SRF should be developed that will include security
2302 clearances, medical examinations and tests, and a thorough program of
2303 education about procedures to be employed in health monitoring as well
2304 as on the risks and requirements for employees. Clear inclusion and
2305 exclusion criteria for employees, based on the requirements of the
2306 certification process, should be developed prior to hiring of personnel.
2307 Baseline medical evaluations of personnel should use the existing
2308 medical evaluation standards appropriate at the time the evaluations are
2309 performed. Since the SRF will be functional for a period of time prior to
2310 the arrival of the Mars samples, monitoring before the arrival of the Mars
2311 samples should include several evaluations over time (a period of two

2312 years has been proposed). Recommended baseline evaluations
2313 include a medical history, physical examination, tests on the person
2314 (e.g., chest X-ray), and tests on samples from the person (e.g., blood
2315 and urine). All testing should be as non-invasive as possible, and
2316 maintain a balance between estimated risks from the Mars samples
2317 and the risks associated with the tests. Test specimens should also be
2318 archived for future comparison, if needed, and may include serum,
2319 lymphocytes, semen and/or hair. In addition, neuropsychological
2320 evaluations using standard testing techniques with well-established
2321 interpretation methods should be administered. Symptom data should
2322 be obtained using standardized instruments available at the time of the
2323 SRF commissioning.²⁷

2324 ➤ *During Mars Sample Handling at the SRF.* A schedule for regular
2325 evaluations of personnel should be established, using the same
2326 evaluation methods adopted for the baseline data collection. Procedures
2327 for standard medical management of personnel illnesses should be
2328 available either on-site or with adequate transportation to a medical
2329 facility, as needed. Intervention should be correlated with exposure, or an
2330 identified risk of exposure, to the Mars samples. If an exposure occurs
2331 and the exposed individual has or develops symptoms, the person
2332 should be transferred to a medical facility with BSL-4 containment
2333 capabilities until proper assessment of the individual is accomplished. If
2334 an exposure occurs and the individual does not have or develop
2335 symptoms, procedures for quarantine of the individual should be
2336 developed with specific guidelines as to the length of quarantine
2337 required if the person remains asymptomatic. If an individual becomes
2338 symptomatic and there is no evidence of an exposure, the individual
2339 should be treated as appropriate for the symptoms, and monitoring
2340 should continue as prescribed by the Draft Protocol.

2341

27. The exact survey instrument has not been identified, but it would be possible to use currently existing surveys, similar to the Millennium Cohort Study (U.S.) or the GAZEL Cohort survey (France), sponsored by the U.S. Department of Defense and INSERM, respectively. Current information about these two surveys, may be found online at: <<http://www.gazel.inserm.fr>> and <<http://www.millenniumcohort.org>>.

- 2342 ➤ *After Completion of Life Detection/Biohazard Testing.* The question of
2343 how long to continue monitoring of SRF personnel has to be addressed.
2344 Certainly, the duration of monitoring will be influenced heavily by the
2345 outcomes of the Life Detection and Biohazard testing. Several factors
2346 may need to be considered in this decision, such as the protection of the
2347 workers versus the protection of the general population. Clearly
2348 articulate decisions will be needed on whether to have lifetime
2349 surveillance for the personnel, or to have a mandatory period followed by
2350 optional reporting (if the risk is determined to be low). Monitoring could
2351 become optional if the samples are deemed safe by the Life Detection
2352 and Biohazard testing. The need for surveillance of relatives or people
2353 living close to the personnel should be considered. A distinction should
2354 be made between monitoring for risk management and the continued
2355 collection of data for a research study. The interpretation of personnel
2356 evaluations may require the use of a control group or population-based
2357 estimations of frequencies of different events. If so, sources for this
2358 information should be specified. Finally, the issue should be addressed
2359 on how to ensure provision of adequate health insurance or services to
2360 support any required long-term monitoring and care for the SRF
2361 personnel.
- 2362 ● *Monitoring at Secondary Sites.* The level of monitoring to be used at
2363 secondary sites receiving and working on portions of the Mars samples
2364 should be based on the results of the Life Detection and Biohazard testing.
2365 If the Mars samples are still potentially hazardous, or their biohazard status
2366 is unknown, several points should be considered in developing a protocol
2367 for monitoring at secondary sites. First, secondary sites should be identified
2368 prior to the arrival of the Mars samples, to allow for pre-certification of
2369 personnel and baseline data gathering. Second, all distributions of sample
2370 materials should be tracked, and procedures for monitoring of containment
2371 at the secondary sites should be developed. Third, consider monitoring
2372 personnel at secondary sites using the same protocols used at the SRF.
2373 The number of additional personnel exclusively located at secondary sites
2374 is expected to be small.

2375 If the Mars samples are deemed safe, either through “sterilization” or by
2376 Biohazard test results, the methods should be used for tracking all sample
2377 distributions and all individuals in contact with the samples. In such a
2378 circumstance, only event reporting is needed.
2379

2380 Database Issues A central database with data analysis capabilities and
2381 procedures should be used for environmental data (baseline, monitoring),
2382 personnel data (baseline, during operations, follow-up), secondary site data, and
2383 sample tracking data. Procedures for regular data analysis and reporting should
2384 be developed. Access to, and confidentiality of, the data should be defined and
2385 assured. Data analysis should distinguish between surveillance and research,
2386 with consideration given to the requirements for ethical review and approval for any
2387 research protocols.
2388

2389 Future Research and Development Needs


- 2390 ● Criteria for inclusion/exclusion of personnel to work at the SRF or at
2391 secondary sites.
- 2392 ● The time frame of personnel monitoring, i.e., “lifetime” versus limited period
2393 (according to hazards).
- 2394 ● If long-term monitoring is implemented, which parameters to monitor on a
2395 long-term basis?
- 2396 ● Need for informed consent for testing and possible long-term monitoring.
- 2397 ● Level of baseline testing and monitoring for secondary site workers as
2398 compared to workers at the SRF.
- 2399 ● Protection of individuals from life-insurance or health-insurance
2400 discrimination.
- 2401 ● Procedures for database management and data analysis, with
2402 consideration of confidentiality and security issues.
- 2403 ● Should monitoring be restricted to relevant public health measures, as
2404 opposed to extending the Draft Protocol to allow for epidemiological
2405 research?
- 2406 ● Level of medical facilities needed at the SRF.

2407 **Summary** Monitoring methods for personnel and the environment should be
2408 developed with consideration given to international regulatory, cultural, and ethical
2409 issues. The radiation and chemical risks are considered to be of low probability
2410 and can be assessed early in the chemical testing procedures to reduce the
2411 monitoring burden. Procedures must be developed for database management
2412 and data analysis, with assurances of confidentiality and security of the data.
2413 Procedures for monitoring personnel should include procedures for education and
2414 certification.

2415

2416 **Personnel Management Considerations in Protocol Implementation**

2417 The staffing of the Sample Receiving Facility(-ies) can be accomplished in a
2418 number of ways. For example, scientists can be recruited to fill permanent
2419 positions at the SRF, or could be selected through a competitive grants program
2420 for work at the SRF, or some combination of the two approaches. Considering the
2421 variety of tasks that must be accomplish during design, construction, and
2422 operation of the facilities, as well as during implementation of the final protocol, it
2423 will be advisable to use a variety of different personnel selection processes.
2424 Personnel should be hired progressively during the development of the project
2425 and the facility(-ies). The functions and responsibilities of the Director's position
2426 will be substantially aided by appropriate committees and advisory groups. In the
2427 event that more than one facility is used, the required methods and procedures
2428 outlined in the Draft Protocol should be applied beyond the SRF to any facility or
2429 site planned to handle martian samples during the implementation of the final
2430 protocol. Because researchers and the public worldwide will have an interest in
2431 returned martian materials, the international character of the program should be
2432 respected throughout the entire process. Figure 7 on the next page presents a
2433 high level schedule and overview of the process from now until the samples are
2434 returned to Earth. One concept of the functions, staffing requirements, and
2435 organization for a Mars Sample Receiving Facility, is further elaborated in Figures
2436 8, 9, and 10. These figures outline staffing needs and proposed organizations at
2437 10-, 5- and 3-years before the arrival of actual samples at the SRF.



2438

2439

2440 Figure 7. Example overall timetable of the required activities to design, build, and
 2441 operate the SRF. The double-headed arrows indicate timing of the staff organization
 2442 described in the subsequent figures (EVT = Experiment Verification Test).
 2443

2444

2445 These proposed management, staffing, and organizational frameworks amount to
 2446 a working hypothesis for the design of the building and operation of the SRF,
 2447 based on the following assumptions:

2448

- The protocol must be fully and successfully tested before the actual
 2449 handling of the martian samples. The exact makeup and sequence of the
 2450 Experiment Verification Tests (EVTs) are TBD.

2451


- It is estimated that a complete EVT will last approximately 6 months and at
 2452 least one complete EVT must be demonstrated successfully before actual
 2453 handling of the returned samples. Thus, the first EVT must begin no later
 2454 than 18 months before the returned samples arrive at the SRF in order to

- 2455 allow enough time to adjust and repeat the EVT, if necessary (at least 9-10
2456 months before experiments begin on actual returned samples).
- 2457 ● These EVTs are consistent with the recommendation of the SSB (1997) and
2458 earlier Workshops in this Series that the SRF be operational two years
2459 before the arrival of the actual Mars samples. These EVTs are part of the
2460 normal operational testing.
 - 2461 ● Based on experiences at other BSL-4 laboratories in the United States and
2462 France, no less than one-year is required to staff and properly train the
2463 technical and scientific personnel.
 - 2464 ● Commissioning of the SRF, which can be performed in parallel with the
2465 staffing and training, will require at least 18 months.
 - 2466 ● In order to accommodate the staffing, training and commissioning
2467 requirements of the SRF, construction of the facility must be finished 3 years
2468 before the actual operations. From past experiences, in France and the
2469 United States, construction of the facility itself will also require 3 years.
 - 2470 ● It is estimated that about 3 years will be needed to develop design
2471 specifications and plans for the SRF, and obtain necessary authorizations
2472 to build the facility. To accommodate all the activities necessary to design,
2473 build and operate an SRF, the entire process must begin fully ten years in
2474 advance of sample return.

2475
2476 To illustrate one approach to staffing and organization that meets facility and
2477 protocol requirements, the text below provides specific details related to the
2478 recommended staffing and organizational plans. It is emphasized that these
2479 scenarios are not fixed requirements of this Draft Protocol, but are intended to
2480 provide a conceptual structure on which to base future organizational and staffing
2481 plans.

2482
2483 10 Years in Advance As soon as the decision is made to build and/or update a
2484 Mars SRF, ~10 years before the actual operations, four positions should be staffed
2485 in order to prepare specifications for future activities and a substantive review of
2486 the design of the facility (see Figure 8). The key positions to be filled 10 years prior

2487 to sample return are the Project Manager/Director, a Director for Administration, a
 2488 Project Scientist/Director for Science, and an Environment, Health, and Safety
 2489 Officer. The Director, who is responsible for the overall sample handling project
 2490 implementation, will have the assistance of an SRF Oversight Committee. This
 2491 Committee will monitor progress and assure compliance of the project with the
 2492 final protocol and with whatever science requirements are to be implemented in
 2493 the Facility. In this example, it is anticipated that the initial Director will have a
 2494 background in scientific facility engineering, and that transition to a Director with a
 2495 science background will occur after construction of the facility is assured. The
 2496



2497

2498 Figure 8. Top-level staffing requirements and structure of the SRF at 10 years prior
 2499 to arrival of the returned sample(s). Permanent positions are in plain boxes;
 2500 committees are in grey boxes. Not all positions are full-time.

2501

2502 Director will be assisted by the Environment, Health, and Safety Officer to ensure
2503 that the actual design requirements related to these critical topics are
2504 implemented properly. A Director for Administration will focus on budget and
2505 staffing issues, and the development of the staffing plan to cover the life of the
2506 project. Additional engineering support (e.g., the Facility Engineer) would be added
2507 as necessary.

2508
2509 The Project Scientist/Director for Science will coordinate the work of scientific
2510 committees and working groups that will develop science specifications and
2511 support the design process for their respective disciplines or areas. Also at this
2512 point in the project, a Communications Officer should be available, at least on a
2513 part-time basis, to ensure attention to risk communications and outreach—
2514 keeping the community informed and identifying and answering questions
2515 regarding the SRF. All communications, plans, and activities at the SRF should be
2516 consistent with those outlined in any comprehensive communication plan
2517 developed for the mission and the Mars exploration program as a whole (see the
2518 section titled "Maintaining and Updating the Protocol," below).

2519
2520 From the beginning of the process, three different kinds of committees should be
2521 installed to help the Directors and Scientific Discipline Heads in overseeing their
2522 changing responsibilities:

- 2523 ● The Science Working Group (SWG) will be charged with helping to guide the
2524 overall project during the construction phase, to provide recommen-
2525 dations and expertise in assuring its compliance with sample scientific
2526 requirements and the final protocol. The members of the SWG will be
2527 chosen from an *ad hoc* set of scientists representing the required
2528 disciplines and expertise. Later, they will be replaced by the Investigators
2529 Working Group, comprised of selected Principal Investigators from an
2530 open competition seeking proposals for sample analysis activities within
2531 the Facility.
- 2532 ● Scientific design committees will be specialized in four disciplines, Life
2533 Detection, Biohazard testing, Physical/Chemical, and Curation, with

2534 members designated by the agencies participating in the mission. These
2535 committees will prepare the design and review and oversee the project to
2536 ensure the facility can operate consistent with the operational aspects of the
2537 planned protocol. As soon as the Scientific Discipline Heads are hired,
2538 these committees will become Discipline Advisory Panels to assist them.


2539 ● Finally, the SRF Oversight Committee will be composed of 12 to 15
2540 members selected by the Program leadership, perhaps with some cross-
2541 membership from the NASA Planetary Protection Advisory Committee and
2542 the French Planetary Protection Committee. These committees will be in
2543 charge of reviewing the overall process and the proposed measures to
2544 comply with the requirements of the final protocol. The Science Oversight
2545 Committee will report to Program Management and the Planetary Protection
2546 Officer, above the level of the Project Manager/Facility Director. However, it is
2547 expected that they will interact directly with that Manager on a regular basis.
2548

2549 Membership on the various committees will be staggered to ensure an
2550 appropriate turnover without losing the “project memory.” Agencies involved with
2551 the SRF should set up jointly an international search committee for recruitment of
2552 the Directors, various functional managers, the Facility Engineer, and the Scientific
2553 Discipline Heads.

2554

2555 5 Years in Advance At roughly midway through the construction of the facility, the
2556 Scientific Discipline Heads should be hired for each required scientific discipline
2557 (see Figure 9 on the next page). These managers will ensure that construction is
2558 completed properly to accommodate the specific needs of their disciplines. With
2559 the help of experts working as part of the scientific working group and discipline
2560 advisory panels, they will complete the general and specific operating procedures
2561 to handle the martian samples and the training program for staff to be hired. At this
2562 point, a Facility Administrative/Staff Manager will also be hired to assist in the
2563 hiring of the technical staff and prepare for future administrative and personnel
2564 needs of the facility.


2565



2566
 2567 Figure 9. Top-level staffing requirements and structure of the SRF at 5 years prior
 2568 to arrival of the returned sample(s). Permanent positions are in plain boxes;
 2569 committees are in grey boxes.
 2570

2571
 2572 3 Years in Advance In order to have a fully operational facility two years before
 2573 samples are returned, the final staffing and training of various operational
 2574 positions must begin three years prior to actual operations (see Figure 10). At this
 2575 time, the required supporting groups, such as an Institutional Bio-Safety
 2576 Committee (IBSC) and an Institutional Animal Care and Use Committee (IACUC),
 2577 will be formed, and staff necessary to support facility operations, administrative
 2578 functions, communications, and safety program implementation will be added,
 2579 Also at this time, it is anticipated that the *ad hoc* Science Working Group (which
 2580 until this time would have dealt with both science issues and issues of planetary

2581 protection protocol compliance), will be supplanted by an Investigators Working
 2582 Group selected through an open solicitation that would provide for scientific
 2583 investigations to be accomplished within the facility. The relationship of these
 2584 selected science investigations to the accomplishment of the protocol objectives
 2585 may be close or distant, depending on the strategy undertaken to implement the
 2586 protocol in its final form.
 2587



2588
 2589 Figure 10. Staffing requirements and structure of the SRF at 3 years prior to arrival
 2590 of the returned sample(s); permanent positions are in plain boxes; committees are in
 2591 grey boxes; stippled boxes indicate an Institutional Bio-Safety Committee (IBSC) and
 2592 an Institutional Animal Care and Use Committee (IACUC).
 2593
 2594

2595 **Future Considerations** Three major issues will require further consideration in the
2596 overall staffing of the SRF.

2597 1. Currently, no one has experience in simultaneous operations or activities in
2598 combined BSL-4 and cleanroom conditions as will be needed for PPL- α
2599 through PPL- δ . The advice of experts from the pharmaceutical or micro-
2600 process industries would be helpful.

2601 2. Details on the optimal staffing mix at the SRF must be considered further. It
2602 is not clear what mix of government employees, semi-permanent staff
2603 employees, outside contractors, and guest scientists will be needed to staff
2604 the facility and implement the final protocol. In planning for facility staffing
2605 and operations, international access and participation should be
2606 considered throughout the process.

2607 3. In order to comply with planetary protection constraints and protocol
2608 requirements, a sustained and adequate budget will be needed throughout
2609 the design, construction, and implementation phases of this project.
2610

2611 **Contingency Planning for Different Protocol Outcomes**

2612 Developing contingency plans for different outcomes of the final protocol will
2613 require anticipating how the scientific community might interpret test results and
2614 react under a variety of possible scenarios following the return of martian
2615 samples. In addition to considering how to interpret possible scientific results, it
2616 will be important to plan how to respond in the face of possible breaches in
2617 containment. Recommended response to various likely scenarios are discussed
2618 below:

2619
2620 **Organic Carbon** It is likely that carbon will be found in sample materials. The
2621 sensitivity of current and future methods will be very high, so that at least some
2622 level of contaminants should be detected, and perhaps carbon compounds from
2623 Mars, as well. The existing base of knowledge on meteorites and other material
2624 collected from space will be useful in providing baseline information to help guide
2625 these investigations. Since the Viking results focused on volatile organics, further
2626 attention to the question is appropriate. *In situ* measurements of non-volatile

2627 organics on missions prior to the sample return mission would be useful to gauge
2628 predictions of anticipated sample organic content.

2629

2630 Extant Life or Biomarkers Positive If extant life or evidence of biomarkers are
2631 detected in the samples, all work on the samples will continue to be done in strict
2632 containment until more definitive data can be gathered (see *Release Criteria* and
2633 *Biohazard Testing* sections, above.) Maximum effort should be made to determine
2634 if any of the positive results are originating from Earth life or Mars life. Information
2635 management will become an issue, both for scientific communication and in
2636 shaping the debate among scientists. It will be important to plan for how and when
2637 initial information, with its attendant uncertainties, should be disseminated to the
2638 public.

2639

2640 Non-Earth Life Confirmed In keeping with the SSB recommendations [SSB 1997],
2641 and the stated release criteria, sample materials will be released from
2642 containment only if they are shown to contain no extraterrestrial life-forms, or they
2643 are sterilized prior to release. If non-terrestrial life is confirmed, a previously
2644 constituted SRF Oversight Committee will need to review the protocol, the steps
2645 taken in support of the protocol, and ongoing provisions for containment. If a
2646 portion of a sample is confirmed as positive for non-terrestrial life, subsequent
2647 testing and analyses on all sample materials will continue in containment. This
2648 means that all physical, chemical, and geological characterization, as well as Life
2649 Detection and Biohazard tests requiring non-sterilized material should continue to
2650 be done in strict containment, either in the SRF or in any other test facilities that
2651 may be used. Experimentation on methods to sterilize samples containing the
2652 newly-discovered life should begin in conjunction with investigations of
2653 appropriate biological culture conditions. Once appropriate biological sterilization
2654 techniques can be validated, detailed plans for distribution of samples can be
2655 developed or revised in order to meet the established or revised scientific
2656 objectives. Management issues will include administrative and technical
2657 procedures for scientific study and curation, as well as informing the public.

2658 Although it is premature to develop specific recommendations at this time, it is
2659 possible to identify issues that will need further discussion in advance of sample
2660 return. The concerns fall into three broad categories: Science and Testing; Facility
2661 and Technological; and Policy and Administrative.

2662

2663 Science and Testing Confirmation of a preliminary discovery of martian life should
2664 require a careful reconsideration of results from many parts of the final protocol,
2665 ranging from a review of preparation, through scanning and testing methods, to
2666 verification of biocontainment materials and sterilization techniques, and a
2667 reassessment of conditions for banking, storage, transportation and curation. If
2668 evidence of any martian life is found, there should be a plan to aggressively
2669 expand the studies with the expectation that there will be multiple, additional life
2670 forms, given that evidence that life can be supported on Mars. In addition, it will be
2671 important to understand the culture and environmental conditions that are required
2672 to maintain and perhaps to grow the new life-form to obtain more material for study
2673 in the lab, and what precautions are needed in the process. Also, it will be
2674 important to review the final protocol to recommend modifications in physical,
2675 geological, and chemical tests of sample materials, adding or deleting tests as
2676 needed.

2677

2678 Facility and Technological Concerns Questions about the adequacy of the SRF to
2679 maintain the new life form must also be addressed, including the possible need to
2680 add equipment, change operations, review emergency plans, or upgrade the
2681 facilities because of what has been found. Concerns about security should also
2682 be reconsidered, especially in view of the potential disruptive activities of any
2683 terrorists or 'radical' groups that may be opposed to sample return. The
2684 advisability of allowing distribution of untested sample material outside the SRF
2685 may need to be reconsidered, as well.

2686

2687 Policy and Administrative Concerns If martian life is detected, both short-and long-
2688 term policy issues will arise. The short-term listing of concerns relates to

2689 procedures regarding access to and distribution of sample materials, as well as
2690 to the publication and review of research findings. The chain of custody of sample
2691 materials will be important in the assessment of data quality, as well as in
2692 addressing the legal requirements of who is allowed to "touch" the sample (or
2693 verifying who has handled the sample appropriately or inappropriately). It will be
2694 critical to incorporate chain-of-custody considerations into the final protocol well in
2695 advance of sample return.

2696

2697 As part of sample return planning, it will be important to develop an organized
2698 communication plan which will lay a strong foundation in public understanding
2699 and acceptance prior to the mission, and allow for an open dialogue with all
2700 sectors of the public. Such a plan should include consideration of the diverse
2701 questions, concerns, and issues likely to be raised, including those related to the
2702 mission and spacecraft operations, the sample return and Biohazard testing, the
2703 administrative and legal matters associated with the effort, and to the potential
2704 implications of discovering extraterrestrial life. Plans should be developed well in
2705 advance in order to avoid a frenzied, reactive mode of communications between
2706 government officials, the scientific community, the mass media, and the public.
2707 Any plan that is developed should avoid a NASA-centric focus by including linkages
2708 with other government agencies, international partners, and external
2709 organizations, as appropriate. It will also be advisable to anticipate the kinds of
2710 questions the public might ask, and to disclose information early and often to
2711 address their concerns, whether scientific or non-scientific.

2712

2713 In the long term, the discovery of extraterrestrial life, whether extant or extinct, *in situ*
2714 or within returned sample materials, will also have implications beyond science
2715 and the SRF *per se*. Such a discovery would likely trigger a review of sample return
2716 missions, and plans for both robotic and human missions. Legal questions could
2717 arise about ownership of the data, or of the entity itself, potentially compounded by
2718 differences in laws between the United States and the countries of international
2719 partners. In any event, ethical, legal and social issues should be considered

2720 seriously. Expertise in these areas should be reflected in the membership on
2721 appropriate oversight committee(s).

2722

2723 *Contradictory/Inconsistent Results* Given the number of techniques, spanning
2724 several scientific disciplines, it is very likely that contradictory or inconsistent
2725 results will be found. Differences in the sensitivity of methods will exist and
2726 confidence in the reliability and level of experimental controls will differ among
2727 procedures. It is important to stress the need for replication of experiments and
2728 duplication of results among multiple sites to add confidence to the results
2729 assessed. In addition, it will be important to follow a strict scientific procedure for
2730 interpreting data and making decisions about sample materials. There is a need
2731 to involve multidisciplinary experts and groups in the overall decision making
2732 process as well as in devising procedures for drawing conclusions, certifying
2733 results, and deciding whether samples are safe enough to be released to lower
2734 containment levels.

2735

2736 *Application of Release Criteria* According to the COMPLEX report on ‘*The*
2737 *Quarantine and Certification of Martian Samples*’ [SSB 2002]:

2738 *“If the samples are shown to be altogether barren of organic matter, to*
2739 *contain no detectable organic carbon compounds and no other evidence of*
2740 *past or present biological activity, untreated aliquots of the samples should*
2741 *be released for study beyond the confines of the Quarantine Facility.”*

2742

2743 The stated goal of the MSHP Workshop Series was to design a protocol to test
2744 returned sample(s) for biohazards and the presence of martian life, to ensure that
2745 a sample is safe to be released without sterilization, for further study. The release
2746 criteria listed in this Draft Protocol are consistent with the cited NRC
2747 recommendation, but this Draft Protocol imposes the additional requirement to
2748 complete Biohazard testing on all samples, taking into account the possibility of
2749 non-carbon-based life. As such, this Draft Protocol is more conservative than the
2750 most recent NRC recommendation [SSB 2002], but justifiably so in terms of what
2751 is known and not known about life elsewhere.

2752 Conversely, arguments have been advanced suggesting that a sterilization step be
2753 added to the protocol for “good measure,” for the release of any materials, even if
2754 the samples are devoid of organic compounds and do not demonstrate any
2755 biohazard. After an evaluation of the arguments advanced regarding this concept,
2756 both pro and con, this additional step is not required by this Draft Protocol. Central
2757 to an understanding of the arguments is the question of risk, i.e., Can *any* protocol
2758 be guaranteed to be absolutely risk-free? If not, what is an acceptable level of risk
2759 (for example, one that approximates the risk from the natural influx of martian
2760 materials into Earth’s biosphere)? And, is there any treatment method that can
2761 eliminate all risks from the returned samples, while preserving them for the
2762 detailed scientific study envisaged by the scientific community? Clearly, the issue
2763 of sterilization will require serious additional attention and research well in
2764 advance of sample return. Likewise, the safety of releasing materials that have
2765 passed both Life Detection and Biohazard testing should be carefully challenged
2766 through a rigorous quality assurance program applied to the completion of the
2767 Draft Protocol.

2768

2769 *Breach of Containment* Anticipating a containment breach and planning for such
2770 an event is an essential element of facility management. The responses to a
2771 breach will depend on where it occurs and what happens. Conceivably, it could
2772 occur in an area with a high population density or in a remote location. The breach
2773 could be a result of an accident or a crime—as a result of activity either outside or
2774 within containment. Required steps on how to handle breaches (based on long
2775 term experience and emergency plans for handling pathogenic biological material
2776 under BSL-3 and BSL-4 containment), are known. Additional information for
2777 responding to breaches and containment problems has been gained through
2778 decades of experience in handling lunar and other extraterrestrial materials.

2779

2780 Clearly, an emergency plan will be needed well in advance to develop
2781 recommended responses to various breach scenarios. The first steps will involve
2782 investigation of the degree of compromise, considering both biosafety and sample

2783 integrity. Full documentation of any breach event will be required as well as
2784 identifying the degree of sample compromise, what organizations or personnel
2785 should be involved in all phases of a response, and how notifications and
2786 communications should be handled. The plan should focus on all aspects of
2787 mitigation, cleanup, and recovery from perspectives of both biosafety and sample
2788 integrity (e.g., decontamination of the area, sample recovery, re-packaging and
2789 labeling as compromised, or destruction if required, etc.).

2790

2791 **Maintaining and Updating the Protocol**

2792 The recent report from the NRC [SSB 2002] recommended:

2793 *“A continuing committee of senior biologists and geochemists that includes*
2794 *appropriate international representation should be formed and charged with*
2795 *reviewing every step of the planning, construction, and employment of the*
2796 *Mars Quarantine Facility. The committee should be formed during the*
2797 *earliest stages of planning for a Mars sample return mission. Members of*
2798 *the committee should also participate in the design of the spacecraft and*
2799 *those portions of the mission profile where biological contamination is a*
2800 *threat.”*

2801


2802 This Draft Protocol refers to the necessary committees, including the SRF
2803 Oversight Committee, and the NASA Planetary Protection Advisory Committee
2804 (PPAC). The protocol implementation and update process will require
2805 establishment of these expert oversight and review committees, re-evaluations of
2806 proposed plans at key points in time before sample return, and open
2807 communication with scientists, international partners, and the public regarding
2808 risks, benefits, and plans. The scope of the task is summarized in Figure 11. A
2809 narrative explanation of recommendations and activities in the process follows.

2810

2811 *Final Scientific and Policy Reviews* Reviews of the Draft Protocol should provide
2812 for the highest degree of scientific scrutiny and evaluation.²⁸ The evaluation should
2813 be conducted jointly by scientific organizations from both the United States and

28. This Protocol was jointly derived by NASA and CNES, reflecting their intention to jointly accomplish the sample return mission. A final protocol should reflect reviews by all of the eventual mission partners.

2814 France (and other countries, as appropriate) to avoid prolonged negotiations and
2815 resolutions that may arise when such reviews are conducted separately. This
2816 review should probably occur at the level of the National Research Council in the
2817 United States, and its equivalent scientific organization in France, whichever is
2818 most appropriate (among the French institutions discussed were Centre National
2819 de la Recherche Scientifique (CNRS), or representatives of various
2820 Etablissements Publics à Caractère Scientifique et Technique (EPST), including –
2821 but not exclusively – CNRS or Académie des Sciences). Final decisions about
2822 which institutions should be involved in scientific reviews are TBD, but should
2823 include NASA's Planetary Protection Advisory Committee, and the French multi-
2824 Ministry-sponsored Planetary Protection Committee.



2825
2826

2827

Figure 11. Protocol update and implementation process.

2828 Clarity of Meaning and Terminology Clarity of meaning is essential to the
2829 implementation of any process especially when the process involves international
2830 agreements. Therefore, absolute consistency should be used in the language for
2831 any documents and charters associated with the eventual final protocol. When the
2832 actual definition of a word or phrase is in dispute, reference should be made to
2833 those definitions or meanings that are standard and accepted when interpreted at
2834 the international level. Clarity in terminology will be especially important when
2835 describing levels of containment to avoid confusion caused by mixing United
2836 States and French definitions of BSL and P4 containment. PPL containment
2837 definitions should be jointly derived to avoid these mixed meanings.

2838

2839 Ethical and Public Reviews Evaluations of the proposal should be conducted both
2840 internal and external to NASA and Centre National d'Etudes Spatiale (CNES) and
2841 the space research communities in the nations participating in the mission. An
2842 ethical review should be conducted at least at the level of the Agencies
2843 participating and these reviews made public early in the process (in France, the
2844 national bioethics committee, Comité Consultatif National d'Ethique pour les
2845 Sciences de la Vie et de la Santé, CCNE, is the appropriate organization). The final
2846 protocol should be announced broadly to the scientific community with a request
2847 for comments and input from scientific societies and other interested
2848 organizations. Broad acceptance at both lay public and scientific levels is essential
2849 to the overall success of this research effort.

2850

2851 Future Modifications to the Protocol When a final protocol has been adopted and
2852 approved by a consensus of appropriate scientific organizations, few changes
2853 should be made to its content. Changes should be made as scientific information,
2854 methodology, and/or technology improve between the time of the approval and the
2855 actual physical implementation of the final protocol within the SRF laboratories.
2856 Changes in methodologies or technologies to be used in implementing the final
2857 protocol may be considered if a proposed change would meet the following
2858 criteria:

- 2859 ● Increases the sensitivity or selectivity of the test,
- 2860 ● Reduces the length of time necessary for a test without a reduction in
- 2861 sensitivity or selectivity,
- 2862 ● Reduces the complexity of the sample handling process,
- 2863 ● Increases the overall safety of the process,
- 2864 ● Reduces the chances of contamination to the sample or the environment,
- 2865 ● Reduces the cost of the process, or
- 2866 ● Represents a new technology or method that has the broad, general
- 2867 acceptance of the scientific community.

2868

2869 Changes to the final protocol should receive appropriate expert review at the same
2870 level as the initial document.

2871

2872 Advisory Committees and Expert Panels Changes in scientific methodology and
2873 instrumentation are inevitable due to the long development time envisaged for this
2874 mission. This necessitates long term, consistent, input and advice from the
2875 external scientific communities of the partners engaged in the mission. To
2876 facilitate this process, a standing Planetary Protection Advisory Committee (PPAC)
2877 is being appointed in the United States to provide input to the NASA Office of Space
2878 Science and the NASA Planetary Protection Officer, and that a similar standing
2879 committee (Planetary Protection Committee, PPC) is being appointed in France.
2880 Both of these committees should provide for the participation of representatives of
2881 governmental regulatory agencies to make use of their particular expertise as well
2882 as to enhance communications among those various agencies, NASA, and CNES.

2883

2884 Standing joint working committees or specialized expert panels should be
2885 appointed (perhaps in cooperation with the SRF's Science Working Group) with
2886 appropriate expertise to provide support and advice to the United States PPAC and
2887 the French PPC in each of three specific areas: technical processes, scientific
2888 procedures, and safety/biosafety issues. To provide the most effective level of
2889 support, these groups should be comprised of members with expertise in a

2890 particular area of concern and organized into individual panels. No expert should
2891 be a member of more than one panel. The overall membership of the committees
2892 and expert panels should be selected to meet the specific needs of the agencies,
2893 and should represent the scientific goals of the agencies and the external science
2894 communities. Their work should aim at providing the respective agencies with
2895 information essential to the success and safety of the Mars sample return
2896 missions. These panels and committees may function jointly or independently
2897 depending on the specific need.

2898

2899 The PPAC and French PPC should receive the annual reports of the three panels,
2900 which will also provide annual written reviews to the NASA Planetary Protection
2901 Officer and, in France, to the appropriate Minister to whom the committee reports.
2902 These reviews should include relevant operational issues and concerns and
2903 provide risk assessments of the technical processes, scientific procedures, and
2904 safety/biosafety plans and processes. These reviews should be made available to
2905 scientific and professional organizations with interests in the mission activities.

2906

2907 Communications Unusual or unprecedented scientific activities are often subject
2908 to extreme scrutiny at both the scientific and political levels. Therefore, a
2909 communication plan must be developed as early as possible to ensure timely,
2910 and accurate dissemination of information to the public about the sample return
2911 mission, and to address concerns and perceptions about associated risks. The
2912 communication plan should be pro-active and designed in a manner that allows
2913 the public and stakeholders to participate in an open, honest dialogue about all
2914 phases of the mission with NASA, policy makers, and international partners. Risk
2915 management and planetary protection information should be balanced with
2916 education/outreach from the scientific perspective about the anticipated benefits
2917 and uncertainties associated with Mars exploration and sample return. The
2918 communication plan should also address how the public and scientific community
2919 will be informed of results and findings during Life Detection and Biohazard
2920 testing, including the potential discovery of extraterrestrial life. Because of the

2921 intense interest likely during initial sample receipt, containment, and testing,
2922 procedures and criteria should be developed in advance for determining when and
2923 how observations or data may be designated as “results suitable for formal
2924 announcement.” Details about the release of SRF information, the management of
2925 the communication plan, and its relationship to the overall communications effort
2926 of the international Mars exploration program should be decided well in advance of
2927 the implementation of this protocol.

2928

2929 Flow Charts and Timelines In order to assure the rational use both of the facilities
2930 and sample materials, development of appropriate flow charts and time lines will
2931 be needed to coordinate the complex series of interrelated procedures. Safety
2932 issues must be prominent at all significant decision points in the process
2933 (e.g., release from containment, and downgrading to lower level of containment).
2934 It is essential to identify the critical points for these decisions in advance so that all
2935 participants understand their timing, and to ensure that such decisions are not
2936 negotiated in haste. Flow diagrams are intended to coordinate complex testing
2937 and inclusion of all required elements, especially those concerning biosafety and
2938 biohazards leading to the sharing of sample material with the external scientific
2939 community. In addition to containing timelines, procedures and processes, flow
2940 charts should also include key decision points for changing the status of the
2941 sample to a less restrictive PPL and proceeding in a particular direction along
2942 branches of the decision tree. Each such chart should incorporate a risk tree and
2943 assessment process.

2944

2945 Workshops/Reviews The need to change schedules and procedures may be
2946 anticipated during the time between now and sample return. To provide assurance
2947 that rules exist between the involved international partners and the scientific
2948 communities, two workshop/reviews should be scheduled prior to sample return
2949 to Earth in order to reaffirm details about process, methodology, safety, and
2950 release criteria. The first review should be conducted at the conclusion of the
2951 facilities design phase to determine if the physical structure meets the scientific

2952 and safety standards as defined within the specifications. In addition, the first
2953 workshop should review the existing procedures that will be conducted within the
2954 facility(ies) to confirm the specific flow chart outlining the approved sequence of
2955 tests and analyses. A second similar workshop/review should occur after the
2956 samples have been collected on Mars, but in advance of their actual return to Earth
2957 for evaluation. Details about who should coordinate these workshop/reviews and
2958 modify schedules or procedures are TBD.

2959
2960 *Preparations and Processes for Decision Making about Release of Samples* It will
2961 be important to make advanced preparations for organized data interpretation and
2962 decision making. These preparations will be especially critical in the event that a
2963 distinctly martian life-form is found within the returned samples. While it is
2964 impossible to develop details of the protocol at this time, it will be crucial to have
2965 considered how decisions will be made, by whom, and based on what principles if
2966 an extraterrestrial life-form is discovered. A specific committee should be
2967 established at least a year ahead of sample return to develop contingency
2968 protocols and processes that will be in place if and when martian life is found and
2969 verified. It is likely that protocol test results will not lead to unanimous decisions in
2970 all instances. It will thus be important to have a review and approval infrastructure
2971 for handling decisions about whether to release sample materials from
2972 containment, or reduce containment to a lower level upon completion of the final
2973 protocol tests. Addressing the overall decision making process in a formal
2974 manner will be critical for drawing conclusions, certifying results, and deciding
2975 whether samples are releasable or not. Any decision to release samples should
2976 involve selected multidisciplinary experts and groups, such as an Interagency
2977 Committee on Back Contamination (ICBC) similar to the one used during the
2978 *Apollo* program. The U.S. PPAC and French PPC should be involved in reporting to
2979 relevant bodies in their respective countries. Details on the structure(s) associated
2980 with decision making are TBD.

2981

2982 The organizational structures, management plans, charters and reporting lines for
2983 many of the proposed committees and groups will need to be developed in the
2984 coming years. Many questions cannot be resolved until additional details on facility
2985 design, operational logistics, mission architecture or anticipated schedules are
2986 made available. Future work should use this Draft Protocol to support the
2987 development of these items.

2988

2989

2990

APPENDIX A: MSHP WORKSHOP SERIES BASIC ASSUMPTIONS

The Mars Sample Handling Protocol (MSHP) Workshop Series was designed to touch on a variety of questions in pursuit of the stated objective, such as: “What types/categories of tests (e.g., biohazard; life detection) should be performed upon the samples? What criteria must be satisfied to demonstrate that the samples do not present a biohazard? What constitutes a representative sample to be tested? What is the minimum allocation of sample material required for analyses exclusive to the Protocol, and what Physical/Chemical analyses are required to complement biochemical or biological screening of sample material? Which analyses must be done within containment and which can be accomplished using sterilized material outside of containment? What facility capabilities are required to complete the Protocol? What is the minimum amount of time required to complete a hazard determination Protocol? By what process should the Protocol be modified to accommodate new technologies that may be brought to practice in the coming years (i.e., from the time that a sample receiving facility would be operational through the subsequent return of the first martian samples?)

To keep the Workshops focused, a set of basic assumptions were provided to guide and constrain deliberations; these assumptions were:

1. Regardless of which mission architecture is eventually selected, samples will be returned from martian sites which were selected based on findings and data from the Mars Surveyor program missions.
2. Samples will be returned *sometime in the next decade*.
3. Samples will not be sterilized prior to return to Earth.
4. The exterior of the Sample Return Canister will be free from contamination by Mars materials.
5. When the Sample Return Canister (SRC) is returned to Earth, it will be opened only in a Sample Receiving Facility (SRF) where samples will

- undergo rigorous testing under containment and quarantine prior to any controlled distribution ('release') for scientific study.
6. The amount of sample to be returned in a SRC is anticipated to be 500-1000 grams.
 7. The sample will likely be a mixture of types including rock cores, pebbles, soil, and atmospheric gases.
 8. The amount of sample used to determine if biohazards are present must be the minimum amount necessary.
 9. Samples must be handled and processed in such a way as to prevent terrestrial (chemical or biological) contamination.
 10. Strict containment of unsterilized samples will be maintained until quarantine testing for biohazards and Life Detection is accomplished. Sub-samples of selected materials may be allowed outside containment only if they are sterilized first.
 11. The SRF will have the capability to accomplish effective sterilization of sub-samples as needed.
 12. The SRF will be operational two years before samples are returned to Earth.
 13. The primary objective of the SRF and protocols is to determine Whether the returned samples constitute a threat to the Earth's biosphere and populations (not science study *per se*) and to contain them until this determination is made.

APPENDIX B: REFERENCES

- Allen, C., *et al.*, "Effects of sterilizing doses of gamma radiation on Mars analog rocks and minerals," *J. Geophysical Research* 104, 27043-27066 (2000).
- Bruch, C. W., R. B. Setlow, and J. D. Rummel, eds., *Mars Sample Handling Protocol Workshop Series, Workshop 2a Final Report*, NASA CP-210924 (2001).
- Carr, Michael H., *et al.*, *Mars Sample Handling and Requirements Panel (MSHARP) Final Report*, NASA Jet Propulsion Lab, Pasadena, California, April 1999, NASA TM-1999-209145 (1999).
- CDC-NIH, *Biosafety in Microbial Laboratories*, 3rd edition, HHS Publication #(CDC) 93-8395, U.S. Government Printing Office, Washington D.C. (1993).
- Daly, M. J., *Dienococcus radiodurans as an Analogue to Extremophile Organisms that may have Survived on Mars*, Appendix A, pp. 67–69 in: Space Studies Board, *The Quarantine and Certification of Martian Samples*, Committee on Planetary and Lunar Exploration (COMPLEX), chaired by John Wood, National Research Council, National Academy Press, Washington D.C. (2002).
- DeVincenzi, D. L., J. Bagby, M. Race, and J. D. Rummel, *Mars Sample Quarantine Protocol Workshop*, NASA Ames Research Center, Moffett Field, California, June 1997, NASA CP-1999-208772 (1999).
- Exobiology Program Office, *An Exobiological Strategy for Mars Exploration*, NASA Headquarters, Washington, D.C., NASA SP-530 (1995).
- Maxwell, J. A., *Rock and Mineral Analysis*, Wiley/Interscience, NY, 584p. (1968).
- NASA, *Biological Contamination Control for Outbound and Inbound Planetary Spacecraft*, NPD 8020.7E, Signed by the NASA Administrator, 19 February 1999.
- Neal, C. R., "Issues involved in a martian sample return: Integrity preservation and the Curation and Analysis Planning Team for Extraterrestrial Materials (CAPTEM) position," *J. Geophys. Res.* 105, 22487-22506 (2000).
- Pollard, E. C., "The Physics of Viruses." Academic Press, New York, N.Y., p. 53 (1953).
- Race, M. S. and J. D. Rummel, eds., *Mars Sample Handling Protocol Workshop Series, Workshop 1 Final Report*, NASA CP-2000-20964 (2000).
- Race, M. S., G. T. A. Kovacs, J. D. Rummel, and S. E. Acevedo, eds., *Mars Sample Handling Protocol Workshop Series, Workshop 2 Final Report*, NASA CP-2001-210923 (2001a).

- Race, M. S., K. H. Nealson, J. D. Rummel, and S. E. Acevedo, eds., *Mars Sample Handling Protocol Workshop Series, Workshop 3 Final Report*, NASA CP-2001-211388 (2001b).
- Race, M. S., D. L. DeVincenzi, J. D. Rummel, and S. E. Acevedo, eds., *Mars Sample Handling Protocol Workshop Series, Workshop 4 Final Report*, NASA CP-2002-211841 (2002).
- Space Studies Board, *Biological Contamination of Mars: Issues and Recommendations*, Task Group on Planetary Protection, chaired by Kenneth Nealson, National Research Council, National Academy Press, Washington D.C. (1992).
- Space Studies Board, *Mars Sample Return: Issues and Recommendations*, Task Group on Issues in Sample Return, chaired by Kenneth Nealson, National Research Council, National Academy Press, Washington D.C. (1997).
- Space Studies Board, *Size Limits of Very Small Microorganisms: Proceedings of a Workshop*, Steering Group for the Workshop on Size Limits of Very Small Microorganisms, National Research Council, National Academy Press, Washington D.C. (1999).
- Space Studies Board, *The Quarantine and Certification of Martian Samples*, Committee on Planetary and Lunar Exploration (COMPLEX), chaired by John Wood, National Research Council, National Academy Press, Washington D.C. (2002).

APPENDIX C: OVERSIGHT AND REVIEW COMMITTEE (ORC)

Joshua Lederberg, Ph.D.
(Committee Co-Chairperson)
President Emeritus
Rockefeller University
1230 York Avenue
New York NY 10021
USA
tel# 212-327-7809
fax# 212-327-8651
jsl@rockvax.rockefeller.edu

Lynn R. Goldman, M.D.
(Committee Co-Chairperson)
Adjunct Professor
Pew Environmental Health Commission
Johns Hopkins School of Public Health
624 N. Broadway, Room 414
Baltimore MD 21205
USA
tel# 410-614-9301
fax# 410-614-8964
lgoldman@jhsp.h.edu

Committee Members:

James R. Arnold, Ph.D.
Department of Chemistry
University of California, San Diego
9500 Gilman Drive
La Jolla CA 92093-0524
USA
tel# 858-534-2908
fax# 858-534-7840
jarnold@ucsd.edu

Anthony S. Fauci, M.D.
Director, National Institute of Allergy and
Infectious Diseases
National Institutes of Health
9000 Rockville Pike
Bethesda MD 20892
USA
tel# 301-496-2263
fax# 301-496-4409
afauci@niaid.nih.gov

Purnell W. Choppin, M.D.
President Emeritus
Howard Hughes Medical Institute
4000 Jones Bridge Road
Chevy Chase MD 20815-6789
USA
tel# 301-215-8554
fax# 301-215-8566
choppinp@hhmi.org

Representing Dr. Fauci:
Carole Heilman, Ph.D.
Director, Division of Microbiology and
Infectious Diseases
6700-B Rockledge Dr., Rm 3142
Bethesda MD 20817
USA
tel# 301-496-1884
cheilman@niaid.nih.gov

Dominique Dormont, M.D.
CEA - Service de Neurovirologie
60 Avenue de la Division Leclerc
BP 6, 92265 Fontenay-aux-Roses
Cedex
FRANCE
tel# 33 01 46 54 81 22
fax# 33 01 46 54 77 26
dormont@dsvidf.cea.fr

Edward McSweegan, Ph.D.
Health Scientist Administrator,
National Institutes of Health
6700-B Rockledge Dr.
Bethesda MD 20892-7630
USA
tel# 301-402-8370
mcsweegan@nih.gov

Committee Members (cont.)

Nina V. Fedoroff, Ph.D.
Director, Life Sciences Consortium
The Pennsylvania State University
519 Wartik Laboratory
University Park PA 16802-5807
USA
tel# 814-863-5717
fax# 814-863-1357
nvf1@psu.edu

Patricia N. Fultz, Ph.D.
Professor of Microbiology
University of Alabama
Bevill Biomedical Research Bldg.
845 South 19th Street
Birmingham AL 35294-2170
tel# 205-934-0790
fax# 205-975-6788
pnf@uab.edu

John Hobbie, Ph.D.
Co-Director,
The Ecosystems Center
Marine Biological Laboratory
7 MBL Street
Woods Hole MA 02543
USA
tel# 508-289-7470
fax# 508-457-1548
jhobbie@mbl.edu

Heinrich D. Holland, Ph.D.
Harvard University
Department of Earth and Planetary
Sciences
20 Oxford Street
Cambridge MA 02138
USA
tel# 617-495-5892
fax# 617-496-4387
holland@eps.harvard.edu

Debra L. Hunt, Dr. P.H.
Director, Biological Safety Division
Occupational and Environmental
Safety Office
Box 3149
Duke University Medical Center
Durham NC 27710
tel# 919-684-8822
fax# 919-681-7509
hunt0009@mc.duke.edu

Stuart A. Kauffman, M.D.
Founder, Bios Group LP
317 Paseo de Peralta
Santa Fe NM 87501
USA
tel# 505-992-6700
fax# 505-988-2229
stu@biosgroup.com

Robert W. McKinney, Ph.D.
Director, Division of Safety
National Institutes of Health
Building 31, Room 1C02
Bethesda MD 20892-2260
USA
tel# 301-496-1357
fax# 301-402-0316
rm130d@nih.gov

Florabel G. Mullick, M.D.
Director, Ctr. for Adv. Pathology
Principal Deputy Director, Armed
Forces Institute of Pathology
6825 16th Street, NW Bldg #54
Washington DC 20306-6000
USA
tel# 202-782-2503
fax# 202-782-7166
mullick@afip.osd.mil

Committee Members (cont.):

Robert Naquet, Ph.D.
Directeur de Recherche Émérite
Institut Alfred Fessard
1 Avenue de la Terrasse
Gif-sur-Yvette 91198 Cedex
FRANCE
tel# 33 1 69 07 61 45
fax# 33 1 69 07 05 38
naquet@iaf.cnrs-gif.fr

Gilbert S. Omenn, M.D., Ph.D.
Exec. Vice Pres. – Medical Affairs
University of Michigan
M7324 Medical Sciences I Bldg.
1301 Catherine Street
Ann Arbor MI 48109-0626
USA
tel# 734-647-9351
fax# 734-647-9739
gomenn@umich.edu

Leslie Orgel, Ph.D.
Chemical Evolution Laboratory
The Salk Institute for Biological
Studies
10010 North Torrey Pines Road
La Jolla CA 92037
USA
tel# 858-453-4100 (x1322)
fax# 858-558-7359
orgel@salk.edu

Mary Jane Osborn, Ph.D.
Professor and Head
Department of Microbiology
Univ. of Connecticut Health Center
263 Farmington Avenue
Farmington CT 06030-3205
USA
tel# 860-679-2318
fax# 860-679-1239
osborn@sun.uhc.edu

Lucy S. Tompkins, M.D., Ph.D.
Professor of Medicine, Microbiology, and
Immunology
Stanford University Medical Center
300 Pasteur Drive Room H1537J
Stanford CA 94305
USA
tel# 650-725-3861
fax# 650-498-2761
lucytomp@stanford.edu

Jean-Didier Vincent, Ph.D.
Director, L'Institut Alfred Fessard
1 Avenue de la Terrasse
Gif-sur-Yvette 91198 Cedex
FRANCE
tel# 33 1 69 82 34 34
fax# 33 1 69 07 05 38
vincent@iaf.cnrs-gif.fr

Robert M. Walker, Ph.D.
Director, McDonnell Center for the
Space Sciences
Department of Physics
Washington University in St. Louis
Campus Box 1105
St. Louis MO 63130
USA
tel# 314-935-6297/6257
fax# 314-935-6219
rmw@howdy.wustl.edu

Consultants to the ORC:

John R. Bagby, Ph.D.
(Apollo Lunar PP Consultant)
5315 Foxtire Lane
Lohman MO 65053
tel# 573-893-5544
fax# 573-751-6041
bagby@computerland.net

Steven J. Dick, Ph.D.
(Historical Consultant)
U.S. Naval Observatory
3450 Massachusetts Avenue, NW
Washington DC 20392-5420
tel# 202-762-0379
fax# 202-762-1489
dick.steve@usno.navy.mil

Kathie L. Olsen, Ph.D.
(NASA Administrator's Liaison)
Chief Scientist, Code AS
NASA Headquarters
300 E Street SW
Washington DC 20546-0001
tel# 202-358-4509
fax# 202-358-3931
kolsen@hq.nasa.gov

John D. Rummel, Ph.D.
(Executive Secretary)
Planetary Protection Officer
NASA Headquarters, Code S
Washington DC 20546
tel# 202-358-0702
fax# 202-358-3097
jrummel@hq.nasa.gov

APPENDIX D: MARS SAMPLE HANDLING PROTOCOL DEVELOPMENT LEADERSHIP GROUP

Ms. Sara E. Acevedo
SETI Institute
2035 Landings Drive
Mt. View CA 94043
tel# 650-960-4539
sacevedo@mail.arc.nasa.gov

Dr. Gregory T. A. Kovacs
Electrical Engineering & Medicine
Stanford University, CISX-202
Stanford CA 94305-4075
tel# 650-725-3637
kovacs@cis.stanford.edu

Dr. Joseph Bielitzki
DARPA/DSO
3701 N. Fairfax Drive
Arlington VA 22203-1714
tel# 703-696-5278
jbielitzki@darpa.mil

Dr. Glenn MacPherson
National Museum of Natural History
Smithsonian Institution
Washington DC 20560-0119
tel# 202-357-2260
glenn@volcano.si.edu

Dr. Carl W. Bruch
684 O'Neil Road
Hudson WI 54016-7839
tel# 715-386-2346
cbruch@isd.net

Dr. Kenneth H. Nealson
Wrigley Professor of Geobiology
University of Southern California
Dept of Earth Sciences, Mail Code 0740
223 Science Hall
Los Angeles CA 90089-0740
tel# 213-821-2271
knealson@usc.edu

Dr. Jean-Louis Counil
CNES
18, Ave Edouard Belin
F-31401 Toulouse Cedex 4
FRANCE
tel# 33-5-61-27-32-36
jean-louis.counil@cnes.fr

Dr. Margaret S. Race
SETI Institute
2035 Landings Drive
Mountain View CA 94043
tel# 925-947-1272
mracemom@aol.com

Dr. Donald L. DeVincenzi
MS 245-1
NASA Ames Research Center
Moffett Field CA 94035-1000
tel# 650-604-5251
ddevincenzi@mail.arc.nasa.gov

Dr. John D. Rummel
Code S
NASA Headquarters
Washington DC 20546
tel# 202-358-0702
jrummel@hq.nasa.gov

Mr. Paul Jackson Schad
US Environmental Protection Agency
c/o Code U
NASA Headquarters
Washington DC 20024
tel# 202-358-0593
pschad@hq.nasa.gov

Dr. Richard B. Setlow
Biology Department, 463
Brookhaven National Laboratory
PO Box 5000
Upton NY 11973-5000
tel# 631-344-3391
setlow@bnl.gov

Mr. Pericles D. Stabekis
The Windermere Group
Suite 201
525 School Street SW
Washington DC 20024
tel# 202-484-8247
pstabeki@hq.nasa.gov

Dr. Allan Treiman
Lunar and Planetary Institute
3600 Bay Area Blvd.
Houston TX 77058
tel# 281-486-2117
treiman@lpi.usra.edu

Dr. Michel Viso
CNES/DPI/E2U
2 place Maurice-Quentin
F-75055 Paris
FRANCE
tel# 33-1-44-76-79-51
Michel.Viso@cnes.fr

Dr. Norman Wainwright
7 MBL Street
Marine Biological Laboratory
Woods Hole MA 02543
tel# 508-289-7343
nwainwri@mbl.edu

APPENDIX E: GLOSSARY OF TERMS AND ACRONYMS

ALH	Alan Hills (Antarctica)
BFP	Blue Fluorescent Protein
BHK cells	A cloned cell line widely used as a viral host, in studies of oncogenic transformation and of cell physiology.
BSL	Biosafety Level
CAPTEM	Curation and Analysis Planning Team for Extraterrestrial Materials (NASA)
CCNE	Comité Consultatif National d’Ethique pour les Sciences de la Vie et de la Santé (French)
CDC	Centers for Disease Control and Prevention (U.S.)
‘cleanliness’	Free from biological or chemical contamination
CNES	Centre National d’Etudes Spatiale (French)
CNRS	Centre National de la Recherche Scientifique (French)
COMPLEX	Committee on Planetary and Lunar Exploration (U.S.)
‘coupons’	Small, regular samples of solid laboratory materials such as plastic
CP	Conference Proceedings (NASA)
D ₃₇	The average radiation dose required to inactivate a live or infectious particle
DNA	Deoxyribonucleic Acid
Eh	Oxidation Potential
EPST	Etablissements Publics à Caractère Scientifique (French)
EVT	Experiment Verification Test
GC/MS	Gas Chromatograph/Mass Spectrometer
GFP	Green Fluorescent Protein
HEPA	High Efficiency Particulate Air (filter)
HHS	Department of Health and Human Services (U.S.)

IACUC	Institutional Animal Care and Use Committee
IBSC	Institutional Bio-Safety Committee
i.c.	Intracranially
ICBC	Interagency Committee on Back Contamination
INSERM	Institut National de la Santé et de la Recherche Médicale (French)
i.p.	Intraperitoneally
IR	Infrared
Knockout mouse	A mouse that is genetically engineered (both alleles of a critically targeted gene are replaced by an inactive allele using homologous recombination) to produce a particular designer alteration whereby a specifically targeted gene becomes inactivated (or "knocked-out")
LAL	<i>Limulus</i> Amebocyte Lysate
LC/MS	Liquid Chromatograph/Mass Spectrometer
LD/BH	Life Detection/Biohazard (Testing)
LD/MS	Laser Desorption Mass Spectroscopy
MeV	Mega Electron Volts
Mrads	Megarads
MS	Mass Spectroscopy
MSHARP	Mars Sample Handling and Requirements Panel (NASA)
MSHP	Mars Sample Handling Protocol
MSR	Mars Sample Return
NAS	National Academy of Science (U.S.)
NASA	National Aeronautics and Space Administration (U.S.)
Nd:YAG	Neodymium-doped:Yttrium Aluminum Garnet (Laser)
NIH	National Institutes of Health (U.S.)
NPD	NASA Policy Directive
NRC	National Research Council (U.S.)

Nude mouse	A mouse that lacks a thymus and, therefore, cannot generate mature T lymphocytes to mount most types of immune responses
PAH	Polycyclic Aromatic Hydrocarbon
'passaging'	A sub-culturing technique
P/C	Physical and Chemical (Testing)
PCR	Polymerase Chain Reaction
<i>per os</i>	Oral administration (e.g., that a drug is to be swallowed)
pH	Measure of hydrogen ion concentration (acidity)
PP	Planetary Protection
PPAC	Planetary Protection Advisory Committee (NASA)
PPC	Planetary Protection Committee (French)
PPL	Planetary Protection Level
rDNA	Ribosomal DNA
'readout'	A measure of potential biohazard effect
'riffle splitter'	A mechanical separation device used for geological samples
RNA	Ribonucleic Acid
'rocklets'	Millimeter-sized rock fragments
SCID	Severely Compromised Immunodeficient
SCID-Hu	Severely Compromised Immunodeficient (human)
'simulant'	Analogue
SP	Special Publication (NASA)
SRC	Sample Return Canister
SRF	Sample Receiving Facility
SSB	Space Studies Board (U.S.)
TBC	To Be Confirmed
TBD	To Be Determined
TEM	Transmission Electron Microscopy
TM	Technical Memorandum (NASA)
TOC	Total Organic Carbon

USAMRIID	U.S. Army Medical Research Institute of Infectious Diseases
USDA	U.S. Department of Agriculture
UV	Ultraviolet
WHO	World Health Organization
'witness plates'	Controls for forward contamination; used to monitor for bioload on spacecraft
XRD	X-ray Diffraction
XRF	X-ray Fluorescence

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188		
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA, 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE		3. DATES COVERED (From - To)	
31/10/02		Conference Proceedings			
4. TITLE AND SUBTITLE			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
A Draft Test Protocol for Detecting Possible Biohazards in Martian Samples Returned to Earth					
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
J. D. Rummel, M.S. Race, D. L. DeVincenzi, P.J. Schad, P.D. Stabekis, M. Viso, and S. E. Acevedo, eds.					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER		
Ames Research Center Moffett Field CA 94035-1000			A-0208825		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING/MONITOR'S ACRONYM(S)		
			11. SPONSORING/MONITORING REPORT NUMBER		
National Aeronautics and Space Administration Washington, D.C. 20546-0001			NASA/CP-2002-211842		
12. DISTRIBUTION/AVAILABILITY STATEMENT					
Unclassified-Unlimited Subject Category 88			Distribution: Standard Availability: NASA CASI (301) 621-0390		
13. SUPPLEMENTARY NOTES					
POC: John D. Rummel, Code S, NASA Headquarters, Washington DC 20546					
14. ABSTRACT					
<p>This document presents the first complete draft of a protocol for detecting possible biohazards in Mars samples returned to Earth; it is the final product of the Mars Sample Handling Protocol Workshop Series, convened in 2000-2001 by NASA's Planetary Protection Officer. The goal of the five-workshop Series was to develop a comprehensive protocol by which returned martian sample materials could be assessed for the presence of any biological hazard(s) while safeguarding the purity of the samples from possible terrestrial contamination. The reference numbers for the proceedings from the five individual Workshops (1, 2, 2a, 3, and 4) are: NASA/CP-2000-20964, NASA/CP-2001-210923, NASA/CP-2001-210924, NASA/CP-2001-211388, NASA/CP-2002-211841.</p>					
15. SUBJECT TERMS					
Planetary protection; Mars sample handling protocol; biohazard testing.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			John D. Rummel
Uncl.	Uncl.	Uncl.	Uncl.	127	19b. TELEPHONE NUMBER (Include area code)
					202-358-0702